



UNSW  
SYDNEY

## MATH1131 Mathematics 1A – Algebra

### Lecture 5: Lengths and the Dot Product

Lecturer: Sean Gardiner – [sean.gardiner@unsw.edu.au](mailto:sean.gardiner@unsw.edu.au)

Based on slides by Jonathan Kress

## Length in $n$ dimensions

Recall the **length** of  $\mathbf{a} \in \mathbb{R}^n$  with

$$\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{pmatrix} = a_1 \mathbf{e}_1 + a_2 \mathbf{e}_2 + \cdots + a_n \mathbf{e}_n$$

is defined to be

$$|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + \cdots + a_n^2}.$$

## Length in $n$ dimensions

Recall the **length** of  $\mathbf{a} \in \mathbb{R}^n$  with

$$\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{pmatrix} = a_1 \mathbf{e}_1 + a_2 \mathbf{e}_2 + \cdots + a_n \mathbf{e}_n$$

is defined to be

$$|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + \cdots + a_n^2}.$$

If  $|\mathbf{a}| = 1$ , we say that  $\mathbf{a}$  is a **unit vector**.

## Length in $n$ dimensions

Recall the **length** of  $\mathbf{a} \in \mathbb{R}^n$  with

$$\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{pmatrix} = a_1 \mathbf{e}_1 + a_2 \mathbf{e}_2 + \cdots + a_n \mathbf{e}_n$$

is defined to be

$$|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + \cdots + a_n^2}.$$

If  $|\mathbf{a}| = 1$ , we say that  $\mathbf{a}$  is a **unit vector**.

### Properties

1.  $|\mathbf{a}|$  is a real number.
2.  $|\mathbf{a}| \geq 0$ .
3.  $|\mathbf{a}| = 0$  if and only if  $\mathbf{a} = \mathbf{0}$ .
4.  $|\lambda \mathbf{a}| = |\lambda| |\mathbf{a}|$  for all  $\lambda \in \mathbb{R}$ .

# Length in $n$ dimensions

## Properties

### Proof of properties

Property 1 follows from the definition of  $\sqrt{\cdot}$ . Since the components  $a_1, \dots, a_n$  are in  $\mathbb{R}$ , we have  $a_1^2 + a_2^2 + \dots + a_n^2 \geq 0$ .

Hence  $|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + \dots + a_n^2}$  is defined and a real number.

# Length in $n$ dimensions

## Properties

### Proof of properties

Property 1 follows from the definition of  $\sqrt{\cdot}$ . Since the components  $a_1, \dots, a_n$  are in  $\mathbb{R}$ , we have  $a_1^2 + a_2^2 + \dots + a_n^2 \geq 0$ .

Hence  $|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + \dots + a_n^2}$  is defined and a real number.

In fact, the definition of  $\sqrt{\cdot}$  says that  $\sqrt{a_1^2 + a_2^2 + \dots + a_n^2} \geq 0$ . This means  $|\mathbf{a}| \geq 0$ , which is Property 2.

# Length in $n$ dimensions

## Properties

### Proof of properties

Property 1 follows from the definition of  $\sqrt{\cdot}$ . Since the components  $a_1, \dots, a_n$  are in  $\mathbb{R}$ , we have  $a_1^2 + a_2^2 + \dots + a_n^2 \geq 0$ .

Hence  $|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + \dots + a_n^2}$  is defined and a real number.

In fact, the definition of  $\sqrt{\cdot}$  says that  $\sqrt{a_1^2 + a_2^2 + \dots + a_n^2} \geq 0$ . This means  $|\mathbf{a}| \geq 0$ , which is Property 2.

For Property 3 we use that  $\sqrt{x} = 0$  if and only if  $x = 0$ . Hence

$$\begin{aligned} |\mathbf{a}| = 0 &\iff a_1^2 + a_2^2 + \dots + a_n^2 = 0 \\ &\iff a_1 = a_2 = \dots = a_n = 0 \\ &\iff \mathbf{a} = \mathbf{0} \end{aligned}$$

# Length in $n$ dimensions

## Properties

### Proof of properties (continued)

For Property 4, take  $\lambda \in \mathbb{R}$ . Since

$$\lambda \mathbf{a} = \begin{pmatrix} \lambda a_1 \\ \lambda a_2 \\ \vdots \\ \lambda a_n \end{pmatrix}$$

we have

$$\begin{aligned} |\lambda \mathbf{a}| &= \sqrt{(\lambda a_1)^2 + (\lambda a_2)^2 + \cdots + (\lambda a_n)^2} \\ &= \sqrt{\lambda^2(a_1^2 + a_2^2 + \cdots + a_n^2)} \\ &= \sqrt{\lambda^2} \sqrt{a_1^2 + a_2^2 + \cdots + a_n^2} \\ &= |\lambda| |\mathbf{a}|. \end{aligned}$$





## Length in $n$ dimensions - Examples

### Example

Find the two unit vectors parallel to  $\mathbf{b} = \begin{pmatrix} 1 \\ 3 \\ 2 \\ 6 \end{pmatrix}$ .

## Length in $n$ dimensions - Examples

### Example

Find the two unit vectors parallel to  $\mathbf{b} = \begin{pmatrix} 1 \\ 3 \\ 2 \\ 6 \end{pmatrix}$ .

$$\text{Here } |\mathbf{b}| = \sqrt{1^2 + 3^2 + 2^2 + 6^2} = \sqrt{50} = 5\sqrt{2}.$$

## Length in $n$ dimensions - Examples

### Example

Find the two unit vectors parallel to  $\mathbf{b} = \begin{pmatrix} 1 \\ 3 \\ 2 \\ 6 \end{pmatrix}$ .

Here  $|\mathbf{b}| = \sqrt{1^2 + 3^2 + 2^2 + 6^2} = \sqrt{50} = 5\sqrt{2}$ .

So the unit vector  $\hat{\mathbf{b}} = \frac{1}{|\mathbf{b}|}\mathbf{b} = \frac{1}{5\sqrt{2}} \begin{pmatrix} 1 \\ 3 \\ 2 \\ 6 \end{pmatrix}$ .

## Length in $n$ dimensions - Examples

### Example

Find the two unit vectors parallel to  $\mathbf{b} = \begin{pmatrix} 1 \\ 3 \\ 2 \\ 6 \end{pmatrix}$ .

Here  $|\mathbf{b}| = \sqrt{1^2 + 3^2 + 2^2 + 6^2} = \sqrt{50} = 5\sqrt{2}$ .

So the unit vector  $\hat{\mathbf{b}} = \frac{1}{|\mathbf{b}|}\mathbf{b} = \frac{1}{5\sqrt{2}} \begin{pmatrix} 1 \\ 3 \\ 2 \\ 6 \end{pmatrix}$ .

The second unit vector that is parallel to  $\mathbf{b}$  is  $-\hat{\mathbf{b}}$ , that is,

$$-\frac{1}{5\sqrt{2}} \begin{pmatrix} 1 \\ 3 \\ 2 \\ 6 \end{pmatrix}.$$

## Length in $n$ dimensions - Examples

### Example

Find a vector of length 5 parallel to  $\mathbf{w} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ .

## Length in $n$ dimensions - Examples

### Example

Find a vector of length 5 parallel to  $\mathbf{w} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ .

Here  $|\mathbf{w}| = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$ .

## Length in $n$ dimensions - Examples

### Example

Find a vector of length 5 parallel to  $\mathbf{w} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ .

Here  $|\mathbf{w}| = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$ .

So the unit vector  $\hat{\mathbf{w}} = \frac{1}{|\mathbf{w}|} \mathbf{w} = \frac{1}{\sqrt{14}} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ .

## Length in $n$ dimensions - Examples

### Example

Find a vector of length 5 parallel to  $\mathbf{w} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ .

Here  $|\mathbf{w}| = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$ .

So the unit vector  $\hat{\mathbf{w}} = \frac{1}{|\mathbf{w}|} \mathbf{w} = \frac{1}{\sqrt{14}} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ .

A vector parallel to  $\mathbf{w}$  with length 5 is therefore given by

$$5\hat{\mathbf{w}} = \frac{5}{\sqrt{14}} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}.$$



# Dot product

## Definition

The **dot product** (or **scalar product**) of two vectors  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$  with

$$\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{pmatrix} \quad \text{and} \quad \mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}$$

is

$$\mathbf{a} \cdot \mathbf{b} = a_1 b_1 + a_2 b_2 + \cdots + a_n b_n = \sum_{k=1}^n a_k b_k.$$

# Dot product

## Examples

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 4 \end{pmatrix} =$$

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 2 \end{pmatrix} =$$

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} -2 \\ 1 \end{pmatrix} =$$

$$\begin{pmatrix} 3 \\ 0 \\ 4 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} =$$

# Dot product

## Examples

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 4 \end{pmatrix} = 1 \times 3 + 2 \times 4 = 11$$

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 2 \end{pmatrix} =$$

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} -2 \\ 1 \end{pmatrix} =$$

$$\begin{pmatrix} 3 \\ 0 \\ 4 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} =$$

# Dot product

## Examples

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 4 \end{pmatrix} = 1 \times 3 + 2 \times 4 = 11$$

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 2 \end{pmatrix} = 1 \times 1 + 2 \times 2 = 5$$

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} -2 \\ 1 \end{pmatrix} =$$

$$\begin{pmatrix} 3 \\ 0 \\ 4 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} =$$

# Dot product

## Examples

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 4 \end{pmatrix} = 1 \times 3 + 2 \times 4 = 11$$

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 2 \end{pmatrix} = 1 \times 1 + 2 \times 2 = 5$$

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} -2 \\ 1 \end{pmatrix} = 1 \times (-2) + 2 \times 1 = 0$$

$$\begin{pmatrix} 3 \\ 0 \\ 4 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} =$$

# Dot product

## Examples

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 4 \end{pmatrix} = 1 \times 3 + 2 \times 4 = 11$$

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 2 \end{pmatrix} = 1 \times 1 + 2 \times 2 = 5$$

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} -2 \\ 1 \end{pmatrix} = 1 \times (-2) + 2 \times 1 = 0$$

$$\begin{pmatrix} 3 \\ 0 \\ 4 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = 3 \times 1 + 0 \times 1 + 4 \times 1 = 7$$

# Properties

## Properties of the dot product

For all vectors  $\mathbf{a}$ ,  $\mathbf{b}$ ,  $\mathbf{c} \in \mathbb{R}^n$  and scalars  $\lambda \in \mathbb{R}$ ,

- $\mathbf{a} \cdot \mathbf{a} = |\mathbf{a}|^2$ , so  $|\mathbf{a}| = \sqrt{\mathbf{a} \cdot \mathbf{a}}$
- $\mathbf{a} \cdot \mathbf{b} \in \mathbb{R}$
- $\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a}$  (commutative law)
- $\mathbf{a} \cdot (\lambda \mathbf{b}) = \lambda(\mathbf{a} \cdot \mathbf{b})$  (associative law of scalar multiplication)
- $\mathbf{a} \cdot (\mathbf{b} + \mathbf{c}) = \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{c}$  (distributive law)

## Exercise

Prove these laws.

# Properties

## Properties of the dot product

For all vectors  $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{R}^n$  and scalars  $\lambda \in \mathbb{R}$ ,

- $\mathbf{a} \cdot \mathbf{a} = |\mathbf{a}|^2$ , so  $|\mathbf{a}| = \sqrt{\mathbf{a} \cdot \mathbf{a}}$
- $\mathbf{a} \cdot \mathbf{b} \in \mathbb{R}$
- $\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a}$  (commutative law)
- $\mathbf{a} \cdot (\lambda \mathbf{b}) = \lambda(\mathbf{a} \cdot \mathbf{b})$  (associative law of scalar multiplication)
- $\mathbf{a} \cdot (\mathbf{b} + \mathbf{c}) = \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{c}$  (distributive law)

## Exercise

Prove these laws.


Note that the dot product is **not** itself associative, since an expression like  $(\mathbf{a} \cdot \mathbf{b}) \cdot \mathbf{c}$  has no sensible meaning.



# Cosine rule for triangles

Consider a triangle in  $\mathbb{R}^n$   
with sides **a**, **b** and  
**c** = **a** − **b**.


Let  $\theta$  be the smaller angle  
between **a** and **b**.



# Cosine rule for triangles

Consider a triangle in  $\mathbb{R}^n$   
with sides **a**, **b** and  
**c** = **a** − **b**.


Let  $\theta$  be the smaller angle  
between **a** and **b**.



# Cosine rule for triangles

Consider a triangle in  $\mathbb{R}^n$   
with sides **a**, **b** and  
**c** = **a** − **b**.

Let  $\theta$  be the smaller angle  
between **a** and **b**.




Applying Pythagoras' Theorem to the shaded triangle:

# Cosine rule for triangles

Consider a triangle in  $\mathbb{R}^n$   
with sides **a**, **b** and  
**c** = **a** - **b**.

Let  $\theta$  be the smaller angle  
between **a** and **b**.




Applying Pythagoras' Theorem to the shaded triangle:

$$|c|^2 = \left(|a| \sin \theta\right)^2 + \left(|a| \cos \theta - |b|\right)^2$$

# Cosine rule for triangles

Consider a triangle in  $\mathbb{R}^n$   
with sides  $\mathbf{a}$ ,  $\mathbf{b}$  and  
 $\mathbf{c} = \mathbf{a} - \mathbf{b}$ .

Let  $\theta$  be the smaller angle  
between  $\mathbf{a}$  and  $\mathbf{b}$ .




Applying Pythagoras' Theorem to the shaded triangle:

$$\begin{aligned} |\mathbf{c}|^2 &= \left(|\mathbf{a}| \sin \theta\right)^2 + \left(|\mathbf{a}| \cos \theta - |\mathbf{b}|\right)^2 \\ &= |\mathbf{a}|^2 \sin^2 \theta + |\mathbf{a}|^2 \cos^2 \theta + |\mathbf{b}|^2 - 2|\mathbf{a}||\mathbf{b}| \cos \theta \end{aligned}$$

# Cosine rule for triangles

Consider a triangle in  $\mathbb{R}^n$   
with sides  $\mathbf{a}$ ,  $\mathbf{b}$  and  
 $\mathbf{c} = \mathbf{a} - \mathbf{b}$ .

Let  $\theta$  be the smaller angle  
between  $\mathbf{a}$  and  $\mathbf{b}$ .




Applying Pythagoras' Theorem to the shaded triangle:

$$\begin{aligned} |\mathbf{c}|^2 &= \left(|\mathbf{a}| \sin \theta\right)^2 + \left(|\mathbf{a}| \cos \theta - |\mathbf{b}|\right)^2 \\ &= |\mathbf{a}|^2 \sin^2 \theta + |\mathbf{a}|^2 \cos^2 \theta + |\mathbf{b}|^2 - 2|\mathbf{a}||\mathbf{b}| \cos \theta \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - 2|\mathbf{a}||\mathbf{b}| \cos \theta \end{aligned}$$

# Cosine rule for triangles

Consider a triangle in  $\mathbb{R}^n$   
with sides **a**, **b** and  
**c** = **a** - **b**.

Let  $\theta$  be the smaller angle  
between **a** and **b**.



Applying Pythagoras' Theorem to the shaded triangle:

$$\begin{aligned} |c|^2 &= \left(|a| \sin \theta\right)^2 + \left(|a| \cos \theta - |b|\right)^2 \\ &= |a|^2 \sin^2 \theta + |a|^2 \cos^2 \theta + |b|^2 - 2|a||b| \cos \theta \\ &= |a|^2 + |b|^2 - 2|a||b| \cos \theta \end{aligned}$$

This is called the **cosine rule**.

## Geometric interpretation

From the cosine rule for triangles,

$$\begin{aligned} 2|\mathbf{a}||\mathbf{b}| \cos \theta \\ = |\mathbf{a}|^2 + |\mathbf{b}|^2 - |\mathbf{a} - \mathbf{b}|^2 \end{aligned}$$



## Geometric interpretation

From the cosine rule for triangles,

$$\begin{aligned} & 2|\mathbf{a}||\mathbf{b}| \cos \theta \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - |\mathbf{a} - \mathbf{b}|^2 \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - \left( (a_1 - b_1)^2 + \cdots + (a_n - b_n)^2 \right) \end{aligned}$$

## Geometric interpretation

From the cosine rule for triangles,

$$\begin{aligned} & 2|\mathbf{a}||\mathbf{b}| \cos \theta \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - |\mathbf{a} - \mathbf{b}|^2 \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - \left( (a_1 - b_1)^2 + \cdots + (a_n - b_n)^2 \right) \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - \left( (a_1^2 + b_1^2 - 2a_1b_1) + \cdots + (a_n^2 + b_n^2 - 2a_nb_n) \right) \end{aligned}$$

## Geometric interpretation

From the cosine rule for triangles,

$$\begin{aligned} & 2|\mathbf{a}||\mathbf{b}|\cos\theta \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - |\mathbf{a} - \mathbf{b}|^2 \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - \left( (a_1 - b_1)^2 + \cdots + (a_n - b_n)^2 \right) \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - \left( (a_1^2 + b_1^2 - 2a_1b_1) + \cdots + (a_n^2 + b_n^2 - 2a_nb_n) \right) \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - \left( |\mathbf{a}|^2 + |\mathbf{b}|^2 - 2\mathbf{a} \cdot \mathbf{b} \right) \end{aligned}$$

# Geometric interpretation

From the cosine rule for triangles,

$$\begin{aligned} & 2|\mathbf{a}||\mathbf{b}|\cos\theta \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - |\mathbf{a} - \mathbf{b}|^2 \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - \left( (a_1 - b_1)^2 + \cdots + (a_n - b_n)^2 \right) \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - \left( (a_1^2 + b_1^2 - 2a_1b_1) + \cdots + (a_n^2 + b_n^2 - 2a_nb_n) \right) \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - \left( |\mathbf{a}|^2 + |\mathbf{b}|^2 - 2\mathbf{a} \cdot \mathbf{b} \right) \\ &= 2\mathbf{a} \cdot \mathbf{b} \end{aligned}$$

## Geometric interpretation


From the cosine rule for triangles,

$$\begin{aligned} & 2|\mathbf{a}||\mathbf{b}| \cos \theta \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - |\mathbf{a} - \mathbf{b}|^2 \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - \left( (a_1 - b_1)^2 + \cdots + (a_n - b_n)^2 \right) \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - \left( (a_1^2 + b_1^2 - 2a_1b_1) + \cdots + (a_n^2 + b_n^2 - 2a_nb_n) \right) \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 - \left( |\mathbf{a}|^2 + |\mathbf{b}|^2 - 2\mathbf{a} \cdot \mathbf{b} \right) \\ &= 2\mathbf{a} \cdot \mathbf{b} \end{aligned}$$

So  $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta$

and  $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|}$

where  $\theta$  is the smaller angle between vectors  $\mathbf{a}$  and  $\mathbf{b}$  joined tail-to-tail:



## Dot product - Examples

### Example

Use

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta$$

to find the smaller angle  $\theta$  between  $\mathbf{a} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ .

## Dot product - Examples

### Example

Use

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta$$

to find the smaller angle  $\theta$  between  $\mathbf{a} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ .

$$\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|}$$

## Dot product - Examples

### Example

Use

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta$$

to find the smaller angle  $\theta$  between  $\mathbf{a} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ .

$$\begin{aligned} \cos \theta &= \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} \\ &= \frac{2 \times 1 + 0 \times 1}{\sqrt{2^2 + 0^2} \sqrt{1^2 + 1^2}} \end{aligned}$$



## Dot product - Examples

### Example

Use

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta$$

to find the smaller angle  $\theta$  between  $\mathbf{a} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ .

$$\begin{aligned} \cos \theta &= \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} \\ &= \frac{2 \times 1 + 0 \times 1}{\sqrt{2^2 + 0^2} \sqrt{1^2 + 1^2}} \\ &= \frac{2}{2\sqrt{2}} = \frac{1}{\sqrt{2}}. \end{aligned}$$

## Dot product - Examples

### Example

Use

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta$$

to find the smaller angle  $\theta$  between  $\mathbf{a} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ .

$$\begin{aligned} \cos \theta &= \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} \\ &= \frac{2 \times 1 + 0 \times 1}{\sqrt{2^2 + 0^2} \sqrt{1^2 + 1^2}} \\ &= \frac{2}{2\sqrt{2}} = \frac{1}{\sqrt{2}}. \end{aligned}$$

So the angle  $\theta$  is given by  $\arccos\left(\frac{1}{\sqrt{2}}\right) = \frac{\pi}{4}$ .

## Dot product - Examples

### Example

Find the smaller angle  $\theta$  between  $\mathbf{a} = \begin{pmatrix} 2 \\ 0 \\ 3 \\ -1 \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \\ 0 \\ 2 \end{pmatrix}$ .

## Dot product - Examples

### Example

Find the smaller angle  $\theta$  between  $\mathbf{a} = \begin{pmatrix} 2 \\ 0 \\ 3 \\ -1 \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \\ 0 \\ 2 \end{pmatrix}$ .

$$\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|}$$

## Dot product - Examples

### Example

Find the smaller angle  $\theta$  between  $\mathbf{a} = \begin{pmatrix} 2 \\ 0 \\ 3 \\ -1 \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \\ 0 \\ 2 \end{pmatrix}$ .

$$\begin{aligned}\cos \theta &= \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} \\ &= \frac{2 \times 1 + 0 \times 1 + 3 \times 0 + (-1) \times 2}{\sqrt{2^2 + 0^2 + 3^2 + (-1)^2} \sqrt{1^2 + 1^2 + 0^2 + 2^2}}\end{aligned}$$

## Dot product - Examples

### Example

Find the smaller angle  $\theta$  between  $\mathbf{a} = \begin{pmatrix} 2 \\ 0 \\ 3 \\ -1 \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \\ 0 \\ 2 \end{pmatrix}$ .

$$\begin{aligned}\cos \theta &= \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} \\ &= \frac{2 \times 1 + 0 \times 1 + 3 \times 0 + (-1) \times 2}{\sqrt{2^2 + 0^2 + 3^2 + (-1)^2} \sqrt{1^2 + 1^2 + 0^2 + 2^2}} \\ &= \frac{0}{\sqrt{14}\sqrt{6}} = 0.\end{aligned}$$

## Dot product - Examples

### Example

Find the smaller angle  $\theta$  between  $\mathbf{a} = \begin{pmatrix} 2 \\ 0 \\ 3 \\ -1 \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \\ 0 \\ 2 \end{pmatrix}$ .

$$\begin{aligned}\cos \theta &= \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} \\ &= \frac{2 \times 1 + 0 \times 1 + 3 \times 0 + (-1) \times 2}{\sqrt{2^2 + 0^2 + 3^2 + (-1)^2} \sqrt{1^2 + 1^2 + 0^2 + 2^2}} \\ &= \frac{0}{\sqrt{14}\sqrt{6}} = 0.\end{aligned}$$

So the angle  $\theta$  is given by  $\arccos(0) = \frac{\pi}{2}$ .

# Dot product

## Theorems

### Theorem

For any two non-zero vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$\mathbf{a} \perp \mathbf{b} \text{ if and only if } \mathbf{a} \cdot \mathbf{b} = 0.$$



# Dot product

## Theorems

### Theorem

For any two non-zero vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$\mathbf{a} \perp \mathbf{b}$  if and only if  $\mathbf{a} \cdot \mathbf{b} = 0$ .

### Proof

# Dot product

## Theorems

### Theorem

For any two non-zero vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$\mathbf{a} \perp \mathbf{b} \text{ if and only if } \mathbf{a} \cdot \mathbf{b} = 0.$$

### Proof

Let  $\mathbf{a}$  and  $\mathbf{b}$  be two non-zero vectors in  $\mathbb{R}^n$

# Dot product

## Theorems

### Theorem

For any two non-zero vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$\mathbf{a} \perp \mathbf{b} \text{ if and only if } \mathbf{a} \cdot \mathbf{b} = 0.$$

### Proof

Let  $\mathbf{a}$  and  $\mathbf{b}$  be two non-zero vectors in  $\mathbb{R}^n$ , and let  $\theta$  be the smaller angle between  $\mathbf{a}$  and  $\mathbf{b}$ .

# Dot product

## Theorems

### Theorem

For any two non-zero vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$\mathbf{a} \perp \mathbf{b} \text{ if and only if } \mathbf{a} \cdot \mathbf{b} = 0.$$

### Proof

Let  $\mathbf{a}$  and  $\mathbf{b}$  be two non-zero vectors in  $\mathbb{R}^n$ , and let  $\theta$  be the smaller angle between  $\mathbf{a}$  and  $\mathbf{b}$ .

If  $\mathbf{a} \perp \mathbf{b}$ , that is, if  $\theta = \frac{\pi}{2}$ , then  $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \frac{\pi}{2} = 0$ .

# Dot product

## Theorems

### Theorem

For any two non-zero vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$\mathbf{a} \perp \mathbf{b} \text{ if and only if } \mathbf{a} \cdot \mathbf{b} = 0.$$

### Proof

Let  $\mathbf{a}$  and  $\mathbf{b}$  be two non-zero vectors in  $\mathbb{R}^n$ , and let  $\theta$  be the smaller angle between  $\mathbf{a}$  and  $\mathbf{b}$ .

If  $\mathbf{a} \perp \mathbf{b}$ , that is, if  $\theta = \frac{\pi}{2}$ , then  $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \frac{\pi}{2} = 0$ .

Conversely, if  $\mathbf{a} \cdot \mathbf{b} = 0$ , then  $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} = 0$ . Hence  $\theta = \frac{\pi}{2}$  and  $\mathbf{a} \perp \mathbf{b}$ . □

# Dot product

## Theorems

### Theorem

For any two non-zero vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$\mathbf{a} \perp \mathbf{b} \text{ if and only if } \mathbf{a} \cdot \mathbf{b} = 0.$$

### Proof

Let  $\mathbf{a}$  and  $\mathbf{b}$  be two non-zero vectors in  $\mathbb{R}^n$ , and let  $\theta$  be the smaller angle between  $\mathbf{a}$  and  $\mathbf{b}$ .

If  $\mathbf{a} \perp \mathbf{b}$ , that is, if  $\theta = \frac{\pi}{2}$ , then  $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \frac{\pi}{2} = 0$ .

Conversely, if  $\mathbf{a} \cdot \mathbf{b} = 0$ , then  $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} = 0$ . Hence  $\theta = \frac{\pi}{2}$  and  $\mathbf{a} \perp \mathbf{b}$ . □

This is where we need that  $\mathbf{a}$  and  $\mathbf{b}$  are non-zero vectors.

# Dot product

## Theorems

### Theorem (Cauchy-Schwarz inequality)

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} \cdot \mathbf{b}| \leq |\mathbf{a}||\mathbf{b}|$$

# Dot product

## Theorems

### Theorem (Cauchy-Schwarz inequality)

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} \cdot \mathbf{b}| \leq |\mathbf{a}||\mathbf{b}|$$

### Proof



# Dot product

## Theorems

### Theorem (Cauchy-Schwarz inequality)

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} \cdot \mathbf{b}| \leq \|\mathbf{a}\| \|\mathbf{b}\|$$

### Proof

Let  $\mathbf{a}$  and  $\mathbf{b}$  be two vectors in  $\mathbb{R}^n$ , and let  $\theta$  be the smaller angle between  $\mathbf{a}$  and  $\mathbf{b}$ .

# Dot product

## Theorems

### Theorem (Cauchy-Schwarz inequality)

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} \cdot \mathbf{b}| \leq \|\mathbf{a}\| \|\mathbf{b}\|$$

### Proof

Let  $\mathbf{a}$  and  $\mathbf{b}$  be two vectors in  $\mathbb{R}^n$ , and let  $\theta$  be the smaller angle between  $\mathbf{a}$  and  $\mathbf{b}$ .

Then  $\mathbf{a} \cdot \mathbf{b} = \|\mathbf{a}\| \|\mathbf{b}\| \cos \theta$ .

# Dot product

## Theorems

### Theorem (Cauchy-Schwarz inequality)

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} \cdot \mathbf{b}| \leq |\mathbf{a}||\mathbf{b}|$$

### Proof

Let  $\mathbf{a}$  and  $\mathbf{b}$  be two vectors in  $\mathbb{R}^n$ , and let  $\theta$  be the smaller angle between  $\mathbf{a}$  and  $\mathbf{b}$ .

Then  $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta$ .

Since  $-1 \leq \cos \theta \leq 1$ , it follows that

$$-|\mathbf{a}||\mathbf{b}| \leq \mathbf{a} \cdot \mathbf{b} \leq |\mathbf{a}||\mathbf{b}|,$$

which means  $|\mathbf{a} \cdot \mathbf{b}| \leq |\mathbf{a}||\mathbf{b}|$ . □

# Triangle inequality

## Triangle inequality (Minkowski's inequality)

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

# Triangle inequality

## Triangle inequality (Minkowski's inequality)

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

This inequality is commonly known as the **triangle inequality** since we can illustrate it as follows:

# Triangle inequality


## Triangle inequality (Minkowski's inequality)

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

This inequality is commonly known as the **triangle inequality** since we can illustrate it as follows:

When we add two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$  we get this triangle.



# Triangle inequality

## Triangle inequality (Minkowski's inequality)


For any two vectors **a** and **b** in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

This inequality is commonly known as the **triangle inequality** since we can illustrate it as follows:

When we add two vectors **a** and **b** in  $\mathbb{R}^n$  we get this triangle.

Look at the lengths of the sides of the triangle.



# Triangle inequality

## Triangle inequality (Minkowski's inequality)


For any two vectors **a** and **b** in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

This inequality is commonly known as the **triangle inequality** since we can illustrate it as follows:

When we add two vectors **a** and **b** in  $\mathbb{R}^n$  we get this triangle.

Look at the lengths of the sides of the triangle.



The distance travelled along **a** and then **b** can never be shorter than the distance travelled along **a + b**, and will only be equal in distance when **a** and **b** point in the same direction.



# Proofs with dot products

## Triangle inequality

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

Proof

# Proofs with dot products

## Triangle inequality

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

## Proof

Let  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ . Using the Cauchy-Schwarz inequality,

# Proofs with dot products

## Triangle inequality

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

## Proof

Let  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ . Using the Cauchy-Schwarz inequality,

$$|\mathbf{a} + \mathbf{b}|^2 = (\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b})$$

# Proofs with dot products

## Triangle inequality

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

## Proof

Let  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ . Using the Cauchy-Schwarz inequality,

$$\begin{aligned} |\mathbf{a} + \mathbf{b}|^2 &= (\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b}) \\ &= \mathbf{a} \cdot \mathbf{a} + \mathbf{b} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{b} + \mathbf{b} \cdot \mathbf{a} \end{aligned}$$

# Proofs with dot products

## Triangle inequality

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

## Proof

Let  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ . Using the Cauchy-Schwarz inequality,

$$\begin{aligned} |\mathbf{a} + \mathbf{b}|^2 &= (\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b}) \\ &= \mathbf{a} \cdot \mathbf{a} + \mathbf{b} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{b} + \mathbf{b} \cdot \mathbf{a} \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 + 2\mathbf{a} \cdot \mathbf{b} \end{aligned}$$

# Proofs with dot products

## Triangle inequality

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

## Proof

Let  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ . Using the **Cauchy-Schwarz inequality**,

$$\begin{aligned} |\mathbf{a} + \mathbf{b}|^2 &= (\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b}) \\ &= \mathbf{a} \cdot \mathbf{a} + \mathbf{b} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{b} + \mathbf{b} \cdot \mathbf{a} \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 + 2\mathbf{a} \cdot \mathbf{b} \\ &\leq |\mathbf{a}|^2 + |\mathbf{b}|^2 + 2|\mathbf{a}||\mathbf{b}| \end{aligned}$$

# Proofs with dot products

## Triangle inequality

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

## Proof

Let  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ . Using the **Cauchy-Schwarz inequality**,

$$\begin{aligned} |\mathbf{a} + \mathbf{b}|^2 &= (\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b}) \\ &= \mathbf{a} \cdot \mathbf{a} + \mathbf{b} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{b} + \mathbf{b} \cdot \mathbf{a} \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 + 2\mathbf{a} \cdot \mathbf{b} \\ &\leq |\mathbf{a}|^2 + |\mathbf{b}|^2 + 2|\mathbf{a}||\mathbf{b}| \\ &= (|\mathbf{a}| + |\mathbf{b}|)^2. \end{aligned}$$

# Proofs with dot products

## Triangle inequality

For any two vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $\mathbb{R}^n$

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|$$

## Proof

Let  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ . Using the **Cauchy-Schwarz inequality**,

$$\begin{aligned} |\mathbf{a} + \mathbf{b}|^2 &= (\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b}) \\ &= \mathbf{a} \cdot \mathbf{a} + \mathbf{b} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{b} + \mathbf{b} \cdot \mathbf{a} \\ &= |\mathbf{a}|^2 + |\mathbf{b}|^2 + 2\mathbf{a} \cdot \mathbf{b} \\ &\leq |\mathbf{a}|^2 + |\mathbf{b}|^2 + 2|\mathbf{a}||\mathbf{b}| \\ &= (|\mathbf{a}| + |\mathbf{b}|)^2. \end{aligned}$$

Taking square roots of both sides gives the claim. □




## Theorem

The angle subtended by a diameter at the circumference of a circle is a right angle.

# Proofs with dot products

## Theorem


The angle subtended by a diameter at the circumference of a circle is a right angle.



# Proofs with dot products

## Theorem


The angle subtended by a diameter at the circumference of a circle is a right angle.



# Proofs with dot products

## Theorem


The angle subtended by a diameter at the circumference of a circle is a right angle.



# Proofs with dot products

## Theorem

The angle subtended by a diameter at the circumference of a circle is a **right angle**.




# Proofs with dot products

## Theorem

The angle subtended by a diameter at the circumference of a circle is a right angle.

## Proof




# Proofs with dot products

## Theorem

The angle subtended by a diameter at the circumference of a circle is a right angle.

## Proof

Let  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  be the vectors as shown in the diagram.



# Proofs with dot products

## Theorem


The angle subtended by a diameter at the circumference of a circle is a right angle.

## Proof

Let  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  be the vectors as shown in the diagram. We need to show that

$$(\mathbf{b} - \mathbf{a}) \cdot (\mathbf{c} - \mathbf{a}) = 0$$

as this means  $(\mathbf{b} - \mathbf{a}) \perp (\mathbf{c} - \mathbf{a})$ .





# Proofs with dot products

## Theorem

The angle subtended by a diameter at the circumference of a circle is a right angle.


## Proof

Let  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  be the vectors as shown in the diagram. We need to show that

$$(\mathbf{b} - \mathbf{a}) \cdot (\mathbf{c} - \mathbf{a}) = 0$$

as this means  $(\mathbf{b} - \mathbf{a}) \perp (\mathbf{c} - \mathbf{a})$ .

Now,  $|\mathbf{a}| = |\mathbf{b}| = |\mathbf{c}|$  and  $\mathbf{c} = -\mathbf{b}$ .



# Proofs with dot products

## Theorem

The angle subtended by a diameter at the circumference of a circle is a right angle.

## Proof


Let **a**, **b** and **c** be the vectors as shown in the diagram. We need to show that

$$(\mathbf{b} - \mathbf{a}) \cdot (\mathbf{c} - \mathbf{a}) = 0$$

as this means  $(\mathbf{b} - \mathbf{a}) \perp (\mathbf{c} - \mathbf{a})$ .

Now,  $|\mathbf{a}| = |\mathbf{b}| = |\mathbf{c}|$  and  $\mathbf{c} = -\mathbf{b}$ . Hence

$$(\mathbf{b} - \mathbf{a}) \cdot (\mathbf{c} - \mathbf{a}) = \mathbf{b} \cdot \mathbf{c} - \mathbf{b} \cdot \mathbf{a} - \mathbf{a} \cdot \mathbf{c} + \mathbf{a} \cdot \mathbf{a}$$



# Proofs with dot products

## Theorem

The angle subtended by a diameter at the circumference of a circle is a right angle.

## Proof


Let  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  be the vectors as shown in the diagram. We need to show that

$$(\mathbf{b} - \mathbf{a}) \cdot (\mathbf{c} - \mathbf{a}) = 0$$

as this means  $(\mathbf{b} - \mathbf{a}) \perp (\mathbf{c} - \mathbf{a})$ .

Now,  $|\mathbf{a}| = |\mathbf{b}| = |\mathbf{c}|$  and  $\mathbf{c} = -\mathbf{b}$ . Hence

$$\begin{aligned}(\mathbf{b} - \mathbf{a}) \cdot (\mathbf{c} - \mathbf{a}) &= \mathbf{b} \cdot \mathbf{c} - \mathbf{b} \cdot \mathbf{a} - \mathbf{a} \cdot \mathbf{c} + \mathbf{a} \cdot \mathbf{a} \\ &= -\mathbf{b} \cdot \mathbf{b} - \mathbf{b} \cdot \mathbf{a} + \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{a}\end{aligned}$$



# Proofs with dot products

## Theorem

The angle subtended by a diameter at the circumference of a circle is a right angle.

## Proof


Let  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  be the vectors as shown in the diagram. We need to show that

$$(\mathbf{b} - \mathbf{a}) \cdot (\mathbf{c} - \mathbf{a}) = 0$$

as this means  $(\mathbf{b} - \mathbf{a}) \perp (\mathbf{c} - \mathbf{a})$ .

Now,  $|\mathbf{a}| = |\mathbf{b}| = |\mathbf{c}|$  and  $\mathbf{c} = -\mathbf{b}$ . Hence

$$\begin{aligned}(\mathbf{b} - \mathbf{a}) \cdot (\mathbf{c} - \mathbf{a}) &= \mathbf{b} \cdot \mathbf{c} - \mathbf{b} \cdot \mathbf{a} - \mathbf{a} \cdot \mathbf{c} + \mathbf{a} \cdot \mathbf{a} \\&= -\mathbf{b} \cdot \mathbf{b} - \mathbf{b} \cdot \mathbf{a} + \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{a} \\&= -|\mathbf{b}|^2 + |\mathbf{a}|^2\end{aligned}$$



# Proofs with dot products

## Theorem

The angle subtended by a diameter at the circumference of a circle is a right angle.

## Proof

Let  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  be the vectors as shown in the diagram. We need to show that

$$(\mathbf{b} - \mathbf{a}) \cdot (\mathbf{c} - \mathbf{a}) = 0$$

as this means  $(\mathbf{b} - \mathbf{a}) \perp (\mathbf{c} - \mathbf{a})$ .

Now,  $|\mathbf{a}| = |\mathbf{b}| = |\mathbf{c}|$  and  $\mathbf{c} = -\mathbf{b}$ . Hence

$$\begin{aligned}(\mathbf{b} - \mathbf{a}) \cdot (\mathbf{c} - \mathbf{a}) &= \mathbf{b} \cdot \mathbf{c} - \mathbf{b} \cdot \mathbf{a} - \mathbf{a} \cdot \mathbf{c} + \mathbf{a} \cdot \mathbf{a} \\&= -\mathbf{b} \cdot \mathbf{b} - \mathbf{b} \cdot \mathbf{a} + \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{a} \\&= -|\mathbf{b}|^2 + |\mathbf{a}|^2 \\&= 0,\end{aligned}$$

which gives the claim. □

