



UNSW  
SYDNEY

MATH1131 Mathematics 1A – Algebra

## Lecture 20: Finding Inverses and Determinants

Lecturer: Sean Gardiner – [sean.gardiner@unsw.edu.au](mailto:sean.gardiner@unsw.edu.au)

Based on slides by Jonathan Kress

# Inverses of $2 \times 2$ matrices

## Definition

The inverse of the general  $2 \times 2$  matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

# Inverses of $2 \times 2$ matrices

## Definition

The inverse of the general  $2 \times 2$  matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

Note that  $A^{-1}$  is defined if and only if  $ad - bc \neq 0$ .

# Inverses of $2 \times 2$ matrices

## Definition

The inverse of the general  $2 \times 2$  matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

Note that  $A^{-1}$  is defined if and only if  $ad - bc \neq 0$ .

If  $ad - bc = 0$ , then the matrix  $A$  has no inverse, that is, it is not invertible or singular.

# Inverses of $2 \times 2$ matrices

## Definition

The inverse of the general  $2 \times 2$  matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

Note that  $A^{-1}$  is defined if and only if  $ad - bc \neq 0$ .

If  $ad - bc = 0$ , then the matrix  $A$  has no inverse, that is, it is not invertible or singular.

For example,

$$\begin{pmatrix} 1 & 2 \\ 3 & 8 \end{pmatrix}^{-1}$$

# Inverses of $2 \times 2$ matrices

## Definition

The inverse of the general  $2 \times 2$  matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

Note that  $A^{-1}$  is defined if and only if  $ad - bc \neq 0$ .

If  $ad - bc = 0$ , then the matrix  $A$  has no inverse, that is, it is not invertible or singular.

For example,

$$\begin{pmatrix} 1 & 2 \\ 3 & 8 \end{pmatrix}^{-1} = \frac{1}{2} \begin{pmatrix} 8 & -2 \\ -3 & 1 \end{pmatrix}.$$

## $2 \times 2$ Determinants

For the  $2 \times 2$  matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

$ad - bc$  is called the **determinant** of  $A$  and is denoted  $\det(A)$  or  $|A|$ .

## $2 \times 2$ Determinants

For the  $2 \times 2$  matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

$ad - bc$  is called the **determinant** of  $A$  and is denoted  $\det(A)$  or  $|A|$ .

Since

$$A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix},$$

we have

**$A$  is invertible if and only if  $\det(A) \neq 0$ .**



## $2 \times 2$ Determinants – Example

### Example

For each of the following matrices, determine if it is invertible and if so, find its inverse.

$$A = \begin{pmatrix} 3 & 9 \\ 2 & 6 \end{pmatrix}$$

$$B = \begin{pmatrix} -3 & -1 \\ 2 & 6 \end{pmatrix}$$

## $2 \times 2$ Determinants – Example

### Example

For each of the following matrices, determine if it is invertible and if so, find its inverse.

$$A = \begin{pmatrix} 3 & 9 \\ 2 & 6 \end{pmatrix}$$

$$B = \begin{pmatrix} -3 & -1 \\ 2 & 6 \end{pmatrix}$$

$$\det(A)$$

## $2 \times 2$ Determinants – Example

### Example

For each of the following matrices, determine if it is invertible and if so, find its inverse.

$$A = \begin{pmatrix} 3 & 9 \\ 2 & 6 \end{pmatrix}$$

$$B = \begin{pmatrix} -3 & -1 \\ 2 & 6 \end{pmatrix}$$

$$\det(A) = 3 \times 6 - 9 \times 2 = 0$$

## $2 \times 2$ Determinants – Example

### Example

For each of the following matrices, determine if it is invertible and if so, find its inverse.

$$A = \begin{pmatrix} 3 & 9 \\ 2 & 6 \end{pmatrix}$$

$$B = \begin{pmatrix} -3 & -1 \\ 2 & 6 \end{pmatrix}$$

$\det(A) = 3 \times 6 - 9 \times 2 = 0$  , so  $A$  is not invertible.

## $2 \times 2$ Determinants – Example

### Example

For each of the following matrices, determine if it is invertible and if so, find its inverse.

$$A = \begin{pmatrix} 3 & 9 \\ 2 & 6 \end{pmatrix}$$

$$B = \begin{pmatrix} -3 & -1 \\ 2 & 6 \end{pmatrix}$$

$\det(A) = 3 \times 6 - 9 \times 2 = 0$  , so  $A$  is not invertible.

$\det(B)$

## $2 \times 2$ Determinants – Example

### Example

For each of the following matrices, determine if it is invertible and if so, find its inverse.

$$A = \begin{pmatrix} 3 & 9 \\ 2 & 6 \end{pmatrix}$$

$$B = \begin{pmatrix} -3 & -1 \\ 2 & 6 \end{pmatrix}$$

$\det(A) = 3 \times 6 - 9 \times 2 = 0$  , so  $A$  is not invertible.

$\det(B) = (-3) \times 6 - (-1) \times 2 = -16$

## $2 \times 2$ Determinants – Example

### Example

For each of the following matrices, determine if it is invertible and if so, find its inverse.

$$A = \begin{pmatrix} 3 & 9 \\ 2 & 6 \end{pmatrix}$$

$$B = \begin{pmatrix} -3 & -1 \\ 2 & 6 \end{pmatrix}$$

$\det(A) = 3 \times 6 - 9 \times 2 = 0$  , so  $A$  is not invertible.

$\det(B) = (-3) \times 6 - (-1) \times 2 = -16 \neq 0$

## $2 \times 2$ Determinants – Example

### Example

For each of the following matrices, determine if it is invertible and if so, find its inverse.

$$A = \begin{pmatrix} 3 & 9 \\ 2 & 6 \end{pmatrix}$$

$$B = \begin{pmatrix} -3 & -1 \\ 2 & 6 \end{pmatrix}$$

$\det(A) = 3 \times 6 - 9 \times 2 = 0$  , so  $A$  is not invertible.

$\det(B) = (-3) \times 6 - (-1) \times 2 = -16 \neq 0$  , so  $B$  is invertible.



## $2 \times 2$ Determinants – Example

### Example

For each of the following matrices, determine if it is invertible and if so, find its inverse.

$$A = \begin{pmatrix} 3 & 9 \\ 2 & 6 \end{pmatrix}$$

$$B = \begin{pmatrix} -3 & -1 \\ 2 & 6 \end{pmatrix}$$

$\det(A) = 3 \times 6 - 9 \times 2 = 0$  , so  $A$  is not invertible.

$\det(B) = (-3) \times 6 - (-1) \times 2 = -16 \neq 0$  , so  $B$  is invertible.

Its inverse is given by the formula:  $B^{-1} = -\frac{1}{16} \begin{pmatrix} 6 & 1 \\ -2 & -3 \end{pmatrix}$ .

## $2 \times 2$ Determinants – Example

### Example

For each of the following matrices, determine if it is invertible and if so, find its inverse.

$$A = \begin{pmatrix} 3 & 9 \\ 2 & 6 \end{pmatrix}$$

$$B = \begin{pmatrix} -3 & -1 \\ 2 & 6 \end{pmatrix}$$

$\det(A) = 3 \times 6 - 9 \times 2 = 0$  , so  $A$  is not invertible.

$\det(B) = (-3) \times 6 - (-1) \times 2 = -16 \neq 0$  , so  $B$  is invertible.

Its inverse is given by the formula:  $B^{-1} = -\frac{1}{16} \begin{pmatrix} 6 & 1 \\ -2 & -3 \end{pmatrix}$ .

Unfortunately we can't easily learn general formulae for the inverses of matrices of size larger than  $2 \times 2$ . Instead, we first need to consider how elementary row operations are related to matrix multiplication.

## Row operations by matrix multiplication

Swapping rows ( $R_i \leftrightarrow R_j$ ) can be achieved by matrix multiplication.

## Row operations by matrix multiplication

**Swapping** rows ( $R_i \leftrightarrow R_j$ ) can be achieved by matrix multiplication.

For example, if

- $A$  is some  $4 \times 4$  matrix,

## Row operations by matrix multiplication

**Swapping** rows ( $R_i \leftrightarrow R_j$ ) can be achieved by matrix multiplication.

For example, if

- $A$  is some  $4 \times 4$  matrix, and
- $E$  is made by swapping rows 2 and 4 of  $I_4$ ,

## Row operations by matrix multiplication

**Swapping** rows ( $R_i \leftrightarrow R_j$ ) can be achieved by matrix multiplication.

For example, if

- $A$  is some  $4 \times 4$  matrix, and
- $E$  is made by swapping rows 2 and 4 of  $I_4$ ,

$$E = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{pmatrix},$$

## Row operations by matrix multiplication

**Swapping** rows ( $R_i \leftrightarrow R_j$ ) can be achieved by matrix multiplication.

For example, if

- $A$  is some  $4 \times 4$  matrix, and
- $E$  is made by swapping rows 2 and 4 of  $I_4$ ,

$$E = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{pmatrix},$$

then  $EA$  is  $A$  with rows 2 and 4 swapped. That is,

$$EA = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 13 & 14 & 15 & 16 \\ 9 & 10 & 11 & 12 \\ 5 & 6 & 7 & 8 \end{pmatrix}.$$

## Row operations by matrix multiplication

A row can be **scaled** ( $R_i \rightarrow \alpha R_i$  for  $\alpha \neq 0$ ) by matrix multiplication.



## Row operations by matrix multiplication

A row can be **scaled** ( $R_i \rightarrow \alpha R_i$  for  $\alpha \neq 0$ ) by matrix multiplication.

For example, if

- $A$  is some  $4 \times 4$  matrix,

## Row operations by matrix multiplication

A row can be **scaled** ( $R_i \rightarrow \alpha R_i$  for  $\alpha \neq 0$ ) by matrix multiplication.

For example, if

- $A$  is some  $4 \times 4$  matrix, and
- $E$  is made by multiplying the third row of  $I_4$  by 2,

## Row operations by matrix multiplication

A row can be **scaled** ( $R_i \rightarrow \alpha R_i$  for  $\alpha \neq 0$ ) by matrix multiplication.

For example, if

- $A$  is some  $4 \times 4$  matrix, and
- $E$  is made by multiplying the third row of  $I_4$  by 2,

$$E = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{pmatrix},$$

## Row operations by matrix multiplication

A row can be **scaled** ( $R_i \rightarrow \alpha R_i$  for  $\alpha \neq 0$ ) by matrix multiplication.

For example, if

- $A$  is some  $4 \times 4$  matrix, and
- $E$  is made by multiplying the third row of  $I_4$  by 2,

$$E = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{pmatrix},$$

then  $EA$  is  $A$  with  $R_3$  replaced by  $2R_3$ . That is,

$$EA = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 18 & 20 & 22 & 24 \\ 13 & 14 & 15 & 16 \end{pmatrix}.$$

## Row operations by matrix multiplication

Rows can **have multiples of other rows added to them** ( $R_i \rightarrow R_i + \alpha R_j$  for  $i \neq j$ ) by matrix multiplication.

## Row operations by matrix multiplication

Rows can **have multiples of other rows added to them** ( $R_i \rightarrow R_i + \alpha R_j$  for  $i \neq j$ ) by matrix multiplication.

For example, if

- $A$  is some  $4 \times 4$  matrix,

## Row operations by matrix multiplication

Rows can **have multiples of other rows added to them** ( $R_i \rightarrow R_i + \alpha R_j$  for  $i \neq j$ ) by matrix multiplication.

For example, if

- $A$  is some  $4 \times 4$  matrix, and
- $E$  is made by adding 3 lots of  $R_4$  to  $R_2$  in  $I_4$ ,

## Row operations by matrix multiplication

Rows can **have multiples of other rows added to them** ( $R_i \rightarrow R_i + \alpha R_j$  for  $i \neq j$ ) by matrix multiplication.

For example, if

- $A$  is some  $4 \times 4$  matrix, and
- $E$  is made by adding 3 lots of  $R_4$  to  $R_2$  in  $I_4$ ,

$$E = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{pmatrix},$$



## Row operations by matrix multiplication

Rows can **have multiples of other rows added to them** ( $R_i \rightarrow R_i + \alpha R_j$  for  $i \neq j$ ) by matrix multiplication.

For example, if

- $A$  is some  $4 \times 4$  matrix, and
- $E$  is made by adding 3 lots of  $R_4$  to  $R_2$  in  $I_4$ ,

$$E = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{pmatrix},$$

then  $EA$  is  $A$  with  $R_2$  replaced by  $R_2 + 3R_4$ . That is,

$$EA = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 44 & 48 & 52 & 56 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{pmatrix}.$$

# Matrix inverses

Suppose  $A$  is a square matrix, and the sequence of row operations that reduces it to **Reduced Row Echelon Form** (RREF) has corresponding matrices  $E_1, E_2, \dots, E_k$ .

# Matrix inverses

Suppose  $A$  is a square matrix, and the sequence of row operations that reduces it to **Reduced Row Echelon Form** (RREF) has corresponding matrices  $E_1, E_2, \dots, E_k$ .

Then

$$E_k \dots E_2 E_1 A = U$$

where  $U$  is some RREF matrix. (**Note that since matrix multiplication is not commutative, we must multiply on the left.**)

# Matrix inverses

Suppose  $A$  is a square matrix, and the sequence of row operations that reduces it to **Reduced Row Echelon Form** (RREF) has corresponding matrices  $E_1, E_2, \dots, E_k$ .

Then

$$E_k \dots E_2 E_1 A = U$$

where  $U$  is some RREF matrix. (Note that since matrix multiplication is not commutative, we must multiply on the left.)

If  $U = I$ , then we know  $A$  is invertible, and in fact

$$A^{-1} = E_k \dots E_2 E_1.$$

## Matrix inverses

Conversely, if  $A$  is invertible, then the system of equations  $A\mathbf{x} = \mathbf{b}$  has a unique solution  $\mathbf{x} = A^{-1}\mathbf{b}$ .

# Matrix inverses

Conversely, if  $A$  is invertible, then the system of equations  $A\mathbf{x} = \mathbf{b}$  has a unique solution  $\mathbf{x} = A^{-1}\mathbf{b}$ .

This means that when  $A$  is reduced to REF, it must have a leading entry in every column.

# Matrix inverses

Conversely, if  $A$  is invertible, then the system of equations  $A\mathbf{x} = \mathbf{b}$  has a unique solution  $\mathbf{x} = A^{-1}\mathbf{b}$ .

This means that when  $A$  is reduced to REF, it must have a leading entry in every column.

Since  $A$  is square, it must also have a leading entry in every row. That is, the leading entries of  $A$  are exactly the diagonal entries of  $A$ .

# Matrix inverses

Conversely, if  $A$  is invertible, then the system of equations  $A\mathbf{x} = \mathbf{b}$  has a unique solution  $\mathbf{x} = A^{-1}\mathbf{b}$ .

This means that when  $A$  is reduced to REF, it must have a leading entry in every column.

Since  $A$  is square, it must also have a leading entry in every row. That is, the leading entries of  $A$  are exactly the diagonal entries of  $A$ .

So when  $A$  is further reduced to RREF, each leading entry will become 1, and all other entries will become 0.



# Matrix inverses

Conversely, if  $A$  is invertible, then the system of equations  $A\mathbf{x} = \mathbf{b}$  has a unique solution  $\mathbf{x} = A^{-1}\mathbf{b}$ .

This means that when  $A$  is reduced to REF, it must have a leading entry in every column.

Since  $A$  is square, it must also have a leading entry in every row. That is, the leading entries of  $A$  are exactly the diagonal entries of  $A$ .

So when  $A$  is further reduced to RREF, each leading entry will become 1, and all other entries will become 0.

That is, the RREF of  $A$  is  $I$ .

# Matrix inverses

Conversely, if  $A$  is invertible, then the system of equations  $A\mathbf{x} = \mathbf{b}$  has a unique solution  $\mathbf{x} = A^{-1}\mathbf{b}$ .

This means that when  $A$  is reduced to REF, it must have a leading entry in every column.

Since  $A$  is square, it must also have a leading entry in every row. That is, the leading entries of  $A$  are exactly the diagonal entries of  $A$ .

So when  $A$  is further reduced to RREF, each leading entry will become 1, and all other entries will become 0.

That is, the RREF of  $A$  is  $I$ .

So whenever  $A$  is invertible, its RREF is  $I$ .

# Matrix inverses

If  $A$  is not invertible, then the equation  $A\mathbf{x} = \mathbf{b}$  has either no solutions or infinitely many solutions.

# Matrix inverses

If  $A$  is not invertible, then the equation  $A\mathbf{x} = \mathbf{b}$  has either no solutions or infinitely many solutions.

In both cases, this means any REF of  $A$  contains fewer leading entries than there are columns.

# Matrix inverses

If  $A$  is not invertible, then the equation  $A\mathbf{x} = \mathbf{b}$  has either no solutions or infinitely many solutions.

In both cases, this means any REF of  $A$  contains fewer leading entries than there are columns.

So  $A$  contains a zero row when reduced to REF (since  $A$  is square).

# Matrix inverses

If  $A$  is not invertible, then the equation  $A\mathbf{x} = \mathbf{b}$  has either no solutions or infinitely many solutions.

In both cases, this means any REF of  $A$  contains fewer leading entries than there are columns.

So  $A$  contains a zero row when reduced to REF (since  $A$  is square).

Any further reduction to RREF will certainly never reach  $I$ , since  $I$  contains no zero rows.

# Matrix inverses

If  $A$  is not invertible, then the equation  $A\mathbf{x} = \mathbf{b}$  has either no solutions or infinitely many solutions.

In both cases, this means any REF of  $A$  contains fewer leading entries than there are columns.

So  $A$  contains a zero row when reduced to REF (since  $A$  is square).

Any further reduction to RREF will certainly never reach  $I$ , since  $I$  contains no zero rows.

So whenever  $A$  is not invertible, its RREF is not  $I$ .

# Matrix inverses

Putting this all together, we now have a way to find the inverse of any square matrix  $A$ .



# Matrix inverses

Putting this all together, we now have a way to find the inverse of any square matrix  $A$ .

- Write the matrix  $A$  alongside the matrix  $I$ , separated by a bar:  $(A|I)$ .

# Matrix inverses

Putting this all together, we now have a way to find the inverse of any square matrix  $A$ .

- Write the matrix  $A$  alongside the matrix  $I$ , separated by a bar:  $(A|I)$ .
- Row-reduce  $A$  to RREF by applying elementary row operations. Simultaneously, apply the same row operations to the matrix right of the bar:  $(A|I) \rightarrow (U|B)$  where  $U$  is in RREF.

# Matrix inverses

Putting this all together, we now have a way to find the inverse of any square matrix  $A$ .

- Write the matrix  $A$  alongside the matrix  $I$ , separated by a bar:  $(A|I)$ .
- Row-reduce  $A$  to RREF by applying elementary row operations. Simultaneously, apply the same row operations to the matrix right of the bar:  $(A|I) \rightarrow (U|B)$  where  $U$  is in RREF.
- If  $U = I$ ,  $A$  is invertible and its inverse is  $B$ .

# Matrix inverses

Putting this all together, we now have a way to find the inverse of any square matrix  $A$ .

- Write the matrix  $A$  alongside the matrix  $I$ , separated by a bar:  $(A|I)$ .
- Row-reduce  $A$  to RREF by applying elementary row operations. Simultaneously, apply the same row operations to the matrix right of the bar:  $(A|I) \rightarrow (U|B)$  where  $U$  is in RREF.
- If  $U = I$ ,  $A$  is invertible and its inverse is  $B$ .
- If  $U \neq I$ ,  $A$  is not invertible.

# Matrix inverses – Examples

## Example

Find the inverse of  $A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 2 \\ 1 & -1 & 1 \end{pmatrix}$ .

# Matrix inverses – Examples

## Example

Find the inverse of  $A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 2 \\ 1 & -1 & 1 \end{pmatrix}$ .

First write  $(A|I)$ , and row-reduce the left matrix:

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 1 & 2 & 0 & 1 & 0 \\ 1 & -1 & 1 & 0 & 0 & 1 \end{array} \right)$$

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 1 & 2 & 0 & 1 & 0 \\ 1 & -1 & 1 & 0 & 0 & 1 \end{array} \right)$$

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 1 & 2 & 0 & 1 & 0 \\ 1 & -1 & 1 & 0 & 0 & 1 \end{array} \right) \xrightarrow[\substack{R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - R_1}]{\hspace{1cm}} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -4 & -2 & 1 & 0 \\ 0 & -3 & -2 & -1 & 0 & 1 \end{array} \right)$$



## Matrix inverses – Examples

$$\begin{pmatrix} 1 & 2 & 3 & | & 1 & 0 & 0 \\ 2 & 1 & 2 & | & 0 & 1 & 0 \\ 1 & -1 & 1 & | & 0 & 0 & 1 \end{pmatrix} \xrightarrow{\substack{R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - R_1}} \begin{pmatrix} 1 & 2 & 3 & | & 1 & 0 & 0 \\ 0 & -3 & -4 & | & -2 & 1 & 0 \\ 0 & -3 & -2 & | & -1 & 0 & 1 \end{pmatrix}$$
$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \begin{pmatrix} 1 & 2 & 3 & | & 1 & 0 & 0 \\ 0 & -3 & -4 & | & -2 & 1 & 0 \\ 0 & 0 & 2 & | & 1 & -1 & 1 \end{pmatrix}$$

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 1 & 2 & 0 & 1 & 0 \\ 1 & -1 & 1 & 0 & 0 & 1 \end{array} \right) \xrightarrow{\substack{R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - R_1}} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -4 & -2 & 1 & 0 \\ 0 & -3 & -2 & -1 & 0 & 1 \end{array} \right)$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -4 & -2 & 1 & 0 \\ 0 & 0 & 2 & 1 & -1 & 1 \end{array} \right)$$

$$\xrightarrow{\substack{R_2 \rightarrow -\frac{1}{3}R_2 \\ R_3 \rightarrow \frac{1}{2}R_3}} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & \frac{4}{3} & \frac{2}{3} & -\frac{1}{3} & 0 \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \end{array} \right)$$

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 1 & 2 & 0 & 1 & 0 \\ 1 & -1 & 1 & 0 & 0 & 1 \end{array} \right) \xrightarrow[\substack{R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - R_1}]{} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -4 & -2 & 1 & 0 \\ 0 & -3 & -2 & -1 & 0 & 1 \end{array} \right)$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -4 & -2 & 1 & 0 \\ 0 & 0 & 2 & 1 & -1 & 1 \end{array} \right)$$

$$\xrightarrow[\substack{R_2 \rightarrow -\frac{1}{3}R_2 \\ R_3 \rightarrow \frac{1}{2}R_3}]{} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & \frac{4}{3} & \frac{2}{3} & -\frac{1}{3} & 0 \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \end{array} \right)$$

$$\xrightarrow[\substack{R_2 \rightarrow R_2 - \frac{4}{3}R_3 \\ R_1 \rightarrow R_1 - 3R_3}]{} \left( \begin{array}{ccc|ccc} 1 & 2 & 0 & -\frac{1}{2} & \frac{3}{2} & -\frac{3}{2} \\ 0 & 1 & 0 & 0 & \frac{1}{3} & -\frac{2}{3} \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \end{array} \right)$$

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 1 & 2 & 0 & 1 & 0 \\ 1 & -1 & 1 & 0 & 0 & 1 \end{array} \right) \xrightarrow{\substack{R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - R_1}} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -4 & -2 & 1 & 0 \\ 0 & -3 & -2 & -1 & 0 & 1 \end{array} \right)$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -4 & -2 & 1 & 0 \\ 0 & 0 & 2 & 1 & -1 & 1 \end{array} \right)$$

$$\xrightarrow{\substack{R_2 \rightarrow -\frac{1}{3}R_2 \\ R_3 \rightarrow \frac{1}{2}R_3}} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & \frac{4}{3} & \frac{2}{3} & -\frac{1}{3} & 0 \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \end{array} \right)$$

$$\xrightarrow{\substack{R_2 \rightarrow R_2 - \frac{4}{3}R_3 \\ R_1 \rightarrow R_1 - 3R_3}} \left( \begin{array}{ccc|ccc} 1 & 2 & 0 & -\frac{1}{2} & \frac{3}{2} & -\frac{3}{2} \\ 0 & 1 & 0 & 0 & \frac{1}{3} & -\frac{2}{3} \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \end{array} \right)$$

$$\xrightarrow{R_1 \rightarrow R_1 - 2R_2} \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & -\frac{1}{2} & \frac{5}{6} & -\frac{1}{6} \\ 0 & 1 & 0 & 0 & \frac{1}{3} & -\frac{2}{3} \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \end{array} \right)$$

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 1 & 2 & 0 & 1 & 0 \\ 1 & -1 & 1 & 0 & 0 & 1 \end{array} \right) \xrightarrow{\dots} \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & -\frac{1}{2} & \frac{5}{6} & -\frac{1}{6} \\ 0 & 1 & 0 & 0 & \frac{1}{3} & -\frac{2}{3} \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \end{array} \right)$$

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 1 & 2 & 0 & 1 & 0 \\ 1 & -1 & 1 & 0 & 0 & 1 \end{array} \right) \xrightarrow{\dots} \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & -\frac{1}{2} & \frac{5}{6} & -\frac{1}{6} \\ 0 & 1 & 0 & 0 & \frac{1}{3} & -\frac{2}{3} \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \end{array} \right)$$

So  $A$  row-reduces to  $I$  in RREF, which means  $A$  is invertible.

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 1 & 2 & 0 & 1 & 0 \\ 1 & -1 & 1 & 0 & 0 & 1 \end{array} \right) \longrightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & -\frac{1}{2} & \frac{5}{6} & -\frac{1}{6} \\ 0 & 1 & 0 & 0 & \frac{1}{3} & -\frac{2}{3} \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \end{array} \right)$$

So  $A$  row-reduces to  $I$  in RREF, which means  $A$  is invertible.

Furthermore, the resultant right-hand matrix is the inverse of  $A$ , so we have found

$$A^{-1} = \left( \begin{array}{ccc} -\frac{1}{2} & \frac{5}{6} & -\frac{1}{6} \\ 0 & \frac{1}{3} & -\frac{2}{3} \\ \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \end{array} \right).$$

# Matrix inverses – Examples

## Example

Try to find the inverse of  $A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$ .



# Matrix inverses – Examples

## Example

Try to find the inverse of  $A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$ .

First write  $(A|I)$ , and row-reduce the left matrix:

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array} \right)$$

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array} \right)$$

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array} \right) \xrightarrow[\substack{R_2 \rightarrow R_2 - 4R_1 \\ R_3 \rightarrow R_3 - 7R_1}]{\phantom{R_2 \rightarrow R_2 - 4R_1}} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & -6 & -12 & -7 & 0 & 1 \end{array} \right)$$

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array} \right) \xrightarrow{\substack{R_2 \rightarrow R_2 - 4R_1 \\ R_3 \rightarrow R_3 - 7R_1}} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & -6 & -12 & -7 & 0 & 1 \end{array} \right)$$

$$\xrightarrow{R_3 \rightarrow R_3 - 2R_2} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 \end{array} \right)$$

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array} \right) \xrightarrow{\substack{R_2 \rightarrow R_2 - 4R_1 \\ R_3 \rightarrow R_3 - 7R_1}} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & -6 & -12 & -7 & 0 & 1 \end{array} \right)$$
$$\xrightarrow{R_3 \rightarrow R_3 - 2R_2} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 \end{array} \right)$$

At this point we know  $A$  will not row-reduce to become  $I$ , since its REF contains a zero row.

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array} \right) \xrightarrow{\substack{R_2 \rightarrow R_2 - 4R_1 \\ R_3 \rightarrow R_3 - 7R_1}} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & -6 & -12 & -7 & 0 & 1 \end{array} \right)$$
$$\xrightarrow{R_3 \rightarrow R_3 - 2R_2} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 \end{array} \right)$$

At this point we know  $A$  will not row-reduce to become  $I$ , since its REF **contains a zero row**.

So  $A$  is not invertible.

## Matrix inverses – Examples

$$\left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array} \right) \xrightarrow{\substack{R_2 \rightarrow R_2 - 4R_1 \\ R_3 \rightarrow R_3 - 7R_1}} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & -6 & -12 & -7 & 0 & 1 \end{array} \right)$$
$$\xrightarrow{R_3 \rightarrow R_3 - 2R_2} \left( \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 \end{array} \right)$$

At this point we know  $A$  will not row-reduce to become  $I$ , since its REF **contains a zero row**.

So  $A$  is not invertible.

It would be useful if we could decide if a matrix is not invertible before carrying out these steps. What we would like to have is something similar to the  $2 \times 2$  determinant, but for larger matrices.

# Determinants

We have seen how to find the determinant of a  $2 \times 2$  matrix. For a  $1 \times 1$  matrix, the determinant is just the single entry itself. We now look at larger matrices.



# Determinants

We have seen how to find the determinant of a  $2 \times 2$  matrix. For a  $1 \times 1$  matrix, the determinant is just the single entry itself. We now look at larger matrices.

## Definition

For a square matrix  $A$ , the  $(i, j)$ th minor, written  $|A_{ij}|$ , is the determinant of the submatrix obtained from  $A$  by deleting row  $i$  and column  $j$ .

# Determinants

We have seen how to find the determinant of a  $2 \times 2$  matrix. For a  $1 \times 1$  matrix, the determinant is just the single entry itself. We now look at larger matrices.

## Definition

For a square matrix  $A$ , the  $(i, j)$ th minor, written  $|A_{ij}|$ , is the determinant of the submatrix obtained from  $A$  by deleting row  $i$  and column  $j$ .

For example, if

$$A = \begin{pmatrix} 1 & 4 & 6 \\ 1 & -8 & 7 \\ 5 & 9 & 0 \end{pmatrix},$$

we have

$$|A_{23}|$$

# Determinants

We have seen how to find the determinant of a  $2 \times 2$  matrix. For a  $1 \times 1$  matrix, the determinant is just the single entry itself. We now look at larger matrices.

## Definition

For a square matrix  $A$ , the  $(i, j)$ th minor, written  $|A_{ij}|$ , is the determinant of the submatrix obtained from  $A$  by deleting row  $i$  and column  $j$ .

For example, if

$$A = \begin{pmatrix} 1 & 4 & 6 \\ 1 & -8 & 7 \\ 5 & 9 & 0 \end{pmatrix},$$

we have

$$|A_{23}|$$

# Determinants

We have seen how to find the determinant of a  $2 \times 2$  matrix. For a  $1 \times 1$  matrix, the determinant is just the single entry itself. We now look at larger matrices.

## Definition

For a square matrix  $A$ , the  $(i, j)$ th minor, written  $|A_{ij}|$ , is the determinant of the submatrix obtained from  $A$  by deleting row  $i$  and column  $j$ .

For example, if

$$A = \begin{pmatrix} 1 & 4 & 6 \\ 1 & -8 & 7 \\ 5 & 9 & 0 \end{pmatrix},$$

we have

$$|A_{23}| = \begin{vmatrix} 1 & 4 \\ 5 & 9 \end{vmatrix}$$

# Determinants

We have seen how to find the determinant of a  $2 \times 2$  matrix. For a  $1 \times 1$  matrix, the determinant is just the single entry itself. We now look at larger matrices.

## Definition

For a square matrix  $A$ , the  $(i, j)$ th minor, written  $|A_{ij}|$ , is the determinant of the submatrix obtained from  $A$  by deleting row  $i$  and column  $j$ .

For example, if

$$A = \begin{pmatrix} 1 & 4 & 6 \\ 1 & -8 & 7 \\ 5 & 9 & 0 \end{pmatrix},$$

we have

$$|A_{23}| = \begin{vmatrix} 1 & 4 \\ 5 & 9 \end{vmatrix} = 1 \times 9 - 5 \times 4 = -11.$$

# Determinants

## Definition

The **determinant** of an  $n \times n$  matrix  $A$  with entries  $a_{ij}$  is written as  **$\det(A)$**  or  **$|A|$** , and is given by:

$$\det(A) = a_{11}|A_{11}| - a_{12}|A_{12}| + a_{13}|A_{13}| - \cdots + (-1)^{n+1}a_{1n}|A_{1n}|$$

# Determinants

## Definition

The **determinant** of an  $n \times n$  matrix  $A$  with entries  $a_{ij}$  is written as  $\det(A)$  or  $|A|$ , and is given by:

$$\det(A) = a_{11}|A_{11}| - a_{12}|A_{12}| + a_{13}|A_{13}| - \cdots + (-1)^{n+1}a_{1n}|A_{1n}|$$

Note that each term contains a determinant of an  $(n-1) \times (n-1)$  matrix, so we need to apply this definition recursively until all the submatrices are at most size  $2 \times 2$ .

# Determinants

## Definition

The **determinant** of an  $n \times n$  matrix  $A$  with entries  $a_{ij}$  is written as  **$\det(A)$**  or  **$|A|$** , and is given by:

$$\det(A) = a_{11}|A_{11}| - a_{12}|A_{12}| + a_{13}|A_{13}| - \cdots + (-1)^{n+1}a_{1n}|A_{1n}|$$

Note that each term contains a determinant of an  $(n-1) \times (n-1)$  matrix, so we need to apply this definition recursively until all the submatrices are at most size  $2 \times 2$ .

Notice in particular that  **$\det(I) = 1$**  no matter the size of  $I$ .



# Determinants

## Definition

The **determinant** of an  $n \times n$  matrix  $A$  with entries  $a_{ij}$  is written as  $\det(A)$  or  $|A|$ , and is given by:

$$\det(A) = a_{11}|A_{11}| - a_{12}|A_{12}| + a_{13}|A_{13}| - \cdots + (-1)^{n+1}a_{1n}|A_{1n}|$$

Note that each term contains a determinant of an  $(n-1) \times (n-1)$  matrix, so we need to apply this definition recursively until all the submatrices are at most size  $2 \times 2$ .

Notice in particular that  $\det(I) = 1$  no matter the size of  $I$ .

This is because

$$\begin{aligned}\det(I_n) &= 1|I_{n-1}| - 0 + \cdots + 0 \\ &= 1|I_{n-2}| - 0 + \cdots + 0 \\ &= \cdots \\ &= 1.\end{aligned}$$

# Determinants – Example

## Example

Find

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix}.$$

# Determinants – Example

## Example

Find

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix}.$$

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix}$$

# Determinants – Example

## Example

Find

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix}.$$

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix} = 6 \det \begin{pmatrix} 2 & 1 \\ 4 & 3 \end{pmatrix} - 8 \det \begin{pmatrix} 5 & 1 \\ 7 & 3 \end{pmatrix} + 9 \det \begin{pmatrix} 5 & 2 \\ 7 & 4 \end{pmatrix}$$

# Determinants – Example

## Example

Find

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix}.$$

$$\begin{aligned} \det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix} &= 6 \det \begin{pmatrix} 2 & 1 \\ 4 & 3 \end{pmatrix} - 8 \det \begin{pmatrix} 5 & 1 \\ 7 & 3 \end{pmatrix} + 9 \det \begin{pmatrix} 5 & 2 \\ 7 & 4 \end{pmatrix} \\ &= 6 \times 2 - 8 \times 8 + 9 \times 6 \end{aligned}$$

# Determinants – Example

## Example

Find

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix}.$$

$$\begin{aligned} \det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix} &= 6 \det \begin{pmatrix} 2 & 1 \\ 4 & 3 \end{pmatrix} - 8 \det \begin{pmatrix} 5 & 1 \\ 7 & 3 \end{pmatrix} + 9 \det \begin{pmatrix} 5 & 2 \\ 7 & 4 \end{pmatrix} \\ &= 6 \times 2 - 8 \times 8 + 9 \times 6 \\ &= 2. \end{aligned}$$

# Determinants

The definition we just saw for the general determinant “expands” along the top row.

In fact, we can also expand along any other row or any column.

# Determinants

The definition we just saw for the general determinant “expands” along the top row.

In fact, we can also expand along any other row or any column.

The determinant is the sum of terms of the form  $(-1)^{i+j} a_{ij} |A_{ij}|$  **along any row or column**.



# Determinants

The definition we just saw for the general determinant “expands” along the top row.

In fact, we can also expand along any other row or any column.

The determinant is the sum of terms of the form  $(-1)^{i+j} a_{ij} |A_{ij}|$  **along any row or column**.

It's easy to remember the sign for each coefficient from the following checkerboard pattern:

$$\begin{pmatrix} + & - & + & - & \cdots \\ - & + & - & + & \cdots \\ + & - & + & - & \cdots \\ - & + & - & + & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

# Determinants – Examples

## Example

Find

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix}$$

by expanding along a row other than the top row or along a column.

# Determinants – Examples

## Example

Find

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix}$$

by expanding along a row other than the top row or along a column.

Expanding down the middle column:

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix}$$

# Determinants – Examples

## Example

Find

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix}$$

by expanding along a row other than the top row or along a column.

Expanding down the middle column:

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix} = -8 \det \begin{pmatrix} 5 & 1 \\ 7 & 3 \end{pmatrix} + 2 \det \begin{pmatrix} 6 & 9 \\ 7 & 3 \end{pmatrix} - 4 \det \begin{pmatrix} 6 & 9 \\ 5 & 1 \end{pmatrix}$$

# Determinants – Examples

## Example

Find

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix}$$

by expanding along a row other than the top row or along a column.

Expanding down the middle column:

$$\begin{aligned} \det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix} &= -8 \det \begin{pmatrix} 5 & 1 \\ 7 & 3 \end{pmatrix} + 2 \det \begin{pmatrix} 6 & 9 \\ 7 & 3 \end{pmatrix} - 4 \det \begin{pmatrix} 6 & 9 \\ 5 & 1 \end{pmatrix} \\ &= (-8) \times 8 + 2 \times (-45) - 4 \times (-39) \end{aligned}$$

## Determinants – Examples

### Example

Find

$$\det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix}$$

by expanding along a row other than the top row or along a column.

Expanding down the middle column:

$$\begin{aligned} \det \begin{pmatrix} 6 & 8 & 9 \\ 5 & 2 & 1 \\ 7 & 4 & 3 \end{pmatrix} &= -8 \det \begin{pmatrix} 5 & 1 \\ 7 & 3 \end{pmatrix} + 2 \det \begin{pmatrix} 6 & 9 \\ 7 & 3 \end{pmatrix} - 4 \det \begin{pmatrix} 6 & 9 \\ 5 & 1 \end{pmatrix} \\ &= (-8) \times 8 + 2 \times (-45) - 4 \times (-39) \\ &= 2. \end{aligned}$$

# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix}$$

# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix}$$

$$\det \begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix}$$



# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix}$$

$$\det \begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix} = 0 - 0 + 2 \det \begin{pmatrix} 4 & 5 & -1 \\ 2 & 0 & -3 \\ 8 & 0 & -5 \end{pmatrix} - 0$$

# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix}$$

$$\begin{aligned} \det \begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix} &= 0 - 0 + 2 \det \begin{pmatrix} 4 & 5 & -1 \\ 2 & 0 & -3 \\ 8 & 0 & -5 \end{pmatrix} - 0 \\ &= 2 \left( -5 \det \begin{pmatrix} 2 & -3 \\ 8 & -5 \end{pmatrix} + 0 - 0 \right) \end{aligned}$$

# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix}$$

$$\begin{aligned} \det \begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix} &= 0 - 0 + 2 \det \begin{pmatrix} 4 & 5 & -1 \\ 2 & 0 & -3 \\ 8 & 0 & -5 \end{pmatrix} - 0 \\ &= 2 \left( -5 \det \begin{pmatrix} 2 & -3 \\ 8 & -5 \end{pmatrix} + 0 - 0 \right) \\ &= 2 \times (-5) \times 14 \end{aligned}$$

# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix}$$

$$\begin{aligned} \det \begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix} &= 0 - 0 + 2 \det \begin{pmatrix} 4 & 5 & -1 \\ 2 & 0 & -3 \\ 8 & 0 & -5 \end{pmatrix} - 0 \\ &= 2 \left( -5 \det \begin{pmatrix} 2 & -3 \\ 8 & -5 \end{pmatrix} + 0 - 0 \right) \\ &= 2 \times (-5) \times 14 = -140. \end{aligned}$$

# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix}$$

$$\begin{aligned} \det \begin{pmatrix} 4 & 5 & 8 & -1 \\ 2 & 0 & -2 & -3 \\ 0 & 0 & 2 & 0 \\ 8 & 0 & -4 & -5 \end{pmatrix} &= 0 - 0 + 2 \det \begin{pmatrix} 4 & 5 & -1 \\ 2 & 0 & -3 \\ 8 & 0 & -5 \end{pmatrix} - 0 \\ &= 2 \left( -5 \det \begin{pmatrix} 2 & -3 \\ 8 & -5 \end{pmatrix} + 0 - 0 \right) \\ &= 2 \times (-5) \times 14 = -140. \end{aligned}$$

We expanded along the third row, and then the second column, because they contained the most zeros.

# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

$$\det \begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

$$\det \begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} = 4 \det \begin{pmatrix} 2 & -2 & -3 \\ 0 & 3 & 0 \\ 0 & 0 & -1 \end{pmatrix} - 0 + 0 - 0$$



# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

$$\begin{aligned} \det \begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} &= 4 \det \begin{pmatrix} 2 & -2 & -3 \\ 0 & 3 & 0 \\ 0 & 0 & -1 \end{pmatrix} - 0 + 0 - 0 \\ &= 4 \left( 2 \det \begin{pmatrix} 3 & 0 \\ 0 & -1 \end{pmatrix} - 0 + 0 \right) \end{aligned}$$

# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

$$\begin{aligned} \det \begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} &= 4 \det \begin{pmatrix} 2 & -2 & -3 \\ 0 & 3 & 0 \\ 0 & 0 & -1 \end{pmatrix} - 0 + 0 - 0 \\ &= 4 \left( 2 \det \begin{pmatrix} 3 & 0 \\ 0 & -1 \end{pmatrix} - 0 + 0 \right) \\ &= 4 \times 2 \times (-3) \end{aligned}$$

# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

$$\begin{aligned} \det \begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} &= 4 \det \begin{pmatrix} 2 & -2 & -3 \\ 0 & 3 & 0 \\ 0 & 0 & -1 \end{pmatrix} - 0 + 0 - 0 \\ &= 4 \left( 2 \det \begin{pmatrix} 3 & 0 \\ 0 & -1 \end{pmatrix} - 0 + 0 \right) \\ &= 4 \times 2 \times (-3) = -24. \end{aligned}$$

# Determinants – Examples

## Example

Find the determinant of

$$\begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

$$\begin{aligned} \det \begin{pmatrix} 4 & 1 & 7 & -1 \\ 0 & 2 & -2 & -3 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} &= 4 \det \begin{pmatrix} 2 & -2 & -3 \\ 0 & 3 & 0 \\ 0 & 0 & -1 \end{pmatrix} - 0 + 0 - 0 \\ &= 4 \left( 2 \det \begin{pmatrix} 3 & 0 \\ 0 & -1 \end{pmatrix} - 0 + 0 \right) \\ &= 4 \times 2 \times (-3) = -24. \end{aligned}$$

We expanded along the first column, and then the first column again, because they contained the most zeros.

# Determinants

The determinant of an **upper triangular matrix** (a matrix with zeros below every diagonal entry) is the product of all its diagonal entries:

# Determinants

The determinant of an **upper triangular matrix** (a matrix with zeros below every diagonal entry) is the product of all its diagonal entries:

$$\begin{aligned} \det \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} & \cdots & a_{1n} \\ 0 & a_{22} & a_{23} & a_{24} & \cdots & a_{2n} \\ 0 & 0 & a_{33} & a_{34} & \cdots & a_{3n} \\ 0 & 0 & 0 & a_{44} & \cdots & a_{4n} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & a_{nn} \end{pmatrix} \\ = a_{11} \det \begin{pmatrix} a_{22} & a_{23} & a_{24} & \cdots & a_{2n} \\ 0 & a_{33} & a_{34} & \cdots & a_{3n} \\ 0 & 0 & a_{44} & \cdots & a_{4n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a_{nn} \end{pmatrix} + 0 + \cdots + 0 \\ = \dots \\ = a_{11} a_{22} a_{33} a_{44} \cdots a_{nn}. \end{aligned}$$