



UNSW
SYDNEY

MATH1131 Mathematics 1A – Algebra


Lecture 8: Triple Scalar Product and the Point-Normal Form

Lecturer: Sean Gardiner – sean.gardiner@unsw.edu.au

Based on slides by Jonathan Kress


Parallelepiped

The 3D version of a parallelogram is called the **parallelepiped**.




Parallelepiped

The 3D version of a parallelogram is called the **parallelepiped**.



Parallelepiped


The 3D version of a parallelogram is called the **parallelepiped**.



$$\text{Volume} = \text{area of base} \times \text{altitude}$$

Parallelepiped


The 3D version of a parallelogram is called the **parallelepiped**.



$$\text{Volume} = \text{area of base} \times \text{altitude}$$

Parallelepiped


The 3D version of a parallelogram is called the **parallelepiped**.



$$\begin{aligned}\text{Volume} &= \text{area of base} \times \text{altitude} \\ &= |\mathbf{b} \times \mathbf{c}| |\mathbf{a}| \cos \theta\end{aligned}$$

Parallelepiped

The 3D version of a parallelogram is called the **parallelepiped**.



$$\begin{aligned}\text{Volume} &= \text{area of base} \times \text{altitude} \\ &= |\mathbf{b} \times \mathbf{c}| |\mathbf{a}| \cos \theta \\ &= |\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})|\end{aligned}$$

Scalar triple product

Definition

The **triple scalar product** of vectors \mathbf{a} , \mathbf{b} , $\mathbf{c} \in \mathbb{R}^3$ is

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}).$$

Scalar triple product

Definition

The **triple scalar product** of vectors \mathbf{a} , \mathbf{b} , $\mathbf{c} \in \mathbb{R}^3$ is

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}).$$

The scalar triple product can also be written as a 3×3 determinant:

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \cdot \begin{pmatrix} b_2 c_3 - b_3 c_2 \\ b_3 c_1 - b_1 c_3 \\ b_1 c_2 - b_2 c_1 \end{pmatrix}$$

Scalar triple product

Definition

The **triple scalar product** of vectors \mathbf{a} , \mathbf{b} , $\mathbf{c} \in \mathbb{R}^3$ is

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}).$$

The scalar triple product can also be written as a 3×3 determinant:

$$\begin{aligned}\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) &= \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \cdot \begin{pmatrix} b_2 c_3 - b_3 c_2 \\ b_3 c_1 - b_1 c_3 \\ b_1 c_2 - b_2 c_1 \end{pmatrix} \\ &= a_1 \begin{vmatrix} b_2 & b_3 \\ c_2 & c_3 \end{vmatrix} - a_2 \begin{vmatrix} b_1 & b_3 \\ c_1 & c_3 \end{vmatrix} + a_3 \begin{vmatrix} b_1 & b_2 \\ c_1 & c_2 \end{vmatrix}\end{aligned}$$

Scalar triple product

Definition

The **triple scalar product** of vectors \mathbf{a} , \mathbf{b} , $\mathbf{c} \in \mathbb{R}^3$ is

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}).$$

The scalar triple product can also be written as a 3×3 determinant:

$$\begin{aligned}\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) &= \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \cdot \begin{pmatrix} b_2 c_3 - b_3 c_2 \\ b_3 c_1 - b_1 c_3 \\ b_1 c_2 - b_2 c_1 \end{pmatrix} \\ &= a_1 \begin{vmatrix} b_2 & b_3 \\ c_2 & c_3 \end{vmatrix} - a_2 \begin{vmatrix} b_1 & b_3 \\ c_1 & c_3 \end{vmatrix} + a_3 \begin{vmatrix} b_1 & b_2 \\ c_1 & c_2 \end{vmatrix} \\ &= \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}\end{aligned}$$

Scalar triple product

Properties

For all $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{R}^3$

- $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \mathbf{b} \cdot (\mathbf{c} \times \mathbf{a}) = \mathbf{c} \cdot (\mathbf{a} \times \mathbf{b})$
- $\mathbf{a} \cdot (\mathbf{c} \times \mathbf{b}) = -\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$
- $\mathbf{a} \cdot (\mathbf{a} \times \mathbf{b}) = 0.$

Scalar triple product

Properties

For all $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{R}^3$

- $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \mathbf{b} \cdot (\mathbf{c} \times \mathbf{a}) = \mathbf{c} \cdot (\mathbf{a} \times \mathbf{b})$
- $\mathbf{a} \cdot (\mathbf{c} \times \mathbf{b}) = -\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$
- $\mathbf{a} \cdot (\mathbf{a} \times \mathbf{b}) = 0.$

Proof

By expanding and rearranging terms,

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$$

$$= a_1(b_2c_3 - b_3c_2) + a_2(b_3c_1 - b_1c_3) + a_3(b_1c_2 - b_2c_1)$$

$$= b_1(c_2a_3 - c_3a_2) + b_2(c_3a_1 - c_1a_3) + b_3(c_1a_2 - c_2a_1) = \mathbf{b} \cdot (\mathbf{c} \times \mathbf{a})$$

$$= c_1(a_2b_3 - a_3b_2) + c_2(a_3b_1 - a_1b_3) + c_3(a_1b_2 - a_2b_1) = \mathbf{c} \cdot (\mathbf{a} \times \mathbf{b})$$

Scalar triple product

Properties

For all $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{R}^3$

- $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \mathbf{b} \cdot (\mathbf{c} \times \mathbf{a}) = \mathbf{c} \cdot (\mathbf{a} \times \mathbf{b})$
- $\mathbf{a} \cdot (\mathbf{c} \times \mathbf{b}) = -\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$
- $\mathbf{a} \cdot (\mathbf{a} \times \mathbf{b}) = 0.$

Proof

Using properties of the cross product (anti-commutativity) and the dot product (associative law of scalar multiplication),

$$\begin{aligned}\mathbf{a} \cdot (\mathbf{c} \times \mathbf{b}) &= \mathbf{a} \cdot (-1)(\mathbf{b} \times \mathbf{c}) \\ &= -\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})\end{aligned}$$

Scalar triple product

Properties

For all $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{R}^3$

- $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \mathbf{b} \cdot (\mathbf{c} \times \mathbf{a}) = \mathbf{c} \cdot (\mathbf{a} \times \mathbf{b})$
- $\mathbf{a} \cdot (\mathbf{c} \times \mathbf{b}) = -\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$
- $\mathbf{a} \cdot (\mathbf{a} \times \mathbf{b}) = 0.$

Proof

Using the first property above and properties of the dot and cross products,

$$\begin{aligned}\mathbf{a} \cdot (\mathbf{a} \times \mathbf{b}) &= \mathbf{b} \cdot (\mathbf{a} \times \mathbf{a}) \\ &= \mathbf{b} \cdot \mathbf{0} \\ &= 0\end{aligned}$$

Scalar triple product – Examples

Example

Find the volume of the parallelepiped with edges given by the vectors

$$\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \quad \begin{pmatrix} -3 \\ -2 \\ 7 \end{pmatrix}, \quad \begin{pmatrix} -4 \\ 3 \\ 7 \end{pmatrix}.$$

Scalar triple product – Examples

Example

Find the volume of the parallelepiped with edges given by the vectors

$$\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \quad \begin{pmatrix} -3 \\ -2 \\ 7 \end{pmatrix}, \quad \begin{pmatrix} -4 \\ 3 \\ 7 \end{pmatrix}.$$

$$\text{Volume} = |\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})|$$

Scalar triple product – Examples

Example

Find the volume of the parallelepiped with edges given by the vectors

$$\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \quad \begin{pmatrix} -3 \\ -2 \\ 7 \end{pmatrix}, \quad \begin{pmatrix} -4 \\ 3 \\ 7 \end{pmatrix}.$$

$$\text{Volume} = |\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})| = \left| \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \cdot \left(\begin{pmatrix} -3 \\ -2 \\ 7 \end{pmatrix} \times \begin{pmatrix} -4 \\ 3 \\ 7 \end{pmatrix} \right) \right|$$

Scalar triple product – Examples

Example

Find the volume of the parallelepiped with edges given by the vectors

$$\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \quad \begin{pmatrix} -3 \\ -2 \\ 7 \end{pmatrix}, \quad \begin{pmatrix} -4 \\ 3 \\ 7 \end{pmatrix}.$$

$$\begin{aligned} \text{Volume} &= |\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})| = \left| \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \cdot \left(\begin{pmatrix} -3 \\ -2 \\ 7 \end{pmatrix} \times \begin{pmatrix} -4 \\ 3 \\ 7 \end{pmatrix} \right) \right| \\ &= \left| \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} -35 \\ -7 \\ -17 \end{pmatrix} \right| \end{aligned}$$

Scalar triple product – Examples

Example

Find the volume of the parallelepiped with edges given by the vectors

$$\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \quad \begin{pmatrix} -3 \\ -2 \\ 7 \end{pmatrix}, \quad \begin{pmatrix} -4 \\ 3 \\ 7 \end{pmatrix}.$$

$$\begin{aligned} \text{Volume} &= |\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})| = \left| \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \cdot \left(\begin{pmatrix} -3 \\ -2 \\ 7 \end{pmatrix} \times \begin{pmatrix} -4 \\ 3 \\ 7 \end{pmatrix} \right) \right| \\ &= \left| \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} -35 \\ -7 \\ -17 \end{pmatrix} \right| \\ &= |-100| \end{aligned}$$

Scalar triple product – Examples

Example

Find the volume of the parallelepiped with edges given by the vectors

$$\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \quad \begin{pmatrix} -3 \\ -2 \\ 7 \end{pmatrix}, \quad \begin{pmatrix} -4 \\ 3 \\ 7 \end{pmatrix}.$$

$$\begin{aligned} \text{Volume} &= |\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})| = \left| \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \cdot \left(\begin{pmatrix} -3 \\ -2 \\ 7 \end{pmatrix} \times \begin{pmatrix} -4 \\ 3 \\ 7 \end{pmatrix} \right) \right| \\ &= \left| \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} -35 \\ -7 \\ -17 \end{pmatrix} \right| \\ &= |-100| \\ &= 100 \end{aligned}$$

Scalar triple product – Examples

Example

Show that the points $A(3, 3, 5)$, $B(1, 0, 1)$, $C(2, 2, 4)$ and $D(2, 1, 2)$ are coplanar.

Scalar triple product – Examples

Example

Show that the points $A(3, 3, 5)$, $B(1, 0, 1)$, $C(2, 2, 4)$ and $D(2, 1, 2)$ are coplanar.

Three vectors all lie in the same plane **if and only if** the parallelepiped spanned by them has zero volume.

Scalar triple product – Examples

Example

Show that the points $A(3, 3, 5)$, $B(1, 0, 1)$, $C(2, 2, 4)$ and $D(2, 1, 2)$ are coplanar.

Three vectors all lie in the same plane **if and only if** the parallelepiped spanned by them has zero volume.

$$\text{Volume} = |\overrightarrow{AB} \cdot (\overrightarrow{AC} \times \overrightarrow{AD})|$$

Scalar triple product – Examples

Example

Show that the points $A(3, 3, 5)$, $B(1, 0, 1)$, $C(2, 2, 4)$ and $D(2, 1, 2)$ are coplanar.

Three vectors all lie in the same plane **if and only if** the parallelepiped spanned by them has zero volume.

$$\text{Volume} = |\overrightarrow{AB} \cdot (\overrightarrow{AC} \times \overrightarrow{AD})| = \left| \begin{pmatrix} -2 \\ -3 \\ -4 \end{pmatrix} \cdot \left(\begin{pmatrix} -1 \\ -1 \\ -1 \end{pmatrix} \times \begin{pmatrix} -1 \\ -2 \\ -3 \end{pmatrix} \right) \right|$$

Scalar triple product – Examples

Example

Show that the points $A(3, 3, 5)$, $B(1, 0, 1)$, $C(2, 2, 4)$ and $D(2, 1, 2)$ are coplanar.

Three vectors all lie in the same plane **if and only if** the parallelepiped spanned by them has zero volume.

$$\begin{aligned}\text{Volume} &= |\overrightarrow{AB} \cdot (\overrightarrow{AC} \times \overrightarrow{AD})| = \left| \begin{pmatrix} -2 \\ -3 \\ -4 \end{pmatrix} \cdot \left(\begin{pmatrix} -1 \\ -1 \\ -1 \end{pmatrix} \times \begin{pmatrix} -1 \\ -2 \\ -3 \end{pmatrix} \right) \right| \\ &= \left| \begin{pmatrix} -2 \\ -3 \\ -4 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix} \right|\end{aligned}$$

Scalar triple product – Examples

Example

Show that the points $A(3, 3, 5)$, $B(1, 0, 1)$, $C(2, 2, 4)$ and $D(2, 1, 2)$ are coplanar.

Three vectors all lie in the same plane **if and only if** the parallelepiped spanned by them has zero volume.

$$\begin{aligned}\text{Volume} &= |\overrightarrow{AB} \cdot (\overrightarrow{AC} \times \overrightarrow{AD})| = \left| \begin{pmatrix} -2 \\ -3 \\ -4 \end{pmatrix} \cdot \left(\begin{pmatrix} -1 \\ -1 \\ -1 \end{pmatrix} \times \begin{pmatrix} -1 \\ -2 \\ -3 \end{pmatrix} \right) \right| \\ &= \left| \begin{pmatrix} -2 \\ -3 \\ -4 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix} \right| \\ &= 0\end{aligned}$$

Scalar triple product – Examples

Example


Show that the points $A(3, 3, 5)$, $B(1, 0, 1)$, $C(2, 2, 4)$ and $D(2, 1, 2)$ are coplanar.

Three vectors all lie in the same plane **if and only if** the parallelepiped spanned by them has zero volume.

$$\begin{aligned}\text{Volume} &= |\overrightarrow{AB} \cdot (\overrightarrow{AC} \times \overrightarrow{AD})| = \left| \begin{pmatrix} -2 \\ -3 \\ -4 \end{pmatrix} \cdot \left(\begin{pmatrix} -1 \\ -1 \\ -1 \end{pmatrix} \times \begin{pmatrix} -1 \\ -2 \\ -3 \end{pmatrix} \right) \right| \\ &= \left| \begin{pmatrix} -2 \\ -3 \\ -4 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix} \right| \\ &= 0\end{aligned}$$

So the four points are coplanar.

Parametric vector form of a plane




Recall: A plane parallel to two (non-parallel) vectors \mathbf{v}_1 and \mathbf{v}_2 , and containing the point with position vector \mathbf{a} , is described by:

$$\mathbf{x} = \mathbf{a} + \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2, \quad \lambda_1, \lambda_2 \in \mathbb{R}.$$

We called this the **parametric vector form** of the plane.

Parametric vector form of a plane




Recall: A plane parallel to two (non-parallel) vectors \mathbf{v}_1 and \mathbf{v}_2 , and containing the point with position vector \mathbf{a} , is described by:

$$\mathbf{x} = \mathbf{a} + \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2, \quad \lambda_1, \lambda_2 \in \mathbb{R}.$$

We called this the **parametric vector form** of the plane.
The linear combinations $\lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2$ are all parallel to the plane.

Parametric vector form of a plane



Recall: A plane parallel to two (non-parallel) vectors \mathbf{v}_1 and \mathbf{v}_2 , and containing the point with position vector \mathbf{a} , is described by:


$$\mathbf{x} = \mathbf{a} + \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2, \quad \lambda_1, \lambda_2 \in \mathbb{R}.$$

We called this the **parametric vector form** of the plane.

The linear combinations $\lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2$ are all parallel to the plane.


The **normal** vector $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2$ is perpendicular to all of these.

Point-normal form of a plane



The **normal** vector $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2$ is perpendicular to the plane.

Point-normal form of a plane




The **normal** vector $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2$ is perpendicular to the plane.

In particular, $\mathbf{x} - \mathbf{a}$ is perpendicular to \mathbf{n} for all \mathbf{x} in the plane, so

$$\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0.$$

Point-normal form of a plane



The **normal** vector $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2$ is perpendicular to the plane.

In particular, $\mathbf{x} - \mathbf{a}$ is perpendicular to \mathbf{n} for all \mathbf{x} in the plane, so

$$\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0.$$

This is called the **point-normal form** of the plane.

It can also be written in the form $\mathbf{n} \cdot \mathbf{x} = \mathbf{n} \cdot \mathbf{a}$.

Cartesian and point-normal forms

Suppose $\mathbf{n} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix}$ is a vector normal to some plane passing through the point $A(a_1, a_2, a_3)$.

Cartesian and point-normal forms

Suppose $\mathbf{n} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix}$ is a vector normal to some plane passing through the point $A(a_1, a_2, a_3)$.

The point-normal form of this plane is:

$$\mathbf{n} \cdot \mathbf{x} = \mathbf{n} \cdot \mathbf{a}$$

Cartesian and point-normal forms

Suppose $\mathbf{n} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix}$ is a vector normal to some plane passing through the point $A(a_1, a_2, a_3)$.

The point-normal form of this plane is:

$$\mathbf{n} \cdot \mathbf{x} = \mathbf{n} \cdot \mathbf{a}$$

$$\begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} \cdot \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$$

Cartesian and point-normal forms

Suppose $\mathbf{n} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix}$ is a vector normal to some plane passing through the point $A(a_1, a_2, a_3)$.

The point-normal form of this plane is:

$$\mathbf{n} \cdot \mathbf{x} = \mathbf{n} \cdot \mathbf{a}$$

$$\begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} \cdot \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$$

$$n_1x + n_2y + n_3z = n_1a_1 + n_2a_2 + n_3a_3$$

Cartesian and point-normal forms

Suppose $\mathbf{n} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix}$ is a vector normal to some plane passing through the point $A(a_1, a_2, a_3)$.

The point-normal form of this plane is:

$$\mathbf{n} \cdot \mathbf{x} = \mathbf{n} \cdot \mathbf{a}$$

$$\begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} \cdot \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$$

$$n_1x + n_2y + n_3z = n_1a_1 + n_2a_2 + n_3a_3$$

or we could write:

$$n_1x + n_2y + n_3z = d, \quad \text{where } d = \mathbf{n} \cdot \mathbf{a}$$

Cartesian and point-normal forms

Suppose $\mathbf{n} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix}$ is a vector normal to some plane passing through the point $A(a_1, a_2, a_3)$.

The point-normal form of this plane is:

$$\mathbf{n} \cdot \mathbf{x} = \mathbf{n} \cdot \mathbf{a}$$

$$\begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} \cdot \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$$

$$n_1x + n_2y + n_3z = n_1a_1 + n_2a_2 + n_3a_3$$

or we could write:

$$n_1x + n_2y + n_3z = d, \quad \text{where } d = \mathbf{n} \cdot \mathbf{a}$$

This is recognisable as a **Cartesian form** of the plane ($ax + by + cz = d$ for some $a, b, c, d \in \mathbb{R}$ with at least one of a, b , and c non-zero).

Cartesian and point-normal forms – Examples

Example

Find a Cartesian equation of the plane with normal $\begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix}$ that passes through the point $(1, 2, 1)$.

Cartesian and point-normal forms – Examples

Example

Find a Cartesian equation of the plane with normal $\begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix}$ that passes through the point $(1, 2, 1)$.

The Cartesian form will be $4x + (-2)y + 3z = d$, where:

$$d = \mathbf{n} \cdot \mathbf{a}$$

Cartesian and point-normal forms – Examples

Example

Find a Cartesian equation of the plane with normal $\begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix}$ that passes through the point $(1, 2, 1)$.

The Cartesian form will be $4x + (-2)y + 3z = d$, where:

$$d = \mathbf{n} \cdot \mathbf{a} = \begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$$

Cartesian and point-normal forms – Examples

Example

Find a Cartesian equation of the plane with normal $\begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix}$ that passes through the point $(1, 2, 1)$.

The Cartesian form will be $4x + (-2)y + 3z = d$, where:

$$d = \mathbf{n} \cdot \mathbf{a} = \begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} = 3.$$

Cartesian and point-normal forms – Examples

Example

Find a Cartesian equation of the plane with normal $\begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix}$ that passes through the point $(1, 2, 1)$.

The Cartesian form will be $4x + (-2)y + 3z = d$, where:

$$d = \mathbf{n} \cdot \mathbf{a} = \begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} = 3.$$

That is, a Cartesian form for the plane is $4x - 2y + 3z = 3$.

Cartesian and point-normal forms – Examples

Example

Write the plane

$$\mathbf{x} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda_1 \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} + \lambda_2 \begin{pmatrix} 5 \\ -4 \\ 1 \end{pmatrix}, \quad \lambda_1, \lambda_2 \in \mathbb{R},$$

in point-normal form and in Cartesian form.

Cartesian and point-normal forms – Examples

Example

Write the plane

$$\mathbf{x} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda_1 \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} + \lambda_2 \begin{pmatrix} 5 \\ -4 \\ 1 \end{pmatrix}, \quad \lambda_1, \lambda_2 \in \mathbb{R},$$

in point-normal form and in Cartesian form.

To find a normal vector \mathbf{n} to the plane, we can take the cross product of the two parallel vectors \mathbf{v}_1 and \mathbf{v}_2 :

$$\mathbf{n} = \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} \times \begin{pmatrix} 5 \\ -4 \\ 1 \end{pmatrix}$$

Cartesian and point-normal forms – Examples

Example

Write the plane

$$\mathbf{x} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda_1 \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} + \lambda_2 \begin{pmatrix} 5 \\ -4 \\ 1 \end{pmatrix}, \quad \lambda_1, \lambda_2 \in \mathbb{R},$$

in point-normal form and in Cartesian form.

To find a normal vector \mathbf{n} to the plane, we can take the cross product of the two parallel vectors \mathbf{v}_1 and \mathbf{v}_2 :

$$\mathbf{n} = \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} \times \begin{pmatrix} 5 \\ -4 \\ 1 \end{pmatrix} = \begin{pmatrix} 11 \\ 13 \\ -3 \end{pmatrix}.$$

Cartesian and point-normal forms – Examples

Example

Write the plane

$$\mathbf{x} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda_1 \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} + \lambda_2 \begin{pmatrix} 5 \\ -4 \\ 1 \end{pmatrix}, \quad \lambda_1, \lambda_2 \in \mathbb{R},$$

in point-normal form and in Cartesian form.

Since we know the plane contains the point $(1, 2, 3)$, we can write a point-normal form as:

Cartesian and point-normal forms – Examples

Example

Write the plane

$$\mathbf{x} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda_1 \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} + \lambda_2 \begin{pmatrix} 5 \\ -4 \\ 1 \end{pmatrix}, \quad \lambda_1, \lambda_2 \in \mathbb{R},$$

in point-normal form and in Cartesian form.

Since we know the plane contains the point $(1, 2, 3)$, we can write a point-normal form as:

$$\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = \begin{pmatrix} 11 \\ 13 \\ -3 \end{pmatrix} \cdot \left(\mathbf{x} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \right) = 0.$$

Cartesian and point-normal forms – Examples

Example

Write the plane

$$\mathbf{x} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda_1 \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} + \lambda_2 \begin{pmatrix} 5 \\ -4 \\ 1 \end{pmatrix}, \quad \lambda_1, \lambda_2 \in \mathbb{R},$$

in point-normal form and in Cartesian form.

Since we know the plane contains the point $(1, 2, 3)$, we can write a point-normal form as:

$$\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = \begin{pmatrix} 11 \\ 13 \\ -3 \end{pmatrix} \cdot \left(\mathbf{x} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \right) = 0.$$

Similarly we can write a Cartesian form as:

$$11x + 13y - 3z = \mathbf{n} \cdot \mathbf{a} = 28.$$

Cartesian and point-normal forms – Examples

Example

Write the plane $2x + 9y - 7z = 21$ in point-normal form.

Cartesian and point-normal forms – Examples

Example

Write the plane $2x + 9y - 7z = 21$ in point-normal form.

One normal vector \mathbf{n} to the plane must be $\mathbf{n} = \begin{pmatrix} 2 \\ 9 \\ -7 \end{pmatrix}$.

Cartesian and point-normal forms – Examples

Example

Write the plane $2x + 9y - 7z = 21$ in point-normal form.

One normal vector \mathbf{n} to the plane must be $\mathbf{n} = \begin{pmatrix} 2 \\ 9 \\ -7 \end{pmatrix}$.

To find a point on the plane, set any two of the unknowns to be 0 and solve for the third unknown.

Cartesian and point-normal forms – Examples

Example

Write the plane $2x + 9y - 7z = 21$ in point-normal form.

One normal vector \mathbf{n} to the plane must be $\mathbf{n} = \begin{pmatrix} 2 \\ 9 \\ -7 \end{pmatrix}$.

To find a point on the plane, set any two of the unknowns to be 0 and solve for the third unknown.

For example, setting $x = y = 0$ gives $-7z = 21$

Cartesian and point-normal forms – Examples

Example

Write the plane $2x + 9y - 7z = 21$ in point-normal form.

One normal vector \mathbf{n} to the plane must be $\mathbf{n} = \begin{pmatrix} 2 \\ 9 \\ -7 \end{pmatrix}$.

To find a point on the plane, set any two of the unknowns to be 0 and solve for the third unknown.

For example, setting $x = y = 0$ gives $-7z = 21 \implies z = -3$.

Cartesian and point-normal forms – Examples

Example

Write the plane $2x + 9y - 7z = 21$ in point-normal form.

One normal vector \mathbf{n} to the plane must be $\mathbf{n} = \begin{pmatrix} 2 \\ 9 \\ -7 \end{pmatrix}$.

To find a point on the plane, set any two of the unknowns to be 0 and solve for the third unknown.

For example, setting $x = y = 0$ gives $-7z = 21 \implies z = -3$.

So $(0, 0, -3)$ is a point on the plane.

Cartesian and point-normal forms – Examples

Example

Write the plane $2x + 9y - 7z = 21$ in point-normal form.

One normal vector \mathbf{n} to the plane must be $\mathbf{n} = \begin{pmatrix} 2 \\ 9 \\ -7 \end{pmatrix}$.

To find a point on the plane, set any two of the unknowns to be 0 and solve for the third unknown.

For example, setting $x = y = 0$ gives $-7z = 21 \implies z = -3$.

So $(0, 0, -3)$ is a point on the plane.

We can therefore write a point-normal form of the plane as:

$$\begin{pmatrix} 2 \\ 9 \\ -7 \end{pmatrix} \cdot \left(\mathbf{x} - \begin{pmatrix} 0 \\ 0 \\ -3 \end{pmatrix} \right) = 0.$$

Summary of plane equations – Vector parametric form

Given a plane in **vector parametric form**:

$$\mathbf{x} = \mathbf{a} + \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2, \quad \lambda_1, \lambda_2 \in \mathbb{R}$$

Summary of plane equations – Vector parametric form

Given a plane in **vector parametric form**:

$$\mathbf{x} = \mathbf{a} + \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2, \quad \lambda_1, \lambda_2 \in \mathbb{R}$$

- **Vectors parallel to the plane** include \mathbf{v}_1 , \mathbf{v}_2 , $2\mathbf{v}_1 - 5\mathbf{v}_2$, etc.

Summary of plane equations – Vector parametric form

Given a plane in **vector parametric form**:

$$\mathbf{x} = \mathbf{a} + \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2, \quad \lambda_1, \lambda_2 \in \mathbb{R}$$

- **Vectors parallel to the plane** include \mathbf{v}_1 , \mathbf{v}_2 , $2\mathbf{v}_1 - 5\mathbf{v}_2$, etc.
- Position vectors for **points on the plane** include \mathbf{a} , $\mathbf{a} + \mathbf{v}_1$, $\mathbf{a} + \mathbf{v}_2$, $\mathbf{a} + 2\mathbf{v}_1 - 5\mathbf{v}_2$, etc.

Summary of plane equations – Vector parametric form

Given a plane in **vector parametric form**:

$$\mathbf{x} = \mathbf{a} + \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2, \quad \lambda_1, \lambda_2 \in \mathbb{R}$$

- **Vectors parallel to the plane** include \mathbf{v}_1 , \mathbf{v}_2 , $2\mathbf{v}_1 - 5\mathbf{v}_2$, etc.
- Position vectors for **points on the plane** include \mathbf{a} , $\mathbf{a} + \mathbf{v}_1$, $\mathbf{a} + \mathbf{v}_2$, $\mathbf{a} + 2\mathbf{v}_1 - 5\mathbf{v}_2$, etc.
- A **normal vector to the plane** is $\mathbf{v}_1 \times \mathbf{v}_2$.

Summary of plane equations – Cartesian form

Given a plane in **Cartesian form**:

$$ax + by + cz = d, \quad a, b, c, d \in \mathbb{R}$$

Summary of plane equations – Cartesian form

Given a plane in **Cartesian form**:

$$ax + by + cz = d, \quad a, b, c, d \in \mathbb{R}$$

- A **normal vector to the plane** is $\begin{pmatrix} a \\ b \\ c \end{pmatrix}$.

Summary of plane equations – Cartesian form

Given a plane in **Cartesian form**:

$$ax + by + cz = d, \quad a, b, c, d \in \mathbb{R}$$

- A **normal vector to the plane** is $\begin{pmatrix} a \\ b \\ c \end{pmatrix}$.
- **Points on the plane** include:
 - $(\frac{d}{a}, 0, 0)$ (if $a \neq 0$)
 - $(0, \frac{d}{b}, 0)$ (if $b \neq 0$)
 - $(0, 0, \frac{d}{c})$ (if $c \neq 0$)

Summary of plane equations – Cartesian form

Given a plane in **Cartesian form**:

$$ax + by + cz = d, \quad a, b, c, d \in \mathbb{R}$$

- A **normal vector to the plane** is $\begin{pmatrix} a \\ b \\ c \end{pmatrix}$.
- **Points on the plane** include:
 - $(\frac{d}{a}, 0, 0)$ (if $a \neq 0$)
 - $(0, \frac{d}{b}, 0)$ (if $b \neq 0$)
 - $(0, 0, \frac{d}{c})$ (if $c \neq 0$)
- **Vectors parallel to the plane** include $\begin{pmatrix} b \\ -a \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ c \\ -b \end{pmatrix}, \begin{pmatrix} -c \\ 0 \\ a \end{pmatrix}$,
and any linear combinations of these.

Summary of plane equations – Point-normal form

Given a plane in **point-normal form**:

$$\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0$$

Summary of plane equations – Point-normal form

Given a plane in **point-normal form**:

$$\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0$$

- A **normal vector to the plane** is \mathbf{n} .

Summary of plane equations – Point-normal form

Given a plane in **point-normal form**:

$$\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0$$

- A **normal vector to the plane** is \mathbf{n} .
- The position vector for **a point on the plane** is \mathbf{a} . To find others, it is easiest to first convert to Cartesian form.

Summary of plane equations – Point-normal form

Given a plane in **point-normal form**:

$$\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0$$


- A **normal vector to the plane** is \mathbf{n} .
- The position vector for **a point on the plane** is \mathbf{a} . To find others, it is easiest to first convert to Cartesian form.
- **Vectors parallel to the plane** are any vectors \mathbf{v} such that $\mathbf{n} \cdot \mathbf{v} = 0$. It is easiest to find these by first converting to Cartesian form.

Shortest distance to a plane

How might we find the shortest distance from a point B to the plane given by $\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0$, or find the point P on the plane that is closest to B ?


Shortest distance to a plane

How might we find the shortest distance from a point B to the plane given by $\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0$, or find the point P on the plane that is closest to B ?




Shortest distance to a plane

How might we find the shortest distance from a point B to the plane given by $\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0$, or find the point P on the plane that is closest to B ?




Shortest distance to a plane

How might we find the shortest distance from a point B to the plane given by $\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0$, or find the point P on the plane that is closest to B ?




Shortest distance to a plane

How might we find the shortest distance from a point B to the plane given by $\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0$, or find the point P on the plane that is closest to B ?




Shortest distance to a plane

How might we find the shortest distance from a point B to the plane given by $\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0$, or find the point P on the plane that is closest to B ?



Shortest distance to a plane


How might we find the shortest distance from a point B to the plane given by $\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0$, or find the point P on the plane that is closest to B ?



Position vector of P : $\mathbf{b} - \text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a}) = \mathbf{b} - \frac{\mathbf{n} \cdot (\mathbf{b} - \mathbf{a})}{|\mathbf{n}|^2} \mathbf{n}$

Shortest distance to a plane

How might we find the shortest distance from a point B to the plane given by $\mathbf{n} \cdot (\mathbf{x} - \mathbf{a}) = 0$, or find the point P on the plane that is closest to B ?



Position vector of P : $\mathbf{b} - \text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a}) = \mathbf{b} - \frac{\mathbf{n} \cdot (\mathbf{b} - \mathbf{a})}{|\mathbf{n}|^2} \mathbf{n}$

Shortest distance: $|\text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a})| = \frac{|\mathbf{n} \cdot (\mathbf{b} - \mathbf{a})|}{|\mathbf{n}|}$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

First find a vector normal to the plane.

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

First find a vector normal to the plane. As before, we can take the cross product of two vectors parallel to the plane. For example,

$$\overrightarrow{PQ} \times \overrightarrow{PR}$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

First find a vector normal to the plane. As before, we can take the cross product of two vectors parallel to the plane. For example,

$$\overrightarrow{PQ} \times \overrightarrow{PR} = \begin{pmatrix} -4 \\ 0 \\ -2 \end{pmatrix} \times \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix}$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

First find a vector normal to the plane. As before, we can take the cross product of two vectors parallel to the plane. For example,

$$\overrightarrow{PQ} \times \overrightarrow{PR} = \begin{pmatrix} -4 \\ 0 \\ -2 \end{pmatrix} \times \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} = 2 \begin{pmatrix} -2 \\ 0 \\ -1 \end{pmatrix} \times 3 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} =$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

First find a vector normal to the plane. As before, we can take the cross product of two vectors parallel to the plane. For example,

$$\overrightarrow{PQ} \times \overrightarrow{PR} = \begin{pmatrix} -4 \\ 0 \\ -2 \end{pmatrix} \times \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} = 2 \begin{pmatrix} -2 \\ 0 \\ -1 \end{pmatrix} \times 3 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = 6 \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix}.$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

First find a vector normal to the plane. As before, we can take the cross product of two vectors parallel to the plane. For example,

$$\overrightarrow{PQ} \times \overrightarrow{PR} = \begin{pmatrix} -4 \\ 0 \\ -2 \end{pmatrix} \times \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} = 2 \begin{pmatrix} -2 \\ 0 \\ -1 \end{pmatrix} \times 3 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = 6 \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix}.$$

Since only the direction of the normal vector is important, we can disregard the length of $\overrightarrow{PQ} \times \overrightarrow{PR}$ and just choose $\mathbf{n} = \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix}$.

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

By choosing \mathbf{a} to be the position vector for the point P , say, we can find the required projected vector as follows:

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

By choosing \mathbf{a} to be the position vector for the point P , say, we can find the required projected vector as follows:

$$\text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a})$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

By choosing \mathbf{a} to be the position vector for the point P , say, we can find the required projected vector as follows:

$$\text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a}) = \frac{(\mathbf{b} - \mathbf{a}) \cdot \mathbf{n}}{|\mathbf{n}|^2} \mathbf{n}$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

By choosing \mathbf{a} to be the position vector for the point P , say, we can find the required projected vector as follows:

$$\begin{aligned}\text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a}) &= \frac{(\mathbf{b} - \mathbf{a}) \cdot \mathbf{n}}{|\mathbf{n}|^2} \mathbf{n} \\ &= \frac{(4 - 1) \times 1 + (-2 - 2) \times 1 + (3 - 3) \times (-2)}{1^2 + 1^2 + (-2)^2} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix}\end{aligned}$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

By choosing \mathbf{a} to be the position vector for the point P , say, we can find the required projected vector as follows:

$$\begin{aligned}\text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a}) &= \frac{(\mathbf{b} - \mathbf{a}) \cdot \mathbf{n}}{|\mathbf{n}|^2} \mathbf{n} \\&= \frac{(4 - 1) \times 1 + (-2 - 2) \times 1 + (3 - 3) \times (-2)}{1^2 + 1^2 + (-2)^2} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix} \\&= -\frac{1}{6} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix}\end{aligned}$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

So the position vector of the closest point X is given by:

$$\mathbf{b} - \text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a})$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

So the position vector of the closest point X is given by:

$$\mathbf{b} - \text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a}) = \begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix} - \frac{-1}{6} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix}$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$.

Also find the point X on the plane that is closest to B .

So the position vector of the closest point X is given by:

$$\mathbf{b} - \text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a}) = \begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix} - \frac{-1}{6} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix} = \begin{pmatrix} 25/6 \\ -11/6 \\ 8/3 \end{pmatrix}$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

So the position vector of the closest point X is given by:

$$\mathbf{b} - \text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a}) = \begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix} - \frac{-1}{6} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix} = \begin{pmatrix} 25/6 \\ -11/6 \\ 8/3 \end{pmatrix}$$

That is, X is the point $(\frac{25}{6}, -\frac{11}{6}, \frac{8}{3})$.

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

So the position vector of the closest point X is given by:

$$\mathbf{b} - \text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a}) = \begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix} - \frac{-1}{6} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix} = \begin{pmatrix} 25/6 \\ -11/6 \\ 8/3 \end{pmatrix}$$

That is, X is the point $(\frac{25}{6}, -\frac{11}{6}, \frac{8}{3})$.

The shortest distance is given by:

$$|\text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a})|$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

So the position vector of the closest point X is given by:

$$\mathbf{b} - \text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a}) = \begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix} - \frac{-1}{6} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix} = \begin{pmatrix} 25/6 \\ -11/6 \\ 8/3 \end{pmatrix}$$

That is, X is the point $(\frac{25}{6}, -\frac{11}{6}, \frac{8}{3})$.

The shortest distance is given by:

$$|\text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a})| = \left| -\frac{1}{6} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix} \right|$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

So the position vector of the closest point X is given by:

$$\mathbf{b} - \text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a}) = \begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix} - \frac{-1}{6} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix} = \begin{pmatrix} 25/6 \\ -11/6 \\ 8/3 \end{pmatrix}$$

That is, X is the point $(\frac{25}{6}, -\frac{11}{6}, \frac{8}{3})$.

The shortest distance is given by:

$$|\text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a})| = \left| -\frac{1}{6} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix} \right| = \frac{1}{6} \sqrt{1^2 + 1^2 + (-2)^2}$$

Shortest distance to a plane – Example

Example

Find the shortest distance between the point $B(4, -2, 3)$ and the plane passing through the points $P(1, 2, 3)$, $Q(-3, 2, 1)$, and $R(4, 5, 6)$. Also find the point X on the plane that is closest to B .

So the position vector of the closest point X is given by:

$$\mathbf{b} - \text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a}) = \begin{pmatrix} 4 \\ -2 \\ 3 \end{pmatrix} - \frac{-1}{6} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix} = \begin{pmatrix} 25/6 \\ -11/6 \\ 8/3 \end{pmatrix}$$

That is, X is the point $(\frac{25}{6}, -\frac{11}{6}, \frac{8}{3})$.

The shortest distance is given by:

$$|\text{proj}_{\mathbf{n}}(\mathbf{b} - \mathbf{a})| = \left| -\frac{1}{6} \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix} \right| = \frac{1}{6} \sqrt{1^2 + 1^2 + (-2)^2} = \frac{1}{\sqrt{6}}$$