

# Lec20: Properties of Determinants

Laure Helme-Guizon (Dr H)

[Laure@unsw.edu.au](mailto:Laure@unsw.edu.au)

Jonathan Kress

[j.kress@unsw.edu.au](mailto:j.kress@unsw.edu.au)

Red-Centre, Rooms 3090 and 3073

2020 Term 1

# Determinants and inverses

Exercise 1. Use  $\det(AB) = \det(A)\det(B)$  to prove that

$$\det(A^{-1}) = \frac{1}{\det(A)}.$$

# Properties of determinants


## Properties of determinants



- (a)  $\det(A^T) = \det(A)$
- (b)  $\det(AB) = \det(A)\det(B)$
- (c)  $R_i \leftarrow R_i + \alpha R_j$  for  $i \neq j$  does not change the determinant.
- (d)  $R_i \leftrightarrow R_j$  for  $i \neq j$  changes the sign of the determinant.
- (e)  $R_i \leftarrow \alpha R_i$  scales the determinant by  $\alpha$ .
- (f) If  $A$  has a zero row or column then  $\det(A) = 0$ .

## Some important consequences:



- (g)  $\det(\alpha A) = \alpha^n \det(A)$  for an  $n \times n$  matrix.  Note that it is  $\alpha^n$  not  $\alpha$
- (h)  $\det(A^{-1}) = 1/\det(A)$ .
- (i) Swapping two columns changes the sign of the determinant.
- (j) Row operations can simplify the calculation of determinants.
- (k)  $A$  is invertible *if and only*  $\det(A) \neq 0$ .
- (l) If one row of  $A$  is a multiple of another row then  $\det(A) = 0$ .
- (m) If one column of  $A$  is a multiple of another column then  $\det(A) = 0$ .

# Determinants

## Exercise 2.

Suppose that  $A$  is a  $3 \times 3$  matrix and  $\det(A) = 5$ . Find  $\det(B)$  if  $B$  is given by

1.  $A^T$ .
2.  $A$  with the first two rows swapped.
3.  $A$  with the first two rows swapped and the last two columns swapped.
4.  $A$  with the second row multiplied by 7.
5.  $2A$ .
6.  $A$  with the second row replaced by 3 times the first row.
7.  $A^{-1}$ .

## Determinants: Checking the rules on an example

Exercise 3. Let  $A = \begin{pmatrix} 1 & 3 \\ 2 & 8 \end{pmatrix}$ .

1. Find  $\det(A)$ .
2. Check that  $R_2 \leftarrow R_2 - 5R_1$  doesn't change the determinant.
3. Check that  $C_2 \leftarrow C_2 + 4C_1$  doesn't change the determinant.
4. Show that  $R_2 \leftarrow 3R_2 + 4R_1$  does change the determinant.

## Determinants: Checking the rules on an example

Exercise 3, continued.  $A = \begin{pmatrix} 1 & 3 \\ 2 & 8 \end{pmatrix}$ .

4. Show that  $R_2 \leftarrow 3R_2 + 4R_1$  does change the determinant.
5. Find the determinant after swapping the rows of  $A$ .
6. Reduce  $A$  to echelon form and find the determinant.

# Determinants and inverses



## Determinants and inverses.

For a square matrix  $A$ , the following are equivalent:

- $\det(A) \neq 0$
- $A$  is invertible
- $A\vec{x} = \vec{0}$  has the unique solution  $\vec{x} = \vec{0}$
- $A\vec{x} = \vec{b}$  has a unique solution for all  $\vec{b} \in \mathbb{R}^n$ .



*The determinant is a detector of invertible matrices*

# Determinants

Exercise 4. Last lecture, we found that  $A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$  does not have an inverse.

Verify this fact by checking that  $\det(A) = 0$ .



# Determinants

Exercise 5. For what values of  $\alpha$  is  $A = \begin{pmatrix} 1 & 2 & -1 \\ 3 & 1 & 0 \\ \alpha & 0 & 1 \end{pmatrix}$  invertible?

# Determinants with Maple

```
> # exercise 5
> with(LinearAlgebra):
> # Enter the matrices columnwise

A := < <1,3,alpha>|<2,1,0>|<-1, 0, 1> >;

A :=  $\begin{bmatrix} 1 & 2 & -1 \\ 3 & 1 & 0 \\ \alpha & 0 & 1 \end{bmatrix}$ 

> # Calculate the determinant of A

Det_of_A := Determinant(A);

Det_of_A :=  $-5 + \alpha$ 
```

# Determinants

Exercise 6. Find  $\begin{vmatrix} 1 & -1 & 0 & 3 \\ 2 & -2 & 6 & -1 \\ 4 & -2 & 1 & 7 \\ 3 & 5 & -7 & 2 \end{vmatrix}$

# Determinants

Exercise 7. Find

$$\begin{vmatrix} 1 & 5 & 4 & 3 \\ 3 & -2 & 12 & -2 \\ 1 & -3 & 4 & 3 \\ 2 & -11 & 8 & 1 \end{vmatrix}$$

# Determinants

Exercise 8. Let  $t \in (-\pi, \pi]$ . For what values of  $t$  is

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 + \cos t & 1 + \sin 2t \\ 1 & 1 + \sin t & 1 + \sin 2t \end{pmatrix}$$

invertible?

# Determinants

Exercise 8, continued. Let  $t \in (-\pi, \pi]$ . For what values of  $t$  is

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 + \cos t & 1 + \sin 2t \\ 1 & 1 + \sin t & 1 + \sin 2t \end{pmatrix}$$

invertible?

# Determinant with Maple

```
> restart;
with(LinearAlgebra):
> # Enter the matrices columnwise
A := < <1,1,1> | <1,1+cos(t),1+sin(t)> | <1,1+sin(2*t),1+sin(2*t)> >;
      A := 
$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 + \cos(t) & 1 + \sin(2t) \\ 1 & 1 + \sin(t) & 1 + \sin(2t) \end{bmatrix}$$

> # Calculate the determinant of A
Det_of_A := Determinant(A);
      Det_of_A :=  $\cos(t) \sin(2t) - \sin(2t) \sin(t)$ 
> # Factorise the determinant of A
factor(Det_of_A);
       $\sin(2t) (\cos(t) - \sin(t))$ 
> # A is invertible unless det(A) = 0
solve(Det_of_A = 0, t);
       $\frac{\pi}{4}, 0$ 
```

# THE END

- Check online that your marks are correctly recorded.
- Read the information about the online final exam on Moodle!
- Make sure you book at time for your exam.
- Past papers with solutions are in the course pack and on Moodle. The 2019 past paper give you an idea of what to expect.
- A link to last semester's exam revision live stream will be posted online.
- Make sure you have a UNSW approved calculator.
- Please complete the myExperience surveys.
- A pre-exam consultation roster will be posted on Moodle.

Good luck!