

School of Mathematics and Statistics

Math1131 Mathematics 1A

CALCULUS LECTURE 4 LIMITS TO INFINITY (PART 2)

Milan Pahor



Copyright 2020 © University of New South Wales and licensors/contributors as identified. All rights reserved. Except where otherwise noted, this work is licensed under CC BY-NC-ND: creativecommons.org/licenses/by-nc-nd/4.0/ The views and opinions expressed in this work are those of the author and do not necessarily reflect the official position of the University of New South Wales. UNSW and UNSW crest are registered trademarks of UNSW Sydney.

MATH1131 CALCULUS

Limit of Functions at Infinity Part 2

 $\lim_{x \to \infty} f(x) = L$ if for every (small) positive real number ϵ there is a (large) real number M with the property that if x > M then $|f(x) - L| < \epsilon$.

Limits of the form $\lim_{x \to a} f(x)$ are best attacked via factorisation.

 $\lim_{x \to a^{-}} f(x)$ is the limit of f(x) as x approaches a from the left

 $\lim_{x \to a^+} f(x)$ is the limit of f(x) as x approaches a from the right

 $\lim_{x \to a} f(x)$ exists if and only if $\lim_{x \to a^{-}} f(x)$ and $\lim_{x \to a^{+}} f(x)$ both exist and are equal.

Just like differentiation from first principles we also have a formal definition of $\lim_{x\to\infty} f(x)$. It is crucial that we understand these fundamental definitions as they provide a solid foundation for the analysis. We can't just keep waving our hands around! If we want you to evaluate a limit formally we will always give you a clear warning.

Definition: $\lim_{x \to \infty} f(x) = L$ if for every (small) positive real number ϵ there is a (large) real number M with the property that if x > M then $|f(x) - L| < \epsilon$.

Discussion

Note that $|f(x) - L| < \epsilon$ simply means that the difference between f(x) and L is really really small. The limit definition is then simply saying that any degree of closeness between f(x) and L can be generated by choosing x to be sufficiently large.

For $\lim_{x \to \infty} f(x)$ to be equal to L the function must eventually (x > M) get into (and stay in!) an ϵ band of L.

Example 1: Consider $\lim_{x \to \infty} \frac{10x - 4}{5x + 7}$.

- a) Show that the value of the limit is L=2.
- b) Find M so that f(x) is within $\frac{1}{1000}$ of its limit whenever x > M.
- c) Does b) prove that $\lim_{x \to \infty} \frac{10x 4}{5x + 7} = 2$?
- d) Prove from the limit definition that $\lim_{x \to \infty} \frac{10x 4}{5x + 7} = 2$.

That is given $\epsilon > 0$ find M with the property that if x > M then $|f(x) - L| < \epsilon$.

- a)
- b)

So $\left|\frac{10x-4}{5x+7}-2\right| \leq \frac{1}{1000}$ provided we choose x>3598.6. Note that we are not really trying to solve $\left|\frac{10x-4}{5x+7}-2\right| \leq \frac{1}{1000}$. We are just trying to find an M with the property that the inequality is true for all x>M. This value of M is not unique and any M>3598.6 would also work


c) NO!!

d)

$$\bigstar \quad M = \frac{18}{5\epsilon} - \frac{7}{5}. \quad \bigstar$$


Note that since the above M depends on ϵ we sometimes write $M(\epsilon) = \frac{18}{5\epsilon} - \frac{7}{5}$.

Let's take a look at the graph for the situation of part b) where $\epsilon = \frac{1}{1000} = 0.001$. First the function and its limit of L = 2 as a horizontal asymptote:



Next a band (red to green) of length $\pm \frac{1}{1000}$ either side of the limit L=2.

Observe below that after M=3598.6 the function is clearly locked within this limit band.




Example 2: Find $\lim_{x \to \infty} \frac{\sin(x)}{x^2 + e^x}$ and prove from the limit definition that your answer is correct. Find a value of M for $\epsilon = \frac{1}{100}$ and sketch the behaviour of the function and its limit

Clearly
$$\lim_{x \to \infty} \frac{\sin(x)}{x^2 + e^x} =$$


So
$$L = 0$$
. We now examine $|f(x) - L| = \left| \frac{\sin(x)}{x^2 + e^x} - 0 \right| =$

So we have $|f(x) - L| < \frac{1}{x^2}$ we therefore only need $\frac{1}{x^2} < \epsilon$. And hence:

Let's take a look at the graph:



Observe below that after M=10 the function is clearly within its $\frac{1}{100}$ limit band although a smaller value of M=5 would also do!!



Note that above we have most definitely **NOT** proven that $\lim_{x \to \infty} \frac{\sin(x)}{x^2 + e^x} = 0!$ The full proof would demand that we use a general small ϵ rather than the specific $\frac{1}{100}$.

Limit of Functions at a Point

There are certain situations where a function fails to be defined **at** a point but it is perfectly happy **near** the point. We then use $\lim_{x \to a} f(x)$ to get a feeling for the behaviour of the function.

Consider the following table of values for $f(x) = \frac{x^2 - 25}{x - 5}$ near x = 5.

x	4.9	4.99	5	5.01	5.1
y	9.9	9.99	?	10.01	10.1

It is clear that the function is trying to get to 10 at x=5 even though it is undefined there. We write

$$\lim_{x \to 5} \frac{x^2 - 25}{x - 5} = 10$$

and say

"The limit as x approaches 5 of $\frac{x^2 - 25}{x - 5}$ is 10".

Example 3: Find $\lim_{x \to 5} \frac{x^2 - 25}{x - 5}$ and hence sketch the function $y = \frac{x^2 - 25}{x - 5}$.



So for limits as x approaches a finite value our main technique is to factorise top and bottom. Always check first that you are facing the indeterminate form " $\frac{0}{0}$ "

Example 4: Evaluate each of the following limits

a)
$$\lim_{x \to 2} \frac{x^2 - 5x + 6}{x^2 - 4}$$

b)
$$\lim_{x \to 1} \frac{x^2 + 1}{x^2 + 4}$$

$$\star$$
 a) $-\frac{1}{4}$ b) $\frac{2}{5}$ \star

In order to define these limits a little more formally we require the concept of a one-sided limit:


 $\lim_{x \to a^{-}} f(x)$ is the limit of f(x) as x approaches a from the left

 $\lim_{x \to a^+} f(x)$ is the limit of f(x) as x approaches a from the right

We can then say that the full limit $\lim_{x \to a} f(x)$ formally exists if and only if $\lim_{x \to a^{-}} f(x)$ and $\lim_{x \to a^{+}} f(x)$ both exist and are equal.

Example 5: Consider the graph of y = f(x) presented in red below:

For each of the following either evaluate the given quantity or explain why it does not exist:



- a) $\lim_{x \to 4^-} f(x)$ b) $\lim_{x \to 4^+} f(x)$
- c) $\lim_{x \to 4} f(x)$ d) $\lim_{x \to 5} f(x)$
- e) f(5) f) $\lim_{x \to 6} f(x)$
- g) $\lim_{x \to 3} f(x)$