



UNSW
SYDNEY

MATH1131 Mathematics 1A – Algebra

Lecture 13: Complex Roots and Powers

Lecturer: Sean Gardiner – sean.gardiner@unsw.edu.au

Based on slides by Jonathan Kress

Powers

Recall that it is easy to find large powers of a complex number if it is first written in exponential polar form.

Powers

Recall that it is easy to find large powers of a complex number if it is first written in exponential polar form.

Example

Calculate $(1 + \sqrt{3}i)^{10}$.

Powers

Recall that it is easy to find large powers of a complex number if it is first written in exponential polar form.

Example

Calculate $(1 + \sqrt{3}i)^{10}$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}}$.

Powers

Recall that it is easy to find large powers of a complex number if it is first written in exponential polar form.

Example

Calculate $(1 + \sqrt{3}i)^{10}$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}}$.

So $(1 + \sqrt{3}i)^{10} = (2e^{i\frac{\pi}{3}})^{10}$

Powers

Recall that it is easy to find large powers of a complex number if it is first written in exponential polar form.

Example

Calculate $(1 + \sqrt{3}i)^{10}$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}}$.

So $(1 + \sqrt{3}i)^{10} = (2e^{i\frac{\pi}{3}})^{10} = 2^{10} e^{i\frac{\pi}{3} \times 10}$

Powers

Recall that it is easy to find large powers of a complex number if it is first written in exponential polar form.

Example

Calculate $(1 + \sqrt{3}i)^{10}$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}}$.

So $(1 + \sqrt{3}i)^{10} = (2e^{i\frac{\pi}{3}})^{10} = 2^{10} e^{i\frac{\pi}{3} \times 10} = 1024e^{i\frac{10\pi}{3}}$

Powers

Recall that it is easy to find large powers of a complex number if it is first written in exponential polar form.

Example

Calculate $(1 + \sqrt{3}i)^{10}$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}}$.

So $(1 + \sqrt{3}i)^{10} = (2e^{i\frac{\pi}{3}})^{10} = 2^{10} e^{i\frac{\pi}{3} \times 10} = 1024e^{i\frac{10\pi}{3}} = 1024e^{-i\frac{2\pi}{3}}$.

Powers

Recall that it is easy to find large powers of a complex number if it is first written in exponential polar form.

Example

Calculate $(1 + \sqrt{3}i)^{10}$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}}$.

So $(1 + \sqrt{3}i)^{10} = (2e^{i\frac{\pi}{3}})^{10} = 2^{10} e^{i\frac{\pi}{3} \times 10} = 1024e^{i\frac{10\pi}{3}} = 1024e^{-i\frac{2\pi}{3}}$.

Since $e^{-i\frac{2\pi}{3}} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$, we have:

Powers

Recall that it is easy to find large powers of a complex number if it is first written in exponential polar form.

Example

Calculate $(1 + \sqrt{3}i)^{10}$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}}$.

So $(1 + \sqrt{3}i)^{10} = (2e^{i\frac{\pi}{3}})^{10} = 2^{10} e^{i\frac{\pi}{3} \times 10} = 1024e^{i\frac{10\pi}{3}} = 1024e^{-i\frac{2\pi}{3}}$.

Since $e^{-i\frac{2\pi}{3}} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$, we have:

$$(1 + \sqrt{3}i)^{10} = 1024e^{-i\frac{2\pi}{3}}$$

Powers

Recall that it is easy to find large powers of a complex number if it is first written in exponential polar form.

Example

Calculate $(1 + \sqrt{3}i)^{10}$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}}$.

So $(1 + \sqrt{3}i)^{10} = (2e^{i\frac{\pi}{3}})^{10} = 2^{10} e^{i\frac{\pi}{3} \times 10} = 1024e^{i\frac{10\pi}{3}} = 1024e^{-i\frac{2\pi}{3}}$.

Since $e^{-i\frac{2\pi}{3}} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$, we have:

$$(1 + \sqrt{3}i)^{10} = 1024e^{-i\frac{2\pi}{3}} = 1024 \left(-\frac{1}{2} - \frac{\sqrt{3}}{2}i \right)$$

Powers

Recall that it is easy to find large powers of a complex number if it is first written in exponential polar form.

Example

Calculate $(1 + \sqrt{3}i)^{10}$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}}$.

So $(1 + \sqrt{3}i)^{10} = (2e^{i\frac{\pi}{3}})^{10} = 2^{10} e^{i\frac{\pi}{3} \times 10} = 1024e^{i\frac{10\pi}{3}} = 1024e^{-i\frac{2\pi}{3}}$.

Since $e^{-i\frac{2\pi}{3}} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$, we have:

$$(1 + \sqrt{3}i)^{10} = 1024e^{-i\frac{2\pi}{3}} = 1024 \left(-\frac{1}{2} - \frac{\sqrt{3}}{2}i \right) = -512 - 512\sqrt{3}i.$$

Roots

Definition

A complex number α is an n^{th} root of $z \in \mathbb{C}$ if

$$\alpha^n = z.$$

Roots

Definition

A complex number α is an n^{th} root of $z \in \mathbb{C}$ if

$$\alpha^n = z.$$

To find n^{th} roots we again prefer to use polar form.

Definition

A complex number α is an n^{th} root of $z \in \mathbb{C}$ if

$$\alpha^n = z.$$

To find n^{th} roots we again prefer to use polar form.

Suppose $z = re^{i\theta}$ and that $\alpha = se^{i\phi}$ is an n^{th} root of z .

Definition

A complex number α is an n^{th} root of $z \in \mathbb{C}$ if

$$\alpha^n = z.$$

To find n^{th} roots we again prefer to use polar form.

Suppose $z = re^{i\theta}$ and that $\alpha = se^{i\phi}$ is an n^{th} root of z .

$$z = re^{i\theta}$$

Definition

A complex number α is an n^{th} root of $z \in \mathbb{C}$ if

$$\alpha^n = z.$$

To find n^{th} roots we again prefer to use polar form.

Suppose $z = re^{i\theta}$ and that $\alpha = se^{i\phi}$ is an n^{th} root of z .

$$z = re^{i\theta} = \alpha^n$$

Definition

A complex number α is an n^{th} root of $z \in \mathbb{C}$ if

$$\alpha^n = z.$$

To find n^{th} roots we again prefer to use polar form.

Suppose $z = re^{i\theta}$ and that $\alpha = se^{i\phi}$ is an n^{th} root of z .

$$\begin{aligned} z = re^{i\theta} &= \alpha^n \\ &= (se^{i\phi})^n \end{aligned}$$

Definition

A complex number α is an n^{th} root of $z \in \mathbb{C}$ if

$$\alpha^n = z.$$

To find n^{th} roots we again prefer to use polar form.

Suppose $z = re^{i\theta}$ and that $\alpha = se^{i\phi}$ is an n^{th} root of z .

$$\begin{aligned} z = re^{i\theta} &= \alpha^n \\ &= (se^{i\phi})^n \\ &= s^n e^{in\phi} \end{aligned}$$

Definition

A complex number α is an n^{th} root of $z \in \mathbb{C}$ if

$$\alpha^n = z.$$

To find n^{th} roots we again prefer to use polar form.

Suppose $z = re^{i\theta}$ and that $\alpha = se^{i\phi}$ is an n^{th} root of z .

$$\begin{aligned} z = re^{i\theta} &= \alpha^n \\ &= (se^{i\phi})^n \\ &= s^n e^{in\phi} \end{aligned}$$

So $s^n = r$ and $n\phi = \theta + 2k\pi$ for some $k \in \mathbb{Z}$.

Roots

Example

Find the 5th roots of 1. That is, find all α such that

$$\alpha^5 = 1.$$

Roots

Example

Find the 5th roots of 1. That is, find all α such that

$$\alpha^5 = 1.$$

In polar form, $1 = e^{i \times 0}$

Roots

Example

Find the 5th roots of 1. That is, find all α such that

$$\alpha^5 = 1.$$

In polar form, $1 = e^{i \times 0} = e^{i \times (0 + 2k\pi)}$ for any $k \in \mathbb{Z}$.

Example

Find the 5th roots of 1. That is, find all α such that

$$\alpha^5 = 1.$$

In polar form, $1 = e^{i \times 0} = e^{i \times (0 + 2k\pi)}$ for any $k \in \mathbb{Z}$.

So if $\alpha^5 = e^{i \times (0 + 2k\pi)}$, we must have $\alpha = e^{i \times \frac{1}{5}(0 + 2k\pi)}$ for some $k \in \mathbb{Z}$.

Example

Find the 5th roots of 1. That is, find all α such that

$$\alpha^5 = 1.$$

In polar form, $1 = e^{i \times 0} = e^{i \times (0 + 2k\pi)}$ for any $k \in \mathbb{Z}$.

So if $\alpha^5 = e^{i \times (0 + 2k\pi)}$, we must have $\alpha = e^{i \times \frac{1}{5}(0 + 2k\pi)}$ for some $k \in \mathbb{Z}$.

Picking $k = 0, 1, 2, 3, \dots$ gives possible arguments:

Example

Find the 5th roots of 1. That is, find all α such that

$$\alpha^5 = 1.$$

In polar form, $1 = e^{i \times 0} = e^{i \times (0 + 2k\pi)}$ for any $k \in \mathbb{Z}$.

So if $\alpha^5 = e^{i \times (0 + 2k\pi)}$, we must have $\alpha = e^{i \times \frac{1}{5}(0 + 2k\pi)}$ for some $k \in \mathbb{Z}$.

Picking $k = 0, 1, 2, 3, \dots$ gives possible arguments:

$$0, \frac{2\pi}{5}, \frac{4\pi}{5}, \frac{6\pi}{5}, \frac{8\pi}{5}, \frac{10\pi}{5}, \dots$$

Example

Find the 5th roots of 1. That is, find all α such that

$$\alpha^5 = 1.$$

In polar form, $1 = e^{i \times 0} = e^{i \times (0 + 2k\pi)}$ for any $k \in \mathbb{Z}$.

So if $\alpha^5 = e^{i \times (0 + 2k\pi)}$, we must have $\alpha = e^{i \times \frac{1}{5}(0 + 2k\pi)}$ for some $k \in \mathbb{Z}$.

Picking $k = 0, 1, 2, 3, \dots$ gives possible arguments:

$$0, \frac{2\pi}{5}, \frac{4\pi}{5}, \frac{6\pi}{5}, \frac{8\pi}{5}, \frac{10\pi}{5}, \dots$$

which are principal arguments:

$$0, \frac{2\pi}{5}, \frac{4\pi}{5}, -\frac{4\pi}{5}, -\frac{2\pi}{5}, 0, \dots$$

Example

Find the 5th roots of 1. That is, find all α such that

$$\alpha^5 = 1.$$

In polar form, $1 = e^{i \times 0} = e^{i \times (0 + 2k\pi)}$ for any $k \in \mathbb{Z}$.

So if $\alpha^5 = e^{i \times (0 + 2k\pi)}$, we must have $\alpha = e^{i \times \frac{1}{5}(0 + 2k\pi)}$ for some $k \in \mathbb{Z}$.

Picking $k = 0, 1, 2, 3, \dots$ gives possible arguments:

$$0, \frac{2\pi}{5}, \frac{4\pi}{5}, \frac{6\pi}{5}, \frac{8\pi}{5}, \frac{10\pi}{5}, \dots$$

which are principal arguments:

$$0, \frac{2\pi}{5}, \frac{4\pi}{5}, -\frac{4\pi}{5}, -\frac{2\pi}{5}, 0, \dots$$

So the 5th roots of 1 are:

$$1, e^{i\frac{2\pi}{5}}, e^{i\frac{4\pi}{5}}, e^{-i\frac{4\pi}{5}}, \text{ and } e^{-i\frac{2\pi}{5}}.$$

Roots

Example

Find the 5th roots of $z = 1 + \sqrt{3}i$.

Roots

Example

Find the 5th roots of $z = 1 + \sqrt{3}i$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}}$

Roots

Example

Find the 5th roots of $z = 1 + \sqrt{3}i$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}} = 2e^{i(\frac{\pi}{3} + 2k\pi)}$ for any $k \in \mathbb{Z}$.

Example

Find the 5th roots of $z = 1 + \sqrt{3}i$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}} = 2e^{i(\frac{\pi}{3}+2k\pi)}$ for any $k \in \mathbb{Z}$.

So if $\alpha^5 = 2e^{i(\frac{\pi}{3}+2k\pi)}$, we have $\alpha = 2^{\frac{1}{5}} e^{i \times \frac{1}{5}(\frac{\pi}{3}+2k\pi)}$ for some $k \in \mathbb{Z}$.

Example

Find the 5th roots of $z = 1 + \sqrt{3}i$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}} = 2e^{i(\frac{\pi}{3}+2k\pi)}$ for any $k \in \mathbb{Z}$.

So if $\alpha^5 = 2e^{i(\frac{\pi}{3}+2k\pi)}$, we have $\alpha = 2^{\frac{1}{5}} e^{i \times \frac{1}{5}(\frac{\pi}{3}+2k\pi)}$ for some $k \in \mathbb{Z}$.

Picking $k = 0, 1, 2, 3, \dots$ gives possible arguments:

Example

Find the 5th roots of $z = 1 + \sqrt{3}i$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}} = 2e^{i(\frac{\pi}{3}+2k\pi)}$ for any $k \in \mathbb{Z}$.

So if $\alpha^5 = 2e^{i(\frac{\pi}{3}+2k\pi)}$, we have $\alpha = 2^{\frac{1}{5}} e^{i \times \frac{1}{5}(\frac{\pi}{3}+2k\pi)}$ for some $k \in \mathbb{Z}$.

Picking $k = 0, 1, 2, 3, \dots$ gives possible arguments:

$$\frac{\pi}{15}, \frac{7\pi}{15}, \frac{13\pi}{15}, \frac{19\pi}{15}, \frac{25\pi}{15}, \frac{31\pi}{15}, \dots$$

Example

Find the 5th roots of $z = 1 + \sqrt{3}i$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}} = 2e^{i(\frac{\pi}{3}+2k\pi)}$ for any $k \in \mathbb{Z}$.

So if $\alpha^5 = 2e^{i(\frac{\pi}{3}+2k\pi)}$, we have $\alpha = 2^{\frac{1}{5}} e^{i \times \frac{1}{5}(\frac{\pi}{3}+2k\pi)}$ for some $k \in \mathbb{Z}$.

Picking $k = 0, 1, 2, 3, \dots$ gives possible arguments:

$$\frac{\pi}{15}, \frac{7\pi}{15}, \frac{13\pi}{15}, \frac{19\pi}{15}, \frac{25\pi}{15}, \frac{31\pi}{15}, \dots$$

which are principal arguments:

$$\frac{\pi}{15}, \frac{7\pi}{15}, \frac{13\pi}{15}, -\frac{11\pi}{15}, -\frac{5\pi}{15}, \frac{\pi}{15}, \dots$$

Example

Find the 5th roots of $z = 1 + \sqrt{3}i$.

In polar form, $1 + \sqrt{3}i = 2e^{i\frac{\pi}{3}} = 2e^{i(\frac{\pi}{3}+2k\pi)}$ for any $k \in \mathbb{Z}$.

So if $\alpha^5 = 2e^{i(\frac{\pi}{3}+2k\pi)}$, we have $\alpha = 2^{\frac{1}{5}} e^{i \times \frac{1}{5}(\frac{\pi}{3}+2k\pi)}$ for some $k \in \mathbb{Z}$.

Picking $k = 0, 1, 2, 3, \dots$ gives possible arguments:

$$\frac{\pi}{15}, \frac{7\pi}{15}, \frac{13\pi}{15}, \frac{19\pi}{15}, \frac{25\pi}{15}, \frac{31\pi}{15}, \dots$$

which are principal arguments:


$$\frac{\pi}{15}, \frac{7\pi}{15}, \frac{13\pi}{15}, -\frac{11\pi}{15}, -\frac{5\pi}{15}, \frac{\pi}{15}, \dots$$

So the 5th roots of $1 + \sqrt{3}i$ are:

$$2^{\frac{1}{5}} e^{i\frac{\pi}{15}}, 2^{\frac{1}{5}} e^{i\frac{7\pi}{15}}, 2^{\frac{1}{5}} e^{i\frac{13\pi}{15}}, 2^{\frac{1}{5}} e^{-i\frac{11\pi}{15}}, \text{ and } 2^{\frac{1}{5}} e^{-i\frac{\pi}{15}}.$$

Roots


Some facts about the n^{th} roots of a complex number z :



Roots

Some facts about the n^{th} roots of a complex number z :


- There are always exactly n different roots.



Roots

Some facts about the n^{th} roots of a complex number z :


- There are always exactly n different roots.
- All the roots have the same modulus $|z|^{\frac{1}{n}}$, and so lie on a circle with centre 0 and radius $|z|^{\frac{1}{n}}$.



Roots

Some facts about the n^{th} roots of a complex number z :


- There are always exactly n different roots.
- All the roots have the same modulus $|z|^{\frac{1}{n}}$, and so lie on a circle with centre 0 and radius $|z|^{\frac{1}{n}}$.
- The roots are evenly spaced around that circle, since adjacent roots are $\frac{2\pi}{n}$ radians apart.



Roots

Some facts about the n^{th} roots of a complex number z :


- There are always exactly n different roots.
- All the roots have the same modulus $|z|^{\frac{1}{n}}$, and so lie on a circle with centre 0 and radius $|z|^{\frac{1}{n}}$.
- The roots are evenly spaced around that circle, since adjacent roots are $\frac{2\pi}{n}$ radians apart.



Roots

Some facts about the n^{th} roots of a complex number z :


- There are always exactly n different roots.
- All the roots have the same modulus $|z|^{\frac{1}{n}}$, and so lie on a circle with centre 0 and radius $|z|^{\frac{1}{n}}$.
- The roots are evenly spaced around that circle, since adjacent roots are $\frac{2\pi}{n}$ radians apart.



Roots

Some facts about the n^{th} roots of a complex number z :


- There are always exactly n different roots.
- All the roots have the same modulus $|z|^{\frac{1}{n}}$, and so lie on a circle with centre 0 and radius $|z|^{\frac{1}{n}}$.
- The roots are evenly spaced around that circle, since adjacent roots are $\frac{2\pi}{n}$ radians apart.



Roots

Some facts about the n^{th} roots of a complex number z :

- There are always exactly n different roots.
- All the roots have the same modulus $|z|^{\frac{1}{n}}$, and so lie on a circle with centre 0 and radius $|z|^{\frac{1}{n}}$.
- The roots are evenly spaced around that circle, since adjacent roots are $\frac{2\pi}{n}$ radians apart.



Roots

It can be useful to find roots by thinking diagrammatically.

Example


Find the 3rd roots of -1 .

Roots

It can be useful to find roots by thinking diagrammatically.

Example

Find the 3rd roots of -1 .




Roots

It can be useful to find roots by thinking diagrammatically.

Example

Find the 3rd roots of -1 .




Roots

It can be useful to find roots by thinking diagrammatically.

Example

Find the 3rd roots of -1 .




Roots

It can be useful to find roots by thinking diagrammatically.

Example

Find the 3rd roots of -1 .




Roots

It can be useful to find roots by thinking diagrammatically.

Example

Find the 3rd roots of -1 .




Roots

It can be useful to find roots by thinking diagrammatically.

Example

Find the 3rd roots of -1 .



So the three 3rd roots of -1 are: -1 , $e^{i\frac{\pi}{3}}$ and $e^{-i\frac{\pi}{3}}$.

Binomial theorem

When calculating small powers of a complex number, it can be useful to apply the **Binomial Theorem**:

Binomial theorem

When calculating small powers of a complex number, it can be useful to apply the **Binomial Theorem**:

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k \quad \text{where} \quad \binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Binomial theorem

When calculating small powers of a complex number, it can be useful to apply the **Binomial Theorem**:

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k \quad \text{where} \quad \binom{n}{k} = \frac{n!}{k!(n-k)!}$$

To find the coefficients $\binom{n}{k}$, use the formula above (recalling $n! = n \times (n-1) \times (n-2) \times \cdots \times 1$), or take the $(k+1)$ th entry in the $(n+1)$ th row of Pascal's triangle:

Binomial theorem

When calculating small powers of a complex number, it can be useful to apply the **Binomial Theorem**:

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k \quad \text{where} \quad \binom{n}{k} = \frac{n!}{k!(n-k)!}$$

To find the coefficients $\binom{n}{k}$, use the formula above (recalling $n! = n \times (n-1) \times (n-2) \times \cdots \times 1$), or take the $(k+1)$ th entry in the $(n+1)$ th row of Pascal's triangle:

$$\begin{array}{ccccccc} & & & & 1 & & & \\ & & & & & & 1 & \\ & & & 1 & & 2 & & 1 \\ & & 1 & & 3 & & 3 & & 1 \\ & 1 & & 4 & & 6 & & 4 & & 1 \\ & & & & & \vdots & & & & \end{array}$$

Binomial theorem

Example

Find $(-2 + i)^7$.

Binomial theorem

Example

Find $(-2 + i)^7$.

The corresponding row in Pascal's triangle is:

1 7 21 35 35 21 7 1

Binomial theorem

Example

Find $(-2 + i)^7$.

The corresponding row in Pascal's triangle is:

1 7 21 35 35 21 7 1

$$(-2 + i)^7$$

Binomial theorem

Example

Find $(-2 + i)^7$.

The corresponding row in Pascal's triangle is:

$$1 \quad 7 \quad 21 \quad 35 \quad 35 \quad 21 \quad 7 \quad 1$$

$$\begin{aligned} &(-2 + i)^7 \\ &= 1 \times (-2)^7 i^0 + 7 \times (-2)^6 i^1 + 21 \times (-2)^5 i^2 + 35 \times (-2)^4 i^3 + \\ &\quad 35 \times (-2)^3 i^4 + 21 \times (-2)^2 i^5 + 7 \times (-2)^1 i^6 + 1 \times (-2)^0 i^7 \end{aligned}$$

Binomial theorem

Example

Find $(-2 + i)^7$.

The corresponding row in Pascal's triangle is:

$$1 \quad 7 \quad 21 \quad 35 \quad 35 \quad 21 \quad 7 \quad 1$$

$$\begin{aligned} &(-2 + i)^7 \\ &= 1 \times (-2)^7 i^0 + 7 \times (-2)^6 i^1 + 21 \times (-2)^5 i^2 + 35 \times (-2)^4 i^3 + \\ &\quad 35 \times (-2)^3 i^4 + 21 \times (-2)^2 i^5 + 7 \times (-2)^1 i^6 + 1 \times (-2)^0 i^7 \\ &= -128 \times 1 + 448 \times i - 672 \times (-1) + 560 \times (-i) + \\ &\quad - 280 \times 1 + 84 \times i - 14 \times (-1) + 1 \times (-i) \end{aligned}$$

Binomial theorem

Example

Find $(-2 + i)^7$.

The corresponding row in Pascal's triangle is:

$$1 \quad 7 \quad 21 \quad 35 \quad 35 \quad 21 \quad 7 \quad 1$$

$$\begin{aligned} &(-2 + i)^7 \\ &= 1 \times (-2)^7 i^0 + 7 \times (-2)^6 i^1 + 21 \times (-2)^5 i^2 + 35 \times (-2)^4 i^3 + \\ &\quad 35 \times (-2)^3 i^4 + 21 \times (-2)^2 i^5 + 7 \times (-2)^1 i^6 + 1 \times (-2)^0 i^7 \\ &= -128 \times 1 + 448 \times i - 672 \times (-1) + 560 \times (-i) + \\ &\quad - 280 \times 1 + 84 \times i - 14 \times (-1) + 1 \times (-i) \\ &= 278 - 29i. \end{aligned}$$

$\cos(n\theta)$ and $\sin(n\theta)$

The Binomial Theorem is particularly useful when expressing $\cos(n\theta)$ and $\sin(n\theta)$ in terms of powers of $\cos \theta$ and $\sin \theta$, and vice versa.

$\cos(n\theta)$ and $\sin(n\theta)$

The Binomial Theorem is particularly useful when expressing $\cos(n\theta)$ and $\sin(n\theta)$ in terms of powers of $\cos \theta$ and $\sin \theta$, and vice versa.

When expressing powers of $\sin \theta$ and $\cos \theta$ in terms of sines and cosines of multiples of θ , it is useful to use the following properties:

$$\begin{aligned}\cos(n\theta) &= \frac{1}{2}(e^{in\theta} + e^{-in\theta}) \\ \sin(n\theta) &= \frac{1}{2i}(e^{in\theta} - e^{-in\theta})\end{aligned}$$

$\cos(n\theta)$ and $\sin(n\theta)$

The Binomial Theorem is particularly useful when expressing $\cos(n\theta)$ and $\sin(n\theta)$ in terms of powers of $\cos \theta$ and $\sin \theta$, and vice versa.

When expressing powers of $\sin \theta$ and $\cos \theta$ in terms of sines and cosines of multiples of θ , it is useful to use the following properties:

$$\begin{aligned}\cos(n\theta) &= \frac{1}{2}(e^{in\theta} + e^{-in\theta}) \\ \sin(n\theta) &= \frac{1}{2i}(e^{in\theta} - e^{-in\theta})\end{aligned}$$

To prove these formulae, consider $z = e^{in\theta} = \cos(n\theta) + i \sin(n\theta)$.

$\cos(n\theta)$ and $\sin(n\theta)$

The Binomial Theorem is particularly useful when expressing $\cos(n\theta)$ and $\sin(n\theta)$ in terms of powers of $\cos \theta$ and $\sin \theta$, and vice versa.

When expressing powers of $\sin \theta$ and $\cos \theta$ in terms of sines and cosines of multiples of θ , it is useful to use the following properties:

$$\begin{aligned}\cos(n\theta) &= \frac{1}{2}(e^{in\theta} + e^{-in\theta}) \\ \sin(n\theta) &= \frac{1}{2i}(e^{in\theta} - e^{-in\theta})\end{aligned}$$

To prove these formulae, consider $z = e^{in\theta} = \cos(n\theta) + i \sin(n\theta)$.

Then

$$\frac{1}{2}(e^{in\theta} + e^{-in\theta})$$

$\cos(n\theta)$ and $\sin(n\theta)$

The Binomial Theorem is particularly useful when expressing $\cos(n\theta)$ and $\sin(n\theta)$ in terms of powers of $\cos \theta$ and $\sin \theta$, and vice versa.

When expressing powers of $\sin \theta$ and $\cos \theta$ in terms of sines and cosines of multiples of θ , it is useful to use the following properties:

$$\begin{aligned}\cos(n\theta) &= \frac{1}{2}(e^{in\theta} + e^{-in\theta}) \\ \sin(n\theta) &= \frac{1}{2i}(e^{in\theta} - e^{-in\theta})\end{aligned}$$

To prove these formulae, consider $z = e^{in\theta} = \cos(n\theta) + i \sin(n\theta)$.

Then

$$\frac{1}{2}(e^{in\theta} + e^{-in\theta}) = \frac{1}{2}(z + \bar{z})$$

$\cos(n\theta)$ and $\sin(n\theta)$

The Binomial Theorem is particularly useful when expressing $\cos(n\theta)$ and $\sin(n\theta)$ in terms of powers of $\cos \theta$ and $\sin \theta$, and vice versa.

When expressing powers of $\sin \theta$ and $\cos \theta$ in terms of sines and cosines of multiples of θ , it is useful to use the following properties:

$$\begin{aligned}\cos(n\theta) &= \frac{1}{2}(e^{in\theta} + e^{-in\theta}) \\ \sin(n\theta) &= \frac{1}{2i}(e^{in\theta} - e^{-in\theta})\end{aligned}$$

To prove these formulae, consider $z = e^{in\theta} = \cos(n\theta) + i \sin(n\theta)$.

Then

$$\frac{1}{2}(e^{in\theta} + e^{-in\theta}) = \frac{1}{2}(z + \bar{z}) = \operatorname{Re}(z)$$

$\cos(n\theta)$ and $\sin(n\theta)$

The Binomial Theorem is particularly useful when expressing $\cos(n\theta)$ and $\sin(n\theta)$ in terms of powers of $\cos \theta$ and $\sin \theta$, and vice versa.

When expressing powers of $\sin \theta$ and $\cos \theta$ in terms of sines and cosines of multiples of θ , it is useful to use the following properties:

$$\begin{aligned}\cos(n\theta) &= \frac{1}{2}(e^{in\theta} + e^{-in\theta}) \\ \sin(n\theta) &= \frac{1}{2i}(e^{in\theta} - e^{-in\theta})\end{aligned}$$

To prove these formulae, consider $z = e^{in\theta} = \cos(n\theta) + i \sin(n\theta)$.

Then

$$\frac{1}{2}(e^{in\theta} + e^{-in\theta}) = \frac{1}{2}(z + \bar{z}) = \operatorname{Re}(z) = \cos(n\theta),$$

$\cos(n\theta)$ and $\sin(n\theta)$

The Binomial Theorem is particularly useful when expressing $\cos(n\theta)$ and $\sin(n\theta)$ in terms of powers of $\cos \theta$ and $\sin \theta$, and vice versa.

When expressing powers of $\sin \theta$ and $\cos \theta$ in terms of sines and cosines of multiples of θ , it is useful to use the following properties:

$$\begin{aligned}\cos(n\theta) &= \frac{1}{2}(e^{in\theta} + e^{-in\theta}) \\ \sin(n\theta) &= \frac{1}{2i}(e^{in\theta} - e^{-in\theta})\end{aligned}$$

To prove these formulae, consider $z = e^{in\theta} = \cos(n\theta) + i \sin(n\theta)$.

Then

$$\frac{1}{2}(e^{in\theta} + e^{-in\theta}) = \frac{1}{2}(z + \bar{z}) = \operatorname{Re}(z) = \cos(n\theta),$$

and

$$\frac{1}{2i}(e^{in\theta} - e^{-in\theta})$$

$\cos(n\theta)$ and $\sin(n\theta)$

The Binomial Theorem is particularly useful when expressing $\cos(n\theta)$ and $\sin(n\theta)$ in terms of powers of $\cos \theta$ and $\sin \theta$, and vice versa.

When expressing powers of $\sin \theta$ and $\cos \theta$ in terms of sines and cosines of multiples of θ , it is useful to use the following properties:

$$\begin{aligned}\cos(n\theta) &= \frac{1}{2}(e^{in\theta} + e^{-in\theta}) \\ \sin(n\theta) &= \frac{1}{2i}(e^{in\theta} - e^{-in\theta})\end{aligned}$$

To prove these formulae, consider $z = e^{in\theta} = \cos(n\theta) + i \sin(n\theta)$.

Then

$$\frac{1}{2}(e^{in\theta} + e^{-in\theta}) = \frac{1}{2}(z + \bar{z}) = \operatorname{Re}(z) = \cos(n\theta),$$

and

$$\frac{1}{2i}(e^{in\theta} - e^{-in\theta}) = \frac{1}{2i}(z - \bar{z})$$

$\cos(n\theta)$ and $\sin(n\theta)$

The Binomial Theorem is particularly useful when expressing $\cos(n\theta)$ and $\sin(n\theta)$ in terms of powers of $\cos \theta$ and $\sin \theta$, and vice versa.

When expressing powers of $\sin \theta$ and $\cos \theta$ in terms of sines and cosines of multiples of θ , it is useful to use the following properties:

$$\begin{aligned}\cos(n\theta) &= \frac{1}{2}(e^{in\theta} + e^{-in\theta}) \\ \sin(n\theta) &= \frac{1}{2i}(e^{in\theta} - e^{-in\theta})\end{aligned}$$

To prove these formulae, consider $z = e^{in\theta} = \cos(n\theta) + i \sin(n\theta)$.

Then

$$\frac{1}{2}(e^{in\theta} + e^{-in\theta}) = \frac{1}{2}(z + \bar{z}) = \operatorname{Re}(z) = \cos(n\theta),$$

and

$$\frac{1}{2i}(e^{in\theta} - e^{-in\theta}) = \frac{1}{2i}(z - \bar{z}) = \operatorname{Im}(z)$$

$\cos(n\theta)$ and $\sin(n\theta)$

The Binomial Theorem is particularly useful when expressing $\cos(n\theta)$ and $\sin(n\theta)$ in terms of powers of $\cos \theta$ and $\sin \theta$, and vice versa.

When expressing powers of $\sin \theta$ and $\cos \theta$ in terms of sines and cosines of multiples of θ , it is useful to use the following properties:

$$\begin{aligned}\cos(n\theta) &= \frac{1}{2}(e^{in\theta} + e^{-in\theta}) \\ \sin(n\theta) &= \frac{1}{2i}(e^{in\theta} - e^{-in\theta})\end{aligned}$$

To prove these formulae, consider $z = e^{in\theta} = \cos(n\theta) + i \sin(n\theta)$.

Then

$$\frac{1}{2}(e^{in\theta} + e^{-in\theta}) = \frac{1}{2}(z + \bar{z}) = \operatorname{Re}(z) = \cos(n\theta),$$

and

$$\frac{1}{2i}(e^{in\theta} - e^{-in\theta}) = \frac{1}{2i}(z - \bar{z}) = \operatorname{Im}(z) = \sin(n\theta).$$

Example

Use De Moivre's Theorem to find formulae for $\cos(4\theta)$ and $\sin(4\theta)$ in terms of powers of $\sin \theta$ and $\cos \theta$.

Example

Use De Moivre's Theorem to find formulae for $\cos(4\theta)$ and $\sin(4\theta)$ in terms of powers of $\sin \theta$ and $\cos \theta$.

De Moivre's Theorem tells us that

$$\cos(4\theta) + i \sin(4\theta) = (\cos \theta + i \sin \theta)^4.$$

Example

Use De Moivre's Theorem to find formulae for $\cos(4\theta)$ and $\sin(4\theta)$ in terms of powers of $\sin \theta$ and $\cos \theta$.

De Moivre's Theorem tells us that

$$\cos(4\theta) + i \sin(4\theta) = (\cos \theta + i \sin \theta)^4.$$

Expanding the right-hand side:

$$\begin{aligned} & (\cos \theta + i \sin \theta)^4 \\ &= \cos^4 \theta + 4i \cos^3 \theta \sin \theta + 6i^2 \cos^2 \theta \sin^2 \theta + 4i^3 \cos \theta \sin^3 \theta + i^4 \sin^4 \theta \\ &= \cos^4 \theta + 4i \cos^3 \theta \sin \theta - 6 \cos^2 \theta \sin^2 \theta - 4i \cos \theta \sin^3 \theta + \sin^4 \theta \end{aligned}$$

Example

Use De Moivre's Theorem to find formulae for $\cos(4\theta)$ and $\sin(4\theta)$ in terms of powers of $\sin \theta$ and $\cos \theta$.

De Moivre's Theorem tells us that

$$\cos(4\theta) + i \sin(4\theta) = (\cos \theta + i \sin \theta)^4.$$

Expanding the right-hand side:

$$\begin{aligned} & (\cos \theta + i \sin \theta)^4 \\ &= \cos^4 \theta + 4i \cos^3 \theta \sin \theta + 6i^2 \cos^2 \theta \sin^2 \theta + 4i^3 \cos \theta \sin^3 \theta + i^4 \sin^4 \theta \\ &= \cos^4 \theta + 4i \cos^3 \theta \sin \theta - 6 \cos^2 \theta \sin^2 \theta - 4i \cos \theta \sin^3 \theta + \sin^4 \theta \end{aligned}$$

So equating real and imaginary parts, we find

$$\cos(4\theta) = \cos^4(\theta) - 6 \cos^2 \theta \sin^2 \theta + \sin^4 \theta,$$

Example

Use De Moivre's Theorem to find formulae for $\cos(4\theta)$ and $\sin(4\theta)$ in terms of powers of $\sin \theta$ and $\cos \theta$.

De Moivre's Theorem tells us that

$$\cos(4\theta) + i \sin(4\theta) = (\cos \theta + i \sin \theta)^4.$$

Expanding the right-hand side:

$$\begin{aligned} & (\cos \theta + i \sin \theta)^4 \\ &= \cos^4 \theta + 4i \cos^3 \theta \sin \theta + 6i^2 \cos^2 \theta \sin^2 \theta + 4i^3 \cos \theta \sin^3 \theta + i^4 \sin^4 \theta \\ &= \cos^4 \theta + 4i \cos^3 \theta \sin \theta - 6 \cos^2 \theta \sin^2 \theta - 4i \cos \theta \sin^3 \theta + \sin^4 \theta \end{aligned}$$

So equating real and imaginary parts, we find

$$\cos(4\theta) = \cos^4(\theta) - 6 \cos^2 \theta \sin^2 \theta + \sin^4 \theta,$$

and

$$\sin(4\theta) = 4 \cos^3 \theta \sin \theta - 4 \cos \theta \sin^3 \theta.$$

$\cos^n \theta$ and $\sin^n \theta$

Example

Write $\sin^3 \theta$ in terms of sines of multiples of θ , and find $\int \sin^3 \theta d\theta$.

Example

Write $\sin^3 \theta$ in terms of sines of multiples of θ , and find $\int \sin^3 \theta d\theta$.

Using $\sin(n\theta) = \frac{1}{2i}(e^{in\theta} - e^{-in\theta})$:

Example

Write $\sin^3 \theta$ in terms of sines of multiples of θ , and find $\int \sin^3 \theta d\theta$.

Using $\sin(n\theta) = \frac{1}{2i}(e^{in\theta} - e^{-in\theta})$:

$$\sin^3 \theta = \left(\frac{1}{2i}(e^{i\theta} - e^{-i\theta}) \right)^3$$

Example

Write $\sin^3 \theta$ in terms of sines of multiples of θ , and find $\int \sin^3 \theta d\theta$.

Using $\sin(n\theta) = \frac{1}{2i}(e^{in\theta} - e^{-in\theta})$:

$$\begin{aligned}\sin^3 \theta &= \left(\frac{1}{2i}(e^{i\theta} - e^{-i\theta}) \right)^3 \\ &= \frac{1}{-8i}(e^{3i\theta} - 3e^{2i\theta}e^{-i\theta} + 3e^{i\theta}e^{-2i\theta} - e^{-3i\theta})\end{aligned}$$

Example

Write $\sin^3 \theta$ in terms of sines of multiples of θ , and find $\int \sin^3 \theta d\theta$.

Using $\sin(n\theta) = \frac{1}{2i}(e^{in\theta} - e^{-in\theta})$:

$$\begin{aligned}\sin^3 \theta &= \left(\frac{1}{2i}(e^{i\theta} - e^{-i\theta}) \right)^3 \\ &= \frac{1}{-8i}(e^{3i\theta} - 3e^{2i\theta}e^{-i\theta} + 3e^{i\theta}e^{-2i\theta} - e^{-3i\theta}) \\ &= -\frac{1}{4} \left(\frac{1}{2i}(e^{3i\theta} - e^{-3i\theta}) - \frac{1}{2i}(3e^{i\theta} - 3e^{-i\theta}) \right)\end{aligned}$$

Example

Write $\sin^3 \theta$ in terms of sines of multiples of θ , and find $\int \sin^3 \theta d\theta$.

Using $\sin(n\theta) = \frac{1}{2i}(e^{in\theta} - e^{-in\theta})$:

$$\begin{aligned}\sin^3 \theta &= \left(\frac{1}{2i}(e^{i\theta} - e^{-i\theta}) \right)^3 \\&= \frac{1}{-8i}(e^{3i\theta} - 3e^{2i\theta}e^{-i\theta} + 3e^{i\theta}e^{-2i\theta} - e^{-3i\theta}) \\&= -\frac{1}{4} \left(\frac{1}{2i}(e^{3i\theta} - e^{-3i\theta}) - \frac{1}{2i}(3e^{i\theta} - 3e^{-i\theta}) \right) \\&= -\frac{1}{4}(\sin(3\theta) - 3\sin \theta)\end{aligned}$$

Example

Write $\sin^3 \theta$ in terms of sines of multiples of θ , and find $\int \sin^3 \theta d\theta$.

Using $\sin(n\theta) = \frac{1}{2i}(e^{in\theta} - e^{-in\theta})$:

$$\begin{aligned}\sin^3 \theta &= \left(\frac{1}{2i}(e^{i\theta} - e^{-i\theta}) \right)^3 \\&= \frac{1}{-8i}(e^{3i\theta} - 3e^{2i\theta}e^{-i\theta} + 3e^{i\theta}e^{-2i\theta} - e^{-3i\theta}) \\&= -\frac{1}{4} \left(\frac{1}{2i}(e^{3i\theta} - e^{-3i\theta}) - \frac{1}{2i}(3e^{i\theta} - 3e^{-i\theta}) \right) \\&= -\frac{1}{4}(\sin(3\theta) - 3\sin \theta)\end{aligned}$$

So

$$\int \sin^3 \theta d\theta$$

Example

Write $\sin^3 \theta$ in terms of sines of multiples of θ , and find $\int \sin^3 \theta d\theta$.

Using $\sin(n\theta) = \frac{1}{2i}(e^{in\theta} - e^{-in\theta})$:

$$\begin{aligned}\sin^3 \theta &= \left(\frac{1}{2i}(e^{i\theta} - e^{-i\theta}) \right)^3 \\&= \frac{1}{-8i}(e^{3i\theta} - 3e^{2i\theta}e^{-i\theta} + 3e^{i\theta}e^{-2i\theta} - e^{-3i\theta}) \\&= -\frac{1}{4} \left(\frac{1}{2i}(e^{3i\theta} - e^{-3i\theta}) - \frac{1}{2i}(3e^{i\theta} - 3e^{-i\theta}) \right) \\&= -\frac{1}{4}(\sin(3\theta) - 3\sin \theta)\end{aligned}$$

So

$$\int \sin^3 \theta d\theta = \int \left(-\frac{1}{4}(\sin(3\theta) - 3\sin \theta) \right) d\theta$$

Example

Write $\sin^3 \theta$ in terms of sines of multiples of θ , and find $\int \sin^3 \theta d\theta$.

Using $\sin(n\theta) = \frac{1}{2i}(e^{in\theta} - e^{-in\theta})$:

$$\begin{aligned}\sin^3 \theta &= \left(\frac{1}{2i}(e^{i\theta} - e^{-i\theta}) \right)^3 \\&= \frac{1}{-8i}(e^{3i\theta} - 3e^{2i\theta}e^{-i\theta} + 3e^{i\theta}e^{-2i\theta} - e^{-3i\theta}) \\&= -\frac{1}{4} \left(\frac{1}{2i}(e^{3i\theta} - e^{-3i\theta}) - \frac{1}{2i}(3e^{i\theta} - 3e^{-i\theta}) \right) \\&= -\frac{1}{4}(\sin(3\theta) - 3\sin \theta)\end{aligned}$$

So

$$\int \sin^3 \theta d\theta = \int \left(-\frac{1}{4}(\sin(3\theta) - 3\sin \theta) \right) d\theta = \frac{1}{12} \cos(3\theta) - \frac{3}{4} \cos \theta.$$