

# Convex set

**line segment** between  $x_1$  and  $x_2$ : all points


$$x = \theta x_1 + (1 - \theta)x_2$$

with  $0 \leq \theta \leq 1$

**convex set**: contains line segment between any two points in the set

$$x_1, x_2 \in C, \quad 0 \leq \theta \leq 1 \quad \implies \quad \theta x_1 + (1 - \theta)x_2 \in C$$

**examples** (one convex, two nonconvex sets)





# Convex combination and convex hull

**convex combination** of  $x_1, \dots, x_k$ : any point  $x$  of the form

$$x = \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_k x_k$$

with  $\theta_1 + \dots + \theta_k = 1$ ,  $\theta_i \geq 0$

**convex hull**  $\text{conv } S$ : set of all convex combinations of points in  $S$




# Convex cone

**conic (nonnegative) combination** of  $x_1$  and  $x_2$ : any point of the form

$$x = \theta_1 x_1 + \theta_2 x_2$$


with  $\theta_1 \geq 0$ ,  $\theta_2 \geq 0$




**convex cone**: set that contains all conic combinations of points in the set

# Hyperplanes and halfspaces

**hyperplane:** set of the form  $\{x \mid a^T x = b\}$  ( $a \neq 0$ )



**halfspace:** set of the form  $\{x \mid a^T x \leq b\}$  ( $a \neq 0$ )



- $a$  is the normal vector
- hyperplanes are affine and convex; halfspaces are convex

# Euclidean balls and ellipsoids


**(Euclidean) ball** with center  $x_c$  and radius  $r$ :

$$B(x_c, r) = \{x \mid \|x - x_c\|_2 \leq r\} = \{x_c + ru \mid \|u\|_2 \leq 1\}$$

**ellipsoid:** set of the form

$$\{x \mid (x - x_c)^T P^{-1} (x - x_c) \leq 1\}$$

with  $P \in \mathbf{S}_{++}^n$  (i.e.,  $P$  symmetric positive definite)



other representation:  $\{x_c + Au \mid \|u\|_2 \leq 1\}$  with  $A$  square and nonsingular


# Convex Functions

## Definition

$f : \mathbf{R}^n \rightarrow \mathbf{R}$  is convex if

$$f(\theta x + (1 - \theta)y) \leq \theta f(x) + (1 - \theta)f(y)$$

for all  $x, y \in \mathbf{dom} f$ ,  $0 \leq \theta \leq 1$



- $f$  is concave if  $-f$  is convex
- $f$  is strictly convex if

$$f(\theta x + (1 - \theta)y) < \theta f(x) + (1 - \theta)f(y)$$

for  $x, y \in \mathbf{dom} f$ ,  $x \neq y$ ,  $0 < \theta < 1$

## Examples on $\mathbf{R}$

convex:

- affine:  $ax + b$  on  $\mathbf{R}$ , for any  $a, b \in \mathbf{R}$
- exponential:  $e^{ax}$ , for any  $a \in \mathbf{R}$
- powers:  $x^\alpha$  on  $\mathbf{R}_{++}$ , for  $\alpha \geq 1$  or  $\alpha \leq 0$
- powers of absolute value:  $|x|^p$  on  $\mathbf{R}$ , for  $p \geq 1$
- negative entropy:  $x \log x$  on  $\mathbf{R}_{++}$

concave:

- affine:  $ax + b$  on  $\mathbf{R}$ , for any  $a, b \in \mathbf{R}$
- powers:  $x^\alpha$  on  $\mathbf{R}_{++}$ , for  $0 \leq \alpha \leq 1$
- logarithm:  $\log x$  on  $\mathbf{R}_{++}$

## Examples on $\mathbf{R}^n$ and $\mathbf{R}^{m \times n}$

affine functions are convex and concave; all norms are convex

### examples on $\mathbf{R}^n$

- affine function  $f(x) = a^T x + b$
- norms:  $\|x\|_p = (\sum_{i=1}^n |x_i|^p)^{1/p}$  for  $p \geq 1$ ;  $\|x\|_\infty = \max_k |x_k|$

### examples on $\mathbf{R}^{m \times n}$ ( $m \times n$ matrices)

- affine function

$$f(X) = \text{tr}(A^T X) + b = \sum_{i=1}^m \sum_{j=1}^n A_{ij} X_{ij} + b$$

- spectral (maximum singular value) norm

$$f(X) = \|X\|_2 = \sigma_{\max}(X) = (\lambda_{\max}(X^T X))^{1/2}$$



# First-order condition


$f$  is **differentiable** if  $\text{dom } f$  is open and the gradient

$$\nabla f(x) = \left( \frac{\partial f(x)}{\partial x_1}, \frac{\partial f(x)}{\partial x_2}, \dots, \frac{\partial f(x)}{\partial x_n} \right)$$

exists at each  $x \in \text{dom } f$

**1st-order condition:** differentiable  $f$  with convex domain is convex iff

$$f(y) \geq f(x) + \nabla f(x)^T (y - x) \quad \text{for all } x, y \in \text{dom } f$$



first-order approximation of  $f$  is global underestimator

## Second-order conditions

$f$  is **twice differentiable** if  $\text{dom } f$  is open and the Hessian  $\nabla^2 f(x) \in \mathbf{S}^n$ ,

$$\nabla^2 f(x)_{ij} = \frac{\partial^2 f(x)}{\partial x_i \partial x_j}, \quad i, j = 1, \dots, n,$$

exists at each  $x \in \text{dom } f$

**2nd-order conditions:** for twice differentiable  $f$  with convex domain

- $f$  is convex if and only if

$$\nabla^2 f(x) \succeq 0 \quad \text{for all } x \in \text{dom } f$$

- if  $\nabla^2 f(x) \succ 0$  for all  $x \in \text{dom } f$ , then  $f$  is strictly convex

## Examples

**quadratic function:**  $f(x) = (1/2)x^T Px + q^T x + r$  (with  $P \in \mathbf{S}^n$ )

$$\nabla f(x) = Px + q, \quad \nabla^2 f(x) = P$$

convex if  $P \succeq 0$

**least-squares objective:**  $f(x) = \|Ax - b\|_2^2$


$$\nabla f(x) = 2A^T(Ax - b), \quad \nabla^2 f(x) = 2A^T A$$

convex (for any  $A$ )

**quadratic-over-linear:**  $f(x, y) = x^2/y$

$$\nabla^2 f(x, y) = \frac{2}{y^3} \begin{bmatrix} y \\ -x \end{bmatrix} \begin{bmatrix} y \\ -x \end{bmatrix}^T \succeq 0$$

convex for  $y > 0$



# Jensen's inequality

**basic inequality:** if  $f$  is convex, then for  $0 \leq \theta \leq 1$ ,

$$f(\theta x + (1 - \theta)y) \leq \theta f(x) + (1 - \theta)f(y)$$

**extension:** if  $f$  is convex, then

$$f(\mathbf{E} z) \leq \mathbf{E} f(z)$$

for any random variable  $z$

basic inequality is special case with discrete distribution

$$\mathbf{prob}(z = x) = \theta, \quad \mathbf{prob}(z = y) = 1 - \theta$$

# Operations that preserve convexity

practical methods for establishing convexity of a function

1. verify definition (often simplified by restricting to a line)
2. for twice differentiable functions, show  $\nabla^2 f(x) \succeq 0$
3. show that  $f$  is obtained from simple convex functions by operations that preserve convexity
  - nonnegative weighted sum
  - composition with affine function
  - pointwise maximum
  - composition

# Positive weighted sum & composition with affine function

**nonnegative multiple:**  $\alpha f$  is convex if  $f$  is convex,  $\alpha \geq 0$

**sum:**  $f_1 + f_2$  convex if  $f_1, f_2$  convex (extends to infinite sums, integrals)

**composition with affine function:**  $f(Ax + b)$  is convex if  $f$  is convex

## examples

- log barrier for linear inequalities

$$f(x) = -\sum_{i=1}^m \log(b_i - a_i^T x), \quad \text{dom } f = \{x \mid a_i^T x < b_i, i = 1, \dots, m\}$$

- (any) norm of affine function:  $f(x) = \|Ax + b\|$

## Pointwise maximum

if  $f_1, \dots, f_m$  are convex, then  $f(x) = \max\{f_1(x), \dots, f_m(x)\}$  is convex

### examples

- piecewise-linear function:  $f(x) = \max_{i=1,\dots,m}(a_i^T x + b_i)$  is convex
- sum of  $r$  largest components of  $x \in \mathbf{R}^n$ :

$$f(x) = x_{[1]} + x_{[2]} + \dots + x_{[r]}$$

is convex ( $x_{[i]}$  is  $i$ th largest component of  $x$ )

proof:

$$f(x) = \max\{x_{i_1} + x_{i_2} + \dots + x_{i_r} \mid 1 \leq i_1 < i_2 < \dots < i_r \leq n\}$$

# Composition with scalar functions

composition of  $g : \mathbf{R}^n \rightarrow \mathbf{R}$  and  $h : \mathbf{R} \rightarrow \mathbf{R}$ :

$$f(x) = h(g(x))$$

$f$  is convex if  $\begin{array}{l} g \text{ convex, } h \text{ convex, } h \text{ nondecreasing} \\ g \text{ concave, } h \text{ convex, } h \text{ nonincreasing} \end{array}$

- proof (for  $n = 1$ , differentiable  $g, h$ )

$$f''(x) = h''(g(x))g'(x)^2 + h'(g(x))g''(x)$$

## examples

- $\exp g(x)$  is convex if  $g$  is convex
- $1/g(x)$  is convex if  $g$  is concave and positive