

Supplemental Material




Figure S1. Representative flow cytometry of LSECs after anti-mouse CD146 column isolation. Gating strategies of LSECs following liver perfusion and LSEC isolation.

Table S1. Deletion Frequency Patterns Following Cas9/sgRNA LNP Treatment.

	-1	-2	-3	-4	>-4	total deletion frequency
Hepatocyte	34.40 ± 5.41 %	4.36 ± 1.29 %	3.01 ± 0.93 %	4.62 ± 1.27 %	14.15 ± 3.40 %	60.54 ± 11.65 %
LSEC	7.68 ± 1.90 %	1.03 ± 0.28 %	0.73 ± 0.14 %	1.38 ± 0.37 %	5.68 ± 1.78 %	16.50 ± 2.96 %

Table S2. Insertion Frequency Patterns Following Cas9/sgRNA LNP Treatment.

	+1	+2	+3	+4	>+4	total insertion frequency
Hepatocyte	3.39 ± 0.91 %	0.83 ± 0.27 %	0.01 ± 0.01 %	0.00	0.00	4.23 ± 1.17 %
LSEC	0.78 ± 0.27 %	0.21 ± 0.10 %	0.01 ± 0.01 %	0.00	0.00	1.01 ± 0.38 %