

Spatio-temporal analysis

Dr Emanuele Giorgi
Lancaster University
`e.giorgi@lancaster.ac.uk`

Toowoomba 21-25 October 2019

Overview

- ▶ Repeated cross-sectional surveys.

Overview

- ▶ Repeated cross-sectional surveys.
- ▶ Analysing reported cases to health facilities.

Geostatistical problems and geostatistical methods

- ▶ **Geostatistical problems:** The object of scientific interest is to make inference on a spatially continuous surface (e.g. disease risk) or some of its properties (e.g. regional prevalence) in a geographical area of interest.
- ▶ **Model-based geostatistics:** Likelihood-based methods of inference that use spatially sparse noisy data to make inference on a spatially continuous surface of interest.


Geostatistical problems and geostatistical methods

- ▶ **Geostatistical problems:** The object of scientific interest is to make inference on a spatially continuous surface (e.g. disease risk) or some of its properties (e.g. regional prevalence) in a geographical area of interest.
- ▶ **Model-based geostatistics:** Likelihood-based methods of inference that use spatially sparse noisy data to make inference on a spatially continuous surface of interest.
- ▶ Geostatistical problems are **NOT** dependent on the data-format (e.g. point-level or areal-level data).

Geostatistical problems: some examples

- ▶ Spatio-temporal interpolation of disease risk.
- ▶ Identification of disease hotspots (i.e. areas where disease risk exceeds policy relevant thresholds).
- ▶ Identification of low transmission areas (i.e. areas where disease risk is below policy relevant thresholds).
- ▶ Estimating the association between risk factors and disease risk metrics.
- ▶ Accounting for the excess of zero-reported cases (spatially structured zero-inflation).
- ▶ Combining data from randomised survey and convenience samples.

The data



The standard model for prevalence

► $\mathcal{D} = \{(x_i, t_i, y_i, n_i) : x_i \in A, i = 1, \dots, N^* = \sum_{i=1}^N n_i\}$

The standard model for prevalence

- ▶ $\mathcal{D} = \{(x_i, t_i, y_i, n_i) : x_i \in A, i = 1, \dots, N^* = \sum_{i=1}^N n_i\}$
- ▶ $d(x, t)$ = explanatory variables.

The standard model for prevalence

- ▶ $\mathcal{D} = \{(x_i, t_i, y_i, n_i) : x_i \in A, i = 1, \dots, N^* = \sum_{i=1}^N n_i\}$
- ▶ $d(x, t)$ = explanatory variables.
- ▶ $S(x, t)$ = unexplained spatial variation.

The standard model for prevalence

- ▶ $\mathcal{D} = \{(x_i, t_i, y_i, n_i) : x_i \in A, i = 1, \dots, N^* = \sum_{i=1}^N n_i\}$
- ▶ $d(x, t)$ = explanatory variables.
- ▶ $S(x, t)$ = unexplained spatial variation.
- ▶ $Z(x, t)$ = unexplained unstructured variation.

The standard model for prevalence

- ▶ $\mathcal{D} = \{(x_i, t_i, y_i, n_i) : x_i \in A, i = 1, \dots, N^* = \sum_{i=1}^N n_i\}$
- ▶ $d(x, t)$ = explanatory variables.
- ▶ $S(x, t)$ = unexplained spatial variation.
- ▶ $Z(x, t)$ = unexplained unstructured variation.
- ▶ A model for the data.

$$\log \left\{ \frac{p(x_i, t_i)}{1 - p(x_i, t_i)} \right\} = d(x_i, t_i)^\top \beta + S(x_i, t_i) + Z(x_i, t_i). \quad (1)$$

The standard model for prevalence

- ▶ $\mathcal{D} = \{(x_i, t_i, y_i, n_i) : x_i \in A, i = 1, \dots, N^* = \sum_{i=1}^N n_i\}$
- ▶ $d(x, t)$ = explanatory variables.
- ▶ $S(x, t)$ = unexplained spatial variation.
- ▶ $Z(x, t)$ = unexplained unstructured variation.
- ▶ A model for the data.

$$\log \left\{ \frac{p(x_i, t_i)}{1 - p(x_i, t_i)} \right\} = d(x_i, t_i)^\top \beta + S(x_i, t_i) + Z(x_i, t_i). \quad (1)$$

- ▶ $E[S(x, t)] = E[Z(x, t)] = 0$; $V[S(x, t)] = \sigma^2$; $V[Z(x, t)] = \tau^2$ and
 $\text{corr}(S(x, t), S(x', t')) = \rho(x, x', t, t'; \theta)$.

The standard model for prevalence

- ▶ $\mathcal{D} = \{(x_i, t_i, y_i, n_i) : x_i \in A, i = 1, \dots, N^* = \sum_{i=1}^N n_i\}$
- ▶ $d(x, t)$ = explanatory variables.
- ▶ $S(x, t)$ = unexplained spatial variation.
- ▶ $Z(x, t)$ = unexplained unstructured variation.
- ▶ A model for the data.

$$\log \left\{ \frac{p(x_i, t_i)}{1 - p(x_i, t_i)} \right\} = d(x_i, t_i)^\top \beta + S(x_i, t_i) + Z(x_i, t_i). \quad (1)$$

- ▶ $E[S(x, t)] = E[Z(x, t)] = 0$; $V[S(x, t)] = \sigma^2$; $V[Z(x, t)] = \tau^2$ and
 $\text{corr}(S(x, t), S(x', t')) = \rho(x, x', t, t'; \theta)$.

- ▶ **Question.** What can we do with this model?

Exploring the residual variation

- ▶ $W(x, t) = S(x, t) + Z(x, t)$

Exploring the residual variation

- ▶ $W(x, t) = S(x, t) + Z(x, t)$
- ▶ $u = \|x - x'\|, v = |t - t'|.$

Exploring the residual variation

- ▶ $W(x, t) = S(x, t) + Z(x, t)$
- ▶ $u = \|x - x'\|$, $v = |t - t'|$.
- ▶ The theoretical STV

$$\begin{aligned}\gamma(u, v; \theta) &= E[\{W(x, t) - W(x', t')\}^2] \\ &= \tau^2 + \sigma^2[1 - \rho(u, v; \theta)].\end{aligned}\tag{2}$$

- ▶ The empirical STV

$$\tilde{\gamma}(u, v) = \frac{1}{2|N(u, v)|} \sum_{(i,j) \in N(u,v)} \{\tilde{Z}(x_i, t_i) - \tilde{Z}(x_j, t_j)\}^2. \tag{3}$$

A test for spatio-temporal correlation

- ▶ $\tilde{Z}(x, t)$ = point estimate from a non-spatial binomial mixed model.

A test for spatio-temporal correlation

- ▶ $\tilde{Z}(x, t)$ = point estimate from a non-spatial binomial mixed model.
- ▶ The algorithm.
 - (i) permute the order of the data, including $\tilde{Z}(x_i, t_i)$, while holding (x_i, t_i) fixed;

A test for spatio-temporal correlation

- ▶ $\tilde{Z}(x, t)$ = point estimate from a non-spatial binomial mixed model.
- ▶ The algorithm.
 - (i) permute the order of the data, including $\tilde{Z}(x_i, t_i)$, while holding (x_i, t_i) fixed;
 - (ii) compute the empirical semi-variogram for $\tilde{Z}(x_i, t_i)$;


A test for spatio-temporal correlation

- ▶ $\tilde{Z}(x, t)$ = point estimate from a non-spatial binomial mixed model.
- ▶ The algorithm.
 - (i) permute the order of the data, including $\tilde{Z}(x_i, t_i)$, while holding (x_i, t_i) fixed;
 - (ii) compute the empirical semi-variogram for $\tilde{Z}(x_i, t_i)$;
 - (iii) repeat(i) and (ii) a large enough number of times, say B ;

A test for spatio-temporal correlation

- ▶ $\tilde{Z}(x, t)$ = point estimate from a non-spatial binomial mixed model.
- ▶ The algorithm.
 - (i) permute the order of the data, including $\tilde{Z}(x_i, t_i)$, while holding (x_i, t_i) fixed;
 - (ii) compute the empirical semi-variogram for $\tilde{Z}(x_i, t_i)$;
 - (iii) repeat (i) and (ii) a large enough number of times, say B ;
 - (iv) use the resulting B empirical semi-variograms to generate 95% tolerance intervals at each of the pre-defined distance bins.

A test for spatio-temporal correlation



The covariance model

- ▶ The Gneiting's spatio-temporal correlation function.

$$\rho(u, v; \theta) = \frac{1}{(1 + v/\psi)^{\delta+1}} \exp \left\{ -\frac{u/\phi}{(1 + v/\psi)^{\xi/2}} \right\}. \quad (4)$$

In the application $\delta = \xi = 0$.

The covariance model

- ▶ The Gneiting's spatio-temporal correlation function.

$$\rho(u, v; \theta) = \frac{1}{(1 + v/\psi)^{\delta+1}} \exp \left\{ -\frac{u/\phi}{(1 + v/\psi)^{\xi/2}} \right\}. \quad (4)$$

In the application $\delta = \xi = 0$.

- ▶ How to choose a covariance model?

The covariance model

- ▶ The Gneiting's spatio-temporal correlation function.

$$\rho(u, v; \theta) = \frac{1}{(1 + v/\psi)^{\delta+1}} \exp \left\{ -\frac{u/\phi}{(1 + v/\psi)^{\xi/2}} \right\}. \quad (4)$$

In the application $\delta = \xi = 0$.

- ▶ How to choose a covariance model?
- ▶ Guiding principle: **parsimony**, e.g. find the simplest model that can explain the variation of the data.

Parameter estimation

- ▶ $[\cdot]$ = “the distribution of \cdot ”.

Parameter estimation

- ▶ $[\cdot]$ = “the distribution of \cdot ”.
- ▶ $\lambda^\top = (\beta^\top, \sigma^2, \theta^\top)$.

Parameter estimation

- ▶ $[\cdot]$ = “the distribution of \cdot ”.
- ▶ $\lambda^\top = (\beta^\top, \sigma^2, \theta^\top)$.
- ▶ The likelihood function.

$$L(\lambda) = [y|\lambda] = \int [W, y|\lambda] dW.$$

- ▶ Bayesian inference.

$$[\lambda|y] = \frac{[\lambda][y|\lambda]}{[y]} = \frac{[\lambda]L(\lambda)}{[y]}.$$

Parameter estimation

- ▶ $[\cdot]$ = “the distribution of \cdot ”.
- ▶ $\lambda^\top = (\beta^\top, \sigma^2, \theta^\top)$.
- ▶ The likelihood function.


$$L(\lambda) = [y|\lambda] = \int [W, y|\lambda] dW.$$

- ▶ Bayesian inference.

$$[\lambda|y] = \frac{[\lambda][y|\lambda]}{[y]} = \frac{[\lambda]L(\lambda)}{[y]}.$$

- ▶ Prior distributions.
 - ▶ $\beta \sim N(0, 10^4)$
 - ▶ $\sigma^2 \sim \text{Uniform}(0, 20)$
 - ▶ $\phi \sim \text{Uniform}(0, 1000)$
 - ▶ $\tau^2/\sigma^2 \sim \text{Uniform}(0, 20)$
 - ▶ $\psi \sim \text{Uniform}(0, 20)$

Estimation: Likelihood-based vs Bayesian inference



Validation of the covariance model


- ▶ Replace (i) with
 - (a) Simulate $W(x_i, t_i)$ at observed locations x_i and times t_i , for $i = 1, \dots, n$, from its marginal multivariate distribution under the assumed model.
 - (b) Conditionally on the simulated values of $W(x_i, t_i)$, simulate binomial data y_i from (1).
 - (c) Compute a point estimates $\tilde{Z}(x_i, t_i)$ using the simulated data.

Validation of the covariance model

- ▶ Replace (i) with
 - (a) Simulate $W(x_i, t_i)$ at observed locations x_i and times t_i , for $i = 1, \dots, n$, from its marginal multivariate distribution under the assumed model.
 - (b) Conditionally on the simulated values of $W(x_i, t_i)$, simulate binomial data y_i from (1).
 - (c) Compute a point estimates $\tilde{Z}(x_i, t_i)$ using the simulated data.
- ▶ Test statistics.

$$T = \sum_{k=1}^K |N(u_k, t_k)| [\tilde{\gamma}(u_k, v_k) - \gamma(u_k, v_k; \theta)]^2. \quad (5)$$

Validation of the covariance model



Spatio-temporal prediction

- ▶ $W^* = \{W(x, t) : x \in \mathcal{X}, t \in \mathcal{T}\}.$

Spatio-temporal prediction

- ▶ $W^* = \{W(x, t) : x \in \mathcal{X}, t \in \mathcal{T}\}.$
- ▶ Likelihood-based inference (plug-in).

$$[W^*|y, \lambda] = \int [W^*, W|y, \lambda] dW$$

Spatio-temporal prediction

- ▶ $W^* = \{W(x, t) : x \in \mathcal{X}, t \in \mathcal{T}\}.$
- ▶ Likelihood-based inference (plug-in).

$$[W^*|y, \lambda] = \int [W^*, W|y, \lambda] dW$$

- ▶ Bayesian inference.

$$[W^*|y] = \int \int [\lambda|y][W|y, \lambda][W^*|W, \lambda] dW d\lambda,$$

Spatio-temporal prediction

- ▶ $W^* = \{W(x, t) : x \in \mathcal{X}, t \in \mathcal{T}\}.$
- ▶ Likelihood-based inference (plug-in).

$$[W^*|y, \lambda] = \int [W^*, W|y, \lambda] dW$$

- ▶ Bayesian inference.

$$[W^*|y] = \int \int [\lambda|y][W|y, \lambda][W^*|W, \lambda] dW d\lambda,$$

- ▶ Likelihood-based inference (accounting for parameter uncertainty).

$$[W^*|y] = \int \int [\hat{\lambda}|y][W|y, \hat{\lambda}][W^*|W, \hat{\lambda}] dW d\hat{\lambda},$$

Targets of prediction

Targets of prediction

- ▶ Exceedance probability surface.

$$\mathcal{R}_I(W^*) = \{r(x, t) = P(p(x, t) > I|y)\}. \quad (6)$$

Targets of prediction

- ▶ Exceedance probability surface.

$$\mathcal{R}_I(W^*) = \{r(x, t) = P(p(x, t) > I|y)\}. \quad (6)$$

- ▶ District-wide average prevalence.

$$p_t(D) = \frac{1}{|D|} \int_D p(x, t) dx, \quad (7)$$

Targets of prediction

- ▶ Exceedance probability surface.


$$\mathcal{R}_I(W^*) = \{r(x, t) = P(p(x, t) > I|y)\}. \quad (6)$$

- ▶ District-wide average prevalence.

$$p_t(D) = \frac{1}{|D|} \int_D p(x, t) dx, \quad (7)$$

[http://fhm-chicas-apps.lancs.ac.uk/shiny/users/giorgi/
mapMalariaSEN/](http://fhm-chicas-apps.lancs.ac.uk/shiny/users/giorgi/mapMalariaSEN/)

Prediction: Likelihood-based vs Bayesian inference



What is a disease outbreak?

- ▶ **WHO:** *“A disease outbreak is the occurrence of cases of disease in excess of what **would normally be expected** in a defined community, geographical area or season. An outbreak may occur in a restricted geographical area, or may extend over several countries. It may last for a few days or weeks, or for several years.”*

What is a disease outbreak?

- ▶ **WHO:** *"A disease outbreak is the occurrence of cases of disease in excess of what **would normally be expected** in a defined community, geographical area or season. An outbreak may occur in a restricted geographical area, or may extend over several countries. It may last for a few days or weeks, or for several years."*
- ▶ A disease outbreak is the occurrence of cases of disease in excess **of a policy-relevant risk threshold** in a defined community, geographical area or season.

- ▶ **Outcome:** Weekly reports of confirmed cases from 142 districts in Mozambique from the first week of 2010 through the ninth week of 2017.

- ▶ **Outcome:** Weekly reports of confirmed cases from 142 districts in Mozambique from the first week of 2010 through the ninth week of 2017.
- ▶ **Population:** WorldPop + 2007 Census Data.

- ▶ **Outcome:** Weekly reports of confirmed cases from 142 districts in Mozambique from the first week of 2010 through the ninth week of 2017.
- ▶ **Population:** WorldPop + 2007 Census Data.
- ▶ **Climatic variables:** FLDAS data are available from NASA at daily temporal resolution and 0.1 degree spatial resolution.
 - ▶ total rainfall (mm)
 - ▶ average weekly temperature ($^{\circ}\text{C}$)
 - ▶ average weekly relative humidity (%)
 - ▶ average weekly surface barometric pressure (hPa)
 - ▶ average weekly saturation vapour pressure deficit (mmHg)

Model formulation

- ▶ i =district; t =week

Model formulation

- ▶ i =district; t =week
- ▶ Y_{it} =random variable of the number of reported cases

Model formulation

- ▶ i =district; t =week
- ▶ Y_{it} =random variable of the number of reported cases
- ▶ d_{it} =vector of explanatory (climatic) variables

Model formulation

- ▶ i =district; t =week
- ▶ Y_{it} =random variable of the number of reported cases
- ▶ d_{it} =vector of explanatory (climatic) variables
- ▶ m_{it} =total number of under 5 children

Model formulation

- ▶ i =district; t =week
- ▶ Y_{it} =random variable of the number of reported cases
- ▶ d_{it} =vector of explanatory (climatic) variables
- ▶ m_{it} =total number of under 5 children
- ▶ $Y_{it}|S_{it} \sim \text{Pois}(m_{it}\lambda_{it})$, such that

$$\log\{\lambda_{it}\} = d_{it}^{\top}\beta + S_{it}$$

Model formulation

- ▶ i =district; t =week
- ▶ Y_{it} =random variable of the number of reported cases
- ▶ d_{it} =vector of explanatory (climatic) variables
- ▶ m_{it} =total number of under 5 children
- ▶ $Y_{it}|S_{it} \sim \text{Pois}(m_{it}\lambda_{it})$, such that

$$\log\{\lambda_{it}\} = d_{it}^{\top}\beta + S_{it}$$

Objective

Spatio-temporal mapping of exceedance probabilities

$$\text{Prob}\{\exp\{S_{it}\} > l | \text{data}\}$$

at a future week t .

Modelling the residual spatio-temporal variation

$$S_{it} = \frac{1}{|\mathcal{D}_i|} \int_{\mathcal{D}_i} S(x, t) dx$$

Modelling the residual spatio-temporal variation

$$S_{it} = \frac{1}{|\mathcal{D}_i|} \int_{\mathcal{D}_i} S(x, t) dx$$

- Spatial covariance structure

$$\text{cov}\{S_{it}, S_{jt}\} = \frac{\sigma^2}{|\mathcal{D}_i||\mathcal{D}_j|} \int_{\mathcal{D}_i} \int_{\mathcal{D}_j} \exp\{-\|x - x'\|/\phi\} dx dx'$$

Modelling the residual spatio-temporal variation

$$S_{it} = \frac{1}{|\mathcal{D}_i|} \int_{\mathcal{D}_i} S(x, t) dx$$

- Spatial covariance structure

$$\text{cov}\{S_{it}, S_{jt}\} = \frac{\sigma^2}{|\mathcal{D}_i||\mathcal{D}_j|} \int_{\mathcal{D}_i} \int_{\mathcal{D}_j} \exp\{-\|x - x'\|/\phi\} dx dx'$$

- $S_t = (S_{1,t}, \dots, S_{142,t})$.

Modelling the residual spatio-temporal variation

$$S_{it} = \frac{1}{|\mathcal{D}_i|} \int_{\mathcal{D}_i} S(x, t) dx$$

- ▶ Spatial covariance structure

$$\text{cov}\{S_{it}, S_{jt}\} = \frac{\sigma^2}{|\mathcal{D}_i||\mathcal{D}_j|} \int_{\mathcal{D}_i} \int_{\mathcal{D}_j} \exp\{-\|x - x'\|/\phi\} dx dx'$$

- ▶ $S_t = (S_{1,t}, \dots, S_{142,t})$.
- ▶ Temporal covariance structure

$$S_t = \rho S_{t-1} + W_t.$$

Models comparison and validation

Models comparison and validation

► $\mathcal{M}_1 : \log\{\lambda_{it}\} = d_{it}^\top \beta$

Models comparison and validation

- ▶ $\mathcal{M}_1 : \log\{\lambda_{it}\} = d_{it}^\top \beta$
- ▶ $\mathcal{M}_2 : \log\{\lambda_{it}\} = \alpha + S_{it}$

Models comparison and validation

- ▶ $\mathcal{M}_1 : \log\{\lambda_{it}\} = d_{it}^\top \beta$
- ▶ $\mathcal{M}_2 : \log\{\lambda_{it}\} = \alpha + S_{it}$
- ▶ $\mathcal{M}_3 : \log\{\lambda_{it}\} = d_{it}^\top \beta + S_{it}$

Models comparison and validation

- ▶ $\mathcal{M}_1 : \log\{\lambda_{it}\} = d_{it}^\top \beta$
- ▶ $\mathcal{M}_2 : \log\{\lambda_{it}\} = \alpha + S_{it}$
- ▶ $\mathcal{M}_3 : \log\{\lambda_{it}\} = d_{it}^\top \beta + S_{it}$
- ▶ Root-mean-square-error (RMSE)

$$\text{RMSE}_t = \sqrt{\frac{1}{142} \sum_{i=1}^{142} (\lambda_{it}^{emp} - \hat{\lambda}_{it})^2}$$

Models comparison and validation


- ▶ $\mathcal{M}_1 : \log\{\lambda_{it}\} = d_{it}^\top \beta$
- ▶ $\mathcal{M}_2 : \log\{\lambda_{it}\} = \alpha + S_{it}$
- ▶ $\mathcal{M}_3 : \log\{\lambda_{it}\} = d_{it}^\top \beta + S_{it}$
- ▶ Root-mean-square-error (RMSE)

$$\text{RMSE}_t = \sqrt{\frac{1}{142} \sum_{i=1}^{142} (\lambda_{it}^{emp} - \hat{\lambda}_{it})^2}$$


- ▶ 95% coverage probabilities (CI)

$$\text{CI}_t = \frac{1}{142} \sum_{i=1}^{142} I(\lambda_{it}^{0.025} < \lambda_{it}^{emp} < \lambda_{it}^{0.975})$$


Results: scale of spatial correlation







Results: validation




Results: exceedance probabilities




Results: exceedance probabilities




Results: exceedance probabilities




CHIBABAVA – 26 weeks




CHIBABAVA – 26 weeks




SANGA – 26 weeks




SANGA – 26 weeks




Results: exceedance probabilities




CHIBABAVA – 26 weeks




CHIBABAVA – 26 weeks



SANGA – 26 weeks




SANGA – 26 weeks




Results: incidence vs exceedance probabilities





Week 1, 2017 – Incidence per 1,000




Week 1, 2017 – EP




Results: exceedance probabilities




Results: exceedance probabilities




CHIBABAVA – 8 weeks




CHIBABAVA – 8 weeks



SANGA – 8 weeks



SANGA – 8 weeks



Conclusions

Conclusions

- ▶ We need to account for unmeasured risk factors (S_{it}).

Conclusions

- ▶ We need to account for unmeasured risk factors (S_{it}).
- ▶ Climatic variables provide information on the long term trends; residual spatio-temporal random effects on short term.

Conclusions

- ▶ We need to account for unmeasured risk factors (S_{it}).
- ▶ Climatic variables provide information on the long term trends; residual spatio-temporal random effects on short term.
- ▶ Exceedance probabilities are a useful tool for detecting outbreaks within at least 4 weeks.

Conclusions

- ▶ We need to account for unmeasured risk factors (S_{it}).
- ▶ Climatic variables provide information on the long term trends; residual spatio-temporal random effects on short term.
- ▶ Exceedance probabilities are a useful tool for detecting outbreaks within at least 4 weeks.
- ▶ Outstanding issue: how to identify relative risk thresholds.

Conclusions

- ▶ We need to account for unmeasured risk factors (S_{it}).
- ▶ Climatic variables provide information on the long term trends; residual spatio-temporal random effects on short term.
- ▶ Exceedance probabilities are a useful tool for detecting outbreaks within at least 4 weeks.
- ▶ Outstanding issue: how to identify relative risk thresholds.
- ▶ Web interface:
<http://www.lancaster.ac.uk/staff/haleac/malaria/>(by Dr Alison Hale)

Conclusions

- ▶ We need to account for unmeasured risk factors (S_{it}).
- ▶ Climatic variables provide information on the long term trends; residual spatio-temporal random effects on short term.
- ▶ Exceedance probabilities are a useful tool for detecting outbreaks within at least 4 weeks.
- ▶ Outstanding issue: how to identify relative risk thresholds.
- ▶ Web interface:
<http://www.lancaster.ac.uk/staff/haleac/malaria/>(by Dr Alison Hale)