

Exercises 6, 12, pp. 52-53.

6. (a) If  $H$  is a subgroup of  $G$ , then for any  $h, h' \in H$ , we have  $h^{-1}h'h \in H$ . Hence  $h^{-1}Hh = H$ , and  $h \in N_G(H)$ . Therefore  $H \leq N_G(H)$ .

If  $H$  is not a subgroup of  $G$ , then multiplication fails so we have no reason to expect  $h^{-1}h'h \in H$ . For example, let

$$H = \left\{ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix} \right\}.$$

Then

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 2 & -5 \\ 0 & 3 \end{pmatrix} \notin H.$$

Hence  $H \not\leq N_G(H)$ .

- (b) ( $\Rightarrow$ ): If  $H$  is abelian, then clearly  $h'h = hh'$  for all  $h, h' \in H$ . Then  $H \leq C_G(H)$ .  
 ( $\Leftarrow$ ): If  $H \leq C_G(H)$ , then for any  $h, h' \in H$ , we have  $h^{-1}h'h = h' \Rightarrow h'h = hh'$ .  
 Hence  $H$  is abelian, as desired.

12. We do each part:

(a) Define

$$p(x_1, \dots, x_4) = 12x_1^5x_2^7x_3 - 18x_2^3x_3 + 11x_1^6x_2x_3^3x_4^{23}.$$

Direct calculation shows that:

$$\begin{aligned} \sigma \cdot p &= (1234) \cdot p \\ &= 12x_2^5x_3^7x_4 - 18x_3^3x_4 + 11x_1^{23}x_2^6x_3x_4^3 \end{aligned}$$

$$\begin{aligned} \tau \cdot (\sigma \cdot p) &= (123) \cdot 12x_2^5x_3^7x_4 - 18x_3^3x_4 + 11x_1^{23}x_2^6x_3x_4^3 \\ &= 12x_1^7x_3^5x_4 - 18x_1^3x_4 + 11x_1x_2^{23}x_3^6x_4^3 \end{aligned}$$

$$\begin{aligned} (\tau \circ \sigma) \cdot p &= (1342) \cdot p \\ &= 12x_1^7x_3^5x_4 - 18x_1^3x_4 + 11x_1x_2^{23}x_3^6x_4^3 \end{aligned}$$

$$\begin{aligned} (\sigma \circ \tau) \cdot p &= (1324) \cdot p \\ &= 12x_2x_3^5x_4^7 - 18x_4^3 + 11x_1^{23}x_2^3x_3^6x_4. \end{aligned}$$

- (b) This definition gives a left group action of  $S_4$  on  $R$ . If  $\sigma, \tau \in S_4$  and  $p \in R$ , then

$$\begin{aligned}\tau \cdot (\sigma \cdot p) &= \tau \cdot (p(x_{\sigma(1)}, \dots, x_{\sigma(4)})) \\ &= p(x_{\tau(\sigma(1))}, \dots, x_{\tau(\sigma(4))}) \\ &= p(x_{(\tau \circ \sigma)(1)}, \dots, x_{(\tau \circ \sigma)(4)}) \\ &= (\tau \circ \sigma) \cdot p.\end{aligned}$$

Hence composition in  $S_4$  is compatible with its action on  $R$ . Clearly  $e \cdot p = p$ . Thus we have satisfied the axioms for a group action, as desired.

- (c) The permutations that stabilize  $x_4$  are the ones that fix 4. The subset of  $S_4$  that does this is:  $\{e, (12), (23), (13), (123), (231)\}$ . Looking at these permutations in cycle notation, clearly they are isomorphic to  $S_3$ . (They are the image of the embedding  $\iota : S_3 \hookrightarrow S_4$ .)
- (d) An element  $\sigma$  stabilizes  $x_1 + x_2$  satisfy  $x_1 + x_2 = x_{\sigma(1)} + x_{\sigma(2)}$ . Hence we must have either  $(\sigma(1), \sigma(2)) = (1, 2)$  or  $(\sigma(1), \sigma(2)) = (2, 1)$ . In the first case,  $\sigma$  fixes 1 and 2, so the possible values are  $\{e, (34)\}$ . In the second case,  $\sigma$  must permute  $(12)$ , so the possible values are  $\{3, (12), (12)(34)\}$ . Letting  $x = (12), y = (34)$ , we see that the stabilizer of  $x_1 + x_2$  is  $\{e, x, y, xy\}$  with  $x^2 = y^2 = e$ . This is clearly the (abelian) group  $\mathbb{Z}_2 \times \mathbb{Z}_2$ .
- (e) If  $\sigma$  stabilizes  $x_1x_2 + x_3x_4$ , then

$$x_{\sigma(1)}x_{\sigma(2)} + x_{\sigma(3)}x_{\sigma(4)} = x_1x_2 + x_3x_4.$$

Then there are two cases:

$$\begin{aligned}(\sigma(1), \sigma(2)) &\in \{(1, 2), (2, 1)\} \wedge (\sigma(3), \sigma(4)) \in \{(3, 4), (4, 3)\}, \\ (\sigma(1), \sigma(2)) &\in \{(3, 4), (4, 3)\} \wedge (\sigma(3), \sigma(4)) \in \{(1, 2), (2, 1)\}\end{aligned}$$

The first case has solutions  $\sigma \in \{e, (12), (34), (12)(34)\}$ . And the second case has solutions  $\sigma \in \{(13)(24), (1324), (1423), (14)(23)\}$ .

To see that these two sets combine to form  $D_8$ , map  $(12) \mapsto s$  and  $r \mapsto (1324)$ . Then we have:

$$\begin{aligned}e &\mapsto e, \quad (1324) \mapsto r, \quad (12)(34) \mapsto r^2, \quad (1423) \mapsto r^3, \\ (12) &\mapsto s, \quad (13)(24) \mapsto sr, \quad (34) \mapsto sr^2, \quad (14)(23) \mapsto sr^3.\end{aligned}$$

It can be checked that this is an isomorphism. Hence the stabilizer of  $x_1x_2 + x_3x_4$  is indeed isomorphic to  $D_8$ .

- (f) Again, for the map to stabilize, we must have either  $x_1 + x_2 = x_{\sigma(1)} + x_{\sigma(2)}$  or

$x_1 + x_2 = x_{\sigma(3)} + x_{\sigma(4)}$ . The same equations follow for  $x_3 + x_4$ . So there are two cases:

$$\begin{aligned} (\sigma(1), \sigma(2)) &\in \{(1, 2), (2, 1)\} \wedge (\sigma(3), \sigma(4)) \in \{(3, 4), (4, 3)\}, \\ (\sigma(1), \sigma(2)) &\in \{(3, 4), (4, 3)\} \wedge (\sigma(3), \sigma(4)) \in \{(1, 2), (2, 1)\} \end{aligned}$$

As we've seen in part (e), this subset is isomorphic to  $D_8$ .

Exercises 16, 17, pp. 65-66.

16. (a) Since  $G$  is finite there can only be a finite amount of subgroups. In particular, there are only a finite amount of subgroups  $\{H_i\}_{i=1}^n$  containing  $H$ . Then any chain  $H \leq H_{i_1} \leq H_{i_2} \leq \dots \leq H_{i_k} \leq G$  is finite, and we may prescribe  $H_{i_k}$  as the maximal subgroup containing  $H$ .
- (b) Suppose  $\langle r \rangle \leq K$ . Then  $|\langle r \rangle| \leq |K|$  while  $|K| \mid |G|$ . But  $\langle r \rangle$  has order  $n$  and  $G$  has order  $2n$ . Hence  $|K|$  can only be  $n$ , in which case  $H = K$ , or  $2n$ , in which case  $K = G$ . This is exactly the definition of  $H$  being maximal, as desired.
- (c) The order of  $x^p$  is  $n/p$ , so  $|\langle x^p \rangle| = n/p$ . If  $K$  contains  $\langle x^p \rangle$ , then  $n/p \leq |K| \Rightarrow n/|K| \leq p$  while  $|K| \mid n \Rightarrow n/|K| = a$  for some  $a \in \mathbb{Z}$ . But the only possible factors of  $p$  are 1 and  $p$ , and  $|K| \neq n$ , so we must have  $a = p$ . Because their orders are equal and one is a subset of the other,  $K = \langle x^p \rangle$ . Hence  $\langle x^p \rangle$  is maximal.
17. (a) The chain  $\mathcal{C}$  is a set of subgroups  $\{H_i\}_{i \in \mathcal{I}}$  on a total order  $\mathcal{I}$  such that  $H_i \leq H_j$  for all  $i \leq j$ .

We first show that if  $x, y \in \bigcup_{i \in \mathcal{I}} H_i = H$ , then

$$xy \in \bigcup_{i \in \mathcal{I}} H_i = H.$$

Since  $x \in H$ , we have  $x \in H_i$  for some  $i \in \mathcal{I}$ . Similarly  $y \in H_j$  for some  $j \in \mathcal{I}$ . Furthermore,  $\mathcal{I}$  is a total order so either  $i \leq j$  or  $i \geq j$ . Without loss of generality assume that  $i \leq j$ , since we could just swap the labels if instead  $j \leq i$ . Then  $H_i \leq H_j$ , so  $x \in H_i \leq H_j$  and  $y \in H_j$  imply  $xy \in H_j \leq H$ .

The other subgroup axioms are straightforward:  $e \in H$  since every  $H_i$  is a subgroup. For any  $x \in H$ ,  $\exists i, x \in H_i \Rightarrow x^{-1} \in H_i \leq H$ .

Hence  $H$  is a subgroup of  $G$ .

- (b) Assume for the sake of contradiction that  $H$  is *not* a proper subgroup, i.e.  $H = G$ . Then each  $g_i$  must lie in some  $H_{\alpha_i}$ . There are only finite  $g_i$ , therefore we can compute the finite maximum  $\max(\alpha_i) = \alpha_j$  for some fixed  $j$ . Then  $H_{\alpha_j}$  is both in

$\mathcal{C}$  and contains each  $g_i$ . Then  $\langle g_1, \dots, g_n \rangle \subset H_{\alpha_j}$ . But  $\langle g_1, \dots, g_n \rangle = G$ ! So  $H_{\alpha_j}$  is not proper, contradicting our assumptions about  $\mathcal{C}$ .

- (c) Part (b) shows that for any chain  $\mathcal{C}$ , the union of all subgroups in the chain  $H$  is an upper bound on  $\mathcal{C}$  that is proper. In other words,  $H \in \mathcal{S}$ , and hence we may apply Zorn's lemma to deduce that  $\mathcal{S}$  contains at least one maximal element. This concludes the proof.

Exercises 1, 18, 24, 40, 41 pp. 85-89.

**Problem 1.** Let  $\varphi : G \rightarrow H$  be a homomorphism and let  $E$  be a subgroup of  $H$ . Prove that  $\varphi^{-1}(E) \leq G$ . If  $E \trianglelefteq H$ , then  $\varphi^{-1}(E) \trianglelefteq G$ . Deduce that  $\ker \varphi \trianglelefteq G$ .

*Proof.* Part 1: We show that  $\varphi^{-1}(E)$  is a subgroup with the subgroup property. Suppose  $g, h \in \varphi^{-1}(E)$ . Then by definition  $\varphi(g), \varphi(h) \in E$ , and since  $E$  is a subgroup, we have in particular  $\varphi(h)^{-1} = \varphi(h^{-1}) \in E$ . Thus  $\varphi(g)\varphi(h^{-1}) = \varphi(gh^{-1}) \in E$ . By definition this means  $gh^{-1} \in \varphi^{-1}(E)$ , which proves that  $\varphi^{-1}(E)$  is indeed a subgroup.

Part 2: Now suppose that  $E \trianglelefteq H$ . To show that  $\varphi^{-1}(E) \trianglelefteq G$ , we have to prove  $gng^{-1} \in \varphi^{-1}(E)$  for all  $n \in \varphi^{-1}(E)$  and  $g \in G$ . By definition,  $\varphi(n) \in E$ , and also the normality of  $E$  implies  $\varphi(g)\varphi(n)\varphi(g)^{-1} \in E$ . Applying the properties of homomorphisms, we have can deduce  $\varphi(gng^{-1}) \in E$ , and so  $gng^{-1} \in \varphi^{-1}(E)$ . Hence  $\varphi^{-1}(E)$  is normal in  $G$ .

Part 3: Immediately from part 2, since  $\{e\} \trianglelefteq H$  and by definition  $\ker \varphi = \varphi^{-1}(e)$ , we have  $\ker \varphi \trianglelefteq G$ .  $\square$

**Problem 18.** Let  $G$  be the quasidihedral group of order 16:

$$G = \langle \sigma\tau \mid \sigma^8 = \tau^2 = 1, \sigma\tau = \tau\sigma^3 \rangle$$

and let  $\overline{G} = G/\langle \sigma^4 \rangle$  be the quotient of  $G$  by the subgroup generated by  $\sigma^4$  (this subgroup is the center of  $G$ , hence is normal).

*Proof.* We do each part in this proof:

- (a) The subgroup  $\langle \sigma^4 \rangle$  has order 2, so Lagrange's theorem implies that  $|\overline{G}| = |G|/|\langle \sigma^4 \rangle| = 16/2 = 8$ .
- (b) We have  $\overline{G} = \{\overline{\tau}^a \overline{\sigma}^b \mid a = 0, 1, b = 0, 1, 2, 3\}$ . This gives 8 elements which cannot be further reduced with the rules  $\overline{\tau}^2 = \overline{\sigma}^4 = 1$  and  $\sigma\tau = \tau\sigma^3$ . Hence these must *exactly* be the elements of  $\overline{G}$ .

(c) Let  $x = \bar{\tau}$  and  $y = \bar{\sigma}$ . The orders can be computed pretty easily:

$$\begin{aligned}
 x^0 y^0 &= 1 \Rightarrow |x^0 y^0| = 1 \\
 x^1 y^0 &= 1 \Rightarrow |x^1 y^0| = |x| = 2 \\
 x^0 y^1 &= 1 \Rightarrow |x^0 y^1| = |y| = 4 \\
 x^1 y^1 &= 1 \Rightarrow xyxy = xxy^3y = 1 \Rightarrow |x^1 y^1| = 2 \\
 x^0 y^2 &= 1 \Rightarrow |x^0 y^2| = 2 \\
 x^1 y^2 &= 1 \Rightarrow |x^1 y^2| = 2 \\
 x^0 y^3 &= 1 \Rightarrow |x^0 y^3| = |y^{-1}| = 4 \\
 x^1 y^3 &= 1 \Rightarrow |x^1 y^3| = |xy^{-1}| = 2
 \end{aligned}$$

(d) Again let  $x = \bar{\tau}$  and  $y = \bar{\sigma}$ . Then

$$\begin{aligned}
 yx &= xy^3 \\
 xy^{-2}x &= xy^2x = xy yx = xyxy^3 = xxy^3y^3 = y^2 \\
 x^{-1}y^{-1}xy &= xy^3xy = xy^2yxy = xy^2xy^3y = xy^2x = y^2
 \end{aligned}$$

(e) Consider the map  $\varphi$  such that  $x \mapsto s$  and  $y \mapsto r$ . Then, looking at (c), clearly  $x$  and  $y$  interact in the same way  $s$  and  $r$  do hence  $\bar{G} \cong D_8$ .

□

**Problem 24.** Prove that if  $N \trianglelefteq G$  and  $H$  is any subgroup of  $G$  then  $N \cap H \trianglelefteq H$ .

*Proof.* Suppose  $N \trianglelefteq G$  and  $H \leq G$ . Let  $n \in H \cap N$  and  $h \in H$ . So  $h \in G$  and  $n \in N$ , for which we deduce that  $hnh^{-1} \in N$ . Also  $n \in H$ , and so  $hnh^{-1} \in H$ . Hence  $hnh^{-1} \in N \cap H$ . Hence  $N \cap H \trianglelefteq H$ .

□

**Problem 40.** Let  $G$  be a group, let  $N$  be a normal subgroup of  $G$  and let  $\bar{G} = G/N$ . Prove that  $\bar{x}$  and  $\bar{y}$  commute in  $\bar{G}$  if and only if  $x^{-1}y^{-1}xy \in N$ .

*Proof.* ( $\Rightarrow$ ): If  $x^{-1}y^{-1}xy \in N$ , then  $(x^{-1}y^{-1}xy)N = N \Rightarrow xyN = Nyx$ . But  $N$  is normal, so we can swap the left and right cosets. Thus  $xNyN = xyN = Nyx = yxN = yN xN$ , as desired.

( $\Leftarrow$ ): If  $xNyN = yNxN$ , then we can just run the argument backwards:

$$xyN = yxN \Rightarrow x^{-1}y^{-1}xyN = N \Rightarrow x^{-1}y^{-1}xy \in N.$$

□

**Problem 41.** Let  $G$  be a group. Prove that  $N = \langle x^{-1}y^{-1}xy \mid x, y \in G \rangle$  is a normal subgroup of  $G$  and  $G/N$  is abelian.

*Proof.*  $N$  is a normal subgroup of  $G$ : Let  $\varphi_g(n) = g^{-1}ng$ . Note that conjugation by  $g$  is a homomorphism. Let  $n \in N$ , which will have the form  $a_1^{\epsilon_1}a_2^{\epsilon_2} \dots a_n^{\epsilon_n}$ , where each  $a_i = x^{-1}y^{-1}xy$  for some  $x, y \in G$  and  $\epsilon_i = \pm 1$ . Now we want to show that  $g^{-1}ng = \varphi_g(n) \in N$  for any  $g \in G$ . Since  $\varphi_g$  is a homomorphism, we have

$$\varphi_g(n) = \varphi_g(a_1)^{\epsilon_1} \varphi_g(a_2)^{\epsilon_2} \dots \varphi_g(a_n)^{\epsilon_n}.$$

Because  $N$  is a subgroup, it suffices now to prove that each  $\varphi_g(a_i) \in N$ . We have

$$\varphi_g(a_i) = \varphi_g(x^{-1}y^{-1}xy) = \varphi_g(x)^{-1}\varphi_g(y)^{-1}\varphi_g(x)\varphi_g(y).$$

The LHS is of the form  $x'^{-1}y'^{-1}x'y'$  for  $x' = \varphi_g(x)$  and  $y' = \varphi_g(y)$ , so it must be in  $N$ . Hence  $\varphi_g(a_i) \in N$ . By extension,  $\varphi_g(n) \in N$ . Therefore  $N$  is normal.  $\square$

*Proof.*  $N$  is abelian: By Exercise 40 we have that  $\bar{x}$  and  $\bar{y}$  commute in  $G/N$  if and only if  $x^{-1}y^{-1}xy \in N$ . But this implies that  $\overline{x^{-1}y^{-1}xy} = 1$  in  $G/N$ . Rearranging gives  $\bar{x}\bar{y} = \bar{y}\bar{x}$ , as desired.  $\square$

Exercise 4, pp. 111.

**Problem 4.** Prove that  $S_n = \langle (12), (123 \dots n) \rangle$  for all  $n \geq 2$ .

*Proof.* It suffices to show that every transposition can be generated from  $x = (12)$  and  $y = (123 \dots n)$ . Indeed, direct calculation shows that we can obtain transpositions of the form  $(i, i+1)$  by conjugating  $y^{i-1}xy^{1-i}$ :

$$\begin{aligned} i &\mapsto 1 \mapsto 2 \mapsto i+1 \\ i+1 &\mapsto 2 \mapsto 1 \mapsto i \\ j &\mapsto j-i+1 \notin \{1, 2\} \mapsto j \quad \forall j \neq i, i+1 \end{aligned}$$

Next, transpositions of the form  $1i$  can be generated recursively using  $(1, i+1) = (1i)(i, i+1)(1i)$ , starting with the base case  $(12)$ . Finally, general transpositions of the form  $(ij)$  can be computed with  $(1i)(1j)(1i)$ , which clearly maps  $i \mapsto j$  and  $j \mapsto i$ . And with all the transpositions, we're done.  $\square$