

Partial Differential Equations

Qualifying Examination

April 10, 1989

Do any 7 problems.

1. Do any three of the following:

(i) State the definition of well-posed problem.

(ii) Write out an example to show the Cauchy problem for heat equation

$$\begin{aligned}u_t &= u_{xx} & x \in \mathbb{R}^1, \quad 0 < t < \infty \\u(0, x) &= \phi(x) \in C^1(\mathbb{R}^1)\end{aligned}$$

is not well-posed.

(iii) Under what additional conditions is the PDE

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2} \quad 0 < t < \infty \quad -\pi < x < \pi$$

well-posed. Explain.

(iv) Is the Neumann problem for Laplace's equation

$$\begin{aligned}\Delta u &= 0 \quad \text{in } \Omega \\ \frac{\partial u}{\partial n} &= f \quad \text{on } \partial\Omega\end{aligned}$$

well-posed where $f \in C^1(\partial\Omega)$ and $\Omega \subset \mathbb{R}^n$ is a bounded smooth domain? Explain.

2. (i) Show $\frac{1}{4\pi r}$ is a weak solution to equation $\Delta u = \delta$, the Dirac delta function, on \mathbb{R}^3 .

(ii) Give two equivalent definitions of Sobolev space $W^{m,p}(\Omega)$.

3. Let $X = \{u \in C^2(\Omega) \cap L^2(\Omega) \mid \frac{\partial u}{\partial n}|_{\partial\Omega} = 0, \int_{\Omega} u = 0\}$. Show that $-\Delta$ is a positive definite operator on X ; $(-\Delta u, u) \geq a \|u\|_{L^2(\Omega)}^2$ for some positive constant a .

4. Solve $\Delta u(x, y) = 0$ on the strip $0 < y < \pi$ with the boundary value

$$u(x, \pi) = u(x, 0) = \begin{cases} 0 & x < 0 \\ 1 & 0 \leq x < \ln 2 \\ 0 & \ln 2 < x \end{cases}$$

(See Fig. 1)

5. A traffic flow can be modeled by equation

$$\begin{cases} \frac{\partial \rho}{\partial t} + u_m \left(1 - \frac{2\rho}{\rho_m}\right) \frac{\partial \rho}{\partial x} = 0 \\ \rho(0, x) = h(x) \end{cases}$$

where $\rho = \rho(t, x)$ is the traffic density,

u_m = maximum permitted speed,

ρ_m = maximum possible density of cars,

$h(x)$ = initial density.

- (i) Find the density $\rho(t, x)$
- (ii) Under what initial density of $h(x)$ may a catastrophe of traffic flow occur at some later time $t > 0$. Explain.

6. Solve the following equation by using the method of separation of variable

$$\begin{cases} \frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + u & 0 < x < \pi, \quad 0 < t \\ u(t, 0) = u(t, \pi) = 0 \\ u(0, x) = x(1 - x) \end{cases}$$

7. State and prove Dirichlet's principle for harmonic functions.

8. Prove the uniqueness of the solution to wave equation of mixed value problem

$$\begin{cases} u_{tt} = C^2 \Delta u & 0 < t \leq T, \quad x \in \Omega \\ \Omega \text{ is bounded domain in } \mathbb{R}^n \\ u(0, x) = f(x) \in C^1(\overline{\Omega}) \\ u_t(0, x) = g(x) \in C^1(\overline{\Omega}) \\ u(x, t)|_{[0, T] \times \partial \Omega} = 0. \end{cases}$$

9. Suppose $P_t(x, y)$ is the heat kernel on a bounded domain Ω in \mathbb{R}^n , i.e.

$$u(t, x) = \int_{\Omega} P_t(x, y) f(y) dy$$

solves equation

$$\begin{cases} u_t = \Delta u & x \in \Omega, \quad 0 < t < T \\ u(0, x) = f(x) \in C^1(\Omega) \\ u(t, x) = 0 & x \in \partial\Omega, \quad 0 \leq t \leq T. \end{cases}$$

Write out the Green's function for Ω in terms of $P_t(x, y)$.

10. The vibration of a beam with infinite length can be described by the following equation

$$\begin{cases} u_{tt} + a^2 \frac{\partial^4 u}{\partial x^4} = 0 \\ u(0, x) = f(x), \quad u_t(0, x) = ag''(x), \quad f, g'' \in L^1(\mathbb{R}^1) \end{cases}$$

where f, g'' are the initial vibration data. Use Fourier transform methods to solve this equation.

HINT: Use the Fourier Cosine Transform:

$$\begin{aligned} \sin b\xi^2 &\mapsto \sqrt{\frac{\pi}{8b}} \left(\cos \frac{x^2}{4b} - \sin \frac{x^2}{4b} \right) \\ \cos b\xi^2 &\mapsto \sqrt{\frac{\pi}{8b}} \left(\cos \frac{x^2}{4b} + \sin \frac{x^2}{4b} \right) \end{aligned}$$

11. (i) Let Ω be a bounded domain in \mathbb{R}^n . State the definition that a function $u \in L^1(\Omega)$ is weakly differentiable. (First order.)
- (ii) We say a function $u \in L^1(\Omega)$ is strongly differentiable if there is a sequence $\{u_m\} \in C^\infty(\Omega)$ and a function $v \in L^1(\Omega)$ such that

$$u_m \rightarrow u, \quad \frac{\partial u_m}{\partial x_i} \rightarrow v \quad \text{in } L^1(\Omega) \quad i = 1, 2, \dots, n.$$

Show that a function u is weakly differentiable if and only if it is strongly differentiable.




Fig. 1