

Algebra Qualifying Exam

Spring 1992

All rings are assumed to have a multiplicative identity, denoted 1. The fields \mathbb{Q} , \mathbb{R} and \mathbb{C} are the fields of *rational*, *real* and *complex* numbers, respectively.

1. Prove that no group of order 120 can be a simple group.
2. Let G be a group, and let H, K be *solvable* subgroups of G with K normal. Prove that HK is a solvable subgroup of G .
3. Let G be a finite group of order greater than 3, and let \mathcal{C} be a conjugacy class of elements in G . If $|\mathcal{C}| = \frac{1}{3}|G|$, show that every element of \mathcal{C} is an element of order 3.
4. Let R be a unique factorization domain in which every prime ideal is maximal. Prove that every prime ideal is principal. (In fact, it turns out that *every* ideal is principal, but you are not asked to prove this.)
5. Let R be a ring. Prove that the following three conditions are equivalent for the left R -module M .

(i) Any increasing chain of submodules

$$M_1 \subseteq M_2 \subseteq \dots,$$

(ii) Any submodule of M is finitely generated.

(iii) Any family of submodules of M has a maximal member with respect to inclusion.

6. Let F be a field and let R be the ring

$$R = \left\{ \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} \mid a, b \in F \right\}.$$

Define the left R -modules

$$M_1 = \left\{ \begin{bmatrix} a \\ 0 \end{bmatrix} \mid a \in F \right\}, M_2 = \left\{ \begin{bmatrix} 0 \\ b \end{bmatrix} \mid b \in F \right\}.$$

(The module action is matrix multiplication.) Prove that M_1 and M_2 are **not** isomorphic as left R -modules.

7. Let V be a vector space of dimension n , and let $1_V \neq T : V \rightarrow V$ be a linear transformation of V . If $\text{im}(T - 1_V) \subseteq \ker(T - 1_V)$, compute the minimal and characteristic polynomials of T on V .
8. Let F_q be the finite field of q elements, and let K be an extension of F_q , of degree n .
 - (a) Prove that the map $\tau_q : K \rightarrow K$, $\tau_q(x) = x^q$ is an automorphism of K , and that F_q is precisely the subfield of fixed elements of τ_q .
 - (b) Compute $\text{Gal}(K/F_q)$.
9. Let $f(x) = x^5 - 2 \in \mathbb{Q}[x]$.
 - (a) Construct a splitting field $K \supseteq \mathbb{Q}$ for $f(x)$ over \mathbb{Q} .
 - (b) If $G = \text{Gal}(K/\mathbb{Q})$, prove that no nonidentity element of G can fix two roots of $f(x)$.