

# Try Bayesian Survival Analysis using the JACC study data (Bayesian Survival Analysis presentations)

Chaochen Wang / Aichi Medical University

2020-01-12

## Parametric Survival Analysis Model (in a Bayesian way)

- We chose Weibull model to describe the survival data we have.
- Weibull model is both a proportional hazard model and an accelerated failure time (AFT) model. The relationship between the acceleration factor and log hazard ratio:

$$\exp(\beta_{PH}^T) = \exp(\kappa \beta_{AFT}^T)$$

Weibull Survival function in a proportional hazard way expression is:

$$S(t|\mathbf{x}) = \exp\{-\lambda t^\kappa e^{\beta_{PH}^T \mathbf{x}}\}$$

Weibull Survival function in an AFT way expression is:

$$S(t|\mathbf{x}) = \exp\{-\lambda t^\kappa e^{\kappa \beta_{AFT}^T \mathbf{x}}\}$$

Acceleration factor is the factor by which time to failure/event is accelerated for  $x_1 = 1$  compared with  $x_1 = 0$

An example of a crude Weibull model using the JAGS program in R:

```
jacc.weibull.model <- function() {  
  for(i in 1:39386){# loop through the subjects  
    # sAge[i] <- (Age[i] - mean(Age[])) / sd(Age[])  
    is.censored[i] ~ dinterval(t[i], c[i])  
    t[i] ~ dweib(shape, lambda[i])  
    lambda[i] <- exp(-mu[i] * shape)  
    mu[i] <- beta[1] + beta[2]*equals(Mlkgre[i], 2) +  
      beta[3]*equals(Mlkgre[i], 3) + beta[4] * equals(Mlkgre[i], 4) +  
      beta[5] * equals(Mlkgre[i], 5) # +  
      # beta[6] * sAge[i]  
  
    ### calculate log-likelihoods  
    # y[i] <- ifelse(is.censored[i], c[i], t[i])  
    # loglik[i] <- log(ifelse(is.censored[i],  
    #                          exp(-lambda[i] * (y[i] ^ shape)),  
    #                          shape * lambda[i] * (y[i] ^ (shape - 1)) * exp(-lambda[i] * (y[i] ^ shape))))  
  }  
  
  ## priors for betas  
  for(j in 1:5){  
    beta[j] ~ dnorm(0, 0.001)  
  }  
  
  ### prior for shape
```

```

shape ~ dgamma(.001, .001)

### Generated values
AFT[2] <- exp(beta[2])
HR[2] <- exp(shape * beta[2])
p.crit[2] <- step(1 - HR[2])

AFT[3] <- exp(beta[3])
HR[3] <- exp(shape * beta[3])
p.crit[3] <- step(1 - HR[3])

AFT[4] <- exp(beta[4])
HR[4] <- exp(shape * beta[4])
p.crit[4] <- step(1 - HR[4])

AFT[5] <- exp(beta[5])
HR[5] <- exp(shape * beta[5])
p.crit[5] <- step(1 - HR[5])
}

```

You can use the following codes to run the above model (careful about the cost of time):

```

library(R2jags) #connect to jags sampling engine in R
jagsfit <- jags.parallel(data = JACCdata,
                        parameters.to.save = c("beta[5]", "HR[5]", "AFT[5]"),
                        #add more variables if you want to check more
                        n.iter = 10000,
                        n.burnin = 2500,
                        n.chains = 3,
                        model.file = jacc.weibull.model)

```

The output of the above crude Weibull model:

```

# Inference for Bugs model at "jacc.weibull.model", fit using jags,
# 3 chains, each with 10000 iterations (first 2500 discarded), n.thin = 7
# n.sims = 3213 iterations saved. Sample size per chain = 1071
# 1. Empirical mean and standard deviation for each variable,
# plus standard error of the mean:
#
#           Mean          SD   Naive SE Time-series SE
# AFT[2]      0.922303  0.064441 0.00113685      0.00420403
# AFT[3]      0.834723  0.051485 0.00090830      0.00362857
# AFT[4]      0.848329  0.052439 0.00092513      0.00401102
# AFT[5]      0.928095  0.041834 0.00073803      0.00336613
# HR[2]       0.890533  0.090586 0.00159811      0.00604246
# HR[3]       0.769913  0.069116 0.00121934      0.00486858

```

```

# HR[4]          0.788205  0.070624  0.00124594    0.00531505
# HR[5]          0.897882  0.058477  0.00103164    0.00468654
# p.crit[2]      0.871460  0.334743  0.00590548    0.01814848
# p.crit[3]      0.996265  0.061008  0.00107630    0.00214585
# p.crit[4]      0.998133  0.043180  0.00076177    0.00097572
# p.crit[5]      0.953626  0.210327  0.00371056    0.02050164
# 2. Quantiles for each variable:
#              2.5%      25%      50%      75%      97.5%
# AFT[2]        0.807308  0.876649  0.918049  0.96495  1.05225
# AFT[3]        0.738423  0.799925  0.832078  0.86719  0.94214
# AFT[4]        0.749645  0.810952  0.847050  0.88448  0.95291
# AFT[5]        0.847796  0.899999  0.927850  0.95594  1.01147
# HR[2]         0.733186  0.825541  0.883106  0.94896  1.07725
# HR[3]         0.644053  0.721892  0.765778  0.81289  0.91643
# HR[4]         0.658787  0.737562  0.785538  0.83614  0.93282
# HR[5]         0.788623  0.857453  0.896668  0.93613  1.01665
# ...

```

*Check model fitting:*




Figure 1: Traceplot for AFT, and HR of daily milk drinkers from crude model (men).




Figure 2: Not overlapping traceplot indicating that model is not converged.




Figure 3: Autocorrelation figures for AFT, and HR of daily milk drinkers from crude model (men).




Figure 4: (bad example) Highly correlated posterior samples indicating that the samples are not independent.

Table 1. Estimated (crude) hazard ratios and acceleration factors for the association of milk intake and risk of stroke mortality using Bayesian MCMC methods in Men of the JACC study data (n = 39,386).

Milk intake	Hazard ratio (HR)					Acceleration factor (AF)			
	Median	Mean (SD)	95% CrI	MCSE	Probability	Median	Mean (SD)	95% CrI	MCSE
Never	-	-	-	-	-	-	-	-	-
1-2 t/Mon	0.88	0.89 (0.09)	(0.73, 1.08)	0.0016	87.15%	0.92	0.92 (0.06)	(0.81, 1.06)	0.0011
1-2 t/Week	0.77	0.77 (0.07)	(0.64, 0.92)	0.0012	99.63%	0.83	0.83 (0.05)	(0.74, 0.94)	0.0009
3-4 t/Week	0.79	0.79 (0.07)	(0.66, 0.93)	0.0012	99.81%	0.85	0.85 (0.05)	(0.75, 0.95)	0.0009
Daily	0.89	0.90 (0.06)	(0.79, 1.02)	0.0010	95.36%	0.93	0.93 (0.04)	(0.85, 1.01)	0.0007

*Note:*

Abbreviations: SD, standard deviation; CrI, credible interval; MCSE, Monte Carlo Standard Error; Probability indicates that p for HR smaller than 1.




Figure 5: Crude model HR posterior density plots.

Table 2. Estimated (age-adjusted) hazard ratios and acceleration factors for the association of milk intake and risk of stroke mortality using Bayesian MCMC methods in Men of the JACC study data (n = 39,386).

Milk intake	Hazard ratio (HR)					Acceleration factor (AF)			
	Median	Mean (SD)	95% CrI	MCSE	Probability	Median	Mean (SD)	95% CrI	MCSE
Never	-	-	-	-	-	-	-	-	-
1-2 t/Mon	0.98	0.98 (0.11)	(0.78, 1.19)	0.0018	57.86%	0.98	0.99 (0.06)	(0.87, 1.11)	0.0011
1-2 t/Week	0.84	0.84 (0.08)	(0.69, 0.99)	0.0014	97.79%	0.90	0.90 (0.05)	(0.80, 0.99)	0.0008
3-4 t/Week	0.86	0.86 (0.08)	(0.71, 1.02)	0.0014	95.92%	0.91	0.91 (0.05)	(0.82, 1.01)	0.0009
Daily	0.75	0.75 (0.05)	(0.67, 0.85)	0.0009	100.00%	0.85	0.85 (0.03)	(0.79, 0.91)	0.0006

*Note:*

Abbreviations: SD, standard deviation; CrI, credible interval; MCSE, Monte Carlo Standard Error; Probability indicates the p for HR smaller than 1.




Figure 6: Age-adjusted HR posterior density plots.