

RESEARCH

Relationships between food groups, eating time slots and diabetes status in adults from the UK National Diet and Nutrition Survey (2008–2017)

Chaochen Wang¹, Suzana Almoosawi² and Luigi Palla^{3,4,5*}

*Correspondence:

Luigi.Palla@uniroma1.it

³Department of Public Health and Infectious Diseases, University of Rome La Sapienza, Piazzale Aldo Moro 5, Rome, 00185 Italy
Full list of author information is available at the end of the article

Abstract

Background: Time of eating has been shown to be associated with diabetes and obesity but less is known about less healthy foods and specific time of their intake during the day (24 hours). We aimed to identify potential relationships between foods and eating time, and see whether these associations may vary by diabetes ~~statuses~~status.

Method: The National Diet and Nutrition Survey (NDNS) including 6802 adults (age ≥ 19 years old) collected 749,026 food recordings by a 4-day-diary. The contingency table cross-classifying 60 food groups with 7 pre-defined eating time slots (6-9am, 9am-12pm, 12-2pm, 2-5pm, 8-10pm, 10pm-6am) ~~were~~was analyzed by Correspondence Analysis (CA). ~~Biplots~~CA biplots displaying the associations were generated for all adults and separately by diabetes ~~statuses~~status (self-reported, pre-diabetes, undiagnosed-diabetes, and non-diabetics) to visually explore the associations between food groups and time of eating across diabetes strata. For selected food groups, odds ratios (OR, 99% confidence intervals, CI) were derived of consuming unhealthy foods at evening/night (8pm-6am) vs. earlier time in the day, by logistic regression models with generalized estimating equations.

Results: The biplots suggested positive associations between evening/night and consumption of puddings, regular soft drinks, sugar confectioneries, chocolates, spirits, beers, ice cream, biscuits, and crisps for all adults in the UK. The OR (99% CIs) of consuming these foods at evening/night were respectively 1.38 (1.03, 1.86), 1.74 (1.47, 2.06), 1.92 (1.38, 2.69), 3.19 (2.69, 3.79), 11.13 (8.37, 14.80), 7.19 (5.87, 8.82), 2.38 (1.79, 3.15), 1.91 (1.67, 2.16), 1.55 (1.27, 1.88) vs. earlier time in the day. Stratified biplots found that sweetened beverages, sugar-confectioneries appeared more strongly associated with evening/night among un-diagnosed diabetics.

Conclusions: Foods consumed in the evening/night time tend to be highly processed, easily accessible, and rich in added sugar or saturated fat. Individuals with undiagnosed diabetes are more likely to consume unhealthy foods at night. Further longitudinal studies are required to ascertain the causal direction of the association between late-eating and diabetes status.

Keywords: chrono-nutrition; time of eating; correspondence analysis; the UK National Diet and Nutrition Survey

Background

The timing of energy intake has been shown to be associated with obesity and diabetes. [1] Specifically, eating late at night or having a late dinner was found to be related to higher risk of obesity [2, 3], hyperglycemia [4], metabolic syndrome [5], diabetes [6], and poorer glycemic control among diabetics [7]. However, the relationship between food choice and the time of food consumption during the day is left largely unknown. Shiftworkers have an increased risk of obesity [8, 9], and diabetes [10], ~~probably~~ possibly due to limited availability of healthy food choice during their night shifts [11, 8]. Identifying those unhealthy foods that might be chosen during late night time would be helpful when guiding people to change their eating habit for the purpose of either weight ~~losing or maintaining~~ loss or glycemic control. Dietary diary recordings from national surveys can provide detailed food choice data for exploration of the relationships between food groups and their time of consumption in the general population.

In this study, we aimed to describe the relationship between food groups and the time of day when they were consumed, and how such relationships may vary by ~~statuses~~ status of type 2 diabetes using the data published by the Rolling Programme of the UK National Diet and Nutrition Survey from 2008 to 2017 as this survey includes diet diaries providing detailed information on the time of day of food intake.

Methods

6802 adults (2810 men and 3992 women) and 749026 food recordings collected by the UK National Diet and Nutrition Survey Rolling Programme (NDNS RP 2008-17) were analyzed in the current study [12]. The sample was randomly drawn from a list of all addresses in the UK, clustered into postcode sectors. Details of the rationale, design and methods of the survey can be found in the ~~previous~~ previously published official study reports [13, 14]. Time of the day was categorized into 7 slots: 6-9 am, 9-12 noon, 12-2 pm, 2-5 pm, 5-8 pm, and 10 pm - 6 am. Foods recorded were classified into 60 standard food groups with 1 to 10 subgroups each: the details are given in Appendix R of the NDNS official report [15]. We focused on the 60 standard food groups in the current analysis. Diabetes ~~statuses were~~ status was defined as: 1) healthy if fasting glucose was lower than 6.10 (mmol/L), hemoglobin A1c (HbA1c) ~~were was~~ less than 6.5 (%), and without self-reported diabetes ~~or under treatment of and treatment for~~ diabetes (n = 2626); 2) pre-diabetic if fasting glucose was lower between 6.10 and 6.99 (inclusive) but without self-reported diabetes ~~or under treatment of and without treatment for~~ diabetes (n = 133); 3) undiagnosed diabetic if either fasting glucose was higher or equal to 7.00 (mmol/L) or HbA1c higher or equal to 6.5 (%) but without self-reported diabetes ~~or under treatment of and treatment for~~ diabetes (n = 99); 4) diabetic if participant had self-reported diabetes or under treatment of diabetes (n = 227). Consequently, the was also a large number of adults whose diabetes statuses could not be confirmed was 3717 (3717 adults of whom 1519 men, and 2198 women) and were kept only whose diabetes status did not fall in one of above categories and could not thus be confirmed; these were retained in the whole sample (unstratified) analyses. In addition, the National Statistics Socio-economic Classification [16] were applied in the survey and therefore

accordingly, the socio-economic ~~classification~~ status for the individuals' household were defined with 8 categories.

Correspondence analysis (CA) [17, 18, 19] was used as a tool for data mining, visualization and hypotheses generation using half of the randomly selected NDNS diary entries data. Specifically, the contingency table generated by cross-tabulating 60 food groups and 7 time slots were analyzed by CA. Through CA, the 60 categories of standard foods and the 7 time slots were projected on ~~two dimensions~~ biplots, i.e., onto two dimensional plots that could jointly contain large percentage of the χ^2 deviation (or inertia) of the contingency table. Biplots that graphically show the association between time of day and food groups were derived for all adults and separately according to their diabetes ~~statuses~~ status. To account for the hierarchical structure of the data (food recorded by the same individuals who lived within the same area/sampling units) and to calculate population average odds ratios (OR), logistic regression models with generalized estimating equations (GEE) were subsequently used to test the associations that were first suggested by visual inspection of biplots generated by CA, using the remaining half of the diary entries data. The marginal ORs and their 99% confidence intervals (CI) were derived of consuming unhealthy food groups (selected by CA, ~~later in the~~ later in the day (8 pm - 6 am, i.e. in the evening and night) compared to earlier in the day (~~between morning and in the morning or~~ afternoon). CA and biplots were conducted and generated by the following packages under R environment [20]: FactoMineR, factoextra, ggplot2, ggrepel [21, 22, 23, 24]. Logistic regression models with GEE were performed with SAS procedure GENMOD [25] adjusted for age, sex, and socio-economic levels, which were deemed the main potential confounders of the associations.

Results

The dataset consisted of 2810 (41.3%) men and 3992 (58.7%) women aged older than or equal to 19 years old with the mean age of 49.9 years (standard deviation, SD = 17.6). Of these individuals 22.6 % ~~of them~~ were current smokers, 24.3 % were past smokers. The average body mass index (BMI) was 27.7 kg/m² (SD = 5.41). Among the food recordings collected (n = 749026), 56.9% were recorded during traditional breakfast (6 am - 9 am: 14.3%), lunch (12 noon - 2 pm: 18.5%), or dinner (5 pm - 8 pm: 24.1%) time slots. Table 1. shows the top 37 food groups that contributed to 90% of the total calories consumed by adults in NDNS RP. These food groups accounted for 478028 of the total diary entries (63.8 %). The random process ~~split~~ split the whole set of food recordings into a hypothesis generating dataset of 374682 and a testing dataset of ~~374344~~ 374344 entries.

Figure 1-5 present the CA biplots that visually summarize the associations between 60 food groups and the time of their consumption in the entire sample and then stratifying by their diabetes ~~statuses~~ status. In Figure 1, the horizontal axis explains 68.9 % of the association structure (inertia) between food and time while the ~~vertical~~ vertical axis reflects 15.3 % of the same relationship. Therefore, a total of 84.2 % of the inertia between food and time were captured in this figure which shows a visual summary of how those two categorical variables are related. Specifically, time slots later than 8 pm are shown in the upper side of the plot closer to ~~foods contained~~ alcoholic products (beers and spirits) or highly processed/energy-condense foods (sugar confectioneries, chocolates, biscuits, regular softdrinks, ice

cream, crisps); ~~time~~ times earlier than noon appear in the left hand side together with typical breakfast foods (cereals, milk, bread, etc.).

To visualize the potentially different associational patterns between food groups choice and time slots according to diabetes status, Figure 2-5 display the CA biplots in subsets of the data. Depending on different diabetes status, these biplots explained ~~between~~ 76.3% ~~to and~~ 84.1% of the inertia in the data. Similarly to the biplot created from the total sample (Figure 1), later time in the day (8 pm and later) are shown in the upper side of each figure and suggested an association with the alcoholic beverages and highly processed or ~~energy-condense~~ energy-dense food groups. Additionally, some food groups and time slots also flagged up associations potentially different by diabetes ~~statuses~~ status. For example, puddings seemed to be ~~shown more~~ closer to later time in the day among undiagnosed diabetics (Figure 4) while for diagnosed diabetic patients (Figure 3) they were closer to traditional dinner time (5 pm to 8 pm) or earlier in the day. Furthermore, sugar confectioneries/chocolates/biscuits/regular soft drinks appeared to be associated with later time in the day (8 pm or later) more strongly among undiagnosed diabetics (Figure 4) than the other participants.

Based on the findings suggested from Figure 1-5, we decided to focus on puddings, regular soft drinks, confectioneries, chocolates, spirits, beers, ice cream, biscuits, crisps as these foods either showed a particularly strong association with time of the day or a different pattern of association across different strata of the survey sample; hence, we tested the following null hypotheses using logistic regression models (adjusted for age, sex, and socio-economic levels) with GEE: that the odds of consuming each selected food at later time of the day (8 pm - 6 am) is the same compared to earlier in the day; and the associations of the above-mentioned food groups and time slots are the same among participants with different diabetes ~~statuses~~ status (i.e. no interaction between the time of food intake and diabetes status). The results are summarized in Table 2.

The listed food groups were found to ~~be consumed during time~~ have higher odds to be consumed between 8 pm and 6 am with higher odds compared to earlier time. The OR (99% CIs) main effects of consuming these foods at evening/night were for puddings 1.38 (1.03, 1.86), for regular soft drinks 1.74 (1.47, 2.06), for sugar confectioneries 1.92 (1.38, 2.69), for chocolates 3.19 (2.69, 3.79), for spirits 11.13 (8.37, 14.80), for beers 7.19 (5.87, 8.82), for ice cream 2.38 (1.79, 3.15), for biscuits 1.91 (1.67, 2.16), for crisps 1.55 (1.27, 1.88) vs. earlier time. Opposite directions of the association for puddings were detected across diabetes ~~statuses~~ status: the ORs (99% CIs) ~~for of~~ consuming puddings at night time (8 pm or later) compared to earlier time were 1.50 (1.10, 2.07), 0.89 (0.16, 4.87), 1.81 (0.41, 7.98), and 0.58 (0.14, 2.43) for healthy, prediabetic, undiagnosed diabetic, and diabetic participants, respectively. Furthermore, undiagnosed diabetic patients were found to have particularly high odds of consuming regular soft drinks (OR: 2.72; 99% CI: 1.44, 5.14), and sugar confectioneries (OR: 13.07; 99%CI: 4.59, 37.24) during night time periods ~~than earlier time in the day and the other participants~~ compared to participants with other diabetes status.

Discussion

The present study described the potential relationships between food groups and time of their consumption in a representative sample from the NDNS RP. ~~All~~ Many unhealthy foods emerged from CA were found to be more likely to be consumed after 8 pm. These included alcoholic/sweetened beverages, chocolates and other foods rich in added sugars and saturated fats such as biscuits and ice cream. Foods chosen in the evening/night time slots tend to be highly processed and easily accessible. Specifically, undiagnosed patients might be at a higher risk of worsening their condition as they were found to have higher odds to choose a number of less healthy foods after 8 pm (sugar confectioneries, regular soft drinks) ~~compared to earlier time~~ than diabetics and non-diabetics. Those foods might need to be targeted when designing intervention to those who might be at risk of being diabetics.

These findings are concerning considering previous research that have indicated that quality of macronutrient intake in the evening is likely to influence fasting glucose levels and glycaemic response to subsequent meals in the morning. [26] More recently, a randomized controlled trial indicated that consuming carbohydrates at dinner irrespective of glycaemic index raised postprandial glucose response to breakfast producing what is known as a second meal effect [27]. Similar observation have been made by Nitta and colleagues who observed that eating sweet snacks post-dinner worsened glycaemic excursions in the evening and at subsequent breakfast ~~[28][28]~~. Added to this is evidence that suggests that the late-night dinners induce post-prandial hyperglycemia in patients with type 2 diabetes and that interventions at this eating occasions can ~~result~~ result in a profound impact on ~~postprandial~~ post-prandial glycaemia. On the balance of this evidence, targeting and improving the timing and quality of foods in evening eating occasions provides a unique opportunity to design intervention to those who might be at risk of being diabetics.

A compelling finding of our study is the observation that diabetes patients were found to be potentially controlling their choice of food groups such as avoiding puddings at night. However, higher odds of consuming alcoholic beverages and energy condensed foods such as chocolates and sugar confectioneries at night among individuals with diabetes suggests that their food choice might need further modifications.

Assessing the relationships between food groups and timing of eating by diabetes status can be considered as a first step towards identifying specific public health targets for behavior change/intervention. This is important as most current public health strategies and dietary recommendations do not provide targeted advice that takes into considerations specific eating occasions while targeted advice is more likely to result in sustainable behavioural change. Our findings are consistent with previous evidence that has found that both sweetened and alcoholic beverages are responsible for large portion of energy consumption at night in other populations [29].

~~An~~ However, an important limitation in this study is the cross-sectional study design. The inability to assess the temporal relationship between timing of food intake and diabetes status means that a cause-effect relationship between time of unhealthy food intake and diabetes status cannot be established. Hence, further prospective studies are warranted to investigate the ~~temporal~~ causal relationship

between diabetes and ~~other disease status in relation to~~ both quality and timing of eating. Moreover, the current study assumes that mis-reporting ~~occured~~ occurred equally amongst all eating occasions. This limitation has been reported by previous ~~research and is a limitation of literature as an important methodological effect of differential misreporting on epidemiological studies in~~ chrono-nutrition in order to suggest possible corrections, e.g., for differential under-reporting at different times of the day (e.g. main meals vs. snack times). [30]

Conclusions

In ~~conclusion~~ summary, our study indicates that foods consumed in the evening/night time tend to be highly processed, easily accessible, and rich in added sugar or saturated fat, whatever the diabetic status. Individuals with undiagnosed diabetes are more likely to consume specific unhealthy foods at night. The survey cross-sectional nature warrants further investigations by longitudinal cohort studies to establish the causal relation between time of eating of unhealthy foods and diabetes.

Acknowledgements

We thank the NDNS team at former MRC Elsie Widdowson Laboratory in Cambridge and NatCen Social Research in London for providing information about data collection and coding.

Funding

This work was supported by Grants-in-Aid for Young Scientists (grant number 19K20199 to C.W.) from the Japan Society for the Promotion of Science (JSPS).

Availability of data and materials

Original data used in this study can be accessed upon request to the UK Data Service (<https://www.ukdataservice.ac.uk>) for academic usage (Study Number: 6533).

Ethics approval and consent to participate

The NDNS RP, funded by Public Health England and the UK Food Standards Agency, is registered with the ISRCTN registry under study ID ISRCTN17261407 and received ethical approval from the Oxfordshire Research Ethics Committee.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

CW, SA, and LP: designed research and had primary responsibility for final content; CW and LP performed statistical analysis; and all authors: wrote the manuscript, read and approved the final manuscript.

Author details

¹Department of Public Health, Aichi Medical University, Nagakute, Aichi, Japan. ²Faculty of Medicine, School of Public Health, Imperial College London, London, UK. ³Department of Public Health and Infectious Diseases, University of Rome La Sapienza, Piazzale Aldo Moro 5, Rome, 00185 Italy. ⁴Department of Medical Statistics, London School of Hygiene & Tropical Medicine, London, UK. ⁵Department of Global Health, School of Tropical Medicine and Global Health, University of Nagasaki, Nagasaki, Japan.

References

1. Almoosawi, S., Vingeliene, S., Karagounis, L., Pot, G.: Chrono-nutrition: a review of current evidence from observational studies on global trends in time-of-day of energy intake and its association with obesity. *Proc Nutr Soc* **75**(4), 487–500 (2016)
2. Xiao, Q., Garaulet, M., Scheer, F.A.: Meal timing and obesity: Interactions with macronutrient intake and chronotype. *International Journal of Obesity* **43**(9), 1701–1711 (2019)
3. Yoshida, J., Eguchi, E., Nagaoka, K., Ito, T., Ogino, K.: Association of night eating habits with metabolic syndrome and its components: a longitudinal study. *BMC Public Health* **18**(1), 1366 (2018)
4. Nakajima, K., Suwa, K.: Association of hyperglycemia in a general Japanese population with late-night-dinner eating alone, but not breakfast skipping alone. *Journal of Diabetes & Metabolic Disorders* **14**(1), 16 (2015)
5. Kutsuma, A., Nakajima, K., Suwa, K.: Potential association between breakfast skipping and concomitant late-night-dinner eating with metabolic syndrome and proteinuria in the Japanese population. *Scientifica* **2014** (2014)
6. Mattson, M.P., Allison, D.B., Fontana, L., Harvie, M., Longo, V.D., Malaisse, W.J., Mosley, M., Notterpek, L., Ravussin, E., Scheer, F.A., *et al.*: Meal frequency and timing in health and disease. *Proceedings of the National Academy of Sciences* **111**(47), 16647–16653 (2014)

7. Sakai, R., Hashimoto, Y., Ushigome, E., Miki, A., Okamura, T., Matsugasaki, M., Fukuda, T., Majima, S., Matsumoto, S., Senmaru, T., et al.: Late-night-dinner is associated with poor glycemic control in people with type 2 diabetes: The kamogawa-dm cohort study. *Endocrine journal*, 17–0414 (2017)
8. Balieiro, L.C.T., Rossato, L.T., Waterhouse, J., Paim, S.L., Mota, M.C., Crispim, C.A.: Nutritional status and eating habits of bus drivers during the day and night. *Chronobiology international* **31**(10), 1123–1129 (2014)
9. Barbadoro, P., Santarelli, L., Croce, N., Bracci, M., Vincitorio, D., Prospero, E., Minelli, A.: Rotating shift-work as an independent risk factor for overweight italian workers: a cross-sectional study. *PLoS One* **8**(5) (2013)
10. Pan, A., Schernhammer, E.S., Sun, Q., Hu, F.B.: Rotating night shift work and risk of type 2 diabetes: two prospective cohort studies in women. *PLoS Med* **8**(12), 1001141 (2011)
11. Bonnell, E.K., Huggins, C.E., Huggins, C.T., McCaffrey, T.A., Palermo, C., Bonham, M.P.: Influences on dietary choices during day versus night shift in shift workers: a mixed methods study. *Nutrients* **9**(3), 193 (2017)
12. MRC Elsie Widdowson Laboratory, NatCen Social Research: National Diet and Nutrition Survey Years 1–8, 2008/09–2015/16 [data collection]. UK Data Service. SN: 6533 (2018). doi:10.5255/ukda-sn-6533-1
13. Bates, B., Lennox, A., Prentice, A., Bates, C.J., Page, P., Nicholson, S., Swan, G.: National Diet and Nutrition Survey: Results from Years 1, 2, 3 and 4 (combined) of the Rolling Programme (2008/2009–2011/2012): A survey carried out on behalf of Public Health England and the Food Standards Agency. Public Health England (2014)
14. Roberts, C., Steer, T., Maplethorpe, N., Cox, L., Meadows, S., Nicholson, S., Page, P., Swan, G.: National Diet and Nutrition Survey: results from years 7 and 8 (combined) of the Rolling Programme (2014/2015–2015/2016) (2018)
15. NatCen Social Research, MRC Elsie Widdowson Laboratory, Univeristy College London. Medical School.: National Diet and Nutrition Survey Years 1–8, 2008/09–2015/16. <http://doi.org/10.5255/UKDA-SN-6533-8> (2018)
16. Rose, D., Pevalin, D.J.: The National Statistics Socio-economic Classification: Origins, Development and Use, (2005)
17. Greenacre, M.: Correspondence Analysis in Practice. Chapman and Hall, New York (2017)
18. Chapman, A.N., Beh, E.J., Palla, L.: Application of correspondence analysis to graphically investigate associations between foods and eating locations. (2017)
19. Palla, L., Chapman, A., Beh, E., Pot, G., Almiron-Roig, E.: Where do adolescents eat less-healthy foods? correspondence analysis and logistic regression results from the uk national diet and nutrition survey. *Nutrients* **12**(8), 2235 (2020)
20. R Core Team: R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria (2019). R Foundation for Statistical Computing. <https://www.R-project.org/>
21. Lê, S., Josse, J., Husson, F.: FactoMineR: A package for multivariate analysis. *Journal of Statistical Software* **25**(1), 1–18 (2008). doi:10.18637/jss.v025.i01
22. Kassambara, A., Mundt, F.: Factoextra: Extract and Visualize the Results of Multivariate Data Analyses. (2019). R package version 1.0.6. <https://CRAN.R-project.org/package=factoextra>
23. Wickham, H.: Ggplot2: Elegant Graphics for Data Analysis. Springer, New York (2016). <https://ggplot2.tidyverse.org>
24. Slowikowski, K.: Ggrepel: Automatically Position Non-Overlapping Text Labels with 'ggplot2'. (2019). R package version 0.8.1. <https://CRAN.R-project.org/package=ggrepel>
25. SAS Institute: SAS 9.4 Language Reference: Concepts. SAS Institute Inc., USA (2013)
26. Wolever, T.M., Jenkins, D.J., Ocana, A.M., Rao, V.A., Collier, G.R.: Second-meal effect: low-glycemic-index foods eaten at dinner improve subsequent breakfast glycemic response. *The American Journal of Clinical Nutrition* **48**(4), 1041–1047 (1988). doi:10.1093/ajcn/48.4.1041. <https://academic.oup.com/ajcn/article-pdf/48/4/1041/24065765/1041.pdf>
27. Halder, S., Egli, L., De Castro, C.A., Tay, S.L., Koh, M.X.N., Darimont, C., Mace, K., Henry, C.J.: High or low glycemic index (gi) meals at dinner results in greater postprandial glycemia compared with breakfast: a randomized controlled trial. *BMJ open diabetes research & care* **8** (2020). doi:10.1136/bmjdr-2019-001099
28. Nitta, A., Imai, S., Kajiyama, S., Miyawaki, T., Matsumoto, S., Ozasa, N., Kajiyama, S., Hashimoto, Y., Tanaka, M., Fukui, M.: Impact of different timing of consuming sweet snack on postprandial glucose excursions in healthy women. *Diabetes & metabolism* **45**, 369–374 (2019). doi:10.1016/j.diabet.2018.10.004
29. Hassen, W.S., Castetbon, K., Tichit, C., Péneau, S., Nechba, A., Ducrot, P., Lampuré, A., Bellisle, F., Hercberg, S., Méjean, C.: Energy, nutrient and food content of snacks in french adults. *Nutrition Journal* **17**(1), 33 (2018)
30. Fayet-Moore, F., McConnell, A., Kim, J., Mathias, K.C.: Identifying eating occasion-based opportunities to improve the overall diets of australian adolescents. *Nutrients* **9** (2017). doi:10.3390/nu9060608

Figures

Figure 1 Biplot of food groups and eating time slots among total sample of NDNS RP.

Figure 2 Biplot of food groups and eating time slots among non-diabetics.

Tables

Figure 3 Biplot of food groups and eating time slots among diabetics.**Figure 4** Biplot of food groups and eating time slots among undiagnosed diabetics.**Figure 5** Biplot of food groups and eating time slots among pre-diabetics.**Table 1** The numbers of food recordings contributed to the total calories consumed by adults in the UK adults. (NDNS RP 2008-2017).

Food group names	n	Calories	Relative Prop	Cal Prop	Cal Cum Prop
Pasta & Rice and other cereals	18353	3512069.99	2.45%	7.36%	7.36%
White Bread	18434	3245641.19	2.46%	6.80%	14.17%
Chips, fried and roast potatoes and potato products	6749	1884058.68	0.90%	3.95%	18.12%
Cakes, buns, sweet pastries, fruit pies	7806	1710594.27	1.04%	3.59%	21.70%
Vegetable (not raw)	51317	1665474.02	6.85%	3.49%	25.19%
Biscuits	13200	1662598.06	1.76%	3.49%	28.68%
Fruit	33903	1641675.02	4.53%	3.44%	32.12%
Miscellaneous unclassified foods	48597	1639024.81	6.49%	3.44%	35.56%
Chicken/turkey	8863	1617820.30	1.18%	3.39%	38.95%
Cheese	10983	1492015.32	1.47%	3.13%	42.07%
Beer lager	8199	1484001.20	1.09%	3.11%	45.19%
Semi-skimmed milk	57611	1302649.72	7.69%	2.73%	47.92%
Potatoes other (in salads and dishes)	10113	1291447.61	1.35%	2.71%	50.62%
Fat spreads	37960	1215278.60	5.07%	2.55%	53.17%
Beef	4987	1124560.42	0.67%	2.36%	55.53%
High fiber breakfast cereals	8215	1072813.73	1.10%	2.25%	57.78%
Whole meal bread	7193	1070695.89	0.96%	2.24%	60.02%
Chocolate	6495	1046112.65	0.87%	2.19%	62.22%
Wine	6967	1027792.96	0.93%	2.15%	64.37%
Brown, granary and wheatgerm bread	6183	1009074.95	0.83%	2.12%	66.48%
Butter	10203	965901.11	1.36%	2.02%	68.51%
Eggs	7554	964769.19	1.01%	2.02%	70.53%
Soft drinks not diet	11387	940516.516	1.52%	1.97%	72.50%
Reduced fat spreads	12620	848834.89	1.68%	1.78%	74.28%
Crisps and savoury snacks	5664	835671.58	0.76%	1.75%	76.04%
Sausages	3025	775004.13	0.40%	1.62%	77.66%
Meat pastries	1979	744639.89	0.26%	1.56%	79.22%
Bacon and ham	8467	738727.49	1.13%	1.55%	80.77%
Yogurt	6776	665484.55	0.90%	1.40%	82.16%
Low-fiber breakfast cereals	4303	560296.32	0.57%	1.17%	83.34%
Nuts and seeds	6259	559873.88	0.84%	1.17%	84.51%
Oily fish	2610	550425.36	0.35%	1.15%	85.67%
Whole Milk	13628	530449.07	1.82%	1.11%	86.78%
White fish, shellfish	1597	498928.82	0.21%	1.05%	87.82%
Puddings	2291	459784.62	0.31%	0.96%	88.79%
Other Milk Cream	6605	434239.37	0.88%	0.91%	89.70%
Pork	1832	420503.76	0.24%	0.88%	90.58%

NDNS RP: National Diet and Nutrition Survey.

Table 2 Odds ratio (99% confidence intervals) for food groups eaten at night (8 pm - 6 am) vs. earlier time in the day, among total and according to different diabetes status, NDNS RP 2008-2017.

Selected food groups	Overall	Healthy	Pre-diabetics	Undiagnosed diabetes	Diabetes
Pudding	1.38 (1.03, 1.86)	1.50 (1.10, 2.07)	0.89 (0.16, 4.87)	1.81 (0.41, 7.98)	0.58 (0.14, 2.43)
Regular soft drink	1.74 (1.47, 2.06)	1.72 (1.43, 2.06)	1.87 (0.97, 3.57)	2.72 (1.44, 5.14)	1.38 (0.65, 2.96)
Sugar Confectionery	1.92 (1.38, 2.69)	1.63 (1.14, 2.32)	2.10 (0.52, 8.46)	13.07 (4.59, 37.24)	5.10 (2.15, 12.09)
Chocolate	3.19 (2.69, 3.79)	3.10 (2.57, 3.73)	4.07 (2.58, 3.73)	2.52 (0.95, 6.66)	5.13 (2.55, 10.30)
Spirit	11.13 (8.37, 14.80)	10.86 (8.01, 14.73)	8.48 (2.26, 31.79)	7.51 (1.99, 5.21)	36.8 (7.36, 183.66)
Beer	7.19 (5.87, 8.82)	7.49 (6.02, 9.34)	4.05 (2.00, 8.20)	7.87 (3.51, 17.63)	6.32 (2.29, 17.47)
Ice Cream	2.38 (1.79, 3.15)	2.45 (1.82, 3.31)	3.32 (0.75, 14.62)	0.98 (0.14, 7.00)	1.65 (0.54, 5.07)
Biscuit	1.91 (1.67, 2.16)	1.78 (1.55, 2.03)	3.51 (2.16, 5.71)	2.75 (1.35, 5.59)	2.44 (1.54, 3.88)
Crisp	1.55 (1.27, 1.88)	1.56 (1.27, 1.92)	1.95 (0.79, 4.78)	1.37 (0.37, 5.12)	1.16 (0.49, 2.75)

Logistic regression models with GEE were adjusted for age, sex, and social-economic levels.

NDNS RP: National Diet and Nutrition Survey.