## Conferences and talks

- 1. R. Turra, Studio del decadimento  $B_s^0 \to D_s^+ D_s^-$ , LHCb Italia, Bologna, January, 2009.
- 2. R. Turra, Misura della sezione d'urto di produzione di fotoni diretti con il rivelatore ATLAS all'LHC, Società Italiana di Fisica (SIF), Bologna, September, 2010.
- 3. R. Turra, Activities in egamma WG, ATLAS Italia, Napoli, May, 2011.
- 4. R. Turra, Search for the Standard Model Higgs Boson in the Decay Mode  $H \to \gamma \gamma$  with ATLAS, Beyond the standard model in particle physics, Quy Nhon (Vietnam), Aug, 2012.

## Posters

- 5. R. Turra, *Measurements of isolated prompt photons in pp collisions with the ATLAS detector*, No. ATL-PHYS-SLIDE-2011-379. Jul, 2011. Presented at PLHC, Perugia 2011.
- 6. R. Turra, *Ricerca di SM Higgs nel canale di decadimento H→gamma gamma in ATLAS*, Apr, 2012. Presented at IFAE, Ferrara 2012.

# **Proceedings**

7. R. Turra, *Measurements of isolated prompt photons in pp collisions with the ATLAS detector*, Tech. Rep. ATL-PHYS-PROC-2011-159, CERN, Geneva, Oct, 2011.

### Selected conf notes

- 8. ATLAS Collaboration, Measurement of the inclusive isolated prompt photon cross section in pp collisions at  $\sqrt{s}=7$  TeV with the ATLAS detector using 35 pb $^{-1}$ , Tech. Rep. ATLAS-CONF-2011-058, CERN, Geneva, Apr. 2011.
- 9. ATLAS Collaboration, *Expected photon performance in the ATLAS experiment*, Tech. Rep. ATL-PHYS-PUB-2011-007, CERN, Geneva, Apr. 2011.
- 10. Search for the Standard Model Higgs boson in the diphoton decay channel with 4.9 fb-1 of ATLAS data at sqrt(s)=7TeV, Tech. Rep. ATLAS-CONF-2011-161, CERN, Geneva, Dec, 2011.
- 11. Observation of an excess of events in the search for the Standard Model Higgs boson in the gamma-gamma channel with the ATLAS detector, Tech. Rep. ATLAS-CONF-2012-091, CERN, Geneva, Jul, 2012.
- 12. Observation of an Excess of Events in the Search for the Standard Model Higgs boson with the ATLAS detector at the LHC, Tech. Rep. ATLAS-CONF-2012-093, CERN, Geneva, Jul, 2012.

### Selected internal notes

- R. Turra et al., Electron and photon reconstruction and identification results from ATLAS at 900 GeV, Tech. Rep. ATL-COM-PHYS-2010-172, CERN, Geneva, Apr, 2010. This COM note supports the CONF note on electron and photon particle identification and reconstruction at 900 GeV (ATLAS-CONF-2010-005).
- 14. R. Turra et al., *Electromagnetic energy scale in-situ calibration and performance: Supporting document for the egamma performance paper*, Tech. Rep. ATL-COM-PHYS-2011-263, CERN, Geneva, Mar, 2011.
- 15. R. Turra et al., Measurement of the inclusive isolated prompt photon cross section in pp collisions at  $\sqrt{s}=7TeV$  with the ATLAS detector, Tech. Rep. ATL-PHYS-INT-2011-013, CERN, Geneva, Mar, 2011.
- 16. R. Turra et al., Measurement of the inclusive isolated prompt photon cross section in pp collisions at  $\sqrt{s}=7$  TeV with the ATLAS detector using 35 pb $^{-1}$ , Tech. Rep. ATL-PHYS-INT-2011-037, CERN, Geneva, Apr, 2011. Supporting document for CERN-PH-EP-2011-115.
- 17. R. Turra et al., *Purity Estimates for the Inclusive Isolated Photons*, Tech. Rep. ATL-PHYS-INT-2011-015, CERN, Geneva, Mar, 2011.

- 18. R. Turra et al., Calculating The Purity of Direct Photon Candidates in ATLAS: Proposed Methods and Plans for Early Data, Tech. Rep. ATL-COM-PHYS-2010-233, CERN, Geneva, May, 2010.
- 19. R. Turra et al., Study of the Di-Photon Backgrounds to the  $H\to\gamma\gamma$  Search with the ATLAS detector at  $\sqrt{s}=$  7 TeV, Tech. Rep. ATL-PHYS-INT-2011-011, CERN, Geneva, Feb, 2011. Supporting document for CERN-PH-EP-2011-088.
- 20. H. HSG1 Working Group, *Background Studies for the Search of Higgs Boson Decaying to Two Photons with 8TeV data*, Tech. Rep. ATL-COM-PHYS-2012-754, CERN, Geneva, Jun, 2012.
- 21. R. Turra et al., Measurement of isolated di-photon cross section in pp collision at  $\sqrt{s} = 7$  TeV with the ATLAS detector, Tech. Rep. ATL-PHYS-INT-2011-071, CERN, Geneva, Sep, 2011. Supporting document for CERN-PH-EP-2011-088.
- 22. R. Turra et al., Search for the Higgs boson in the diphoton final state with 38 pb $^{-1}$  of data recorded by the ATLAS detector at  $\sqrt{s}$ =7 TeV, Tech. Rep. ATL-PHYS-INT-2011-024, CERN, Geneva. Mar. 2011.

#### **Publications**

157 publications as ATLAS author, I've been mainly involved in [23, 24, 25, 26, 27, 24, 28]

- 23. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the inclusive isolated prompt photon cross-section in pp collisions at sqrt(s)= 7 TeV using 35 pb-1 of ATLAS data, Phys.Lett. **B706** (2011) 150-167, arXiv:1108.0253 [hep-ex].
- 24. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the isolated di-photon cross-section in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Rev. **D85** (2012) 012003, arXiv:1107.0581 [hep-ex].
- 25. ATLAS Collaboration Collaboration, G. Aad et al., Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC, arXiv:1207.7214 [hep-ex].
- 26. ATLAS Collaboration Collaboration, A. Collaboration et al., Search for a fermiophobic Higgs boson in the diphoton decay channel with the ATLAS detector, arXiv:1205.0701 [hep-ex].
- 27. ATLAS Collaboration Collaboration, G. Aad et al., Search for the Standard Model Higgs boson in the diphoton decay channel with 4.9 fb-1 of pp collisions at sqrt(s)=7 TeV with ATLAS, Phys.Rev.Lett. 108 (2012) 111803, arXiv:1202.1414 [hep-ex].
- 28. ATLAS Collaboration Collaboration, G. Aad et al., Search for the Standard Model Higgs boson in the two photon decay channel with the ATLAS detector at the LHC, Phys.Lett. **B705** (2011) 452–470, arXiv:1108.5895 [hep-ex].
- 29. ATLAS Collaboration Collaboration, G. Aad et al., Search for a supersymmetric partner to the top quark in final states with jets and missing transverse momentum at sqrt(s) = 7 TeV with the ATLAS detector, arXiv:1208.1447 [hep-ex].
- 30. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of WZ production in proton-proton collisions at sqrt(s) = 7 TeV with the ATLAS detector, arXiv:1208.1390 [hep-ex].
- 31. ATLAS Collaboration Collaboration, G. Aad et al., Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using 4.7  $fb^{-1}$  of sqrt(s) = 7 TeV proton-proton collision data, arXiv:1208.0949 [hep-ex].
- 32. ATLAS Collaboration Collaboration, G. Aad et al., Time-dependent angular analysis of the decay  $B_s \to J/\psi \phi$  and extraction of  $\Delta \Gamma_s$  and the CP-violating weak phase  $\phi_s$  by ATLAS, arXiv:1208.0572 [hep-ex].
- 33. ATLAS Collaboration Collaboration, G. Aad et al., Underlying event characteristics and their dependence on jet size of charged-particle jet events in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, arXiv:1208.0563 [hep-ex].
- 34. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of charged-particle event shape variables in sqrt(s) = 7 TeV proton-proton interactions with the ATLAS detector, arXiv:1207.6915 [hep-ex].

- 35. ATLAS Collaboration Collaboration, G. Aad et al., Search for magnetic monopoles in sqrt(s) = 7 TeV pp collisions with the ATLAS detector, arXiv:1207.6411 [hep-ex].
- 36. ATLAS Collaboration Collaboration, G. Aad et al., Measurements of top quark pair relative differential cross-sections with ATLAS in pp collisions at sqrt(s) = 7 TeV, arXiv:1207.5644 [hep-ex].
- 37. ATLAS Collaboration Collaboration, G. Aad et al., Search for top and bottom squarks from gluino pair production in final states with missing transverse energy and at least three b-jets with the ATLAS detector, arXiv:1207.4686 [hep-ex].
- 38. ATLAS Collaboration Collaboration, G. Aad et al., A search for  $t\bar{t}$  resonances in lepton+jets events with highly boosted top quarks collected in pp collisions at  $\sqrt{s}=7$  TeV with the ATLAS detector, arXiv:1207.2409 [hep-ex].
- 39. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the  $\Lambda_b$  lifetime and mass in the ATLAS experiment, arXiv:1207.2284 [hep-ex].
- 40. ATLAS Collaboration Collaboration, G. Aad et al., Combined search for the Standard Model Higgs boson in pp collisions at  $\sqrt{s}=7$  TeV with the ATLAS detector, arXiv:1207.0319 [hep-ex].
- 41. ATLAS Collaboration Collaboration, G. Aad et al., Search for the Standard Model Higgs boson produced in association with a vector boson and decaying to a b-quark pair with the ATLAS detector, arXiv:1207.0210 [hep-ex].
- 42. ATLAS Collaboration Collaboration, G. Aad et al., Search for the Higgs boson in the  $H \to WW \to l\nu jj$  decay channel at sqrt(s) = 7 TeV with the ATLAS detector, arXiv:1206.6074 [hep-ex].
- 43. ATLAS Collaboration Collaboration, G. Aad et al., Search for the Standard Model Higgs boson in the H to tau+ tau- decay mode in sqrt(s) = 7 TeV pp collisions with ATLAS, arXiv:1206.5971 [hep-ex].
- 44. ATLAS Collaboration Collaboration, G. Aad et al., ATLAS measurements of the properties of jets for boosted particle searches, arXiv:1206.5369 [hep-ex].
- 45. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the b-hadron production cross section using decays to  $D^*\mu^-X$  final states in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, arXiv:1206.3122 [hep-ex].
- 46. ATLAS Collaboration Collaboration, G. Aad et al., Search for a Standard Model Higgs boson in the mass range 200-600 GeV in the  $H \to ZZ \to l^+l^-q\bar{q}$  decay channel, arXiv:1206.2443 [hep-ex].
- 47. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of event shapes at large momentum transfer with the ATLAS detector in pp collisions at sqrt(s) = 7 TeV, arXiv:1206.2135 [hep-ex].
- 48. ATLAS Collaboration Collaboration, G. Aad et al., Hunt for new phenomena using large jet multiplicities and missing transverse momentum with ATLAS in 4.7  $fb^{-1}$  of sqrt(s) = 7 TeV proton-proton collisions, arXiv:1206.1760 [hep-ex].
- 49. ATLAS Collaboration Collaboration, G. Aad et al., Search for the Standard Model Higgs boson in the  $H \to WW(*) \to l\nu l\nu$  decay mode with 4.7  $fb^{-1}$  of ATLAS data at sqrt(s) = 7 TeV, arXiv:1206.0756 [hep-ex].
- 50. ATLAS Collaboration Collaboration, G. Aad et al., A search for flavour changing neutral currents in top-quark decays in pp collision data collected with the ATLAS detector at sqrt(s) = 7 TeV, arXiv:1206.0257 [hep-ex].
- 51. ATLAS Collaboration Collaboration, G. Aad et al., Search for a Standard Model Higgs boson in the  $H \to ZZ \to l^+ l^- \nu \bar{\nu}$  decay channel using 4.7 fb<sup>-1</sup> of sqrt(s) = 7 TeV data with the ATLAS detector, arXiv:1205.6744 [hep-ex].
- 52. ATLAS Collaboration Collaboration, G. Aad et al., Evidence for the associated production of a W boson and a top quark in ATLAS at sqrt(s) = 7 TeV, arXiv:1205.5764 [hep-ex].
- 53. ATLAS Collaboration Collaboration, G. Aad et al., A search for ttbar resonances with the ATLAS detector in 2.05  $fb^{-1}$  of proton-proton collisions at sqrt(s) = 7 TeV, Eur.Phys.J. C72 (2012) 2083, arXiv:1205.5371 [hep-ex].

- 54. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the t-channel single top-quark production cross section in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, arXiv:1205.3130 [hep-ex].
- 55. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of W gamma and Z gamma production cross sections in pp collisions at sqrt(s) = 7 TeV and limits on anomalous triple gauge couplings with the ATLAS detector, arXiv:1205.2531 [hep-ex].
- 56. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the W boson polarization in top quark decays with the ATLAS detector, JHEP 1206 (2012) 088, arXiv:1205.2484 [hep-ex].
- 57. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the top quark pair cross section with ATLAS in pp collisions at sqrt(s) = 7 TeV using final states with an electron or a muon and a hadronically decaying  $\tau$  lepton, arXiv:1205.2067 [hep-ex].
- 58. ATLAS Collaboration Collaboration, G. Aad et al., Search for the resonances in proton-proton collisions at sqrt(s) = 7 TeV with the ATLAS detector, arXiv:1205.1016 [hep-ex].
- 59. ATLAS Collaboration Collaboration, G. Aad et al., Search for lepton flavour violation in the emu continuum with the ATLAS detector in sqrt(s) = 7 TeV pp collisions at the LHC, Eur.Phys.J. C72 (2012) 2040, arXiv:1205.0725 [hep-ex].
- 60. ATLAS Collaboration Collaboration, G. Aad et al., Search for scalar top quark pair production in natural gauge mediated supersymmetry models with the ATLAS detector in pp collisions at sqrt(s) = 7 TeV, arXiv:1204.6736 [hep-ex].
- 61. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of  $\tau$  polarization in  $W \to \tau \nu$  decays with the ATLAS detector in pp collisions at sqrt(s) = 7 TeV, Eur.Phys.J. **C72** (2012) 2062, arXiv:1204.6720 [hep-ex].
- 62. ATLAS Collaboration Collaboration, G. Aad et al., Search for supersymmetry in events with three leptons and missing transverse momentum in sqrt(s) = 7 TeV pp collisions with the ATLAS detector, Phys.Rev.Lett. 108 (2012) 261804, arXiv:1204.5638 [hep-ex].
- 63. ATLAS Collaboration Collaboration, G. Aad et al., Search for TeV-scale gravity signatures in final states with leptons and jets with the ATLAS detector at sqrt(s) = 7 TeV, arXiv:1204.4646 [hep-ex].
- 64. ATLAS Collaboration Collaboration, G. Aad et al., Search for supersymmetry with jets, missing transverse momentum and at least one hadronically decaying tau lepton in proton-proton collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys. Lett. **B714** (2012) 197–214, arXiv:1204.3852 [hep-ex].
- 65. ATLAS Collaboration Collaboration, G. Aad et al., Search for charged Higgs bosons decaying via  $H^+ \to \tau \nu$  in top quark pair events using pp collision data at sqrt(s) = 7 TeV with the ATLAS detector, JHEP 1206 (2012) 039, arXiv:1204.2760 [hep-ex].
- 66. ATLAS Collaboration Collaboration, G. Aad et al., Search for resonant WZ production in the WZ to I nu I' I' channel in sqrt(s) = 7 TeV pp collisions with the ATLAS detector, Phys. Rev. **D85** (2012) 112012, arXiv:1204.1648 [hep-ex].
- 67. ATLAS Collaboration Collaboration, G. Aad et al., Search for pair production of a new quark that decays to a Z boson and a bottom quark with the ATLAS detector, Phys.Rev.Lett. (2012), arXiv:1204.1265 [hep-ex].
- 68. ATLAS Collaboration Collaboration, G. Aad et al., Search for the decay  $B_s^0 \to \mu\mu$  with the ATLAS detector, Phys.Lett. **B713** (2012) 387–407, arXiv:1204.0735 [hep-ex].
- 69. ATLAS Collaboration Collaboration, G. Aad et al., Search for events with large missing transverse momentum, jets, and at least two tau leptons in 7 TeV proton-proton collision data with the ATLAS detector, Phys.Lett. **B714** (2012) 180–196, arXiv:1203.6580 [hep-ex].
- 70. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the WW cross section in sqrt(s) = 7 TeV pp collisions with the ATLAS detector and limits on anomalous gauge couplings, Phys.Lett. B712 (2012) 289–308, arXiv:1203.6232 [hep-ex].
- 71. ATLAS Collaboration Collaboration, G. Aad et al., Search for supersymmetry in pp collisions at sqrt(s) = 7 TeV in final states with missing transverse momentum and b-jets with the ATLAS detector, Phys.Rev. **D85** (2012) 112006, arXiv:1203.6193 [hep-ex].

- 72. ATLAS Collaboration Collaboration, G. Aad et al., Search for gluinos in events with two same-sign leptons, jets and missing transverse momentum with the ATLAS detector in pp collisions at sqrt(s) = 7 TeV, Phys.Rev.Lett. 108 (2012) 241802, arXiv:1203.5763 [hep-ex].
- 73. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the top quark mass with the template method in the  $t\bar{t} \to lepton + jets$  channel using ATLAS data, Eur.Phys.J. **C72** (2012) 2046, arXiv:1203.5755 [hep-ex].
- 74. ATLAS Collaboration Collaboration, G. Aad et al., Search for heavy neutrinos and right-handed W bosons in events with two leptons and jets in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, Eur.Phys.J. C72 (2012) 2056, arXiv:1203.5420 [hep-ex].
- 75. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of  $t\bar{t}$  production with a veto on additional central jet activity in pp collisions at sqrt(s) = 7 TeV using the ATLAS detector, Eur.Phys.J. C72 (2012) 2043, arXiv:1203.5015 [hep-ex].
- 76. ATLAS Collaboration Collaboration, G. Aad et al., Jet mass and substructure of inclusive jets in sqrt(s) = 7 TeV pp collisions with the ATLAS experiment, JHEP 1205 (2012) 128, arXiv:1203.4606 [hep-ex].
- 77. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the charge asymmetry in top quark pair production in pp collisions at sqrt(s) = 7 TeV using the ATLAS detector, Eur.Phys.J. C72 (2012) 2039, arXiv:1203.4211 [hep-ex].
- 78. ATLAS Collaboration Collaboration, G. Aad et al., Observation of spin correlation in  $t\bar{t}$  events from pp collisions at sqrt(s) = 7 TeV using the ATLAS detector, Phys.Rev.Lett. **108** (2012) 212001, arXiv:1203.4081 [hep-ex].
- 79. ATLAS Collaboration Collaboration, G. Aad et al., Determination of the strange quark density of the proton from ATLAS measurements of the  $W \to l\nu$  and  $Z \to ll$  cross sections, Phys.Rev.Lett. 109 (2012) 012001, arXiv:1203.4051 [hep-ex].
- 80. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of inclusive two-particle angular correlations in pp collisions with the ATLAS detector at the LHC, JHEP 1205 (2012) 157, arXiv:1203.3549 [hep-ex].
- 81. ATLAS Collaboration Collaboration, G. Aad et al., Search for second generation scalar leptoquarks in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, arXiv:1203.3172 [hep-ex].
- 82. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the production cross section of an isolated photon associated with jets in proton-proton collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Rev. **D85** (2012) 092014, arXiv:1203.3161 [hep-ex].
- 83. ATLAS Collaboration Collaboration, G. Aad et al., Forward-backward correlations and charged-particle azimuthal distributions in pp interactions using the ATLAS detector, JHEP 1207 (2012) 019, arXiv:1203.3100 [hep-ex].
- 84. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the azimuthal anisotropy for charged particle production in  $\sqrt(s_{NN})=2.76$  TeV lead-lead collisions with the ATLAS detector, Phys.Rev. **C86** (2012) 014907, arXiv:1203.3087 [hep-ex].
- 85. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the polarisation of W bosons produced with large transverse momentum in pp collisions at sqrt(s) = 7 TeV with the ATLAS experiment, Eur.Phys.J. C72 (2012) 2001, arXiv:1203.2165 [hep-ex].
- 86. ATLAS Collaboration Collaboration, G. Aad et al., Search for a light Higgs boson decaying to long-lived weakly-interacting particles in proton-proton collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Rev.Lett. 108 (2012) 251801, arXiv:1203.1303 [hep-ex].
- 87. ATLAS Collaboration Collaboration, G. Aad et al., Single hadron response measurement and calorimeter jet energy scale uncertainty with the ATLAS detector at the LHC, arXiv:1203.1302 [hep-ex].
- 88. ATLAS Collaboration Collaboration, G. Aad et al., Search for new particles decaying to ZZ using final states with leptons and jets with the ATLAS detector in sqrt(s) = 7 Tev proton-proton collisions, Phys.Lett. B712 (2012) 331–350, arXiv:1203.0718 [hep-ex].

- 89. ATLAS Collaboration Collaboration, G. Aad et al., Search for FCNC single top-quark production at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Lett. **B712** (2012) 351–369, arXiv:1203.0529 [hep-ex].
- 90. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the azimuthal ordering of charged hadrons with the ATLAS detector, arXiv:1203.0419 [hep-ex].
- 91. ATLAS Collaboration Collaboration, G. Aad et al., Search for down-type fourth generation quarks with the ATLAS detector in events with one lepton and hadronically decaying W bosons, Phys.Rev.Lett. 109 (2012) 032001, arXiv:1202.6540 [hep-ex].
- 92. ATLAS Collaboration Collaboration, G. Aad et al., Search for same-sign top-quark production and fourth-generation down-type quarks in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, JHEP 1204 (2012) 069, arXiv:1202.5520 [hep-ex].
- 93. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the cross section for top-quark pair production in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector using final states with two high-pt leptons, JHEP 1205 (2012) 059, arXiv:1202.4892 [hep-ex].
- 94. ATLAS Collaboration Collaboration, G. Aad et al., Search for anomaly-mediated supersymmetry breaking with the ATLAS detector based on a disappearing-track signature in pp collisions at sqrt(s) = 7 TeV, Eur.Phys.J. **C72** (2012) 1993, arXiv:1202.4847 [hep-ex].
- 95. ATLAS Collaboration Collaboration, G. Aad et al., Search for pair-produced heavy quarks decaying to Wq in the two-lepton channel at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Rev. **D86** (2012) 012007, arXiv:1202.3389 [hep-ex].
- 96. ATLAS Collaboration Collaboration, G. Aad et al., Search for pair production of a heavy quark decaying to a W boson and a b quark in the lepton+jets channel with the ATLAS detector, Phys.Rev.Lett. **108** (2012) 261802, arXiv:1202.3076 [hep-ex].
- 97. ATLAS Collaboration Collaboration, G. Aad et al., Search for the Standard Model Higgs boson in the decay channel  $H \to ZZ(*) \to 4l$  with 4.8  $fb^{-1}$  of pp collision data at sqrt(s) = 7 TeV with ATLAS, Phys.Lett. **B710** (2012) 383-402, arXiv:1202.1415 [hep-ex].
- 98. ATLAS Collaboration Collaboration, G. Aad et al., Combined search for the Standard Model Higgs boson using up to 4.9 fb-1 of pp collision data at sqrt(s) = 7 TeV with the ATLAS detector at the LHC, Phys.Lett. **B710** (2012) 49–66, arXiv:1202.1408 [hep-ex].
- 99. ATLAS Collaboration Collaboration, G. Aad et al., Search for decays of stopped, long-lived particles from 7 TeV pp collisions with the ATLAS detector, Eur.Phys.J. **C72** (2012) 1965, arXiv:1201.5595 [hep-ex].
- 100. ATLAS Collaboration Collaboration, G. Aad et al., Search for excited leptons in proton-proton collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Rev. **D85** (2012) 072003, arXiv:1201.3293 [hep-ex].
- 101. ATLAS Collaboration Collaboration, G. Aad et al., Rapidity gap cross sections measured with the ATLAS detector in pp collisions at sqrt(s) = 7 TeV, Eur.Phys.J. C72 (2012) 1926, arXiv:1201.2808 [hep-ex].
- 102. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the top quark pair production cross-section with ATLAS in the single lepton channel, Phys.Lett. **B711** (2012) 244–263, arXiv:1201.1889 [hep-ex].
- 103. ATLAS Collaboration Collaboration, G. Aad et al., Study of jets produced in association with a W boson in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Rev. **D85** (2012) 092002, arXiv:1201.1276 [hep-ex].
- 104. ATLAS Collaboration Collaboration, G. Aad et al., Search for anomalous production of prompt like-sign muon pairs and constraints on physics beyond the Standard Model with the ATLAS detector, Phys.Rev. **D85** (2012) 032004, arXiv:1201.1091 [hep-ex].
- 105. ATLAS Collaboration Collaboration, G. Aad et al., *Jet energy measurement with the ATLAS detector in proton-proton collisions at sqrt(s) = 7 TeV*, arXiv:1112.6426 [hep-ex].
- 106. ATLAS Collaboration Collaboration, G. Aad et al., *Measurement of inclusive jet and dijet production in pp collisions at sqrt(s) = 7 TeV using the ATLAS detector*, Phys. Rev. **D86** (2012) 014022, arXiv:1112.6297 [hep-ex].

- 107. ATLAS Collaboration Collaboration, G. Aad et al., Search for heavy vector-like quarks coupling to light quarks in proton-proton collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Lett. **B712** (2012) 22–39, arXiv:1112.5755 [hep-ex].
- 108. ATLAS Collaboration Collaboration, G. Aad et al., Observation of a new  $\chi_b$  state in radiative transitions to  $\Upsilon(1S)$  and  $\Upsilon(2S)$  at ATLAS, Phys. Rev. Lett. **108** (2012) 152001, arXiv:1112.5154 [hep-ex].
- 109. ATLAS Collaboration Collaboration, G. Aad et al., Search for first generation scalar leptoquarks in pp collisions at sqrts=7 TeV with the ATLAS detector, Phys.Lett. **B709** (2012) 158–176, arXiv:1112.4828 [hep-ex].
- 110. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of  $D^{*+/-}$  meson production in jets from pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Rev. **D85** (2012) 052005, arXiv:1112.4432 [hep-ex].
- 111. ATLAS Collaboration Collaboration, G. Aad et al., Search for contact interactions in dilepton events from pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Lett. **B712** (2012) 40–58, arXiv:1112.4462 [hep-ex].
- 112. ATLAS Collaboration Collaboration, G. Aad et al., Search for scalar bottom pair production with the ATLAS detector in pp Collisions at sqrts = 7 TeV, Phys.Rev.Lett. 108 (2012) 181802, arXiv:1112.3832 [hep-ex].
- 113. ATLAS Collaboration Collaboration, G. Aad et al., Search for production of resonant states in the photon-jet mass distribution using pp collisions at sqrt(s) = 7 TeV collected by the ATLAS detector, Phys.Rev.Lett. 108 (2012) 211802, arXiv:1112.3580 [hep-ex].
- 114. ATLAS Collaboration Collaboration, G. Aad et al., Search for the Higgs boson in the  $H \to WW^{(*)} \to l\nu l\nu$  decay channel in pp collisions at sqrts = 7 TeV with the ATLAS detector, Phys.Rev.Lett. 108 (2012) 111802, arXiv:1112.2577 [hep-ex].
- 115. ATLAS Collaboration Collaboration, G. Aad et al., Search for Extra Dimensions using diphoton events in 7 TeV proton-proton collisions with the ATLAS detector, Phys.Lett. **B710** (2012) 538–556, arXiv:1112.2194 [hep-ex].
- 116. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the WZ production cross section and limits on anomalous triple gauge couplings in proton-proton collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Lett. B709 (2012) 341–357, arXiv:1111.5570 [hep-ex].
- 117. ATLAS Collaboration Collaboration, G. Aad et al., Search for Diphoton Events with Large Missing Transverse Momentum in 1  $fb^{-1}$  of 7 TeV Proton-Proton Collision Data with the ATLAS Detector, Phys.Lett. **B710** (2012) 519–537, arXiv:1111.4116 [hep-ex].
- 118. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the production cross section for Z/gamma\* in association with jets in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Rev. **D85** (2012) 032009, arXiv:1111.2690 [hep-ex].
- 119. ATLAS Collaboration Collaboration, G. Aad et al., Kshort and Lambda production in pp interactions at sqrt(s) = 0.9 and 7 TeV measured with the ATLAS detector at the LHC, Phys.Rev. **D85** (2012) 012001, arXiv:1111.1297 [hep-ex].
- 120. ATLAS Collaboration Collaboration, G. Aad et al., Search for strong gravity signatures in same-sign dimuon final states using the ATLAS detector at the LHC, Phys.Lett. **B709** (2012) 322–340, arXiv:1111.0080 [hep-ex].
- 121. ATLAS Collaboration Collaboration, G. Aad et al., A study of the material in the ATLAS inner detector using secondary hadronic interactions, JINST 7 (2012) P01013, arXiv:1110.6191 [hep-ex].
- 122. ATLAS Collaboration Collaboration, G. Aad et al., Searches for supersymmetry with the ATLAS detector using final states with two leptons and missing transverse momentum in sqrts = 7 TeV proton-proton collisions, Phys.Lett. **B709** (2012) 137–157, arXiv:1110.6189 [hep-ex].
- 123. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the ZZ production cross section and limits on anomalous neutral triple gauge couplings in proton-proton collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Rev.Lett. 108 (2012) 041804, arXiv:1110.5016 [hep-ex].

- 124. ATLAS Collaboration Collaboration, G. Aad et al., *Electron performance measurements with the ATLAS detector using the 2010 LHC proton-proton collision data*, Eur.Phys.J. **C72** (2012) 1909, arXiv:1110.3174 [hep-ex].
- 125. ATLAS Collaboration Collaboration, G. Aad et al., Search for Massive Colored Scalars in Four-Jet Final States in sqrts=7 TeV proton-proton collisions with the ATLAS Detector, Eur.Phys.J. C71 (2011) 1828, arXiv:1110.2693 [hep-ex].
- 126. Atlas Collaboration Collaboration, G. Aad et al., Search for new phenomena in final states with large jet multiplicities and missing transverse momentum using sqrt(s)=7 TeV pp collisions with the ATLAS detector, JHEP 1111 (2011) 099, arXiv:1110.2299 [hep-ex].
- 127. Atlas Collaboration Collaboration, G. Aad et al., *Performance of the ATLAS Trigger System in 2010*, Eur.Phys.J. **C72** (2012) 1849, arXiv:1110.1530 [hep-ex].
- 128. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the inclusive and dijet cross-sections of b-jets in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, Eur.Phys.J. C71 (2011) 1846, arXiv:1109.6833 [hep-ex].
- 129. ATLAS Collaboration Collaboration, G. Aad et al., Search for supersymmetry in final states with jets, missing transverse momentum and one isolated lepton in sqrts = 7 TeV pp collisions using  $1\ fb^{-1}$  of ATLAS data, Phys.Rev. **D85** (2012) 012006, arXiv:1109.6606 [hep-ex].
- 130. ATLAS Collaboration Collaboration, G. Aad et al., Search for squarks and gluinos using final states with jets and missing transverse momentum with the ATLAS detector in sqrt(s) = 7 TeV proton-proton collisions, Phys.Lett. **B710** (2012) 67–85, arXiv:1109.6572 [hep-ex].
- 131. ATLAS Collaboration Collaboration, G. Aad et al., Search for the Standard Model Higgs boson in the decay channel  $H \to ZZ^{(*)} \to 4l$  with the ATLAS detector, Phys.Lett. **B705** (2011) 435–451, arXiv:1109.5945 [hep-ex].
- 132. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the jet fragmentation function and transverse profile in proton-proton collisions at a center-of-mass energy of 7 TeV with the ATLAS detector, Eur.Phys.J. C71 (2011) 1795, arXiv:1109.5816 [hep-ex].
- 133. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the inclusive W+- and Z/gamma cross sections in the electron and muon decay channels in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Rev. **D85** (2012) 072004, arXiv:1109.5141 [hep-ex].
- 134. ATLAS Collaboration Collaboration, G. Aad et al., Search for New Phenomena in ttbar Events With Large Missing Transverse Momentum in Proton-Proton Collisions at sqrt(s) = 7 TeV with the ATLAS Detector, Phys.Rev.Lett. 108 (2012) 041805, arXiv:1109.4725 [hep-ex].
- 135. ATLAS Collaboration Collaboration, G. Aad et al., Search for the Higgs boson in the  $H \to WW \to l\nu jj$  decay channel in pp collisions at sqrts = 7 TeV with the ATLAS detector, Phys.Rev.Lett. 107 (2011) 231801, arXiv:1109.3615 [hep-ex].
- 136. ATLAS Collaboration Collaboration, G. Aad et al., Search for a Standard Model Higgs boson in the  $H \to ZZ \to l^+ l^- \nu \bar{\nu}$  decay channel with the ATLAS detector, Phys.Rev.Lett. **107** (2011) 221802, arXiv:1109.3357 [hep-ex].
- 137. ATLAS Collaboration Collaboration, G. Aad et al., Search for a heavy neutral particle decaying into an electron and a muon using  $1\ fb^{-1}$  of ATLAS data, Eur.Phys.J. **C71** (2011) 1809, arXiv:1109.3089 [hep-ex].
- 138. ATLAS Collaboration Collaboration, G. Aad et al., Search for displaced vertices arising from decays of new heavy particles in 7 TeV pp collisions at ATLAS, Phys.Lett. **B707** (2012) 478–496, arXiv:1109.2242 [hep-ex].
- 139. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the cross-section for b-jets produced in association with a Z boson at sqrt(s)=7 TeV with the ATLAS detector, Phys.Lett. B706 (2012) 295-313, arXiv:1109.1403 [hep-ex].
- 140. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the cross section for the production of a W boson in association with b-jets in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Lett. **B707** (2012) 418–437, arXiv:1109.1470 [hep-ex].
- 141. ATLAS Collaboration Collaboration, G. Aad et al., Measurements of the electron and muon inclusive cross-sections in proton-proton collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Lett. **B707** (2012) 438–458, arXiv:1109.0525 [hep-ex].

- 142. ATLAS Collaboration Collaboration, G. Aad et al., Search for New Physics in the Dijet Mass Distribution using 1  $fb^{-1}$  of pp Collision Data at sqrt(s) = 7 TeV collected by the ATLAS Detector, Phys.Lett. **B708** (2012) 37–54, arXiv:1108.6311 [hep-ex].
- 143. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the Transverse Momentum Distribution of W Bosons in pp Collisions at sqrt(s) = 7 TeV with the ATLAS Detector, Phys.Rev. **D85** (2012) 012005, arXiv:1108.6308 [hep-ex].
- 144. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the centrality dependence of the charged particle pseudorapidity distribution in lead-lead collisions at  $\sqrt{s_{NN}}=2.76$  TeV with the ATLAS detector, Phys.Lett. **B710** (2012) 363–382, arXiv:1108.6027 [hep-ex].
- 145. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the pseudorapidity and transverse momentum dependence of the elliptic flow of charged particles in lead-lead collisions at  $\sqrt{s_{NN}}=2.76$  TeV with the ATLAS detector, Phys.Lett. **B707** (2012) 330–348, arXiv:1108.6018 [hep-ex].
- 146. ATLAS Collaboration Collaboration, G. Aad et al., *Performance of Missing Transverse Momentum Reconstruction in Proton-Proton Collisions at 7 TeV with ATLAS*, Eur.Phys.J. **C72** (2012) 1844, arXiv:1108.5602 [hep-ex].
- 147. ATLAS Collaboration Collaboration, G. Aad et al., Search for a heavy Standard Model Higgs boson in the channel  $H \to ZZ \to llqq$  using the ATLAS detector, Phys.Lett. **B707** (2012) 27–45, arXiv:1108.5064 [hep-ex].
- 148. ATLAS Collaboration Collaboration, G. Aad et al., A measurement of the ratio of the W and Z cross sections with exactly one associated jet in pp collisions at sqrt(s) = 7 TeV with ATLAS, Phys.Lett. B708 (2012) 221–240, arXiv:1108.4908 [hep-ex].
- 149. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the W to tau nu Cross Section in pp Collisions at sqrt(s) = 7 TeV with the ATLAS experiment, Phys.Lett. **B706** (2012) 276–294, arXiv:1108.4101 [hep-ex].
- 150. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the top quark pair production cross section in pp collisions at sqrt(s) = 7 TeV in dilepton final states with ATLAS, Phys.Lett. **B707** (2012) 459-477, arXiv:1108.3699 [hep-ex].
- 151. ATLAS Collaboration Collaboration, G. Aad et al., *Measurement of the Z to tau tau Cross Section with the ATLAS Detector*, Phys.Rev. **D84** (2011) 112006, arXiv:1108.2016 [hep-ex].
- 152. ATLAS Collaboration Collaboration, G. Aad et al., Search for dilepton resonances in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Rev.Lett. 107 (2011) 272002, arXiv:1108.1582 [hep-ex].
- 153. ATLAS Collaboration Collaboration, G. Aad et al., Search for a heavy gauge boson decaying to a charged lepton and a neutrino in 1 fb-1 of pp collisions at sqrt(s) = 7 TeV using the ATLAS detector, Phys.Lett. **B705** (2011) 28-46, arXiv:1108.1316 [hep-ex].
- 154. ATLAS Collaboration Collaboration, G. Aad et al., *Inclusive search for same-sign dilepton signatures in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector*, JHEP **1110** (2011) 107, arXiv:1108.0366 [hep-ex].
- 155. ATLAS Collaboration Collaboration, G. Aad et al., Search for neutral MSSM Higgs bosons decaying to  $\tau^+\tau^-$  pairs in proton-proton collisions at sqrt(s) = 7 TeV with the ATLAS detector, Phys.Lett. **B705** (2011) 174–192, arXiv:1107.5003 [hep-ex].
- 156. ATLAS Collaboration Collaboration, G. Aad et al., *Properties of jets measured from tracks in proton-proton collisions at center-of-mass energy sqrt(s) = 7 TeV with the ATLAS detector*, Phys.Rev. **D84** (2011) 054001, arXiv:1107.3311 [hep-ex].
- 157. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the transverse momentum distribution of Z/gamma\* bosons in proton-proton collisions at sqrt(s)=7 TeV with the ATLAS detector, Phys.Lett. B705 (2011) 415–434, arXiv:1107.2381 [hep-ex].
- 158. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of multi-jet cross sections in proton-proton collisions at a 7 TeV center-of-mass energy, Eur.Phys.J. C71 (2011) 1763, arXiv:1107.2092 [hep-ex].

- 159. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of dijet production with a veto on additional central jet activity in pp collisions at sqrt(s)=7 TeV using the ATLAS detector, JHEP 1109 (2011) 053, arXiv:1107.1641 [hep-ex].
- 160. ATLAS Collaboration Collaboration, G. Aad et al., Search for Diphoton Events with Large Missing Transverse Energy with 36  $pb^{-1}$  of 7 TeV Proton-Proton Collision Data with the ATLAS Detector, Eur.Phys.J. **C71** (2011) 1744, arXiv:1107.0561 [hep-ex].
- 161. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the Upsilon(1S) Production Cross-Section in pp Collisions at sqrt(s) = 7 TeV in ATLAS, Phys.Lett. **B705** (2011) 9–27, arXiv:1106.5325 [hep-ex].
- 162. ATLAS Collaboration Collaboration, G. Aad et al., Search for new phenomena with the monojet and missing transverse momentum signature using the ATLAS detector in sqrt(s) = 7 TeV proton-proton collisions, Phys.Lett. **B705** (2011) 294–312, arXiv:1106.5327 [hep-ex].
- 163. ATLAS Collaboration Collaboration, G. Aad et al., Search for Heavy Long-Lived Charged Particles with the ATLAS detector in pp collisions at sqrt(s) = 7 TeV, Phys.Lett. **B703** (2011) 428–446, arXiv:1106.4495 [hep-ex].
- 164. ATLAS Collaboration Collaboration, G. Aad et al., Limits on the production of the Standard Model Higgs Boson in pp collisions at sqrt(s) =7 TeV with the ATLAS detector, Eur.Phys.J. C71 (2011) 1728, arXiv:1106.2748 [hep-ex].
- 165. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of Wgamma and Zgamma production in proton-proton collisions at sqrt(s)=7 TeV with the ATLAS Detector, JHEP 1109 (2011) 072, arXiv:1106.1592 [hep-ex].
- 166. ATLAS Collaboration Collaboration, G. Aad et al., *Measurement of the WW cross section in sqrt(s) = 7 TeV pp collisions with ATLAS*, Phys.Rev.Lett. **107** (2011) 041802, arXiv:1104.5225 [hep-ex].
- 167. ATLAS Collaboration Collaboration, G. Aad et al., Search for Contact Interactions in Dimuon Events from pp Collisions at sqrt(s) = 7 TeV with the ATLAS Detector, Phys.Rev. **D84** (2011) 011101, arXiv:1104.4398 [hep-ex].
- 168. ATLAS Collaboration Collaboration, G. Aad et al., Search for pair production of first or second generation leptoquarks in proton-proton collisions at sqrt(s)=7 TeV using the ATLAS detector at the LHC, Phys.Rev. **D83** (2011) 112006, arXiv:1104.4481 [hep-ex].
- 169. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the differential cross-sections of inclusive, prompt and non-prompt J/psi production in proton-proton collisions at sqrt(s) = 7 TeV, Nucl.Phys. **B850** (2011) 387–444, arXiv:1104.3038 [hep-ex].
- 170. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the Inelastic Proton-Proton Cross-Section at  $\sqrt{s}=7$  TeV with the ATLAS Detector, Nature Commun. 2 (2011) 463, arXiv:1104.0326 [hep-ex].
- 171. ATLAS Collaboration Collaboration, G. Aad et al., Search for an excess of events with an identical flavour lepton pair and significant missing transverse momentum in sqrts = 7 TeV proton-proton collisions with the ATLAS detector, Eur.Phys.J. C71 (2011) 1647, arXiv:1103.6208 [hep-ex].
- 172. ATLAS Collaboration Collaboration, G. Aad et al., Search for supersymmetric particles in events with lepton pairs and large missing transverse momentum in  $\sqrt{s}=7$  TeV proton-proton collisions with the ATLAS experiment, Eur.Phys.J. **C71** (2011) 1682, arXiv:1103.6214 [hep-ex].
- 173. ATLAS Collaboration Collaboration, G. Aad et al., Search for high mass dilepton resonances in pp collisions at  $\sqrt{s}=7$  TeV with the ATLAS experiment, Phys.Lett. **B700** (2011) 163–180, arXiv:1103.6218 [hep-ex].
- 174. ATLAS Collaboration Collaboration, G. Aad et al., Search for a heavy particle decaying into an electron and a muon with the ATLAS detector in  $\sqrt{s}=7$  TeV pp collisions at the LHC, Phys.Rev.Lett. 106 (2011) 251801, arXiv:1103.5559 [hep-ex].
- 175. ATLAS Collaboration Collaboration, G. Aad et al., Search for supersymmetry in pp collisions at sqrts = 7TeV in final states with missing transverse momentum and b-jets, Phys.Lett. B701 (2011) 398-416, arXiv:1103.4344 [hep-ex].

- 176. ATLAS Collaboration Collaboration, G. Aad et al., Search for New Physics in Dijet Mass and Angular Distributions in pp Collisions at  $\sqrt{s}=7$  TeV Measured with the ATLAS Detector, New J.Phys. **13** (2011) 053044, arXiv:1103.3864 [hep-ex].
- 177. ATLAS Collaboration Collaboration, G. Aad et al., Measurement of the Muon Charge Asymmetry from W Bosons Produced in pp Collisions at  $\sqrt{s}=7$  TeV with the ATLAS detector, Phys.Lett. B701 (2011) 31-49, arXiv:1103.2929 [hep-ex].
- 178. ATLAS Collaboration Collaboration, G. Aad et al., Search for stable hadronising squarks and gluinos with the ATLAS experiment at the LHC, Phys.Lett. **B701** (2011) 1–19, arXiv:1103.1984 [hep-ex].
- 179. ATLAS Collaboration Collaboration, G. Aad et al., Search for high-mass states with one lepton plus missing transverse momentum in proton-proton collisions at  $\sqrt{s}=7$  TeV with the ATLAS detector, Phys.Lett. **B701** (2011) 50-69, arXiv:1103.1391 [hep-ex].