

Probability and Measure Solutions

WispyAbyss

December 9, 2024

Contents

Forward	1
Chapter 1.1 - Borel's Normal Number Theorem	1
Notes	1
The Unit Interval	1
Solutions	2

Forward

This document will contain notes and solutions corresponding to Probability and Measure, Third Edition, by Patrick Billingsley [amazon].

Chapter 1.1 - Borel's Normal Number Theorem

Notes

For a complete understanding of probability, you need to understand an infinite number of events as well as a finite number of events. We try and present why that must be so here.

The Unit Interval

We take the length of an interval $I = (a, b] = b - a$. Note, for A a disjoint set of intervals in $(0, 1]$, we have that $P(A)$ is well defined. If B is a similar disjoint set, and is disjoint from A , $P(A + B) = P(A) + P(B)$ is well defined as well. Note - we haven't defined anything for intersections yet. These

definitions can also directly stem from the Riemann integral of step functions.

The unit interval can give the probability that a single particle is emitted in a unit interval of time. Or a single phone call comes in. However, it can also model an infinite coin toss. This is done as follows - for $\omega \in (0, 1]$, define:

$$\omega = \sum_{n=1}^{\infty} \frac{d_n(\omega)}{2^n}$$

Where $d_n(\omega)$ is 0 or 1, and comes from the binary expansion of ω . We take ω as the non terminating representation. Note, we were particular when we defined intervals as half inclusive. Examine the set of ω for which $d_i(\omega) = u_i$ for $i = 1, \dots, n$, $u_i \in \{0, 1\}$. We have that:

$$\sum_{i=1}^n \frac{u_i}{2^i} < \omega \leq \sum_{i=1}^n \frac{u_i}{2^i} + \sum_{i=n+1}^{\infty} \frac{1}{2^i}$$

We cannot have the lower extreme value, as this would imply ω takes on its terminating binomial representation, which is what we said we would not do. This is our first taste, I guess, of measure 0 sets, we we still have:

$$\mathbb{P}[\omega : d_i(\omega) = u_i, i = 1, \dots, n] = \frac{1}{2^n}$$

Note, probabilities of various familiar events can be written down immediately. Ultimately, note, however, each probability is the sum of disjoint dyadic intervals of various ranks k . Ie, all the events are still well defined by our probability definition above. We have:

$$\mathbb{P}\left[\omega : \sum_{i=1}^n d_i(\omega) = k\right] = \binom{n}{k} \frac{1}{2^n}$$

Solutions