

# Probability and Measure Solutions

WispyAbyss

January 30, 2025

## Contents

<b>Forward</b>	<b>2</b>
<b>Section 1 - Borel's Normal Number Theorem</b>	<b>3</b>
Notes	3
The Unit Interval	3
The Weak Law of Large Numbers	4
The Strong Law of Large Numbers	6
The Measure Theory of Diophantine Approximation	9
Problems	11
1.1 Infinite Independent Events on a Discrete Space (are im- possible)	11
1.2 Normal Numbers and Complements are Dense in $(0, 1]$	12
1.3 Trifling Set Properties	12
1.4 Trifling Sets In Base $r$	13
1.5 Cantor Set Is Trifling, Uncountable, and Perfect	16
1.6 Alternate $S_n$ Integral Value Proof	17
1.7 Vieta's Formula	19
1.8 Differences between the Weak and Strong Law Is Uniform Convergence	21
1.9 Nowhere Dense Set Existence and Properties	22
1.10 Normal Numbers are of the First Category	23
1.11 Diophantine Approximation Properties	24
<b>Section 2 - Probability Measures</b>	<b>26</b>
Notes	26
Problems	33
2.1	33
2.2 Union and Intersection Equality Property	34

2.3 Equivalent and Non Equivalent Field Definitions . . . . .	35
2.4 Unions of increasing Fields are Fields, but not true for $\sigma$ fields . . . . .	35
2.5 Generated Field and Explicit Definition . . . . .	35
2.6 Comparing Fields and $\sigma$ fields . . . . .	35
2.7 Extending a field/ $\sigma$ field by $\{H\}$ . . . . .	36
2.8 $\sigma$ algebra for class where $A^c$ is countable union of class elements . . . . .	36
2.9 Equivalent $\sigma$ algebra definition . . . . .	36
2.10 Classes Generating The Discrete Field are Large . . . . .	37
2.11 Countably Generated Fields . . . . .	38
2.12 Cardinality of Fields and $\sigma$ Fields . . . . .	39
2.13 Probability Measure on Finite and CoFinite Sets . . . . .	41
2.14 First Category $\sigma$ Field . . . . .	42
2.15 Differences between $B_0$ and $C_0$ . . . . .	43
2.18 Stochastic Arithmetic . . . . .	44
2.19 Nonatomic Probability Spaces . . . . .	50
2.21 Generating Sigma Algebras "From the Inside" . . . . .	53
2.22 . . . . .	54
<b>Section 3 - Existence and Extension</b>	<b>55</b>
Notes . . . . .	55
Construction of the Extension . . . . .	55
Uniqueness of the Extension and the $\pi - \lambda$ Theorem . . . . .	59
Lebesgue Measure on the Unit Interval . . . . .	62
Problems . . . . .	65
3.1 Finite vs. Countable Additivity . . . . .	65
3.2 Redefining the Inner and Outer Measure . . . . .	66
3.3 Countable Additivity and the Outer Measure . . . . .	69
3.6 Extension for a Finitely Additive Probability Measure on a Field . . . . .	70
3.7 Finitely Additive Field Extension . . . . .	73
3.8 Finitely Additive Field Extension to Power Set Field . . . .	75

## Forward

This document will contain notes and solutions corresponding to Probability and Measure, Third Edition, by Patrick Billingsley [amazon].

A note on how I will do the questions. I want to answer every question,

but there really are a lot of them. So, I think I will tackle them this way. To get through a chapter, I'll go through every question that is in the back of the book (as they are the most important, and might be needed at a later time). Then, I'll go maybe over 3-4 more. Then, I'll answer one question I skipped from the previous chapters.

## Section 1 - Borel's Normal Number Theorem

### Notes

For a complete understanding of probability, you need to understand an infinite number of events as well as a finite number of events. We try and present why that must be so here.

### The Unit Interval

We take the length of an interval  $I = (a, b] = b - a$ . Note, for  $A$  a disjoint set of intervals in  $(0, 1]$ , we have that  $P(A)$  is well defined. If  $B$  is a similar disjoint set, and is disjoint from  $A$ ,  $P(A + B) = P(A) + P(B)$  is well defined as well. Note - we haven't defined anything for intersections yet. These definitions can also directly stem from the Riemann integral of step functions.

The unit interval can give the probability that a single particle is emitted in a unit interval of time. Or a single phone call comes in. However, it can also model an infinite coin toss. This is done as follows - for  $\omega \in (0, 1]$ , define:

$$\omega = \sum_{n=1}^{\infty} \frac{d_n(\omega)}{2^n}$$

Where  $d_n(\omega)$  is 0 or 1, and comes from the binary expansion of  $\omega$ . We take  $\omega$  as the non terminating representation. Note, we were particular when we defined intervals as half inclusive. Examine the set of  $\omega$  for which  $d_i(\omega) = u_i$  for  $i = 1, \dots, n$ ,  $u_i \in \{0, 1\}$ . We have that:

$$\sum_{i=1}^n \frac{u_i}{2^i} < \omega \leq \sum_{i=1}^n \frac{u_i}{2^i} + \sum_{i=n+1}^{\infty} \frac{1}{2^i}$$

We cannot have the lower extreme value, as this would imply  $\omega$  takes on its terminating binomial representation, which is what we said we would not do. This is our first taste, I guess, of measure 0 sets, we we still have:

$$\mathbb{P}[\omega : d_i(\omega) = u_i, i = 1, \dots, n] = \frac{1}{2^n}$$

Note, probabilities of various familiar events can be written down immediately. Ultimately, note, however, each probability is the sum of disjoint dyadic intervals of various ranks  $k$ . Ie, all the events are still well defined by our probability definition above. We have:

$$\mathbb{P} \left[ \omega : \sum_{i=1}^n d_i(\omega) = k \right] = \binom{n}{k} \frac{1}{2^n}$$

All these results have been for finitely many components of  $d_i(\omega)$ . What we are interested in, however, is properties of the entire sequence of  $\omega = (d_1(\omega), d_2(\omega), \dots)$ .

## The Weak Law of Large Numbers

What I like about this chapter, is to me - it *emphasizes* the connection between the *structure of real numbers*, and probability. At the end of the day - probability can be seen as just extracting properties of *frequency* over the real numbers, to be understood as probabilistic statements. However, with just our basic real numbers - we can't really prove a lot of properties about infinite things. That is when measure theory comes in later. However, for now, we look at what we can prove - and that starts with the weak law of large numbers. We have:

**Theorem 1.1 - The Weak Law of Large Numbers** For each  $\epsilon$ :

$$\lim_{n \rightarrow \infty} \mathbb{P} \left[ \omega : \left| \frac{1}{n} \sum_{i=1}^n d_i(\omega) - \frac{1}{2} \right| \geq \epsilon \right] = 0$$

Probabilistically - this is saying that if  $n$  is large, then there is a small probability that the fraction/relative *frequency* of heads in  $n$  tosses will deviate much from  $1/2$ . Think about it as a statement over the real numbers as well - it is also interesting. Ultimately, the intervals containing  $\omega$  that do not satisfy the above are getting smaller and smaller and smaller. We formalize this with the following concept:

As  $d_i(\omega)$  are constant over each dyadic interval of rank  $n$  if  $i \leq n$ , the sums  $\sum_{i=1}^n d_i(\omega)$  are also constant over rank  $n$ . Thus, the set in the theorem is just a disjoint union of dyadic intervals of rank  $n$ . Note - the theorem is saying, that the total weight given to those intervals gets smaller and smaller as  $n$  goes to infinity.

Now, we go over how to prove the theorem. It relies on rademacher variables:

$$r_n(\omega) = 2d_n(\omega) - 1$$

These are  $\pm 1$  when  $d_n = 1/0$ . Note, these have the same "being constant on dyadic intervals" properties as  $d_n(\omega)$ . We define:

$$s_n(\omega) = \sum_{i=1}^n r_i(\omega)$$

And so, our theorem is equivalent to proving:

$$\lim_{n \rightarrow \infty} \mathbb{P} \left[ \omega : \left| \frac{1}{n} s_n(\omega) \right| \geq \epsilon \right] = 0$$

Note, rademacher functions also have interpretations, probabilistically, of random walks and such. With these variables, we can ultimately find properties, going all the way to:

$$\int_0^1 s_n^2(\omega) d\omega = n$$

However, what interests me is the following: Chebyshev's Lemma, but as a property of the real numbers. We have:

**Lemma - Chebyshev's Inequality** If  $f$  is a nonnegative step function, then  $[\omega : f(\omega) \geq \alpha]$  is for  $\alpha > 0$  a finite union of intervals, and:

$$\mathbb{P}[\omega : f(\omega) \geq \alpha] \leq \frac{1}{\alpha} \int_0^1 f(\omega) d\omega$$

Proof: Note, it is all just properties of step functions. Let  $c_j$  correspond to the step intervals  $(x_{j-1}, x_j]$ , and let  $\sum'$  be the sum over  $c_j \geq \alpha$ . Then, we have quite easily:

$$\int_0^1 f(\omega) d\omega = \sum c_j(x_j - x_{j-1}) \geq \sum' c_j(x_j - x_{j-1}) \geq \sum' \alpha(x_j - x_{j-1}) = \alpha \mathbb{P}[\omega : f(\omega) \geq \alpha]$$

Thus, we have Chebyshev's inequality, and with it, we can easily prove the Weak Law of Large Numbers. However - it is important to note - these are *properties over the real numbers*, as much as they are probabilistic properties.

## The Strong Law of Large Numbers

Just to first formalize some terms - the frequency of 1 in  $\omega$  is  $\sum_{i=1}^n d_i(\omega)$ , the relative frequency is that number normalized, ie  $\frac{1}{n} \sum_{i=1}^n d_i(\omega)$ , and the asymptotic relative frequency is the limit. We can derive, with some technical tools outside of discrete probability theory, results on the set:

$$N = \left[ \omega : \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n d_i(\omega) = 1/2 \right]$$

We call this the set of normal numbers  $N$ . The tools themselves are the concepts of negligibility. A set  $A$  is negligible if for every  $\epsilon > 0$ , there is a countable number of intervals (not necessarily disjoint) such that:

$$A \subset \bigcup_k I_k \quad \sum_k I_k = \sum_k b_k - a_k < \epsilon$$

For one - I like to note here interpretations. Essentially - if  $A$  is negligible, it is a practical impossibility that  $\omega$  randomly drawn will lie within  $A$ . And if  $A^c$  is negligible, it is a practical certainty that  $\omega$  randomly drawn will lie within  $A$ . These are just how they should be understood - and these understandings are reasonable, as the total "length" that  $A$  takes up can be understood to be incredibly incredibly small.

Some properties of negligibility - note, these are the standard properties, stemming from infinite sums  $(1/2^k)$  summing to values less than  $\epsilon$ . Individual points are negligible, and so to thus are countable sets. So to are countable unions of countable sets.

With these properties - we understand that the property of our model not including  $\omega$  with a terminating sequence (all 0 ending) is not a short coming. These  $\omega$  form a countable set - and so, they can be considered negligible.

**Theorem 1.2** The set of normal numbers  $N$  has negligible complement.

**Proof** As an aside - we note that this proof is stronger than just the negligibility properties we noted above. This is because  $N^c$  is not countable. The set of  $d_i(\omega) = 1$  unless  $i$  is a multiple of 3 clearly belongs to  $N$  - as for each  $n$ ,  $n^{-1} \sum_{i=1}^n d_i(\omega) \geq 2/3$ . However, note this set is uncountable (diagonalization argument).

Note, the proof relies on equivalently defining  $N$  as:

$$N = \left[ \omega : \lim_{n \rightarrow \infty} \frac{1}{n} s_n(\omega) = 0 \right]$$

Then, we can again make use of Chebyshev's Inequality (step function version) to find that:

$$\mathbb{P}[\omega : |s_n(\omega)| \geq n\epsilon] \leq \frac{1}{n^4 \epsilon^4} \int_0^1 s_n^4(\omega) d\omega = \frac{n + 3n(n-1)}{n^4 \epsilon^4} \leq \frac{3}{n^2 \epsilon^4}$$

Where the last step is just via an in depth (but simple) investigation of the integrals of multiplications of rademacher variables. With this property, we can find that if  $A_n = [\omega : |n^{-1} s_n(\omega)| \geq \epsilon_n]$ , then we have a sequence of  $\epsilon_n$  such that  $P(A_n) \leq 3\epsilon_n^{-4} n^{-2}$ , and we can find such a sequence such that:

$$\sum_n \mathbb{P}[A_n] < \infty$$

The final step to proving the theorem is noting that:

$$\bigcap_{n=m}^{\infty} A_n^c \subset N \implies N^c \subset \bigcup_{n=m}^{\infty} A_n$$

Which will ultimately prove the theorem. Note - a lot of details are left out, but I do not consider them important. You should be able to fill in. These are just the major strokes, outlining the proof. It essentially hinges on our integral value, and the relationship between  $A_n$  and the set of normal number  $N$ . qed.

So, we have  $N^c$  is negligible. But, can we have that  $N$  itself is negligible? Well, we could say no - using our "practically impossible" notions, and noting that for  $\omega \in [0, 1]$  randomly drawn, it must be in  $[0, 1]$ , and  $N^c \cup N = [0, 1]$ . But, that is not rigorous. And so, the following theorem will give us our initial basis of *measure*, and also help us note that  $N$  is not negligible.

### Theorem 1.3 - Lebesgue Measure Starting Point

1. If  $\bigcup_k I_k \subset I$ , and the  $I_k$  are disjoint, then  $\sum_k |I_k| \leq |I|$
2. If  $I \subset \bigcup I_k$  (the  $I_k$  need not be disjoint), then  $|I| \leq \sum_k |I_k|$
3. If  $\bigcup I_k = I$ , and the  $I_k$  are disjoint, then  $|I| = \sum_k |I_k|$

Note, this Theorem is true for countably infinite intervals as well. **Proof:** Note that the third part follows directly from (1) and (2). We start with the finite cases. For (1), we can prove by induction on the number of intervals  $n$ . It is clearly true for  $n = 1$ , and it is a fairly simple induction hypothesis to prove in general. We similarly have the same for (2).

The difficult part comes when going to infinite intervals. For (1), it is a simple limit, ie:

$$\sum_k |I_k| = \lim_{n \rightarrow \infty} \sum_{k=1}^n |I_k|$$

Note, each sum is less than  $|I|$ , as the finite case to 1 applies for each finite sum. And so, the inequality can be expanded to the limit. However - we can't do that for (2). Ultimately, the difference between the two cases is the inclusion of unions. We note:

$$\bigcup_k I_k \subset I \implies \bigcup_{k=1}^n I_k \subset I$$

Ie, the inclusion is true for every subset. However, we *do not* necessarily have:

$$I \subset \bigcup_k I_k \implies I \subset \bigcup_{k=1}^n I_k$$

Note that in the following way:  $I = (a, b]$ . We have that  $I_i = (a + 1/i, b]$ . We do indeed have that:

$$I \subset \bigcup_k I_k$$

As if you take  $x \in I$ ,  $a < x \leq b$ , and so we must have for  $i$  large enough,  $a + 1/i < x \leq b$ , and so  $x \in I_i$ . However, note that the inclusion is not actually true for a specific finite subunion. So, we need to take a different strategy to prove the infinite case. This comes from dealing with *open covers of compact spaces*, and relying on the Heine-Borel theorem, which says that intervals  $[a, b]$  are indeed compact. In this case, we are able to bridge between infinite unions and finite unions - as we can take a finite sub cover of an open cover on compact spaces. We prove the theorem essentially for  $[a + \epsilon, b]$ , that:

$$|I| - \epsilon = b - (a + \epsilon) \leq \sum_k |I_k| + \epsilon$$

However, as the  $\epsilon$  is arbitrary, we can conclude the fact for the infinite case as well. qed.



Note - this implies that  $N$  is not negligible. As it was,  $[0, 1]$  would be negligible, but that is incorrect by the above, as any open covering must have total sum at least 1, ie, the total sum is not smaller than arbitrary  $\epsilon$ .

## The Measure Theory of Diophantine Approximation

This section is just additional, so my notes here are sparse. However, I do read through it, and record the theorems, plus some notes I have on them.

**Theorem 1.4** If  $x$  is irrational, there are infinitely many irreducible fractions  $p/q$  such that:

$$\left| x - \frac{p}{q} \right| < \frac{1}{q^2}$$

Honestly, this proof is so good. I like it a lot - it is pretty clever. However, I don't just want to copy it down here - it is in the book. I'm not sure if there is any broad message I can glean from it - just that, it is a property of the real numbers. It just hinges on the following fact (which itself is pretty difficult to prove), that for every  $Q$  positive integer, there is an integer  $q < Q$  and corresponding  $p$  such that:

$$\left| x - \frac{p}{q} \right| < \frac{1}{qQ} \leq \frac{1}{q^2}$$

Note, this is true for  $x$  rational or irrational. However, we have an infinite number of such irreducible fractions for the irrational case, and the contradiction derived in the book is nice as well. Anyway - read the book for this. qed.

Anyway, the above essentially means that, apart from a negligible set of  $x$ , each real number has an infinite set of irreducible rationals such that the bounds in Theorem 1.4 are true. We now consider a generalization - when can we tighten the inequality in Theorem 1.4 - Consider:

$$\left| x - \frac{p}{q} \right| < \frac{1}{q^2 \varphi(q)}$$

Let  $A_\varphi$  consist of the real  $x$  for which the above has infinitely many irreducible solutions. Under what conditions on  $\varphi$  will  $A_\varphi$  have negligible complement? Note that if  $\varphi(q) < 1$ , then the condition is weaker than Theorem 1.4, and so  $A_\varphi$  has negligible complement immediately. It becomes interesting if  $\varphi(q) > 1$ . We will later prove the theorem:

**Theorem 1.5** Suppose that  $\varphi$  is positive and nondecreasing. If:

$$\sum_q \frac{1}{q\varphi(q)} = \infty$$

Then  $A_\varphi$  has negligible complement. We will prove this later, but we can now prove:

**Theorem 1.6** Suppose that  $\varphi$  is positive. If

$$\sum_q \frac{1}{q\varphi(q)} < \infty$$

Then  $A_\varphi$  is negligible.

We will go over the proof soon for this theorem. However - just note what the theorems are saying. Note that in the second -  $\varphi(q)$  must be growing quite quickly. We need the denominator to be quite large, so that the infinite sum is ultimately finite. However, in theorem 1.5, we don't want the  $\varphi(q)$  to be too large, lest the sum actually does become finite. Ultimately - both theorems are conditions on how  $\varphi(q)$  grows. Which, ultimately does make sense. If  $\varphi(q)$  grows to large - it becomes unreasonable to expect our condition to hold infinitely many times. If I ever encounter such situations, where I might want to examine the growth of a function  $\varphi$  - I think examining whether the infinite sum of  $1/\varphi$  equals infinity or not is often a good property that is related to the growth of a function.

**Proof of Theorem 1.6** I'll give the full proof here, as it is interesting to me, and rather short. We want to show that  $A_\varphi$  is negligible. Well, given that the sum is finite, there is a  $q_0$  large enough such that the tail sum  $\sum_{q \geq q_0} \frac{1}{q\varphi(q)} < \epsilon/4$ . If  $x \in A_\varphi$ , then our definition holds for some  $q \geq q_0$ , and as  $0 < x < 1$ , we have that the corresponding  $p$  lies in  $0 \leq p \leq q$ . Thus, we have that:

$$A_\varphi \subset \bigcup_{q \geq q_0} \bigcup_{p=0}^q \left( \frac{p}{q} - \frac{1}{q^2\varphi(q)}, \frac{p}{q} + \frac{1}{q^2\varphi(q)} \right]$$

Which stems from every  $x \in A_\varphi$  being in the right expression, given that  $x$  is within one of the intervals on the right by the property we just described. Now, we have a covering interval - we just need to find the length of it. Note, by assumption, we have all of our  $q$  must satisfy  $q \geq 1$  (or, we can add that in). And so, we have the sum of the intervals is:

$$\sum_{q \geq q_0} \sum_{p=0}^q \frac{2}{q^2\varphi(q)} = \sum_{q \geq q_0} \frac{2(q+1)}{q^2\varphi(q)} \leq \sum_{q \geq q_0} \frac{2(q+q)}{q^2\varphi(q)} \leq \sum_{q \geq q_0} \frac{4}{q\varphi(q)} < \epsilon$$

And thus,  $A_\varphi$  is negligible. qed.

## Problems

### 1.1 Infinite Independent Events on a Discrete Space (are impossible)

1. As for why the existence of an infinite sequence of independent events each with probability  $1/2$  in a discrete probability space would make the section superfluous - I think this is because, in the section, we *rely* on the uncountability of the real numbers. This allows us to make notions like negligible, which helps us make Borel's Number Theorem (the Strong Law of Large numbers). We could then just handle infinite cases with a countable, discrete space - which would make the section unnecessarily in depth ("superfluous").

As for why a discrete space cannot have an infinite sequence of independent events. Note, we can partition the space into sets  $A_1 \cap A_2$ ,  $A_1 \cap A_2^c$ ,  $A_1^c \cap A_2$ , and  $A_1^c \cap A_2^c$ . Note that each has probability  $2^{-2}$ , by independence. And so, each countable point  $\omega$  belongs to one of these sets, and  $P(\omega) \leq 2^{-2}$ . We can continue on for arbitrary  $2^k$  partitions, each of probability at most  $2^{-k}$ . Thus, we find that  $P(\omega) = 0$  for each point. This is a contradiction, as:

$$\sum P(\omega) = 1 \neq 0 = \sum_{\omega} 0 = \sum P(\omega)$$

2. This portion draws the same contradiction as above, namely,  $P(\omega) = 0$  for all  $\omega$ . Each  $\omega$  belongs to a sequence of  $A_1, A_2, A_3^c$ , something like that. Let  $t_i = p_i/1 - p_i$  that corresponds to  $\omega \in A_i$  or  $\omega \in A_i^c$ . We find:

$$P(\omega) \leq \prod_{i=1}^n t_i \leq \exp \left( - \sum_{i=1}^n (1 - t_i) \right)$$

Where the second step notes a property for  $t_i \in [0, 1]$ . Note, it is clear that the above is bounded by:

$$\leq \exp \left( - \sum_{i=1}^n \alpha_i \right)$$

If  $\sum_n \alpha_n$  diverges, the above goes to 0, and we can conclude:

$$P(\omega) = 0$$

Which again, draws out our contradiction. qed.

## 1.2 Normal Numbers and Complements are Dense in $(0, 1]$

Show that  $N$  and  $N^c$  are dense in  $(0, 1]$ . Recall, the definition of *dense* is that  $N$  is dense in  $(0, 1]$  if for each  $x \in (0, 1]$ , and each interval  $J$  containing  $x$ , there is a  $y \in N$  such that  $y \in J$ .

Take  $\omega \in (0, 1]$ . We note  $\omega$  has some form:

$$(d_1(\omega), d_2(\omega), \dots)$$

Note, the problem is equivalent to saying that if  $\omega \in N$ , can we find  $x \in N^c$  arbitrarily close to  $\omega$ , and vice versa, for  $\omega \in N^c$  and  $x \in N$ . We first assume  $\omega \in N$ . We can easily find an  $x \in N^c$ , such that  $x$  is arbitrarily close to  $\omega$ . Just take the first  $k$  elements matching, so that we are within  $\frac{1}{2^k}$  of  $\omega$ , and continue with ones. Clearly, such an  $x$  can be arbitrarily close to  $\omega$ , and within  $N^c$ . So,  $N^c$  is clearly dense.

For the other direction, take  $\omega \in N^c$ , and now, again, match  $x$  on the first  $k$  elements of the binary expansion. For the remainder, oscillate between 1 and 0. Clearly,  $x \in N$ , and arbitrarily close to  $\omega$ . qed.

## 1.3 Trifling Set Properties

**Definition:** Define a set  $A$  to be *trifling* if for each  $\epsilon$  there exists a *finite* sequence of intervals  $I_k$  satisfying that they cover  $A$  and interval sum less than  $\epsilon$ . Recall, from Calculus on Manifolds - this is essentially content 0.

1. A trifling set is also negligible. This must be clear - take the remaining infinite intervals as ones that sum up to less than a small enough  $\epsilon'$ .
2. Show that the *closure* of a trifling set is also trifling. Recall, the closure is all points that are not exterior to  $A$  - exterior meaning that they have open neighborhoods not intersecting  $A$ . Well, take a finite covering less than  $\epsilon/2$  of  $A$ . We define:

$$A \subset \bigcup_{k=1}^n I_k \quad \sum_{k=1}^n I_k < \frac{\epsilon}{2} \quad I'_k = \left( a_k - \frac{\epsilon}{2^{k+2}}, b_k + \frac{\epsilon}{2^{k+2}} \right]$$

$$\implies \sum_{k=1}^n I'_k < \epsilon$$

Note that  $I'_k$  covers  $\overline{A}$ . Take  $x \in \overline{A}$ . By definition, we have  $(x - \frac{\epsilon}{2^{n+2}}, x + \frac{\epsilon}{2^{n+2}})$  intersects  $A$ , and so coincides with some  $I_k$ , and so must be contained within  $I'_k$ . Thus, it is clear that  $I'_k$  covers  $\overline{A}$ , and so  $\overline{A}$  is trifling as well.

3. The rationals in  $(0, 1]$  are bounded and negligible (being countable), but not trifling. Assume we have a covering of the rationals that is finite and sums to less than  $\epsilon$ . Note, we can take the covering to be restricted to  $(0, 1]$ , as all the rationals are in  $(0, 1]$ . For  $\epsilon < 1$  - Theorem 1.3.2 implies that these intervals do not cover all of  $(0, 1]$ . Note, if they don't cover a rational, we are done. We now note that these sets must not cover some interval of non negligible length - if they covered every such interval, their sum would be 1. This interval contains a rational, which contradicts the set being covering.
4. Show that the closure of a negligible set may not be negligible. Again, the closure of the rationals in  $(0, 1]$  is  $(0, 1]$ , which is not negligible.
5. Show that finite unions of trifling sets are trifling, but that this can fail for countable unions. Fail for countable unions - take the union of each rational, which is a countable union of trifling singleton sets - by the above, this is negligible, and not trifling. Note, for finite unions of trifling sets - say  $k$  such sets - take the covering of size  $\frac{\epsilon}{2^k}$ , and note that the union of these intervals is a finite covering of total length less than  $\epsilon$ .

#### 1.4 Trifling Sets In Base $r$

1. First thing to note. We can look at  $A_r(i)$  as iteratively removing intervals from  $(0, 1]$ , where step  $k$  corresponds to removing the numbers whose expansions do not contain  $i$  for the first  $k - 1$  digits, but contain  $i$  at digit  $k$ . At step  $k$ , we remove  $(r - 1)^{k-1}$  intervals (corresponding to the  $r - 1$  possible digits in the first  $k - 1$  spaces) of length  $\frac{1}{r^k}$  (corresponding to the length of the interval starting with  $i \in [r - 1]$  in the  $k$ th entry going to  $i + 1$  in the  $k$ th entry). We find, that the total length of the disjoint intervals removed is:

$$\sum_{k=1}^{\infty} \frac{(r-1)^{k-1}}{r^k} = \frac{1}{r} \sum_{k=1}^{\infty} \frac{(r-1)^{k-1}}{r^{k-1}} = \frac{1}{r} \sum_{k=0}^{\infty} \frac{(r-1)^k}{r^k} = \frac{1}{r} * \frac{1}{1/r} = 1$$

Where the second to last step is the sum of a geometric series. So, at the very least, we have that it could be possible that  $A_r(i)$  is trifling.

Here is how it is trifling. We *know* that a finite amount of points is trifling. As the above sum equals 1 - if we go far enough, the amount removed will be arbitrarily close to 1. Say, within  $\epsilon/2$  of 1. And so, the remaining *intervals* that are uncovered must have at most a total

length of  $\epsilon/2$ . We can cover those intervals with the intervals themselves. Frankly, I think that is enough. Note, of course, at each step we remove an interval that looks like  $(a, b]$ . And so, the remaining intervals should be of the form  $(c, d]$  as well. Everything should be nice, as the intervals in our iteration are disjoint. I literally think that is it. qed.

2. We want to find a trifling set  $A$  such that every point in the unit interval can be represented in the form  $x + y$  with  $x$  and  $y$  in  $A$ .
3. Let  $A_r(i_1, \dots, i_k)$  consist of the numbers in the unit interval in whose base  $r$  expansion the digits  $i_1, \dots, i_k$  nowhere appear consecutively in that order. Show that it is *trifling*.

The first observation I have made: if we have that  $i_1, \dots, i_k$  are all equal, whereas  $j_1, \dots, j_k$  are an arbitrary sequence of digits, we have that:

$$|A_r(j_1, \dots, j_k)| \leq |A_r(i_1, \dots, i_k)|$$

The reason is the following: for the first  $n$  digits of the base  $r$  expansion, there are more numbers without  $i_1, \dots, i_k$  appearing consecutively than there are numbers without  $j_1, \dots, j_k$  appearing consecutively. Consider the following example: Base 3, with  $n = 3$ . We have the following possible sequences:

000	001	002	010	011	012	020	021	022
100	101	102	110	111	112	120	121	122
200	201	202	210	211	212	220	221	222

Consider the count of sequences above without 11. There are 22 such sequences. Now, count the sequences without 12. There are 21 such sequences. Note, above, we have that the sequence 111 has 11 at the start, and 11 at the end, but only takes up one entry. However, we have that 12X and Y12 can never be the same, and so there are two such sequences taken up. We can expand this concept in general - when  $i_1, \dots, i_k$  are all equal, we get the most *collisions* between sequences with digits  $i_1, \dots, i_k$  starting at the possible  $n - k + 1$  starting points. And so, if we find that  $A_r(i_1, \dots, i_k)$  is trifling for  $i_1 = \dots = i_k$ , we can conclude that  $A_r(j_1, \dots, j_k)$  is trifling in general.

To be honest, I am going to skip this one, because I am getting nowhere.

However, here is the work I've done so far, for what it is worth. I have made the following definitions:

$S_{k,n} = \{\text{The length } n \text{ sequences that contain the digit } d \text{ repeated } k \text{ times}\}$

$A_{t,n} = \{\text{The length } n \text{ sequences where } d \text{ repeated } k \text{ times first appears at pos } t\}$

And so, with these definitions, we have:

$$S_{k,n} = \bigcup_{t=1}^{n-k+1} A_{t,n} \implies |S_{k,n}| = \sum_{t=1}^{n-k+1} |A_{t,n}|$$

Where the first step is just definitional, as if  $d$  is repeated  $k$  times, that subsequence first starts at position 1, 2, or up to position  $n - k + 1$ . The second step comes from noting that the sets  $A_{t,n}$  are disjoint - if the sequence first starts at position  $t$ , it *does not* start at position  $t' \neq t$ . And so, if we can find that  $\lim_{n \rightarrow \infty} \frac{|S_{k,n}|}{r^n} = 1$ , then for  $n$  large enough, it will equal  $1 - \epsilon$ , and we can take a *finite* number of intervals to cover the remaining intervals of total length  $\epsilon$  that are not represented in the *finite* union of intervals  $\bigcup_{t=1}^{n-k+1} A_{t,n}$ . The difficult part is actually finding what the above is in terms of numbers. However, I now note that:

$$|A_{t,n}| = r^{n-(t+k-1)} \cdot 1^k \cdot (r-1) \cdot (r^{t-2} - |S_{k,t-2}|)$$

This comes from examining what each of the possible digits in our expansion of  $x \in A_{t,n}$  can be. The remaining  $n - (t + k - 1)$  after the digits  $d$  repeated  $k$  times starting at position  $t$  can be anything we want. This gives us our first element in the product.  $1^k$  refers to the  $k$  digits starting at  $t$  must all be  $d$ .  $(r-1)$  refers to position  $t-1$  - that *must* be any number other than  $d$ . If it is  $d$ , then we get  $x \in A_{t-1,n}$ , which is incorrect. Finally, the remaining first  $t-2$  entries can be any sequence at all, except for a sequence of  $d$  repeated  $k$  times. The count of those sequences is removed from the total  $r^{t-2}$  sequences possible. And so, we find that essentially, both sides are equal. If we make any sequence described on the right side, we have that it is within  $A_{t,n}$ . And, any sequence in  $A_{t,n}$  can be described on the right side. And so now, the work remains to just simplify the calculation of  $|S_{k,n}|$ . Note, we could actually calculate this value by a recursive algorithm. That might help us.

Big Note: The calculation of  $|A_{t,n}|$  assumes somethings, like the existence of position  $t-1$  (which is not there if  $t=1$ ) or  $t > 2$  for  $r^{t-2}$ .

Just make sure to keep these exceptions in mind. Anyway. We make a hand wavy assumption - that  $r^n - |S_{k,n}| > r^{k-1}$ . Note,  $r$  is some base  $n$  digit, and so  $r^{k-1}$  is just some constant. And while we want to eventually prove that the ratio is equal to 1 in limit - ultimately, there will always be some constant distance between  $r^n$  and  $|S_{k,n}|$ . And this constant will continue to grow to infinity. Anyway, for  $n$  large enough, I think it is clear. And so, being hand wavy, and including all terms, although they might not be present, we have:

$$\begin{aligned}\lim_{n \rightarrow \infty} \frac{|S_{k,n}|}{r^n} &\approx \lim_{n \rightarrow \infty} \frac{\sum_{t=1}^{n-k+1} r^{n-(t+k-1)} \cdot (r-1) \cdot (r^{t-2} - |S_{k,t-2}|)}{r^n} \\ &= \lim_{n \rightarrow \infty} \frac{r^{n-k+1}(r-1) \sum_{t=1}^{n-k+1} r^{-t} \cdot (r^{t-2} - |S_{k,t-2}|)}{r^n} \\ &= r^{-k+1}(r-1) \lim_{n \rightarrow \infty} \sum_{t=1}^{n-k+1} r^{-t} \cdot (r^{t-2} - |S_{k,t-2}|)\end{aligned}$$

Now, making use of our hand waviness, we have that:

$$\geq r^{-k+1}(r-1) \lim_{n \rightarrow \infty} \sum_{t=1}^{n-k+1} r^{-t} r^{k-1} = (r-1) \lim_{n \rightarrow \infty} \sum_{t=1}^{n-k+1} r^{-t}$$

Now, we can make use of our geometric series, and have that the above equals:

$$= (r-1) * \frac{1}{r-1} = 1$$

And so yes, the limit does indeed equal 1. Unraveling everything we said above, this allows us to conclude that  $A_r(i_1, \dots, i_k)$  is indeed trifling. Now, there is a saying, that a monkey typing at random for infinity will ultimately write Shakespeare. Well, we can let every word be a digit in some base 10 million language. Ultimately, the probability that a monkey *does not* type our specific Shakespeare sequence is indeed 0, as the amount of "worlds" where the monkey types at random, but does not hit our  $A_r(i_1, \dots, i_k)$  digit sequence is 0.

### 1.5 Cantor Set Is Trifling, Uncountable, and Perfect

1. The Cantor set  $C$  can be defined as the closure of  $A_3(1)$ . Show that  $C$  is uncountable but trifling. First, note uncountable. This can be done by the diagonalization argument. Take any list of numbers in  $A_3(1)$ , which are also in  $C$ . We can make a new number in  $A_3(1)$ , but not in



the list, by taking the  $i$ th entry, and switching the 0 for 2 or 2 for 0. Thus,  $A_3(1)$  is clearly uncountable, and so must be  $C$ .

Now, we note that  $A_3(1)$  is trifling. Take any finite covering of  $A_3(1)$  by intervals with total length less than  $\epsilon$ , then covers  $A_3(1)$ . Note that we can extend each interval by some length of  $\epsilon/2^{i+3}$  on each edge, and this should cover all of  $C$  as well. This is because any element within the closure of  $A_3(1)$ , can also be viewed as within the closure of all the intervals, and so we can extend the interval lengths a bit. Note, this would apply for every trifling set (in  $\mathbb{R}$ , not sure about  $\mathbb{R}^n$  with covering half open rectangles, but I think the idea could be extended).

2. From  $[0, 1]$ , remove the open middle third  $(1/3, 2/3)$ . From the remainder, a union of two closed intervals, removed the open middle thirds  $(1/9, 2/9)$  and  $(7/9, 8/9)$ . Show that  $C$  is what remains when this process continues ad infimum.

Note, this is the standard definition of  $C$ . We have to show this equals our closure definition above. I make the note - the  $n$ th step is the closure of points in  $[0, 1]$  such that the base 3 representation does not contain 1 in the first  $n$  digits. This, I think, is clear. And so, taking the process to infinity, we can conclude that  $C$  is the closure of the set where none of the digits are 1.

3. Show that  $C$  is perfect. A set is *perfect* if it is closed and for each  $x$  in  $A$  and positive  $\epsilon$ , there is a  $y$  in  $A$  such that  $0 < |x - y| < \epsilon$ .

Note,  $C$  is closed, as it is the closure of a set. Now, take  $x \in C$  and  $\epsilon > 0$ . We can find the corresponding  $y$  just by matching say the first  $n$  digits of  $x$ , and then flipping the  $n + 1$  digit from 0 to 2 or vice versa, and then taking any random 0 or 2 for the remaining digit. Note,  $0 < |x - y| < \epsilon$  if  $n$  is large enough. Note - I guess this doesn't apply for the points in the closure. For a point  $x \in C$  in the closure, we can find a  $y$  in  $A_3(1)$  within  $\epsilon/2$  of  $x$ , by the limit definition of the closure. Now, take  $z$  within  $\epsilon/2$  of  $y$  by changing a digit. We now know that  $z \neq x$ , and the property applies. qed.

## 1.6 Alternate $S_n$ Integral Value Proof

We first show the derivative property. We have that:

$$M(t) = \int_0^1 e^{ts_n(\omega)} d\omega$$

We have that  $f(\omega, t) = e^{ts_n(\omega)}$ . Note, we can make use of *Leibnitz's rule* to take the derivative of  $M(t)$  under the integral. See problem 3-32 in Calculus on Manifolds by Spivak. However, this presupposes that  $f$  is continuous. I get around this by the following: note that  $s_n(\omega)$  has only finite points of discontinuity, when we switch from one dyadic interval of rank  $n$  to the next. We can split  $\int_0^1$  so that we ignore those points of discontinuity, and only integrate where  $s_n(\omega)$  is continuous (and constant). Note, the points of discontinuity have content 0, and so the integral on  $[0, 1]$  is equal to the integral on the set not including those points of discontinuity. Thus, we have:

$$M'(t) = \int_0^1 D_2(e^{ts_n(\omega)})(0) d\omega = \int_0^1 s_n(\omega) e^{ts_n(\omega)} d\omega \implies M'(0) = \int_0^1 s_n(\omega) d\omega$$

Now, we can repeat this operation a finite amount of times, successive differentiation under the integral, to clearly find that:

$$M^{(k)}(0) = \int_0^1 s_n(\omega)^k d\omega$$

Now, noting again that  $s_n(\omega)$  is constant on the  $2^n$  dyadic intervals, we have that  $M(t)$  is actually easy to evaluate. For each of those  $2^n$  intervals,  $s_n$  is some sum of the form  $\pm 1 \pm 1 \cdots \pm 1$ , and so we have that:

$$M(t) = \frac{1}{2^n} \sum_{i=1}^{2^n} \exp(t(\pm 1 \pm 1 \cdots \pm 1))$$

We note that this can be broken down with a binomial coefficient. If we have that  $k$  of the  $\pm 1$  are  $-1$ , then the value of the sum is  $n - 2k$ . And so, based off of how many possible sequences of the  $\pm 1$  that contain  $k$   $-1$ , we have that the above can be expressed as:

$$\begin{aligned} &= \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \exp(t(n - 2k)) = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \exp(t(n - k - k)) \\ &= \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \exp(t(n - k) - t(k)) = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \exp(t)^{n-k} \exp(-t)^k \end{aligned}$$

By the Binomial Theorem, the above equals:

$$= \frac{1}{2^n} (\exp(t) + \exp(-t))^n = \left( \frac{e^t + e^{-t}}{2} \right)^n = (\cosh t)^n$$

Where the last step is just an identity. Now, for a new proof of 1.16:

$$\int_0^1 s_n(\omega) d\omega = M'(0) = n \cosh^{n-1}(0) \sinh(0) = n * 1 * 0 = 0$$

Now, for a new proof of 1.18:

$$\int_0^1 s_n^2(\omega) d\omega = M''(0) = n [(n-1) \cosh^{n-2}(0) \sinh^2(0) + \cosh^n(0)] = n * 1 = n$$

Finally, for a new proof of 1.28, we just have to take the fourth derivative, and plug in 0. I will not be going over the steps, but using derivative calculator, we get the fourth derivative at 0 is:

$$\int_0^1 s_n^4(\omega) d\omega = M'''(0) = n(3n-2) = 3n^2 - 2n$$

Which is the final property. qed.

## 1.7 Vieta's Formula

We first find a similar property to the above. We examine:

$$\int_0^1 \exp \left[ i \sum_{k=1}^n a_k r_k(\omega) \right] d\omega$$

We note that the summation is constant on the  $2^n$  dyadic intervals of rank  $n$ . In which case, the integral becomes a summation, that looks like:

$$= \frac{1}{2^n} \sum \exp [i (\pm a_1 \pm a_2 \pm \dots \pm a_n)]$$

Now, we extract the first  $+a_1$  term and  $-a_1$  term from the exponential:

$$= \frac{1}{2^n} \exp(ia_1) \sum \exp [i (\pm a_2 \pm \dots \pm a_n)] + \frac{1}{2^n} \exp(-ia_1) \sum \exp [i (\pm a_2 \pm \dots \pm a_n)]$$

We note that by symmetry, both the summations are equal, and so the above actually equals:

$$= \frac{1}{2^n} (\exp(ia_1) + \exp(-ia_1)) \sum \exp [i (\pm a_2 \pm \dots \pm a_n)]$$

We can continue this process  $n$  times, and then split the  $2^{-n}$ , to find:

$$\int_0^1 \exp \left[ i \sum_{k=1}^n a_k r_k(\omega) \right] d\omega = \prod_{k=1}^n \frac{e^{ia_k} + e^{-ia_k}}{2}$$

Using the cos identity, we find:

$$\int_0^1 \exp \left[ i \sum_{k=1}^n a_k r_k(\omega) \right] d\omega = \prod_{k=1}^n \cos(a_k)$$

We let  $a_k = t2^{-k}$ . We note that  $\sum_{k=1}^{\infty} r_k(\omega)2^{-k} = 2\omega - 1$  - this is because in each entry, we have  $r_k(\omega) = 2d_k(\omega) - 1$ , and  $\omega = \sum_{k=1}^{\infty} d_k(\omega)2^{-k}$ . We can apply the  $2\omega - 1$  operations (as it is continuous and passes the summation limit), to derive the above. We thus have:

$$\lim_{n \rightarrow \infty} \int_0^1 \exp \left[ i \sum_{k=1}^n a_k r_k(\omega) \right] d\omega = \int_0^1 \exp \left[ i \lim_{n \rightarrow \infty} \sum_{k=1}^n a_k r_k(\omega) \right] d\omega = \int_0^1 \exp [ti(2\omega - 1)] d\omega$$

Note, we don't have theorems to pass the limit through the integral yet. However, I believe this will be by the monotone convergence theorem - the partial sums should be non decreasing, given the exponential being nonnegative. Anyway, we don't have to prove that here. We now note this is an integral from 0 to 1 - whose value is:

$$-\frac{ie^{it(2x-1)}}{2t} = \frac{-i^2}{t} \left( \frac{e^{it} - e^{-it}}{2i} \right) = \frac{\sin(t)}{t}$$

Taking the limit on the other side, we can conclude:

$$\frac{\sin(t)}{t} = \prod_{k=1}^{\infty} \cos \frac{t}{2^k}$$

We can now derive Vieta's Formula:

$$\frac{2}{\pi} = \frac{\sqrt{2}}{2} \frac{\sqrt{2 + \sqrt{2}}}{2} \frac{\sqrt{2 + \sqrt{2 + \sqrt{2}}}}{2} \dots$$

We recall the half angle formula for cos:

$$\cos \frac{\theta}{2} = \sqrt{\frac{1 + \cos(\theta)}{2}}$$

We can use this to make an induction argument that:

$$\cos \left( \frac{\pi}{2^{2^k}} \right) = \frac{\sqrt{2 + \sqrt{2 + \dots + \sqrt{2}}}}{2}$$

For the base case of  $k = 1$ , we have:

$$\cos\left(\frac{\pi}{2} \frac{1}{2}\right) = \sqrt{\frac{1 + \cos(\pi/2)}{2}} = \sqrt{\frac{1}{2}} = \frac{\sqrt{2}}{2}$$

Now, for arbitrary  $k$ , we have:

$$\begin{aligned} \cos\left(\frac{\pi}{2} \frac{1}{2^k}\right) &= \cos\left(\frac{1}{2} \frac{\pi}{2} \frac{1}{2^{k-1}}\right) = \sqrt{\frac{1 + \cos\left(\frac{\pi}{2} \frac{1}{2^{k-1}}\right)}{2}} = \sqrt{\frac{1 + \frac{\sqrt{2 + \sqrt{2 + \dots + \sqrt{2}}}}{2}}{2}} \\ &= \sqrt{\frac{2 + \sqrt{2 + \sqrt{2 + \dots + \sqrt{2}}}}{4}} = \frac{\sqrt{2 + \sqrt{2 + \dots + \sqrt{2}}}}{2} \end{aligned}$$

And so, by our identity, we find:

$$\frac{2}{\pi} = \frac{\sin(\pi/2)}{\pi/2} = \prod_{k=1}^{\infty} \cos\left(\frac{\pi}{2} \frac{1}{2^k}\right) = \frac{\sqrt{2}}{2} \frac{\sqrt{2 + \sqrt{2}}}{2} \frac{\sqrt{2 + \sqrt{2 + \sqrt{2}}}}{2} \dots$$

And we have found Vieta's formula. qed.

## 1.8 Differences between the Weak and Strong Law Is Uniform Convergence

A number  $\omega$  is normal in base 2 if and only if for each positive  $\epsilon$  there exists an  $n_0(\epsilon, \omega)$  such that  $|n^{-1} \sum_{i=1}^n d_i(\omega) - 1/2| < \epsilon$  for all  $n$  exceeding  $n_0(\epsilon, \omega)$ . That is just the definition. Theorem 1.2 concerns the entire dyadic expansion (ie, the complement of the set of normal numbers whose infinite sum equals  $1/2$  has probability 0), whereas Theorem 1.1 concerns only the beginning sequence (ie, the limit of probability, of the dyadic expansion numbers whose first  $n$  partial sum is not close to  $1/2$ , is 0). Identify the difference by showing that for  $\epsilon < 1/2$  the  $n_0(\epsilon, \omega)$  above cannot be the same for all  $\omega$  in  $N$  - in other words,  $n^{-1} \sum_{i=1}^n d_i(\omega)$  converges to  $1/2$  for all  $\omega$  in  $N$ , but not uniformly. But see problem 13.9.

Well - noting that for  $\epsilon < 1/2$ , the  $n_0(\epsilon, \omega)$  cannot be the same for all  $\omega$  in  $N$ , means finding  $\omega$  in  $N$  for which the  $n$  is different. We can take just  $\omega$  where the first  $k$  are all 0, and then the remaining digits alternate between 0 and 1. Note, that the first  $n$  for which that sum is actually close to  $1/2$  (when normalized by  $n^{-1}$ ), must increase to  $\infty$ . And so yes, we have convergence to  $1/2$  for  $\omega$  in  $N$ , but not uniform convergence. And so, while Theorem 1.1

can rely on this non uniform convergence (ie, the sets are getting smaller, because ultimately, we reach that  $n$  value for all  $\omega$ ), we cannot rely on that for Theorem 1.2.

Looking at Problem 13.9 - it looks like, however, we might be able to get uniform convergence on subsets of  $N$ . Perhaps this is how we can actually prove Theorem 1.2. Anyway, I like to just note that the underlying difference in the two Theorems comes from an underlying difference in just real number theory - uniform vs non uniform convergence.

### 1.9 Nowhere Dense Set Existence and Properties

1. Show that a trifling set is nowhere dense. A set  $E$  is nowhere dense in  $B$  if each open interval  $I$  contains some interval  $J$  that does not meet  $E$ . Use the previous problems (1.3 b) and theorems (1.3ii) to prove this.

I am going to assume we have a trifling set  $E$  in the real line. We take an open interval of the real line  $I$ . We want to find some interval  $J$  which our trifling set  $E$  does not meet. Assume  $I = (a, b]$ , and so take  $\epsilon = (b - a)/2$ . By trifling, there is a finite interval cover of  $E$  with total length less than  $\epsilon$ . Assume there is no interval  $J$  in  $I$  that  $E$  does not meet. Ie, all subintervals  $J$  meet  $E$ . This means that all subintervals  $J$  are contained within the finite cover. This is because every point inside of  $J$  has an open neighborhood around it that meets  $E$ , and so that open neighborhood meets the interval - and if the neighborhood is small enough, the original point is contained within the interval as well (I guess, this can be, via the closure is trifling as well).

And so, every subinterval of  $I$  is contained any finite open cover of  $E$  - this is a contradiction, as this implies every subinterval of  $I$  is vanishing in length. Some subintervals of  $I$  are of length  $(b - a)/2$ , which is not vanishing. qed.

2. Let  $B = \bigcup_n (r_n - 2^{-n-2}, r_n + 2^{-n-2}]$ , where  $r_1, r_2, \dots$  is an enumeration of the rationals in  $(0, 1]$ . Show that  $(0, 1] - B$  is nowhere dense but not trifling or even negligible.

We first show nowhere dense. Take a subinterval  $I$  of  $(0, 1]$ . Note that  $I$  must contain some rational  $r_n$ , as the rationals are dense. We let  $J$  be one of the intervals from the intersection of  $(r_n - 2^{-n-2}, r_n + 2^{-n-2}] \cap I$ . Note that  $J \subset I$ , and  $J \subset B$ . And so,  $J \cap (0, 1] - B = \emptyset$ . And so  $B$

must be nowhere dense.

We now note that  $(0, 1] - B$ , however, is not negligible (and thus not trifling). We have that:

$$|B| \leq \sum_{n=1}^{\infty} |(r_n - 2^{-n-2}, r_n + 2^{-n-2})| = \sum_{n=1}^{\infty} \frac{1}{2^{n+1}} = 1/2$$

And so, we must have that:

$$|(0, 1] - B| \geq 1 - 1/2 = 1/2$$

If we had any cover, whose total length is less than  $\epsilon < 1/2$ , we would get a contradiction. qed.

3. Take a compact negligible set. Take an infinite open interval cover whose total length is less than  $\epsilon$ . By compactness, a finite subset of those sets covers the compact negligible set as well. And so, the set is trifling.

### 1.10 Normal Numbers are of the First Category

**Definition: Set of the First Category** A set of the first category is a set that is a countable union of nowhere dense sets. This is a topological notion of smallness - similar to first countable, I guess. This is similar to the metric notion of smallness, negligibility. Neither condition implies the other.

1. Show that the non negligible set  $N$  of normal numbers is of the first category by proving that:

$$A_m = \bigcap_{n=m}^{\infty} [\omega : |n^{-1}s_n(\omega)| < 1/2]$$

Is nowhere dense, and  $N \subset \bigcup_m A_m$ . Note, we can prove  $A_m$  is nowhere dense in the following way - if the complement is dense, and each point in that dense set has an interval around it contained in the complement, then  $A_m$  is nowhere dense. This is similar to the above problem. Note, this implies nowhere dense, as every interval  $I$  has a point in the complement (by the complement being dense), and this point has an interval around it that is not contained in  $A_m$ .

We have that:

$$A_m^c = \bigcup_{n=m}^{\infty} [\omega : |n^{-1}s_n(\omega)| \geq 1/2]$$

We note - each point  $\omega \in A_m^c$  has an interval around it in  $A_m^c$ . This is just the dyadic interval -  $\omega \in A_m^c \implies \omega \in [\omega : |n^{-1}s_n(\omega)| \geq 1/2]$ . Every point in the same rank  $n$  dyadic interval as  $\omega$  must also belong to  $A_m^c$ . Now, we just have to find such an  $\omega \in I$ . Note, we can just find some dyadic interval of rank  $n$  that is contained in  $I$ . Then, we can examine:

$$[\omega : |4n+1^{-1}s_{4n+1}(\omega)| \geq 1/2]$$

We can take  $\omega$  such that the first  $n$  bits place  $\omega$  in the dyadic interval of rank  $n$  - then, the remaining bits can be 1, and so  $\omega \in I$ , and  $\omega \in [\omega : |(4n+1)^{-1}s_{4n+1}(\omega)| \geq 1/2] \implies \omega \in A_m^c$ . Note, this is because  $(4n+1)^{-1}s_{4n+1}(\omega) \geq \frac{3n+1-n}{4n+1} = \frac{2n+1}{4n} \geq 1/2$ . Thus,  $A_m^c$  is nowhere dense.

The final part is to prove  $N \subset \bigcup_m A_m$ . Well,  $\omega \in N$  essentially implies for  $m$  large enough,  $\omega \in A_m$ . By the limit definition, we have that at some point,  $\frac{1}{n}s_n(\omega)$  remains within  $\epsilon$  of 0. Just set  $\epsilon = 1/2$ , and we have it.

2. According to a famous theorem of Baire, a nonempty interval is *not* of the first category. Use this fact to prove that the negligible set  $N^c = (0, 1] - N$  is not of the first category.

Proof by contradiction. Assume  $N^c$  is of the first category. Then,  $N$  and  $N^c$  are both of the first category. Note, the countable union of sets of the first category are of the first category (stemming from the countable union of countable sets is still countable), and so this would imply  $[0, 1] = N \cup N^c$  is of the first category as well. This is a contradiction of the Theorem by Baire. Thus,  $N^c$  is not of the first category. qed.

### 1.11 Diophantine Approximation Properties

Prove:

1. If  $x$  is rational, (1.33) has only finitely many irreducible solutions. Recall, (1.33) is the formula:

$$\left| x - \frac{p}{q} \right| < \frac{1}{q^2}$$

Well, we have  $x$  is some irreducible  $\frac{p'}{q}$ . Thus, we have:

$$\left| \frac{p'}{q'} - \frac{p}{q} \right| = \frac{|p'q - q'p|}{q'q}$$



We note that  $|p'q - q'p| \geq 1$ . This is integer multiplication and subtraction, which results in an integer. The only case when this is less than 1 is when it equals 0 - or  $p'q = pq'$ . By different rationals, we can only have one of  $q = q'$  or  $p = p'$ . If we are in that case, divide and not equal. If  $q \neq q'$  and  $p \neq p'$ , then the above is equal only if  $p' = q'$  and  $q = p$  - in which case we are not irreducible. Anyway, this implies that:

$$\left| \frac{p'}{q'} - \frac{p}{q} \right| \geq \frac{1}{q'q}$$

And so, the number of rationals  $\frac{p}{q}$  that satisfy  $\frac{1}{q^2} \geq \frac{1}{q'q}$  is finite.

2. Suppose that  $\varphi(q) \geq 1$  and (1.35) holds for infinitely many pairs  $p, q$  but only for finitely many relatively prime ones. Then  $x$  is rational.

Just as a reminded, we have that (1.35) is:

$$\left| x - \frac{p}{q} \right| < \frac{1}{q^2 \varphi(q)}$$

We have that by Theorem 1.4, if  $x$  is irrational, there are infinitely many irreducible fractions  $p/q$  such that:

$$\left| x - \frac{p}{q} \right| < \frac{1}{q^2}$$

Well, frankly, I don't understand what the "relatively prime" wording here is. Like, if  $x$  is rational - we just set  $p/q$  as  $x$ , and we have infinitely many *reducible* rational fractions that satisfy 1.35. I am just going to take the statement as "finitely many irreducible fractions satisfy 1.35." If  $\varphi(q) = 1$ , it is clear that this implies  $x$  is rational - this is by Theorem 1.4, which says there are infinitely many irreducible solutions for irrational  $x$ .

Now, we consider  $\varphi(q) > 1$ . And, well, what I think we want to note is the following. By part 1 - a rational will always have finitely many solutions, for the  $p/q$  that satisfy  $\frac{1}{q^2 \varphi(q)} > \frac{1}{q'q}$ . So, the only distinction to make is that if  $x$  is *irrational* - well, then:

$$\left| x - \frac{p}{q} \right| < \frac{1}{q^2 \varphi(q)}$$

Either has infinitely many, or 0 solutions. Suppose irrational  $x$  has finitely many solutions to the equation:

$$\left| x - \frac{p}{q} \right| < \frac{1}{q^2 \varphi(q)}$$

I think we can make a contradiction like in Theorem 1.4. Suppose we had finitely many irreducible solutions -  $p_1/q_1, \dots, p_m/q_m$ . Then,  $|x - p_k/q_k|$  is positive, and we can take  $Q$  where  $Q^{-1}$  is smaller than each of the differences. Anyway, I don't actually want to do this.

3. If  $\varphi$  goes to infinity too rapidly, then  $A_\sigma$  is negligible (Theorem 1.6). But however rapidly  $\varphi$  goes to infinity, show that  $A_\varphi$  is nonempty, even uncountable. Note, this one has a solution in the book. I don't want to really go over it - I read the section as it was interesting, but these problems aren't something I'm interested in.

## Section 2 - Probability Measures

### Notes

**Spaces** We first go over  $\Omega$  - a space of points, where each  $\omega \in \Omega$  is a possible result or outcome. What is interesting to me here is - the fact that  $\Omega$  will be interesting from the point of view of *geometry and analysis*, as well as the point of view of probability. In fact, we will probably be able to understand probability properties at times, because of the geometry/analysis properties of the underlying space. Read the section for some examples.

A subset of  $\Omega$  is an *event*, and an element  $\omega \in \Omega$  is a sample point.

**Assigning Probabilities** This section discusses assigning probabilities to events (subsets) in our space  $\Omega$ , in a way such that we will be able to get useful properties out of it. It goes over the unit interval  $(0, 1]$ , and how we have assigned probabilities to disjoint (countable) unions of intervals. We could extend this to say negligible (metric notion) sets are probability 0. However - how can we be sure we have assigned probabilities to all useful sets? The successful procedure, generally, is to assign probabilities to such a large amount of sets, that any set we could possibly think of would be covered (or, not think of, but usefully use).

Another interesting point - can we not just assign a probability to every subset? Well, we will find later, no, we cannot, as this will remove an important property that we need from the spaces - countable additivity. We will go over this later. Given this - we will need to restrict ourselves to a subclass of the class of all subsets of a space.

**Classes of Sets** This section essentially outlines what our class of subsets should and shouldn't have, for it to satisfy some of the properties we noted above, and to be able to express some of the stuff we were doing previously.

We can express the set of normal numbers as a countable union and intersection of disjoint intervals. This is done by expressing the limit definition ( $\epsilon/\delta$ ) in terms of a countable intersection and union. It is pretty simple. However, this tells us - if we want a systematic treatment of section 1 - we need a class of sets that contains the intervals, and is closed under the formation of countable unions and intersections.

A second interesting thing to note is - the singleton  $\{x\}$  is a countable intersection  $\bigcap_n (x - n^{-1}, x]$ . And so, if a class contains singletons, and is closed under *arbitrary unions* - our class is essentially all the subsets of  $\Omega$ . As the theory does not apply to such an extensive class - in the uncountable  $\Omega$  case - our attention needs to be restricted to countable set theory operations.

**Definition - Fields and  $\sigma$  Fields** are sets of subsets of  $\Omega$  with the following properties:

1.  $\Sigma \in F$
2.  $A \in F \implies A^c \in F$
3.  $A, B \in F$  implies  $A \cup B \in F$  for a field, or countable unions for a  $\sigma$  field

Note, via DeMorgan's law, you have that the above definitions are equivalent for finite/countable intersections.

### Examples

1. The finite disjoint unions of subintervals on  $\Omega = (0, 1]$  is a field, denoted  $B_0$ . It is not, however, a  $\sigma$  field, as it doesn't contain singletons (not a subinterval of the form  $(\cdot)$ ), which is a countable intersection of intervals of the form  $(\cdot]$ .
2. Finite and cofinite (ie,  $A^c$  is finite) sets of  $\Omega$  are a field. If  $\Omega$  is finite, then they are a  $\sigma$  field as well (as  $F$  would be all subsets). If  $\Omega$  is not finite -  $F$  is not a  $\sigma$  field.
3. Countable and cocountable sets of  $\Omega$  are a  $\sigma$  field. If  $\Omega$  is uncountable, there are sets that  $F$  misses, however (via the axiom of choice).

**Definition - Generated Sigma Field** For a class of sets  $A$ , the smallest  $\sigma$  field containing  $A$  is called  $\sigma(A)$ . It is the (non empty) intersection of  $\sigma$  fields containing  $A$ . Note, any arbitrary intersection (even uncountable) of  $\sigma$  fields is a  $\sigma$  field - just a definitional property. Some other properties of generated  $\sigma$  fields are:

1.  $A \subset \sigma(A)$
2.  $\sigma(A)$  is a  $\sigma$  field
3. If  $A \subset G$  and  $G$  is a  $\sigma$  field,  $\sigma(A) \subset G$
4. If  $F$  is a  $\sigma$  field,  $\sigma(F) = F$
5. If  $A \subset A'$  then  $\sigma(A) \subset \sigma(A')$
6. If  $A \subset A' \subset \sigma(A)$ , then  $\sigma(A) = \sigma(A')$ .

**Example - Borel Sets** Let  $I$  be the subintervals of  $\Omega = (0, 1]$ , and let  $B = \sigma(I)$ . Note that  $I \subset B_0 \subset B$ , and so  $\sigma(B_0) = B$ . Note that elements of  $B$  are called the Borel sets. Note, it contains open sets on the unit interval (intersection of contained rational intervals).

**Probability Measures** A set function is a real-valued function defined on some class of subsets of  $\Omega$ . A set function  $P$  on a field  $F$  is a *probability measure* if it satisfies the conditions:

1.  $0 \leq P(A) \leq 1$  for  $A \in F$
2.  $P(\emptyset) = 0, P(\Omega) = 1$
3. Countable additivity.

Note, countable additivity implies finite additivity. Note, the conditions can be seen as slightly redundant (ie, they can be relaxed, and still imply the same things).

**Probability Space** If  $F$  is a  $\sigma$  field on  $\Omega$ , and  $P$  is a probability measure on  $F$ , the triple  $(\Omega, F, P)$  is called a *probability measure space*, or simply *probability space*. A support for  $P$  is any  $F$  set  $A$  for which  $P(A) = 1$ .

**Discrete Probability Space** If  $\Omega$  is countable, and  $p(\omega)$  is a nonnegative function on  $\Omega$ , which sums to 1 for all  $\omega$ . Then,  $P(A) = \sum_{\omega \in A} p(\omega)$  is a probability measure, and so  $(\Omega, F, P)$  is a probability space. This forms the basis for discrete probability theory.

An interesting note: why do we call it discrete probability theory, but not *countable probability theory*? Well, I think the reason for this is the following. As noted earlier, and within my measure theory notes - we cannot have a *discrete* probability space on an uncountable space. This is because, any measure which assigns a value to each subset of an uncountable space, cannot have countable additivity. And so - a *discrete probability space* implies that the space  $\Omega$  must be *countable*.

**Massed Discrete Probability Space** This just refers to the notion that  $\Omega$  need not be countable - however, there are finitely/countably many  $\omega_k$  with corresponding masses  $m_k$ , such that  $P(A) = \sum_{\omega_k \in A} m_k$  for  $A$  in  $F$ . Here,  $P$  is discrete, but the space need not be. We can write  $P(A) = \sum_k m_k I_A(\omega_k)$ .

For  $P$  probability measure on a field  $F$ , we can easily prove the following properties:

1.  $P(A) \leq P(B)$  if  $A \subset B$
2.  $P(A^c) = 1 - P(A)$
3.  $P(A) + P(B) = P(A \cup B) + P(A \cap B)$
4.  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
5. Inclusion-exclusion formula, which is just induction on the above formula.
6. Finite subadditivity
7. Continuity from below: If  $A_n$  and  $A$  lie in  $F$  and  $A_n \uparrow A$ , then  $P(A_n) \uparrow P(A)$ . Note, the  $\uparrow$  notation implies inclusion and total union is  $A$ , whereas implies monotone increasing up to  $P(A)$  in the limit.
8. Continuity from above: If  $A_n$  and  $A$  lie in  $F$  and  $A_n \downarrow A$ , then  $P(A_n) \downarrow P(A)$ .
9. Countable subadditivity, which stems from the two easily proved above properties.

10. Finite additivity and  $A_n \downarrow \emptyset$  implies  $P(A_n) \downarrow 0$  implies countable additivity.

**Lebesgue Measure on the Unit Interval** In this section - we essentially say that our measure  $|I| = |(a, b]| = b - a = \lambda(I)$  is a *finitely additive* probability measure on  $B_0$ , the set of finite unions of disjoint intervals  $I$ . However, we are also able to extend this - and say that  $\lambda$  is a *countably additive probability measure* on our *field* of  $B_0$ . This involves using Theorem 1.3 multiple times, where we concluded that if  $A = \bigcup_k I_k$  for disjoint  $I_k$ , then:

$$\lambda(A) = \sum_k \lambda(I_k)$$

However, the tricky bit comes in extending this to  $A = \bigcup_k A_k$  for  $A_k \in B_0$ . However, at the end of the day, each of the  $A$  and  $A_k$  can be expressed as a *finite union of disjoint intervals*, and so we can just break down the summations to where we can use Theorem 1.3 iii on a countable union of disjoint intervals. None of these theorems are particularly ground breaking.

However, I like the note at the end - consider what we might have to do to *extend*  $\lambda$  to the Borel sets  $B = \sigma(B_0)$ , and hopefully prove that  $\lambda$  is still countably additive on such a set. Note, to prove finite additivity on a field - we just had to rely on basic elementary properties of the real number system. However, to prove countable additivity on our field - we had to rely on something deeper about the real numbers - namely, *compactness of  $[a, b]$* . To extend to  $B$ , however, and still prove countable additivity - we will need some *new property*, ultimately.

**Sequence Space** Note, everything here has to do with  $S$ , a *finite* set of points, regarded as *outcomes* of a simple observation, or experiment. Note - we can extend this to *multiple repeated experiments* in the following way. However, for me - I often had trouble with understanding how a *sequence of events* could be independent. Well - if we have  $S$  is a finite possible outcomes of events, we could just allow our  $\sigma$  algebra to be the *sequence space* - and then, different coordinates *should be independent*. Note, I haven't really proved that statement, but it seems to me to be intuitive enough.

Let  $\Omega = S^\infty$  be the space of all infinite sequences:

$$\omega = (z_1(\omega), z_2(\omega), \dots)$$

of elements of  $S$ :  $z_k(\omega) \in S$  for all  $\omega \in S^\infty$  and  $k \geq 1$ . The sequence above can be viewed as the result of repeating infinitely often the simple experiment

represented by  $S$ . For  $S = \{0, 1\}$ , the space  $S^\infty$  is closely related to the unit interval. Note, I think this is the case for  $S = \{0, 1, \dots, n-1\}$  as well - just take  $S$  to be some base  $n$  representation of the numbers, and we can again identify  $S^\infty$  with the unit interval. See problems 1.4 and 1.5, for example.

**Definition - Coordinate Functions** Note that each  $z_k(\cdot)$  is a *coordinate function*, or *natural projection*, or just *projection function*. Such a function takes an element in  $S^\infty$ , and returns one in  $S$ . Let  $S^n = S \times \dots \times S$  be the Cartesian product of  $n$  copies of  $S$  - it consists of the  $n$  long sequences  $(u_1, \dots, u_n)$  of elements of  $S$ . For such a sequence, the set:

$$\{\omega : (z_1(\omega), \dots, z_n(\omega)) = (u_1, \dots, u_n)\}$$

Represents the event that the first  $n$  repetitions of the experiment give the outcomes  $u_1, \dots, u_n$  in sequence.

**Definition: Cylinder of Rank  $n$**  is a set of the form:

$$A = \{\omega : (z_1(\omega), \dots, z_n(\omega)) \in H\}$$

Note - colloquially, I like to say - cylinders of rank  $n$  are just *restrictions* on the first  $n$  experiments. Note that  $H \subset S^n$ , and that if  $H$  is nonempty,  $A$  is also nonempty. The previous example, where  $|H| = 1$ , is called a *thin cylinder*.

Define  $C_0$  as the class of cylinders of all ranks. We have that  $C_0$  is a *field*.  $S^\infty$  and  $\emptyset$  are within  $C_0$ , for  $H = S^n, \emptyset$ . If  $H$  is replaced by  $S^n - H$ , we have  $C_0$  is closed under complements. We also have closed under finite unions - note just that the union of two cylinders of rank  $n \leq m$  is just a restriction on the first  $m$  experiments, and is also a cylinder.

**Definition: Probability Measure on the Field of Cylinders** Let  $p_u$ ,  $u \in S$  be probabilities on  $S$  - nonnegative and summing to 1. Define a set function  $P$  on  $C_0$  in this way:

$$P(A) = \sum_{(u_1, \dots, u_n) \in H} p_{u_1} \cdots p_{u_n}$$

We have that  $P$  is a *probability measure*, with *finite additivity*.  $P(S^\infty) = P(A) = 1$ , where  $A$  is the cylinder with  $H = S^n$ . Clearly,  $0 \leq P(A) \leq 1$ , and

$P(\emptyset) = 0$ . We just need finite additivity - which can also be proved easily, but I won't give it here, just read the book. This is called a *product measure*, given that the definition just is a product of the individual probabilities on  $S$ .

**Theorem 2.3** Every finitely additive probability measure on the field  $C_0$  of cylinders in  $S^\infty$  is in fact countably additive. Note - this theorem is easy to prove, which we will, after stating the following lemma.

**Lemma:** If  $A_n \downarrow A$ , where the  $A_n$  are nonempty cylinders, then  $A$  is nonempty. Using this lemma, we quickly prove Theorem 2.3. Recall, we had that if  $A_n \downarrow \emptyset$  implies that  $P(A_n) \downarrow 0$ , for a finitely additive probability measure, then the probability measure is countably additive as well. Take  $A_n \downarrow \emptyset$ . Note, each  $A_n$  is a cylinder in  $C_0$ . Assume the lemma is true. Now, assume that  $A_n$  does *not* converge to 0. Then,  $P(A_n) \geq P(\emptyset) \geq \epsilon > 0$  for some  $\epsilon$ . But, this implies that  $A_n$  is nonempty. By the lemma, this makes the assumption  $A_n \downarrow \emptyset$  a contradiction, and so we must have  $P(A_n) \downarrow 0$ . Thus, we indeed have that if  $A_n \downarrow \emptyset$ , we must have  $P(A_n) \downarrow 0$ , and by our previous fact, the finitely additive probability measure on our field  $C_0$  is indeed countably additive. qed.

**Proof of Lemma** Note, the proof of the lemma in the book is interesting. It is essentially a diagonalization argument, and taking subsequences of sequences that contain an element repeating an infinite number of times. It becomes easy to note that the  $\omega$  containing these elements appearing infinite number of times is within the intersection of each  $A_n$ , and so  $A$  is nonempty. However, I want to try and prove the more general argument.

By Tychonoff's theorem, we have that the Cartesian product of countable compact topological spaces is compact as well, with the product topology. Note, we can take the discrete topology on  $S$ , and as  $S$  is finite, we have that it must be compact as well. For any open cover of  $S$ , we can take a finite subcover, by identifying each finite number of elements with one of the sets in the open cover that contains it. And so, we have that  $S^\infty$  must be compact as well, by Tychonoff's theorem.

Note, for a proof of Tychonoff's theorem, see my Topology notes. We now note that each  $A_n \in C_0$  is a set within the product topology as well. Each  $A_n$  can be viewed as the intersection of open cylinders as defined in my topology book (the only difference being that there, open cylinders are restrictions on



one coordinate, rather than restrictions on the first  $n$  coordinates). And so, the  $A_n$  in question in the theorem are also elements of the topological space, and so any cover of  $S^\infty$  by sets like  $A_n$  must have a finite subcover. And so, the proof of the lemma comes down to the proof of the following fact: For  $A_1 \supseteq A_2 \supseteq A_3 \supseteq \cdots$  closed nonempty sets on a compact topological space, we must have that:

$$A = \bigcap_{n=1}^{\infty} A_n$$

is nonempty. Proof by contradiction. Assume that  $A = \emptyset$ . Then, we must have that  $A_1^c, A_2^c, \dots$  is an open cover of  $S^\infty$ . By compactness, there is a finite subcover, which we will just identify (wlog) the sets  $A_1^c, A_2^c, \dots, A_n^c$ . Thus, we have:

$$S^\infty = A_1^c \cup \cdots \cup A_n^c$$

However, as  $A_1 \supseteq A_2 \supseteq \cdots \supseteq A_n$ , we must have that  $A_n^c \supseteq \cdots \supseteq A_2^c \supseteq A_1^c$ , which tells us:

$$S^\infty = A_n^c \implies A_n = \emptyset$$

This contradicts  $A_n$  being nonempty, and so, we must have  $A$  is nonempty. qed.

**Extended Sequence Measure** In Chapter 3, we will learn how to extend a countably additive probability measure on a field  $F$  to a countably additive probability measure on the sigma field  $\sigma(F)$ . The term *probability measure* more accurately refers to the extended  $P$ . We have that  $C_0$  is the field above, and we let  $C = \sigma(C_0)$ . Thus,  $(S^\infty, C, P)$  will be a probability space that we will look at later, an important one.

## Problems

### 2.1

Define  $x \vee y = \max(x, y)$ , and for a collection  $\{x_\alpha\}$  define:

$$\bigvee_{\alpha} x_{\alpha} = \sup_{\alpha} x_{\alpha}$$

Similarly, define  $x \wedge y = \min(x, y)$ , and for a collection  $\{x_\alpha\}$  define:

$$\bigwedge_{\alpha} x_{\alpha} = \inf_{\alpha} x_{\alpha}$$

Prove that  $I_{A \cup B} = I_A \vee I_B$ ,  $I_{A \cap B} = I_A \wedge I_B$ ,  $I_{A^c} = 1 - I_A$ , and  $I_{A \Delta B} = |I_A - I_B|$ , in the sense that there is equality at each point of  $\Omega$ . Show that  $A \subset B$  if and only if  $I_A \leq I_B$  pointwise. Check the equation:

$$x \wedge (y \wedge z) = (x \wedge y) \vee (x \wedge z)$$

And deduce the distributive law  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ . By similar arguments prove that:

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C) \quad A \Delta C \subset (A \Delta B) \cup (B \Delta C)$$

$$\left( \bigcup_n A_n \right)^c = \bigcap_n A_n^c \quad \left( \bigcap_n A_n \right)^c = \bigcup_n A_n^c$$

I think, the first thing to note is - what is the  $I_A$  notation? It is just the indicator notation.

1.  $I_{A \cup B} = I_A \vee I_B$  is clear, as if a point is in either  $A$  or  $B$ , the maximum between 0 and 1 is 1.
2.  $I_{A \cap B} = I_A \wedge I_B$  also clear
3.  $I_{A^c} = 1 - I_A$  Also clear
4.  $I_{A \Delta B} = |I_A - I_B|$  If  $\omega$  is in  $A \Delta B$ , it is in one, and not the other. Thus,  $|I_A - I_B| = |1| = |-1| = 1$ .
5.  $A \subset B$  if and only if  $I_A \leq I_B$  pointwise. If  $x \in A$  and  $x \in B$ , then  $1 \leq 1$ . If  $x \in A$ , but  $x \notin B$ , we have  $1 \not\leq 0$ .
6. The remaining points are clear. Identify the  $\wedge$  or  $\vee$  with the corresponding set operation.

## 2.2 Union and Intersection Equality Property

We have:

$$U_k = \bigcup (A_{i_1} \cap \cdots \cap A_{i_k}) \quad I_k = \bigcap (A_{i_1} \cup \cdots \cup A_{i_k})$$

Where union and intersection are over all  $k$  tuples satisfying  $1 \leq i_1 \leq \cdots \leq i_k \leq n$ . We note that  $U_k = I_{n-k+1}$ . This is via the pigeon hold principle - take  $x \in U_k$ , note that it is in some  $A_{i_1} \cap \cdots \cap A_{i_k}$  intersection. Now, note that  $x \in I_{n-k+1}$ , as each of unions being intersected has  $n - k + 1$  unique sets out of the  $n$  total sets - so, it is only missing  $k - 1$  sets. So, each of the unions must include one of the  $A_{i_1}, \dots, A_{i_k}$ , and  $x \in I_{n-k+1}$ . Similar in the other direction.

### 2.3 Equivalent and Non Equivalent Field Definitions

1. We have that  $A^c = \Omega - A \in F$ . We note that  $A \cup B = \Omega - (A^c - B) \in F$ .
2. Let  $\Omega$  be four points,  $F$  the empty set,  $\Omega$ , and all six of the two-point sets. Note, complements and disjoint unions will not get us a set of size 3, which means that  $F$  is not a field.

### 2.4 Unions of increasing Fields are Fields, but not true for $\sigma$ fields

1. Note, any two sets ultimately belong to the same  $F_i$ , and so their union is within the union. Clearly complements are maintained. Let  $\Omega = \bigcup F_n$ , and so  $\Omega$  and  $\emptyset$  are also within the big union.
2. Let  $F_n$  be the discrete  $\sigma$  field on  $\{1, \dots, n\}$ . Note,  $\bigcup_n F_n$  does not contain  $\{1, 2, \dots\}$ , which can be expressed as the countable union of singletons.

### 2.5 Generated Field and Explicit Definition

1. To show that  $f(A)$  is a field - note, the discrete field is a field, so the intersection is non empty. Note that clearly unions and complements of items in the intersection remain within the intersection.
2. Show that the given representation is indeed a field that contains  $A$ , and so  $f(A)$  is equivalent to the representation. Note, this is just set theory stuff. See solutions for the complement, represented as a disjoint intersection of unions.

### 2.6 Comparing Fields and $\sigma$ fields

1. Note that  $f(A)$  must contain finite and cofinite fields, as any finite field is the finite union of the points, and we can take the complement of those to get any cofinite set. Note, the field of cofinite and finite sets contains  $A$ , and so we must have it is equal to  $A$ .
2. Clearly  $f(A) \subset \sigma(A)$ , as  $\sigma(A)$  is within the intersection definition of  $f(A)$ . If  $A$  is finite, we have that  $f(A) = \sigma(A)$ , as  $\sigma(A) \subset f(A)$ , as all unions can be represented as finite unions. Finally, we have  $A \subset f(A) \subseteq \sigma(A)$ , which implies  $\sigma(f(A)) = \sigma(A)$ .
3. If  $A$  is countable, then  $f(A)$  is countable, as each set can be represented as noted in 2.5. There are a countable number of such representations

(as the countable product of countable sets is surjective onto the representation).

## 2.7 Extending a field/ $\sigma$ field by $\{H\}$

Let  $H$  be a set outside of  $F$ , where  $F$  is a field (or  $\sigma$  field). Show that the  $(\sigma)$  field generated by  $F \cup \{H\}$  consists of sets of the form:

$$(H \cap A) \cup (H^c \cap B) \quad A, B \in F$$

Well, first note that the above is a  $(\sigma)$  field containing  $F \cup \{H\}$ . Clearly contains  $F$  and  $H$ , and the emptyset and  $\Omega$ . Now note closed under complements:

$$[(H \cap A) \cup (H^c \cap B)]^c = (H \cap A^c) \cup (H^c \cap B)$$

Now, note closed under unions:

$$\bigcup_k (H \cap A_k) \cup (H^c \cap B_k) = (H \cap \bigcup_k A_k) \cup (H^c \cap \bigcup_k B_k)$$

Finally, note that any  $\sigma$  algebra containing  $F \cup \{H\}$  must contain the one above. And so yes, the set above is the  $(\sigma)$  field generated by  $F \cup \{H\}$ .

## 2.8 $\sigma$ algebra for class where $A^c$ is countable union of class elements

Note that the smallest class over  $A$  that contains countable unions must be a  $\sigma$  algebra containing  $A$ , so this class  $K$  satisfies:

$$\sigma(A) \subseteq K$$

Finally, we note that  $K \subseteq \sigma(A)$ , as every element in  $K$  must be within  $\sigma(A)$ , by definition.

## 2.9 Equivalent $\sigma$ algebra definition

Define:

$$G = \{B \in \sigma(A) \mid \exists \text{ a countable subclass } A_B \text{ of } A \text{ such that } B \in \sigma(A_B)\}$$

Note that  $G = \sigma(A)$ . This can be done by showing

1. We have that  $A \subset G$ . We have that for  $B \in A$ ,  $B \in \sigma(B)$ , where  $A_B = B$  is a countable subclass of  $A$ .

2.  $G$  is a  $\sigma$  algebra. First, note that  $\emptyset, \Omega \in G$ , as  $\emptyset, \Omega \in \sigma(B)$  for  $B \in A$ . We have closed under countable unions, as for  $B_1, B_2, \dots \in G$ , we have that the union is within  $\sigma(A_{B_1} \cup A_{B_2} \cup \dots)$ , where  $A_{B_1} \cup A_{B_2} \cup \dots$  is a countable subclass of  $A$ . Finally, we have for  $B \in G$ , we must have that  $B^c \in \sigma(A_B) \implies B \in G$ .
3. We have  $\sigma(A) \subseteq G$  by the above two points.
4. We have that  $G \subseteq \sigma(A)$ , by definition, as  $G$  only consists of elements within  $\sigma(A)$ .

Thus, every  $B \in \sigma(A) \implies B \in G$ . Thus, every  $B \in \sigma(A)$  has a countable subclass  $A_B$  of  $A$  such that  $B \in \sigma(A_B)$ . qed.

## 2.10 Classes Generating The Discrete Field are Large

1. Show that if  $\sigma(A)$  contains every subset of  $\Omega$ , then for each pair  $\omega$  and  $\omega'$  of distinct points in  $\Omega$  there is in  $A$  a subset  $A$  such that  $I_A(\omega) \neq I_A(\omega')$ .

We first note the following: subsets of  $\Omega$  containing both  $\omega$  and  $\omega'$ , or neither, is a  $\sigma$  field  $F$ .  $F$  contains both the empty set and  $\Omega$ .  $F$  is closed under complements (as the complement of a set containing both is neither, and vice versa).  $F$  is closed under unions (union of both and neither contains both, union of both and both contains both, union of neither and neither contains neither).

We now examine  $\sigma(B)$ , where  $B$  is subsets of  $A$  that contain both  $\omega$  and  $\omega'$ , or neither. Note that  $\sigma(B)$  must be contained within the  $\sigma$  field noted above, and so  $\sigma(B) \neq 2^\Omega$ . Thus, we must have that  $\sigma(B)$  is strictly smaller than  $\sigma(A)$ , which implies  $B$  is strictly smaller than  $A$ , and so  $A$  must contain sets that have either  $\omega$  and no  $\omega'$ , or vice versa.

2. Show that the reverse implication holds if  $\Omega$  is countable.

The reverse implication is that if  $\Omega$  is countable, and  $A$  is such that for every pair  $\omega$  and  $\omega'$ , there is a subset  $A$  such that  $I_A(\omega) \neq I_A(\omega')$ , then  $\sigma(A) = 2^\Omega$ . This can be noted via forming  $\{\omega\}$  as a countable intersection of sets within  $\sigma(A)$ . Note that for every pair  $\omega, \omega'$ , we have a set that contains one and not the other, and via complements, we can always find a set  $A_{\omega'} \in A$  such that  $\omega \in A_{\omega'}$ , but  $\omega' \notin A_{\omega'}$ . By a countable union, we must have:

$$\{\omega\} = \bigcap_{\omega' \in \Omega} A_{\omega'}$$

Thus,  $\sigma(A) = 2^\Omega$ .

3. Show by example that the reverse implication need not hold for uncountable  $\Omega$ .

Well, we can have that  $\Omega = [0, 1]$ , and  $A$  is the class of all single intervals. This is essentially the example gone over in Chapters 1 and 2. Note,  $\sigma(A) = B$ , the Borel  $\sigma$  algebra. At the end of Chapter 3, we will note that  $B \neq 2^\Omega$ , or all possible subsets.

## 2.11 Countably Generated Fields

A  $\sigma$  field is *countable generated*, or *separable*, if it is generated by some countable class of sets.

1. Show that the  $\sigma$  field  $B$  of Borel sets is countable generated. Well, we have that it is generated by intervals with rational endpoints. There are countably many such intervals, there being a 1-1 correspondence with  $\mathbb{Q} \times \mathbb{Q}$ . Note that  $\sigma(\mathbb{Q} \times \mathbb{Q})$  contains all singletons, as we can express every singleton in  $[0, 1]$  as a countable intersection of rational intervals - as we can approach every point from below and above by rational numbers in  $[0, 1]$ . Thus,  $\sigma(\mathbb{Q} \times \mathbb{Q}) = B$ .
2. Show that the  $\sigma$  field of Example 2.4 is countable generated if and only if  $\Omega$  is countable.

In example 2.4, we have the countable and co-countable sets sigma field  $F$ . We assume that  $\Omega$  is uncountable, and  $F$  is countably generated. Thus,  $F$  must be generated by a countable number of singletons - namely, the countable singletons in all generating sets (or their complements - the generating sets must be countable or co-countable). We let  $\Omega_0$  be the union of these countable sets. We note that  $F$  must consist of sets  $B$  and  $B \cup \Omega_0^c$ , with  $B \subset \Omega_0$ . Clearly, each  $B \subset \Omega_0$  is within  $F$ , and so is each  $B \cup \Omega_0^c$ , as  $(B \cup \Omega_0^c)^c = B^c \cap \Omega_0$ , which is also countable.

Now, as for why all sets in  $F$  are of the form  $B$  or  $B \cup \Omega_0^c$ , for  $B \subset \Omega_0$ . Note first that these sets form a  $\sigma$  field. We have that  $B^c = (\Omega_0 - B) \cup \Omega_0^c$ , which is in the field. We have that  $(B \cup \Omega_0^c)^c = B^c \cap \Omega_0 \subset \Omega_0$ , which is also in the field, so it is closed under complements. We have that  $\emptyset$  is in the field, and so by closed under complements, so is  $\Omega$ . Finally, it is closed under unions - clearly  $B_1 \cup B_2$  and  $B_1 \cup \Omega_0^c \cup B_2 \cup \Omega_0^c$  are all within the field, and this can be extended

countably many times. Finally, note that  $B_1 \cup (B_2 \cup \Omega_0^c) = B_1 \cup B_2 \cup \Omega_0^c$  is within the field as well, and can be extended countably many times.

And so, we have defined a field. Note, this field contains all the sets within the generating set, and so all the countable and co-countable sets must be of the form defined above. Now, we find our contradiction. Note that the singletons within  $\Omega_0^c$  are countable sets. However, these singletons are not with  $\Omega_0$ , and they are not of the form  $B \cup \Omega_0^c$ . Note,  $\Omega_0^c$  is nonempty, as  $\Omega$  is uncountable. And so, we have a contradiction - namely, being that the countable generating sets cannot form every countable and co-countable set.

So, if  $\Omega$  is uncountable,  $F$  is *not* countably generated. If  $\Omega$  is countable - we must have that  $F$  is countable generated. Take all the singletons that are in the countable or co-countable sets. This must be a countable amount of singletons, as  $\Omega$  is countable. They generate all the countable and co-countable sets. Note, actually, if  $\Omega$  is countable, the countable and co-countable sets are just the power sets. And so, the singletons generate all sets, and  $F$ .

3. Suppose that  $F_1$  and  $F_2$  are  $\sigma$  fields,  $F_1 \subset F_2$ , and  $F_2$  is countable generated. Show by example that  $F_1$  may not be countably generated.

Let  $F_2$  consist of the Borel sets in  $\Omega = (0, 1]$ , and let  $F_1$  consist of the countable and co-countable sets in  $(0, 1]$ . Note that  $F_1$  cannot be countably generated, by the previous problem.  $F_2$  is countably generated by part 1. Finally,  $F_1 \subset F_2$ . Each countable set is a countable union of singletons, which is in  $F_2$ . The complement of each co-countable set is a countable union of singletons, which is in  $F_2$ , and so is the complement. Thus, we indeed have  $F_1 \subset F_2$ ,  $F_2$  countably generated, by  $F_1$  not. qed.

## 2.12 Cardinality of Fields and $\sigma$ Fields

Show that a  $\sigma$  field cannot be countably infinite - its cardinality must be finite, or else at least that of the continuum (the real numbers). Show by example that a field can be countably infinite.

Note, we have examples for finite and uncountably infinite  $\sigma$  fields. Assume a  $\sigma$  field is countably infinite. Then, we can list every set in the  $\sigma$  field

like so:

$$A_1, A_2, A_3, \dots$$

We will  $\omega \in \{\pm 1\}^{\mathbb{N}}$ . We define:

$$B^\omega = \bigcap_{n \in \mathbb{N}} A_n^{\omega_n}$$

Where  $\omega_n$  is the  $n$ th bit in  $\omega$ . Note,  $-1$  is just the complement. If  $\gamma \neq \omega$ , then we must have:

$$B^\omega \cap B^\gamma = \emptyset$$

As they differ on including some  $A_n$  and  $A_n^c$  in their intersections. We also note that:

$$A_n = \bigcup_{\omega: \omega_n=1} B^\omega$$

As because we must have that  $A_1^{\omega_1} \cap \dots \cap A_{n-1}^{\omega_{n-1}} \cap \widehat{A_n} \cap \dots$  partitions all of  $\Omega$  (think about it this way - for each of the sets listed in the intersection,  $x \in \Omega$  is in either  $A_i$ , or  $A_i^c$ .  $x$  would be in the intersection that makes the right choice for each  $i$ ).

So now, we consider the set  $\{B^\omega : \omega \in \{\pm 1\}^{\mathbb{N}}\}$ . If there are only a finite amount of different  $B^\omega$  - this would imply that there are only a finite amount of distinct  $A_n$ . So, there is an infinite family of  $\omega$  for which the  $B^\omega$  are different. We have for some index set,  $\omega_i, i \in I, B^{\omega_i} \neq B^{\omega_j}$ . Let  $C_n := B^{\omega_n}$ . Note, we just assume that the index set is countable. So, what is the reason of the above mess - we had that each of the  $A_i$  were *distinct*. However, what we get from the  $C_n$ , is that these are countably infinite sets that are *disjoint*. Now, we define:

$$\Gamma : \{0, 1\}^{\mathbb{N}} \rightarrow F, (\gamma) \rightarrow \bigcup_{\gamma_n=1} C_n$$

Note: the map of  $\Gamma$  is *injective*. As the  $C_n$  were disjoint, we see that each set in the disjoint union contains *different elements*. Finally, we note that all possible subsets of  $\{0, 1\}^{\mathbb{N}}$  is uncountable (as we can identify one point in  $[0, 1]$  with each element in  $\{0, 1\}^{\mathbb{N}}$ , this is all of chapter 1). And so, we have that if the  $\sigma$  algebra is not finite, it has cardinality of *at least the real line*.

The steps are actually simple. 1. Find countably many disjoint sets. 2. Create uncountable sets in the  $\sigma$  algebra via disjoint unions. qed.



### 2.13 Probability Measure on Finite and CoFinite Sets

1. The probability measure we define is that  $P(A) = 0$  if  $A$  is finite and  $P(A) = 1$  if  $A$  is cofinite. Note, the measure is not well defined if  $\Omega$  is finite, but we assume that  $\Omega$  is infinite. We note that it is countably additive - for finite  $A_1, \dots, A_n$ , we have the union is finite, and so:

$$P\left(\bigcup A_n\right) = 0 = \sum P(A_n)$$

I now note - we cannot have disjoint cofinite sets. Assume that  $A$  and  $B$  are cofinite. Thus,  $A^c$  and  $B^c$  are finite - this is because  $\Omega - A^c - B^c$  are both non empty, and  $\Omega - A^c - B^c \subset A, \Omega - A^c - B^c \subset B$ . Finally, we note for a disjoint union of  $n - 1$  finite sets and a cofinite set, we must have that their complement is cofinite, and so:

$$P\left(\bigcup A_n\right) = 1 = \sum P(A_n)$$

2. Show that  $P$  is not countably additive if  $\Omega$  is countably infinite.

So, in the case that  $\Omega$  is countably infinite - the countable union of finite sets can result in a cofinite set ( $A_n = \{\omega_1\}$ ). And so:

$$P\left(\bigcup A_n\right) = 1 \neq 0 = \sum P(A_n)$$

3. Show that  $P$  is countably additive if  $\Omega$  is uncountable.

We just need countably additive in the case that the *countable union remains within our field*. Note - the countable union of finite sets cannot be cofinite. This union would mean  $A$  is countable, and  $A^c$  is finite, which would imply that  $\Omega = A \cup A^c$  is countable. And so, we can only have a countable union that results in a finite set  $A$ . And so:

$$P\left(\bigcup A_n\right) = P(A) = 0 = \sum P(A_n)$$

4. Now let  $F$  be the  $\sigma$  field consisting of the countable and the cocountable sets in an uncountable  $\Omega$ , and define  $P$  analogously. Show that  $P$  is countably additive.

Union of countable countable sets is still countable. No such union of disjoint cocountable sets. Union of countable countable sets with one cocountable set is cocountable. qed.

## 2.14 First Category $\sigma$ Field

In  $(0, 1]$  let  $F$  be the class of sets that either (i) are of the first category (countable union of sets that are nowhere dense, which are sets  $E$  for which each interval  $I$  contains an interval  $J$  that does not meet  $E$ ) or (ii) have complement of the first category. Show that  $F$  is a  $\sigma$  field.

The empty set is of the first category, and the complement of  $(0, 1]$  (the empty set) is of the first category. Clearly complements of sets in  $F$  are still in  $F$ , as this is definitional. Finally, we consider countable unions of sets in  $F$ . If each set is first countable, the countable union of a countable amount of nowhere dense sets is still a countable union of nowhere dense sets, and so the countable union of first countable sets is first countable.

Now, we assume at least one of the sets in our countable union has a complement in the first category. Note, the complement of the countable union must be contained within the complement of our first set, and so the complement of the countable union is in the first category as well. Thus, the countable union in this case is also within  $F$ . In all cases,  $F$  contains countable unions of elements of  $F$ . Thus, we can conclude  $F$  is a  $\sigma$  field.

For  $A$  in  $F$ , take  $P(A)$  to be 0 in case (i) and 1 in case (ii). Show that  $P$  is countably additive.

Countably additive only needs to apply for disjoint sets. For disjoint sets in case (i) - the countable union of such disjoint sets is still in case (i), and so  $P(\bigcup A_i) = 0 = \sum 0 = \sum P(A_i)$ .

Now, we note that we cannot have 2 disjoint sets in case (ii). Assume  $A$  and  $B$  are disjoint, and the complements of  $A$  and  $B$  are both first countable. We note that for any two disjoint sets - we must have  $\Omega = A^c \cup B^c$ . Take  $x \in \Omega$ . Assume  $x \notin A^c$ . Thus,  $x \in A \implies x \notin B \implies x \in B^c$ . So,  $x \in A^c$  or  $x \in A$ , in which case,  $x \in B^c$ . However, note that  $A^c \cup B^c$  is still first countable, which implies that  $(0, 1]$  is first countable. However, in an earlier problem, we noted that by Baire, a nonempty interval is *not* of the first category. And so, this is a contradiction, and we cannot have disjoint  $A$  and  $B$  whose complements are first countable.

So, the only remaining case for countably additive is one set of case (ii) and a countable amount of sets of case (i) which are all disjoint. As noted

above - the union of such sets is case (ii), and we have:

$$P\left(\bigcup A_i\right) = 1 = 1 + 0 = P(A_1) + \sum_{i=2}^{\infty} P(A_i) = \sum P(A_i)$$

And thus  $P$  is countably additive. qed.

### 2.15 Differences between $B_0$ and $C_0$

On the field  $B_0$  in  $(0, 1]$  defined  $P(A)$  to be 1 or 0 according as there does or does not exist some positive  $\epsilon_A$  (depending on  $A$ ) such that  $A$  contains the interval  $\left(\frac{1}{2}, \frac{1}{2} + \epsilon_A\right]$ . Show that  $P$  is finitely but not countably additive.

No such example is possible for the field  $C_0$  in  $S^\infty$  (as by Theorem 2.3, every finitely additive probability measure on the field  $C_0$  must be countably additive. This shows the opposite for  $B_0$ ).

We let sets of case (i) be those for which  $P(A) = 0$ , and those of case (ii) be those for which  $P(A) = 1$ . Note, for finitely additive - we are essentially just doing the same shit we did above. Two sets of case (i) - their finite union must also be of case (i). As for why this is the case:

Assume that  $A$  and  $B$  are of case (i), but  $A \cup B$  is of case (ii), ie, there is some positive  $\epsilon_{A \cup B}$  such that:

$$\left(\frac{1}{2}, \frac{1}{2} + \epsilon_{A \cup B}\right] \subseteq A \cup B$$

We will show that this creates a contradiction. For both  $A$  and  $B$ , as they are of case (i), for every  $\epsilon > 0$ , there is a point  $x_{\epsilon A}$  and  $x_{\epsilon B}$  such that:

$$x_{\epsilon A}, x_{\epsilon B} \in \left(\frac{1}{2}, \frac{1}{2} + \epsilon\right] \quad x_{\epsilon A} \notin A \quad x_{\epsilon B} \notin B$$

For the entire interval to be contained, we must have for every  $\epsilon < \epsilon_{A \cup B}$ , we also have:

$$x_{\epsilon A} \in B \quad x_{\epsilon B} \in A$$

This creates a contradiction. As  $A$  and  $B$  are within  $B_0$ , they must be finite unions of disjoint intervals. Note, a point itself is not actually within  $B_0$ , and so these points  $x_{\epsilon A}$  must be contained within some interval of  $B$ . However, this entire interval is not within  $\left(\frac{1}{2}, \frac{1}{2} + \epsilon\right]$ , and so it has some starting point

$1/2 < \epsilon'_B$ , and  $A$  similarly has a starting point  $1/2 < \epsilon'_A$ . We can take a new  $\epsilon = \min(\epsilon'_A, \epsilon'_B)$ , and find new points that must not be contained in  $A$  and  $B$ , but create corresponding intervals in the other set, and so on. This can be done a countably infinite number of times. This process thus implies that  $A$  and  $B$  must both be disjoint unions of countable sets - which contradicts  $A, B \in B_0$ . Thus, we must have that  $A \cup B$  must be of case (i). Thus, we find:

$$P(A \cup B) = 0 = P(A) + P(B)$$

We can use induction to prove for any finite disjoint union of case (i) sets, the measure is finitely countable. Now, we consider case (ii). Like above, it is clear that there is no disjoint union of case (ii) sets. That is because, we just take the minimum of  $\epsilon_A, \epsilon_B$ , and clearly, both intervals are in both sets. So, the final case we consider is a case (ii) set with multiple case (i) sets. Again, this finite disjoint union is of case (ii), and so:

$$P(A \cup B_1 \cup \dots \cup B_n) = 1 = 1 + 0 + \dots + 0 = P(A) + P(B_1) + \dots + P(B_n)$$

The final point to make is that  $P$  is not countably additive. We can define:

$$A_n = \left( \frac{1}{2} + \frac{1}{n+2}, 1 \right]$$

Each  $A_n$  is clearly of case (i). However, the countable union of  $A_n$  is  $(1/2, 1]$ , and so is of case (ii). Thus, we do not have countable additivity, as this would give us a contradiction that  $1 = 0$ . Thus,  $P$  is not countably additive.

I think the most important part is why this is interesting. We noted that if  $S = \{0, 1\}$ , then  $C_0$  is essentially a field on  $(0, 1]$ , as we can identify each of the infinite sequences with a point in  $(0, 1]$ . And, initially, the sets in the field  $C_0$  look like they contain intervals and their unions. However, there must be something substantially different about the field and/or measure on  $C_0$ , when compared with  $B_0$ .

## 2.18 Stochastic Arithmetic

Define a set function  $P_n$  on the class of all subsets of  $\Omega = \{1, 2, \dots\}$  by:

$$P_n(A) = \frac{1}{n} \# [m : 1 \leq m \leq n, m \in A]$$

among the first  $n$  integers, the proportion that lie in  $A$  is just  $P_n(A)$ . Then  $P_n$  is a discrete probability measure (a probability measure defined on a discrete space). The set  $A$  has *density*:

$$D(A) = \lim_n P_n(A)$$

provided this limit exists. Note, density makes sense, as we are essentially getting a ratio of how many numbers are in  $A$ , compared with  $\Omega$  (as  $P_n(\Omega) = 1$  for all cases). Let  $\mathcal{D}$  be the class of sets having density.

(a) Show that  $D$  is finitely but not countably additive on  $\mathcal{D}$ . Take  $A, B \in \mathcal{D}$  disjoint. We have that:

$$D(A \cup B) = \lim_n P_n(A \cup B) = \lim_n P_n(A) + P_n(B) = \lim_n P_n(A) + \lim_n P_n(B) = D(A) + D(B)$$

Note implicit above is that  $\mathcal{D}$  contains finite disjoint unions. Note, we have additivity of  $P_n$  on *disjoint* sets, but it is not true for non-disjoint sets. We show not countably additive by a counter example. Define  $A_i = \{i\}$ . We have that:

$$D(A_i) = \lim_n P_n(A_i) = \lim_n \frac{1}{n} = 0$$

However,  $\bigcup A_i = \Omega$ . Thus:

$$D\left(\bigcup A_i\right) = D(\Omega) = 1 \neq 0 = \sum D(A_i)$$

Note in the above example, we do have that the countable disjoint union is within  $\mathcal{D}$ .

(b) Show that  $\mathcal{D}$  contains the empty set and  $\Omega$  and is closed under the formation of complements, proper differences, and finite disjoint unions, but is not closed under the formation of countable disjoint unions or of finite unions that are not disjoint.

In part (a) we concluded closed under finite disjoint unions. We now consider complements. Take  $A \in \mathcal{D}$ . By definition, we have that the limit exists and

$$\lim_n P_n(A) = \lim_n x_n = x$$

We note that  $P_n(A^c) = 1 - x_n$ . This is because,  $A$  contains some  $i$  elements between 1 and  $n$ . We have that  $A^c$  must contain  $n - i$  elements between 1 and  $n$ . Thus,  $P_n(A^c) = \frac{n-i}{n} = 1 - \frac{i}{n} = 1 - P_n(A) = 1 - x_n$ . Thus, we find:

$$\lim_n P_n(A^c) = \lim_n 1 - x_n = 1 - \lim_n x_n = 1 - x$$

Thus,  $A^c \in \mathcal{D}$ . We now consider proper differences. Take  $A, B \in \mathcal{D}$ . We note:

$$A \setminus B = A \cap B^c = (A^c \cup B)^c$$

Recall that the *proper* difference implies that  $B \subset A$ . Thus,  $A^c \in \mathcal{D}$ ,  $A^c \cap B \in \mathcal{D}$  being a disjoint union, and finally  $(A^c \cup B)^c \in \mathcal{D}$ .

Now, we consider the not closed examples. We will try and find counter examples. I first outline a set that is not within  $\mathcal{D}$ . We need the  $x_n$  values to oscillate, so that the limit does not exist. I define such an  $A$  like so. If we have:

$$P_n(A) = \frac{1}{4} = \frac{n/4}{n}$$

We can make  $P_{4n}(A) = \frac{3}{4}$ , by setting  $11n/4$  of the final  $3n$  entries as 1:

$$P_{4n}(A) = \frac{n/4 + 11n/4}{4n} = \frac{3}{4}$$

Similarly, if we have:

$$P_n(A) = \frac{3}{4} = \frac{3n/4}{n}$$

We can make  $P_{4n}(A) = \frac{1}{4}$ , by setting  $n/4$  of the final  $3n$  entries as 1:

$$P_{4n}(A) = \frac{3n/4 + n/4}{4n} = \frac{1}{4}$$

In such a way, we build up  $A$ . We start with  $A = \{4\}$ . Thus, we have:

$$P_4(A) = \frac{1}{4}$$

Then, we add  $A = \{4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16\}$ , such that:

$$P_{16}(A) = \frac{3}{4}$$

We continue with the process defined above, adding the numbers that correspond to the final entries in the added  $3n$  spots. Thus, we have:

$$\lim P_n(A)$$

Does not exist, as it oscillates between  $1/4$  and  $3/4$ . We have that  $\mathcal{D}$  is not closed under the formation of countable disjoint unions. We have that:

$$A = \{4\} \cup \{5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16\} \cup \dots$$

Note that each set in our countable union has a finite amount of elements - and so for each subset  $B$ ,  $P_n(B) = 0$ . However,  $\bigcup B = A \notin \mathcal{D}$ .

We now consider finite unions that are not disjoint. In this case, we define a new subset, as noted in the solutions. We let  $A$  be the set of even integers. We define  $C_k = \{m : v_k < m \leq v_{k+1}\}$ . Note, for  $v_1 = 0$ ,  $\bigcup C_k = \Omega$ . Finally, we define  $B$  as:

$$B = (A \cap (C_1 \cup C_3 \cup \dots)) \cup (A^c \cap (C_2 \cup C_4 \cup \dots))$$

We thus have that:

$$A \cap B = A \cap (C_1 \cup C_3 \cup \dots)$$

We note that if  $v_k$  increases rapidly enough, then the set  $A \cap B$  has no density. Note, consider just the sets  $A \cap C_{2i-1}$ , which in a disjoint union, would form  $A \cap B$ . We have that:

$$P_{v_{2k-1}}(A \cap B) = \frac{1}{v_{2k-1}} \left| \bigcup_{i=1}^k A \cap C_{2i-1} \right|$$

$$P_{v_{2k}}(A \cap B) = \frac{1}{v_{2k}} \left| \bigcup_{i=1}^k A \cap C_{2i-1} \right|$$

Note, this is because going from  $v_{2k-1}$  to  $v_{2k}$ , we are not including any of the ones included in  $C_{2k}$ . Thus, we have:

$$\frac{P_{v_{2k-1}}}{P_{v_{2k}}} = \frac{v_{2k-1}}{v_{2k}}$$

Note, if  $A \cap B$  has a density  $D(A \cap B)$ , then we must have that:

$$\lim_{k \rightarrow \infty} \frac{P_{v_{2k-1}}}{P_{v_{2k}}} = \frac{D(A \cap B)}{D(A \cap B)} = 1$$

Thus, if we have the limit of the ratio  $\frac{v_{2k-1}}{v_{2k}}$  does not equal 1, then that implies  $A \cap B$  must have no density. Such a sequence would be  $v_k = 2^k$ , in which case the ratio is 2. Now, as noted above, this set is a union of countable singletons, each with density 0, so like our above example - this shows that  $\mathcal{D}$  is not closed under countable unions. However, this example also helps us with finite non disjoint unions. However, we also note that:

$$D(B) = \frac{1}{2}$$

Because, by in large,  $B$  contains one of every two points in the pair  $(x_i, x_i+1)$ . We also have that  $A \in \mathcal{D}$ , as clearly  $D(A) = 1/2$  as well. By the above points,

we have that  $A^c, B^c \in \mathcal{D}$ . However, if we assume that  $\mathcal{D}$  is closed under non disjoint unions, we have:

$$A^c \cup B^c = (A \cap B)^c \in \mathcal{D} \implies A \cap B \in \mathcal{D}$$

Which is a contradiction.

(c) Let  $\mathcal{M}$  consist of the periodic sets  $M_a = \{ka : k = 1, 2, \dots\}$ . Observe that:

$$P_n(M_a) = \frac{1}{n} \left\lfloor \frac{n}{a} \right\rfloor \rightarrow \frac{1}{a} = D(M_a)$$

Show that the field  $f(\mathcal{M})$  generated by  $\mathcal{M}$  is contained in  $\mathcal{D}$ . Show that  $D$  is completely determined on  $f(\mathcal{M})$  by the value it gives for each  $a$  to the event that  $m$  is divisible by  $a$ .

First, we note that by problem 2.5,  $f(\mathcal{M})$  is the class of sets of the form:

$$\bigcup_{i=1}^m \bigcap_{j=1}^{n_i} A_{ij} \quad \text{such that} \quad A_{ij} \in \mathcal{M} \quad \text{or} \quad A_{ij}^c \in \mathcal{M} \quad \text{and the } m \text{ sets } \bigcap_{j=1}^{n_i} A_{ij} \quad \text{are disjoint}$$

Note that by closed under finite disjoint unions, if each  $\bigcap_{j=1}^{n_i} A_{ij} \in \mathcal{D}$ , then  $f(\mathcal{M})$  is completely contained in  $\mathcal{D}$ . We note that:

$$M_a \cap M_b = M_{lcm(a,b)}$$

Take  $x \in M_a \cap M_b$ . Note that  $lcm(a, b)$  is the smallest integer that is divisible by both  $a$  and  $b$ . Recall the fundamental theorem of arithmetic - *wiki*. Go over the proofs - they are not too difficult. Every number has a unique prime factorization. Thus, we have:

$$a = p_1^{e_1} \cdots p_n^{e_n} \quad b = p_1^{f_1} \cdots p_n^{f_n}$$

Where  $p_1, \dots, p_n$  is a list from 1 to a prime larger than  $a$  and  $b$  (infinite primes, one exists). We note that:

$$lcm(a, b) = p_1^{\max(e_1, f_1)} \cdots p_n^{\max(e_n, f_n)}$$

Note, the rhs is clearly divisible by  $a$  and  $b$  - and if we decrease one of the powers by 1, it is not divisible by the two. And so, it is clearly equal, by definition (ie, you cannot have a smaller prime factorization that is divisible by  $a$  and  $b$ ). Note that  $x = ka$  and  $x = k'b$ . Thus, the prime factorization of  $x$  must include *at least*  $p_i^{\max(e_i, f_i)}$  for each  $i$ , and thus  $x = k''lcm(a, b)$ , and



$x \in M_{lcm(a,b)}$ . The other direction is trivial.

What is the point of this? It shows that  $\mathcal{M}$  is closed under intersections, and as  $\mathcal{M} \subset \mathcal{D}$ , we have that:

$$\bigcap_{j=1}^{n_i} A_{ij} \in \mathcal{D} \implies f(\mathcal{M}) \subset \mathcal{D}$$

Now, this implies that  $D$  on  $f(\mathcal{M})$  is determined by  $D$  on  $\mathcal{M}$ . For each intersection, take the  $lcm$  to find the value of  $D$  on the intersection. Then, just add up the fractions.

(d) Assume that  $\sum p^{-1}$  diverges - this is the sum over all primes. Prove that  $D$ , although finitely additive, is not countably additive on the field  $f(\mathcal{M})$ .

Note, finite additivity is given, by finite additivity on all of  $\mathcal{D}$ , and  $f(\mathcal{M}) \subset \mathcal{D}$ . We now make use of the solutions. We define  $B_l = M_a - \bigcup_{1 < p \leq l} M_{ap}$  for some  $a \in \Omega$ . We note that  $B_l \in f(M)$  - by the definition of a field, it is closed under finite unions, complements, and finite intersections, and these operations are used to form  $B_l$  out of sets originally in  $f(M)$ . We now note:

$$D(B_l) = D(M_a) - D\left(\bigcup_{1 < p \leq l} M_{ap}\right)$$

Note that  $\mathcal{D}$  is closed under proper differences, and for  $p > 1$ , we have  $M_{ap} \subset M_a$ . We also note why the above equation is true - for  $B \subset A$ , both in  $D$ , we have that:

$$D(A-B) = D(A \cap B^c) = D((A^c \cup B)^c) = 1 - D(A^c \cup B) = 1 - D(A^c) - D(B) = D(A) - D(B)$$

Where, each equality above we have proved in the preceding questions, as we have that  $A^c$  and  $B$  are disjoint. Thus, we have:

$$D(B_l) = \frac{1}{a} - D\left(\bigcup_{1 < p \leq l} M_{ap}\right)$$

Now, we note that  $D$  is a finite probability measure on  $f(M)$ . Thus, we can apply the inclusion-exclusion formula, to find:

$$D(B_l) = \frac{1}{a} - \sum_{1 < p \leq l} \frac{1}{ap} + \sum_{1 < p < q \leq l} \frac{1}{apq} - \dots$$

$$\implies D(B_l) \leq \frac{1}{a} - \sum_{1 \leq p \leq l} \frac{1}{ap}$$

Note - the implication is given in the notes. However, I do note that:

$$\frac{1}{apq} \sum_{q < k \leq l} \frac{1}{apqk} = \frac{1}{apq} \sum_{q < k \leq l} \frac{1}{k}$$

It is hard to compare with the next term in the inclusion-exclusion principle. We are given in the question that  $\sum \frac{1}{k}$  diverges - and so it must be greater than one, correct? I guess, you can split it up, for every  $apq$  term, like:

$$-\frac{1}{apq} \sum_{m=1}^{l-q} \sum_{q < k_1 < k_2 < \dots < k_m \leq l} \left[ \frac{1}{k_1} - \frac{1}{k_1 k_2} + \dots + \frac{1}{k_1 k_2 \dots k_m} \right]$$

I have spent too much time on this question. I will skip it.

## 2.19 Nonatomic Probability Spaces

A probability measure space  $(\Omega, \mathcal{F}, P)$  is *nonatomic* if  $P(A) > 0$  implies that there exists a  $B$  such that  $B \subset A$  and  $0 < P(B) < P(A)$  ( $A$  and  $B$  in  $\mathcal{F}$ , of course).

Note: An initial observation I had is that  $\mathcal{F}$  cannot contain finite sets - as then we always have the existence of a finite subset with nonzero probability, and we can continue recursively until a single point (which is where I guess the name *nonatomic* comes from), which doesn't satisfy the *nonatomic condition*.

1. Assuming the existence of Lebesgue measure  $\lambda$  on  $\mathcal{B}$  (ie, we have a probability space  $((0, 1], \mathcal{B}, \lambda)$ ), prove that it is nonatomic.

We first considered proving that the sets satisfying the nonatomic condition in  $\mathcal{B}$  contained the intervals and where a field, and thus equal to  $\mathcal{B}$ . However, a simpler way is by examining the following function for  $A$  with  $\lambda(A) > 0$ :

$$f(x) = \lambda(A \cap (0, x])$$

We first note that  $f(0) = 0$  and  $f(1) = P(A)$ . We now note that  $f$ , being a measure, is continuous from below. Ie, take a sequence  $x_n$ , with  $x_0 = 0$ , and  $x_n \rightarrow 1$ , non decreasing. We note that:

$$B_n = A \cap (0, x_n] \implies B_n \subseteq B_{n+1} \implies \bigcup_n B_n = A$$

We also note that as  $A \in \mathcal{B}$  and  $(0, x_n] \in \mathcal{B}$ , each  $B_n \in \mathcal{B}$ , and so is the countable union clearly. By continuity from below we have:

$$\lim_n f(x_n) = \lim_n \lambda(B_n) = \lambda(A) = P(A)$$

Or, even better. We let  $a$  be the infimum of points in  $A$ , and  $b$  the supremum - note, not equal and well defined, as  $A$  has more than 2 points (a finite set is Lebesgue measure 0). We let  $x_n$  start at  $a$ , and approach  $b$ . We still have  $\lambda(B_0) = 0$ . Now, we assume that for every arbitrary sequence of such  $x_n$ , there is no intermediate value - ie, we either have  $\lambda(B_n) = 0$ , or  $\lambda(B_n) = P(A)$ . This must be a contradiction. If it is true for every sequence - there must be a single point  $c \in [a, b)$  where the switch happens - ie, if  $x_n \geq c$ , then  $P(B_n) = P(A)$ , and 0 otherwise. If there were two such  $c$ , we could form a sequence with  $x_n$  between the  $c$ , and get a contradiction from the limit being nondecreasing. However, the existence of such a  $c$  would be a contradiction for the following reasons - it would imply  $P(\{c\}) = P(A) > 0$ . This is because the above would imply for all  $x_n \in (c, b)$ , we have:

$$P(A \cap (c, x_n]) = P(A)$$

And we could define  $C_n = A \cap (c - \frac{1}{n}, x_n]$  for  $x_n$  decreasing to  $c$ ,  $\{c\} = \bigcap C_n$ , and then probability from above would give  $P(\{c\}) = P(A)$ .

Thus, there must be some  $x_n$  satisfying:

$$0 < P(B_n) < P(A)$$

And thus  $\mathcal{B}$  is *nonatomic*.

2. Show that in the nonatomic case that  $P(A) > 0$  and  $\epsilon > 0$  imply that there exists a  $B$  such that  $B \subset A$  and  $0 < P(B) < \epsilon$ .

I think, this follows if there is some  $P(A) = \epsilon$ . Or, we have that as a probability measure space is a  $\sigma$  algebra, we assume some nonempty  $P(A)$ , and then we have the existence of  $0 < P(B) < P(A)$ . By the probability of a complement, we have either  $P(B) \leq P(A)/2$ , or  $P(A - B) \leq P(A)/2$  (both  $B$  and  $A - B$  being in the sigma field). We can continue iteratively to  $\epsilon$ .

3. Show in the nonatomic case that  $0 \leq x \leq P(A)$  implies that there exists a  $B$  such that  $B \subset A$  and  $P(B) = x$ .

By the proof in part A - as we take the  $x$  from 0 to 1 -  $f(x)$  must be *continuous*. It is clear that  $f(x)$  is increasing. Assume that it is not continuous. Then, we have some  $x$  where a *jump discontinuity* is made. In such a case - we can just make use of the above argument, to assign a nonzero probability to a singleton  $x$ , namely  $P(\{x\}) > 0$ . In which case, we have a contradiction in the nonatomic case, as noted at the start. No finite set can have nonzero probability.

Note, just more details. The above argument relies on  $\Omega$  going from 0 to some number  $t$ . It works for the Borel sets, however the hint gives us something more general. Assume the existence of some  $x$ . Inductively define class  $\mathcal{H}_n$  in the following way:

$$\mathcal{H}_0 = \{\emptyset\} \quad \mathcal{H}_n = \left\{ H : H \subset A - \bigcup_{k < n} H_k, P\left(\bigcup_{k < n} H_k\right) + P(H) \leq x \right\}$$

We define  $H_k$  above in the following way. We let  $h_k = \sup \{P(H) : H \in \mathcal{H}_k\}$ , and we have that  $H_k$  is some set in  $\mathcal{H}_k$  with  $P(H_k) > h_k - \frac{1}{k}$ .

We first note that  $P\left(\bigcup_{k < n} H_k\right) \leq x$ . By definition, we have that:

$$P(H_{n-1}) + P\left(\bigcup_{k < n-1} H_k\right) \leq x \implies P\left(\bigcup_{k < n} H_k\right) \leq x$$

As it is also a disjoint union, by definition, we have the implication above. We also have that each  $\mathcal{H}_n$  is nonempty. If we are in the case where  $P\left(\bigcup_{k < n} H_k\right) = x$ , we are in the clear, as we have found a subset of  $A$  satisfying what we need. However, if  $P\left(\bigcup_{k < n} H_k\right) < x$ , we have that  $\epsilon = x - P\left(\bigcup_{k < n} H_k\right) < P(A)$ , and we make use of part (b) to find at least one set that satisfies being within  $\mathcal{H}_n$ . Finally, we have that finding  $P(H_n) > h_n - \frac{1}{n}$  does exist - as, by the supremum, it is either obtained, or we reach it in the limit.

And so, the iteration is well defined. We have that  $H_1, H_2, \dots$  is a *strictly increasing* number of disjoint sets. We have that their countable union (if we must continue that long) is within  $\mathcal{F}$ . We that the unions:

$$\bigcup_{k=1}^n H_k$$

Are strictly increasing. We have by disjoint union:

$$P\left(\bigcup_{k=1}^n H_k\right) = \sum_{k=1}^n P(H_k) > \sum_{k=1}^n h_k - \frac{1}{k}$$

Note, I don't like the hanging  $1/k$  term, as it goes to infinity. However, we can note that:

$$h_k \in \left(x - \sum_{t=1}^{k-1} h_t - \frac{1}{t}, x\right]$$

We just need a condition to prove  $h_k$  grows quick enough. I think this might come from noting that if we exclude members of previous classes of  $\mathcal{H}_n$ , then we can ensure that the remaining  $H$  to define  $\mathcal{H}_n$  are big enough, so that  $h_t$  grows quick enough.

4. Show in the nonatomic case that if  $p_1, p_2, \dots$  are nonnegative and add to 1, then  $A$  can be decomposed into sets  $B_1, B_2$  such that  $P(B_n) = p_n P(A)$ .

Well, first we define the subset  $B_1 = p_1 P(A)$ , then we look at  $A - B_1$ , which still must contain a set of size  $p_2 P(A)$ , and so on. Easy enough.

## 2.21 Generating Sigma Algebras "From the Inside"

1. Suppose that  $\mathcal{A} = \{A_1, A_2, \dots\}$  is a countable partition of  $\Omega$ . Recall that  $\mathcal{A}_0 = \mathcal{A}$  and  $\mathcal{A}_n = \mathcal{A}_{n-1}^*$ , where  $H^*$  consists of the sets in  $H$ , the complements of sets in  $H$ , and the finite and countable unions of sets in  $H$ .

So,  $\mathcal{A}_1 = \mathcal{A}_0^*$  is the sets in  $\mathcal{A}_0$ , the complements of sets in  $\mathcal{A}_0$ , and the finite and countable unions of sets in  $\mathcal{A}_0$ . Show that  $\mathcal{A}_1$  coincides with  $\sigma(\mathcal{A})$ . This is a case where  $\sigma(\mathcal{A})$  can be constructed "from the inside."

Well - coincides is essentially  $\mathcal{A}_1 = \sigma(\mathcal{A})$ . This is true if  $\mathcal{A}_1$  is a  $\sigma$  field, as  $\mathcal{A}_1$  contains  $\mathcal{A}$ . For  $B \in \mathcal{A}_1$ , we note  $B^c \in \mathcal{A}_1$  if  $B$  is in the element of  $\mathcal{A}_0$  case. If  $B$  is a complement or countable union of elements in  $\mathcal{A}_0$  - note that  $B^c$  is just another countable union of elements in  $\mathcal{A}_0$ , in which case  $B^c \in \mathcal{A}_1$ .

Now, we quickly note  $\Omega$  is in  $\mathcal{A}_1$ , and the complement of  $\Omega$  (being an empty union). Finally, we note  $\mathcal{A}_1$  is closed under countable unions

- every element of  $\mathcal{A}_1$  can be expressed as a finite or countable union of elements of  $\mathcal{A}_0$  (as seen with the complement case), and a countable amount of countable/finite unions is still a countable union of elements in  $\mathcal{A}_0$ . Thus, we have that  $\mathcal{A}_1$  is a  $\sigma$  algebra, and  $\mathcal{A}_1 \supseteq \sigma(\mathcal{A})$ . We also have that  $\mathcal{A}_1 \subseteq \sigma(\mathcal{A})$ , again noting that every element of  $\mathcal{A}_1$  is a countable union of elements of  $\mathcal{A}$ . Thus, we can conclude:

$$\mathcal{A}_1 = \sigma(\mathcal{A})$$

2. Show that the set of normal numbers lies in  $\mathcal{I}_6$ . Recall,  $\mathcal{I}$  is the class of subintervals of  $\Omega$ . We recall that the set of normal numbers is of the form:

$$N = \bigcap_{k=1}^{\infty} \bigcup_{m=1}^{\infty} \bigcap_{n=m}^{\infty} [\omega : |n^{-1}s_n(\omega)| < k^{-1}]$$

Where we also recall that each  $[\omega : |n^{-1}s_n(\omega)| < k^{-1}]$  is a finite disjoint union of intervals. Note,  $\mathcal{I}_{n-1}^*$  is also closed under countable intersections. So, we have that  $\mathcal{I}_1$  would contain sets of the form  $[\omega : |n^{-1}s_n(\omega)| < k^{-1}]$ ,  $\mathcal{I}_2$  would have intersections of those,  $\mathcal{I}_3$  would have unions of those, and  $\mathcal{I}_4$  would have intersections of those. Thus, the normals are indeed within  $\mathcal{I}_6$ , give or take.

3. Show that  $\mathcal{H}^* = \mathcal{H}$  if and only if  $\mathcal{H}$  is a  $\sigma$  field. Show that  $\mathcal{I}_{n-1}$  is strictly smaller than  $\mathcal{I}_n$  for all  $n$ .

Well - clearly if  $\mathcal{H}$  is a  $\sigma$  field, then  $\mathcal{H}^* = \mathcal{H}$ , given that countable unions and complements are already contained. Now, assume that  $\mathcal{H}^* = \mathcal{H}$ . Then, for any element in  $A \in \mathcal{H}$ , we know its complement is already in  $\mathcal{H}$ , and any list of elements in  $\mathcal{H}$ , the union is in  $\mathcal{H}$ . Thus, we do have the if and only if.

Assume that  $\mathcal{I}_{n-1} = \mathcal{I}_n$  for all  $n$ . Then,  $\mathcal{I}_{n-1} = \mathcal{B}$ , as it is a  $\sigma$  algebra containing the intervals (note how both  $\supseteq$  and  $\subseteq$  are obtained). Note, this is a contradiction, as we had shown in the chapter that  $\bigcup_n \mathcal{I}_n \neq \mathcal{B}$ . Thus, as  $\mathcal{I}_n \supseteq \mathcal{I}_{n-1}$ , and  $\mathcal{I}_n \neq \mathcal{I}_{n-1}$ , we can conclude that  $\mathcal{I}_{n-1} \subset \mathcal{I}_n$ .

## 2.22

Extend (2.27) to the infinite ordinals  $\alpha$  by defining  $\mathcal{A}_\alpha = \left( \bigcup_{\beta < \alpha} \mathcal{A}_\beta \right)^*$ . Show that, if  $\Omega$  is the first uncountable ordinal, then:

$$\bigcup_{\alpha < \Omega} \mathcal{A}_\alpha = \sigma(\mathcal{A})$$

Show that, if the cardinality of  $\mathcal{A}$  does not exceed that of the continuum ( $\mathbb{R}$ ), then the same is true of  $\sigma(\mathcal{A})$ . Thus  $\mathcal{B}$  has the power of the continuum. I will skip this for now, as I don't really know ordinals too well.

## Section 3 - Existence and Extension

### Notes

The main theorem of the chapter is the following:

**Theorem 3.1 - Probability Measure Extension Theorem** A probability measure on a field has a unique extension to the generated  $\sigma$  field.

### Construction of the Extension

**Outer Measure** I like the discussion on the intuition for defining the *outer measure* and *inner measure*. Say we have  $P$  a probability measure on field  $\mathcal{F}_0$ . We want to define  $P$  on  $\sigma(\mathcal{F}_0)$  - but it doesn't need to be restricted to  $\sigma(\mathcal{F}_0)$ . The outer measure is defined for each subset  $A$  of  $\Omega$  as:

$$P^*(A) = \inf \sum_n P(A_n) \quad \text{such that} \quad A \subset \bigcup_n A_n$$

Note - as  $\mathcal{F}_0$  is a field on  $\Omega$ , it contains  $\Omega$ , so we could take our finite cover as  $\Omega$  itself, at least. So, the infimum is always non-empty. As for intuition - we have  $P$  on  $\mathcal{F}_0$  - to extend  $P$  to sets outside of  $\mathcal{F}_0$  - using what we already have - we just have to approximate  $A$  by sets in  $\mathcal{F}_0$ . The covering is an approximation, and the infimum takes the best such approximation.

**Inner Measure** Note how "from the inside" applies to the inner measure:

$$P_*(A) = 1 - P^*(A^c)$$

The covering of  $A^c$  will perhaps cover more than  $A^c$ , and so the complement of that will be slightly less than  $A$ . I like considering whether we can take  $P_*(A) = \sup \sum_n P(A_n)$  of  $A_n$  unions inside of  $A$ . However, just practically - this would imply normal numbers  $N$  have inner measure of 0 (as there are no nonempty intervals in  $\mathcal{B}_0$  inside of  $N$ ).

**$P^*$  Measurable** So,  $P^*(A)$  holds for every set. Which sets do we want our measure to actually apply to? We would want it to apply to sets where our probability measure properties hold - so perhaps sets where:

$$P^*(A) + P^*(A^c) = 1$$

Note, this should always be the case by countable additivity and  $P(\Omega) = 1$ . And so, we take this as our condition. Note, this is *equivalent* to:

$$P^*(A) = P_*(A)$$

We also note that it will be helpful to apply the more stringent condition:

$$P^*(A \cap E) + P^*(A^c \cap E) = P^*(E)$$

For every set  $E$  - the previous condition held with  $E = \Omega$ , as it would turn out  $P^*(\Omega) = 1$ . In a later problem, we will see that all three conditions are equivalent anyway - see problem 3.2. And so, we just for now, define and say a set  $A$  is called  *$P^*$ -Measurable* if the last condition, with  $E$ , holds. Note, it is the most stringent, and so if we have that  $P^*$ -Measurable sets contain  $\sigma(\mathcal{F}_0)$ , it should be enough anyway. We use the variable  $\mathcal{M}$  to define the class of  $P^*$  measurable sets.

**Properties of  $P^*$  and  $\mathcal{M}$**  We will need these properties to prove that  $\mathcal{M}$  contains  $\sigma(\mathcal{F}_0)$  and that the restriction of  $P^*$  to  $\sigma(\mathcal{F}_0)$  is our required (unique) extension of  $P$ .

1.  $P^*(\emptyset) = 0$
2.  $P^*$  is nonnegative:  $P^*(A) \geq 0$  for every  $A \subset \Omega$
3.  $P^*$  is monotone:  $A \subset B$  implies  $P^*(A) \leq P^*(B)$
4.  $P^*$  is countably subadditive:  $P^*(\cup_n A_n) \leq \sum_n P^*(A_n)$

1 is obvious, as the union of empty sets contains  $\emptyset$ . 2 is obvious, as we already said the infimum is nonempty, and each element is nonnegative (being a sum of  $P$  nonnegative). 3 is also obvious, as any cover of  $B$  covers  $A$ , and so all sums in  $B$ 's infimum are in  $A$ 's. The final part is countable subadditivity. This is the normal  $\epsilon/2^n$  proof:

For each  $A_n$ , take a cover of  $\mathcal{F}_0$  sets  $B_{nk}$ , where  $\sum_k P(B_{nk}) < P^*(A_n) + \epsilon/2^{-n}$ . Note, it is possible to find by infimum definition. Note that all  $B_{nk}$  cover the total union - so, by the definition of infimum:

$$P^*(\cup_n A_n) \leq \sum_{n,k} P(B_{nk}) < \sum_n P^*(A_n) + \epsilon/2^{-n} = \epsilon + \sum_n P^*(A_n)$$

This is true for all  $\epsilon > 0$ , and so 4 follows.



**Lemma 1 - The class  $\mathcal{M}$  is a field** First, we note by finite subadditivity above:

$$P^*(E) \leq P^*(A \cap E) + P^*(A^c \cap E)$$

And so, to prove that  $A$  is in  $\mathcal{M}$ , we just need to prove that:

$$P^*(A \cap E) + P^*(A^c \cap E) \leq P^*(E)$$

It is clear that  $\Omega \in \mathcal{M}$ , and  $\mathcal{M}$  is closed under complements. The final step is to prove  $\mathcal{M}$  is closed under finite unions. Note, it is simple to prove that:

$$P^*(E) \geq P^*((A \cup B) \cap E) + P^*((A \cup B)^c \cap E)$$

Using the  $E$  condition being true on  $A$  and  $B$  (and taking  $(B \cap E)$  as the set  $E$ , and so on) and subadditivity. qed.

**Lemma 2 -  $P^*$  Condition on Countable Disjoint Sets in  $\mathcal{M}$**  If  $A_1, A_2, \dots$  is a finite or infinite sequence of disjoint  $\mathcal{M}$ -sets, then for each  $E \subset \Omega$ :

$$P^*\left(E \cap \left(\bigcup_k A_k\right)\right) = \sum_k P^*(E \cap A_k)$$

Proof: For finite disjoint sets, we use induction. Clearly true for  $n = 1$ . For  $n = 2$ , it is also easy to prove. If  $A_1 \cup A_2 = \Omega$ , then  $A_2 = A_1^c$ , and it is just our condition. Otherwise:

$$P^*(E \cap (A_1 \cup A_2)) = P^*(A_1 \cap E \cap (A_1 \cup A_2)) + P^*(A_1^c \cap E \cap (A_1 \cup A_2))$$

Where the last step again makes use of our condition for  $E = E \cap (A_1 \cup A_2)$ . The above equals:

$$= P^*(E \cap A_1) + P^*(E \cap A_2)$$

Where we have that  $A_1^c \cap E \cap (A_1 \cup A_2) = E \cap A_2$  via disjointness. Ie,  $A_2 \subseteq A_1^c$ , so  $x \in A_2 \implies x \in A_1^c$ . With the  $n = 2$  case, we can imply our inductive hypothesis and find the formula is true for any finite disjoint set of unions. For the infinite case - note:

$$P^*\left(E \cap \left(\bigcup_k A_k\right)\right) \geq \sum_{k=1}^n P^*(E \cap A_k)$$

This is by monotonicity of  $P^*$ . We can take  $n \rightarrow \infty$  on the RHS, and so we can indeed conclude:

$$P^*\left(E \cap \left(\bigcup_k A_k\right)\right) \geq \sum_k P^*(E \cap A_k)$$

And the other inequality  $\leq$  follows from countable subadditivity. qed.

**Lemma 3 - The class  $\mathcal{M}$  is a  $\sigma$  field, and  $P^*$  restricted to  $\mathcal{M}$  is countably additive** Suppose that  $A_1, A_2, \dots$  are disjoint  $\mathcal{M}$  sets with union  $A$ . Since  $F_n = \bigcup_{k=1}^n A_k$  lies in  $\mathcal{M}$ , we have:

$$P^*(E) = P^*(E \cap F_n) + P^*(E \cap F_n^c)$$

By applying lemma 2 and monotonicity, we thus find:

$$P^*(E) \geq \sum_{k=1}^n P^*(E \cap A_k) + P^*(E \cap A^c)$$

We can take  $n$  to infinity, and using Lemma 2 again, we find:

$$P^*(E) \geq P^*(E \cap A) + P^*(E \cap A^c)$$

Thus, the countable disjoint union is also within  $\mathcal{M}$ . Note, any countable union can be expressed as a countable disjoint union - thus  $\mathcal{M}$  is a  $\sigma$  field. Countable additivity of  $P^*$  on  $\mathcal{M}$  follows again from Lemma 3.2, taking  $E = \Omega$ . qed.

Note - Lemmas 1, 2, and 3 only used properties (1) through (4) we defined above, along with an  $\Omega$ ,  $P^*$  being defined on all subsets of  $\Omega$ , and the condition that defined  $\mathcal{M}$ . Now, we can bring it back to comparing with  $P$  on  $\mathcal{F}_0$  and extending to  $P^*$  on  $\sigma(\mathcal{F}_0)$ .

**Lemma 4 - If  $P^*$  is defined by the outer measure, then  $\mathcal{F}_0 \subset \mathcal{M}$**  Note, this just requires  $\mathcal{M}$  containing every set in  $\mathcal{F}$ . Take  $A \in \mathcal{F}_0$ . Given  $E$  and  $\epsilon$ , choose  $\mathcal{F}_0$  sets  $A_n$  such that:

$$E \subset \bigcup_n A_n \quad \text{and} \quad \sum_n P(A_n) \leq P^*(E) + \epsilon$$

Recall - the goal is to show  $\mathcal{F}_0 \subset \mathcal{M}$ , and so we need to show for our  $E \subset \Omega$  defined above:

$$P^*(A \cap E) + P^*(A^c \cap E) \leq P^*(E)$$

Define  $B_n = A_n \cap A$  and  $C_n = A_n \cap A^c$ .  $B_n$  and  $C_n$  are in  $\mathcal{F}_0$  by field properties. Also,  $E \cap A \subset \bigcup_n B_n$ , and  $E \cap A^c \subset \bigcup_n C_n$ . By first  $P^*$  outer measure definition, and then by finite additivity of  $P$  on  $\mathcal{F}_0$ , we have:

$$P^*(A \cap E) + P^*(A^c \cap E) \leq \sum_n P(B_n) + \sum_n P(C_n) = \sum_n P(A_n) \leq P^*(E) + \epsilon$$

As this is true for all  $\epsilon > 0$ , we have that our necessary and sufficient condition is true, and thus  $A \in \mathcal{F}_0 \implies A \in \mathcal{M}$ . qed.

**Lemma 5 - If  $P^*$  is defined by the outer measure, then  $P^*(A) = P(A)$  for  $A \in \mathcal{F}_0$**  Clearly, we have that  $P^*(A) \leq P(A)$ , as we can just take a countable union of  $A$  and emptysets. Now, take any union where  $A \subset \cup_n A_n$ . By countable subadditivity and monotonicity of  $P$  on  $\mathcal{F}_0$ , we have:

$$P(A) = P(A \cap \cup_n A_n) = P(\cup_n A \cap A_n) \leq \sum_n P(A \cap A_n) \leq \sum_n P(A_n)$$

Thus, the infimum clearly takes the value of  $P(A)$ . qed.

**Proof of Extension in Theorem 3.1** Suppose  $P^*$  is defined as our outer measure from a (countably additive) probability measure  $P$  on the field  $\mathcal{F}_0$ . Let  $\mathcal{F} = \sigma(\mathcal{F}_0)$ . By Lemma 3 and 4:

$$\mathcal{F}_0 \subseteq \mathcal{F} \subseteq \mathcal{M} \subseteq 2^\omega$$

By Lemma 5, we have  $P^*(\Omega) = P(\Omega) = 1$ . By Lemma 3,  $P^*$  when restricted to  $\mathcal{M}$  is a probability measure there, and thus  $P^*$  restricted to  $\mathcal{F}$  is also clearly a probability measure on that class as well. By Lemma 5,  $P^*$  agrees with  $P$  on  $\mathcal{F}$ , and thus it is our required extension.

### Uniqueness of the Extension and the $\pi - \lambda$ Theorem

We first give the following definitions:

**$\pi$  System** A class  $\mathcal{P}$  of subsets of  $\Omega$  is a  $\pi$  system if it is closed under the formation of finite intersections:  $A, B \in \mathcal{P} \implies A \cap B \in \mathcal{P}$

**$\lambda$  System** A class  $\mathcal{L}$  of subsets of  $\Omega$  is a  $\lambda$  system if it contains  $\Omega$ , and is closed under complements and finite and countable disjoint intersections:

1.  $\Omega \in \mathcal{L}$
2.  $A \in \mathcal{L} \implies A^c \in \mathcal{L}$
3.  $A_1, A_2, \dots \in \mathcal{L}$  disjoint  $\implies \cup_n A_n \in \mathcal{L}$

Note: As to why these are useful. We ultimately want to prove uniqueness of an extension. We have that an extension is equal on a field - which is also a  $\pi$  system. And ultimately, the *essential* property of the probability measure is its countable additivity, which is defined for a countable disjoint union. These countable disjoint unions are contained within  $\lambda$ .

And so, how we can prove uniqueness can go something like this: if we have a probability measure on a  $\pi$  system, and we can extend it to a  $\lambda$  system (that is somehow related to a  $\sigma$  field), and the measure agrees on the  $\lambda$  system as well - then, the probability measure is unique!

That is a "high flying" statement I would say. Also, it is much more general than what we need it for - like, why should we start with a  $\pi$  system, and not just a field? Well, the generality will be useful later. And so, we shall go and prove portions of that statement:

**Lemma 6  $\pi$  and  $\lambda$  implies  $\sigma$**  A class that is both a  $\pi$  system and a  $\lambda$  system is a  $\sigma$  field. **Proof:** Easy enough, as any countable union can be expressed as a countable disjoint union, using finite intersections. qed. This gives us the connection to a  $\sigma$  field.

**Theorem 3-2:  $\pi - \lambda$  Theorem** If  $\mathcal{P}$  is a  $\pi$  system and  $\mathcal{L}$  is a  $\lambda$  system, then  $\mathcal{P} \subset \mathcal{L}$  implies  $\sigma(\mathcal{P}) \subset \mathcal{L}$ .

**Proof:** Let  $\mathcal{L}_0$  be the  $\lambda$  system generated by  $\mathcal{P}$ , the intersection of all containing  $\lambda$  systems. It is easy to prove intersections of  $\lambda$  systems are  $\lambda$  systems (essentially the same as the proof for  $\sigma(\cdot)$  and nonempty by  $2^\Omega$ ). Thus, we have  $\mathcal{P} \subset \mathcal{L}_0 \subset \mathcal{L}$  by definition. If  $\mathcal{L}$  can be shown to be a  $\pi$  system, then it will follow by Lemma 6 that  $\mathcal{L}_0$  is a  $\sigma$  field. Thus, we *must* have:

$$\mathcal{P} \subseteq \sigma(\mathcal{P}) \subseteq \mathcal{L}_0 \subseteq \mathcal{L}$$

Thus, the theorem follows if  $\mathcal{L}_0$  is a  $\pi$  system as well.

For each  $A \subset \Omega$ , let  $\mathcal{L}_A$  be the class of sets  $B$  such that  $A \cap B \in \mathcal{L}_0$ . If  $A \in \mathcal{P}$ , or  $A \in \mathcal{L}_0$ , then  $\mathcal{L}_A$  is a  $\lambda$  system. Note,  $A \in \mathcal{P} \implies A \in \mathcal{L}_0$ , and so we just prove it for the second case. First,  $\mathcal{L}_A$  contains  $\Omega$ , as  $A \cap \Omega = A \in \mathcal{L}_0$ . We now recall that the conditions:

1.  $\mathcal{L}_A$  contains  $\Omega$
2.  $\mathcal{L}_A$  contains proper differences
3.  $\mathcal{L}_A$  contains disjoint countable unions

Is equivalent to a  $\lambda$  system - as the complement of an element of the set can be expressed as a proper difference. We have the first point. For the second, take  $B_1, B_2 \in \mathcal{L}_A$  with  $B_1 \subset B_2$ . We have:

$$(A \cap B_2) - (A \cap B_1) = A \cap (B_2 - B_1) \in \mathcal{L}_0$$

By the proper difference property of  $\mathcal{L}_0$ , and thus  $B_2 - B_1 \in \mathcal{L}_1$ , fulfilling point 2. The final point is the third. Let  $B_n$  be disjoint  $\mathcal{L}_A$  sets.  $\mathcal{L}_0$  contains  $A \cap B_n$ , which are disjoint, and so  $\mathcal{L}_0$  contains  $\bigcup_n A \cap B_n = A \cap (\bigcup_n B_n)$ , and thus  $\mathcal{L}_A$  contains disjoint unions.

Thus, we have that  $\mathcal{L}_A$  is a  $\lambda$  system. If  $A \in \mathcal{P}$  and  $B \in \mathcal{P}$ , then  $A \cap B \in \mathcal{P} \subset \mathcal{L}_0$ , which is equivalent to  $B \in \mathcal{L}_A$ . Thus,  $A \in \mathcal{P} \implies P \subset \mathcal{L}_A$ , and by minimality, we have:

$$\mathcal{L}_0 \subseteq \mathcal{L}_A$$

This is where it gets tricky. By the above -  $A \in \mathcal{P}$  and  $B \in \mathcal{L}_0$  implies  $B \in \mathcal{L}_0 \subseteq \mathcal{L}_A$ . As this is symmetric - it implies  $A \in \mathcal{L}_B$ . Thus,  $B \in \mathcal{L}_0$  implies  $\mathcal{P} \subset \mathcal{L}_B$  (as for every  $A \in \mathcal{P}$ ,  $A \in \mathcal{L}_B$ ). Thus, again by minimality, we have if  $B \in \mathcal{L}_0$ :

$$\mathcal{L}_0 \subseteq \mathcal{L}_B$$

We essentially used "symmetry" of the  $\mathcal{L}_C$  definition to go from  $A \in \mathcal{P}$  implies  $\mathcal{L}_0 \subseteq \mathcal{L}_A$ , directly to  $B \in \mathcal{L}_0$  implies  $\mathcal{L}_0 \subseteq \mathcal{L}_B$ . Finally -  $B \in \mathcal{L}_0$ , and  $C \in \mathcal{L}_0$ , together imply  $C \in \mathcal{L}_B$ , or  $B \cap C \in \mathcal{L}_0$  by definition. Therefore,  $\mathcal{L}_0$  is a  $\pi$  system, and the theorem follows. qed.

Uniqueness of our extended probability measure  $P^*$  is thus a result of the  $\pi - \lambda$  theorem:

**Theorem 3-3 Uniqueness of Extensions** Suppose that  $P_1$  and  $P_2$  are probability measures on  $\sigma(\mathcal{P})$ , where  $\mathcal{P}$  is a  $\pi$  system. If  $P_1, P_2$  agree on  $\mathcal{P}$ , then they agree on  $\sigma(\mathcal{P})$ .

Let  $\mathcal{L}$  be the class of sets  $A$  in  $\sigma(\mathcal{P})$  such that  $P_1(A) = P_2(A)$ . Clearly,  $\Omega \in \mathcal{L}$ . If  $A \in \mathcal{L}$ , then  $P_1(A^c) = 1 - P_1(A) = 1 - P_2(A) = P_2(A^c)$ , and so  $\mathcal{L}$  is closed under complements (note, we have the formula by being a probability measure). If  $A_n$  are disjoint sets in  $\mathcal{L}$ , then:

$$P_1\left(\bigcup_n A_n\right) = \sum_n P_1(A_n) = \sum_n P_2(A_n) = P_2\left(\bigcup_n A_n\right)$$

Thus,  $\mathcal{L}$  is also closed under disjoint countable unions, and so  $\mathcal{L}$  is a  $\lambda$  system. Thus, as  $\mathcal{P} \subset \mathcal{L}$ , the  $\pi - \lambda$  theorem tells us that  $\sigma(\mathcal{P}) \subseteq \mathcal{L}$ , which implies that  $P_1$  and  $P_2$  agree on  $\sigma(\mathcal{P})$  as well. qed.

And so - we have proved the uniqueness of the extension, as for any two probability measures on  $\sigma(\mathcal{F}_0)$  that agree on  $\mathcal{F}_0$ , agree on the  $\pi$  system  $\mathcal{F}_0$ ,

and thus agree on the extended sigma algebra.

Note - as stated above, our Theorems are pretty general. We could have replaced  $\pi$  system every where with field, as a field has all the properties of a  $\pi$  system. All in total, we have proved:

**Theorem 3.1 - Probability Measure Extension Theorem** A probability measure on a field has a unique extension to the generated  $\sigma$  field.

**Monotone Classes** Are classes that are closed under the formation of monotone unions and intersections (like in continuity from above and below). That's it. There is another close relative of the  $\pi - \lambda$  theorem for these classes:

**Theorem 3.4 - Halmos's Monotone Class Theorem** If  $\mathcal{F}_0$  is a field and  $\mathcal{M}$  is a monotone class, then  $\mathcal{F}_0 \subset \mathcal{M} \implies \sigma(\mathcal{F}_0) \subset \mathcal{M}$ . **Proof:** Very similar to the  $\pi - \lambda$  theorem, in that it relies on first defining a minimal monotone class  $m(\mathcal{F}_0)$  containing  $\mathcal{F}_0$  - and then finding a simpler condition to proving that it is a  $\sigma$  field and thus contains  $\sigma(\mathcal{F}_0)$ . That simpler condition is proving that  $m(\mathcal{F}_0)$  is a field - which clearly implies it is a  $\sigma$  field.

Proving that  $m(\mathcal{F}_0)$  is closed under complements is easy. However - proving it is closed under finite unions is more difficult. The argument, however - is the same in structure as the  $\pi - \lambda$  theorem. It involves a "switching" statement, to prove that the class of sets in  $m(\mathcal{F}_0)$  that are closed under unions are also a monotone class containing  $\mathcal{F}_0$ , and thus  $m(\mathcal{F}_0)$  is contained in that class. Thus, we have  $m(\mathcal{F}_0)$  is a field, and unraveling all the way back proves the theorem. qed.

### Lebesgue Measure on the Unit Interval

Recall the unit interval  $(0, 1]$  with the field  $\mathcal{B}_0$  of finite disjoint unions of subintervals. In Theorem 2.2, we have a probability measure  $\lambda$  on  $\mathcal{B}_0$ , where  $\lambda$  assigns intervals a measure equal to their length. Thus, by Theorem 3.1,  $\lambda$  extends to  $\mathcal{B}$ , the extended  $\lambda$  being the Lebesgue measure. The probability space  $((0, 1], \mathcal{B}, \lambda)$  will thus be the basis of a lot of the probability theory we will go over in the remaining sections of the first chapter. An interesting note: as the intervals in  $(0, 1]$  form a  $\pi$  system generating  $\mathcal{B}$ ,  $\lambda$  is the *only* probability measure on  $\mathcal{B}$  that assigns to each interval its length as its measure.

**Example 1 - Subintervals around rationals** We can assign to each rational  $r_1, r_2, \dots \in (0, 1)$  an interval  $I_n$  of length  $\epsilon/2^n$ . Consider  $A = \cup_n I_n$ . By subadditivity,  $\lambda(A) < \epsilon$ . Note,  $A$  is dense, and note each interval  $I$  must intersect an  $I_n$ , so  $\lambda(A \cap I) > 0$ . Note that  $B = (0, 1) - A$  satisfies  $\lambda(B) > 1 - \epsilon$  - however, no matter how close to 1 the measure of  $B$  is,  $B$  will always be nowhere dense (all intervals contain a rational and thus intersect  $B^c$ ).

**Example 2 -  $k$  Repeated Sequences** This example gives a set with real-world probabilistic interpretations that has similar properties to the set in the previous example. Let  $d_n(\omega)$  be the  $n$ th digit in the dyadic expansion of  $\omega$  - like in the first section. Define:

$$A_n = \{\omega \in (0, 1] : d_i(\omega) = d_{n+i}(\omega) = d_{2n+i}(\omega), i = 1, \dots, n\}$$

Then, we define:

$$A = \bigcup_{n=1}^{\infty} A_n$$

Probabilistically,  $A$  corresponds to the event that in an infinite sequence of tosses of a coin, some finite initial segment is immediately duplicated twice over. From  $\lambda(A_n) = 2^n \cdot 2^{-3n}$  (consider  $3n$  spaces, we have any choice for the first  $n$  spaces, and then fixed choices for the final  $2n$  spaces) (also, note that the dyadic cylinders can be expressed as disjoint interval unions, and so they do have a  $\lambda$  value), we have  $0 < \lambda(A) \leq 1/3$ .  $A$  is dense in the unit interval - we can find an element in  $A$  by taking a midpoint of some interval, and then repeating the sequence again, which remains in the interval. It's measure can be made less than  $1/3$ , by requiring the initial segment is duplicated  $k$  times with  $k$  being large.

And so, we have a real world example of a dense set, but with very very small probability.

**Completeness** A probability measure space  $(\Omega, \mathcal{F}, P)$  is *complete* if  $A \subset B, B \in \mathcal{F}$ , and  $P(B) = 0$  together imply  $A \in \mathcal{F}$  (and hence,  $P(A) = 0$  by monotonicity).

Suppose that  $(\Omega, \mathcal{F}, P)$  is an arbitrary probability space. Define  $P^*$  for  $\mathcal{F}_0 = \mathcal{F} = \sigma(\mathcal{F}_0)$  - note, even though  $\mathcal{F}$  is a  $\sigma$  field,  $P^*$  is still defined for *all subsets of  $\Omega$*  via the outer measure. Consider the  $\sigma$  field  $\mathcal{M}$  of  $P^*$  measurable sets (ones for which  $P^*(A) + P^*(A^c) = 1$ ). The arguments outlines above tell

us that  $(\Omega, \mathcal{M}, P^*)$  is a probability measure space. If  $P^*(B) = 0$  and  $A \subset B$ , then:

$$P^*(A \cap E) + P^*(A^c \cap E) \leq P^*(B) + P^*(E) = P^*(E) \implies A \in \mathcal{M}$$

Thus,  $(\Omega, \mathcal{M}, P^*)$  is a *complete probability measure space*. We note: In any probability space, it is therefore possible to enlarge the  $\sigma$  field and extend the measure in such a way as to get a complete space.

Suppose that  $((0, 1], \mathcal{B}, \lambda)$  is completed in this way. The sets in the completed  $\sigma$  field  $\mathcal{M}$  are called *Lebesgue Sets*, and  $\lambda$  extended to  $\mathcal{M}$  is still called the Lebesgue measure.

**Nonmeasurable Sets** Above, we have focused on what sets are in  $\mathcal{B}$ , and finding more measurable sets. Now, we find a set that is *outside of*  $\mathcal{B}$ . We need the following definitions:

**Definition: Set Shifts/Translations** For  $x, y \in (0, 1]$ , take  $x \oplus y$  to be  $x + y$  or  $x + y - 1$  according as  $x + y$  lies in  $(0, 1]$  or not - this is the circle group, essentially. Define  $A \oplus x = \{a \oplus x : a \in A\}$ . Let  $\mathcal{L}$  be the class of Borel sets  $A$  such that  $A \oplus x$  is a Borel set and  $\lambda(A \oplus x) = \lambda(A)$ . Then  $\mathcal{L}$  is a  $\lambda$  system containing the intervals, and so  $\mathcal{B} \subset \mathcal{L}$  by the  $\pi - \lambda$  theorem. Note, it is clear that:

1. The intervals are contained in  $\mathcal{L}$ .
2. The intervals are a  $\pi$  system.
3. Complements are in  $\mathcal{L}$ , as we can take  $\lambda(A^c \oplus x) = 1 - \lambda(A \oplus x) = 1 - \lambda(A) = \lambda(A^c)$ .
4. Disjoint unions are in  $\mathcal{L}$ :

$$\lambda \left( \bigcup_n A_n \oplus x \right) = \lambda \left( \bigcup_n (A_n \oplus x) \right) = \sum_n \lambda(A_n \oplus x) = \sum_n \lambda(A_n) = \lambda \left( \bigcup_n A_n \right)$$

And so, we do indeed have that  $\mathcal{B} \subset \mathcal{L}$  by the  $\pi - \lambda$  theorem, and  $A \in \mathcal{B}$  implies that  $A \oplus x \in \mathcal{B}$ , and  $\lambda(A \oplus x) = \lambda(A)$ . In this sense -  $\lambda$  is translation invariant on  $\mathcal{B}$ .



**Vitali Sets** Define  $x$  and  $y$  to be equivalent ( $x \sim y$ ) if  $x \oplus r = y$  for some rational  $r$  in  $(0, 1]$ . Let  $H$  be a subset of  $(0, 1]$  consisting of exactly one representative point from each equivalence class - such a set exists via the *axiom of choice*. Note - I believe that  $H$  is uncountable, as I think there are uncountably such equivalence classes - but, I'm not sure, and I'm not gonna try and prove it here.

Now, consider the countably many sets  $H \oplus r$  for rational  $r$ . These sets are disjoint - as no two distinct points of  $H$  are equivalent, and so if  $H \oplus r_1$  and  $H \oplus r_2$  share the point  $h_1 \oplus r_1$  and  $h_2 \oplus r_2$ , then  $h_1 \sim h_2$ , then  $h_1, h_2 \in H$  satisfy  $h_1 \sim h_2$ , which contradicts the definition of  $H$ .

Note that each point of  $(0, 1]$  lies in one of these sets, because  $H$  has a representative from each equivalence class, and each point can be expressed as  $h \oplus r$  for some  $h \in H$  and  $r \in \mathbb{Q}$ . Thus:

$$(0, 1] = \bigcup_r (H \oplus r)$$

Now, we prove that  $H$  is outside of  $\mathcal{B}$  via a contradiction. Assume that  $H \in \mathcal{B}$  - then, translation invariance implies each  $H \oplus r$  is in  $\mathcal{B}$ . By countable additivity, we have:

$$1 = \lambda((0, 1]) = \lambda\left(\bigcup_r (H \oplus r)\right) = \sum_r \lambda(H \oplus r) = \sum_r \lambda(H)$$

If  $\lambda(H) = 0$ , we have a contradiction. If  $\lambda(H) > 0$ , we have a contradiction. Thus, we have a contradiction in all cases, and  $H$  is outside of  $\mathcal{B}$ .  $H$  is called a *Vitali Set*.

## Problems

### 3.1 Finite vs. Countable Additivity

1. In the proof of Theorem 3.1 the assumed finite additivity of  $P$  is used twice and the assumed countable additivity of  $P$  is used once. Where? Finite additivity is used once in Lemma 4 to prove the condition for  $A \in \mathcal{F}_0 \implies A \in \mathcal{M}$ . In Lemma 5, we use countable additivity (via countable subadditivity) to prove  $P^*(A) = P(A)$ , and finite subadditivity via monotonicity.
2. Show by example that a finitely additive probability measure on a field may not be countably subadditive. Show in fact that if a finitely additive probability measure is countably subadditive, then it is necessarily

countably additive as well.

We do the second part first. By countable subadditivity, we have:

$$P\left(\bigcup_n A_n\right) \leq \sum_n P(A_n)$$

Assume  $A_n$  are disjoint. Recall - monotonicity *only* requires finite additivity. And so, we have:

$$\begin{aligned} \sum_{n=1}^k P(A_n) \leq P(A) &\implies \lim_{k \rightarrow \infty} P(A_n) \leq P(A) \implies P\left(\bigcup_n A_n\right) \geq \sum_n P(A_n) \\ &\implies P\left(\bigcup_n A_n\right) = \sum_n P(A_n) \end{aligned}$$

For an example, consider problem 2-15 (which we did above). This is a  $P$  that is finitely but not countably additive.

3. Suppose Theorem 2.1 (Continuity From Above, Below, and Countable Subadditivity) were weakened by strengthening its hypothesis to the assumption that  $\mathcal{F}$  is not a  $\sigma$ -field. Why would this weakened result not suffice for the proof of Theorem 3.1?

In Lemma 2 - we make use of countable subadditivity to prove the Lemma 2 property - we cannot prove Theorem 3.1 without first concluding that  $\mathcal{M}$  is a  $\sigma$  field, which relies on Lemma 2. Also, in Lemma 5, we rely on countable subadditivity to prove  $P^*(A) = P(A)$ .

### 3.2 Redefining the Inner and Outer Measure

Let  $P$  be a probability measure on a field  $\mathcal{F}_0$  and for every subset  $A$  of  $\Omega$  define  $P^*(A)$  by the outer measure. Denote also by  $P$  the extension of  $P$  to  $\mathcal{F} = \sigma(\mathcal{F}_0)$ .

1. Show that:

$$P^*(A) = \inf [P(B) : A \subset B, B \in \mathcal{F}]$$

And:

$$P_*(A) = \sup [P(C) : C \subset A, C \in \mathcal{F}]$$

And show that the infimum and supremum are always achieved.

For the first case - note that:

$$P^*(A) = \inf \left[ \sum_n P(A_n) : A \subseteq \cup A_n, A_n \in \mathcal{F}_0 \right] \geq \inf [P(B) : A \subset B, B \in \mathcal{F}]$$

As for why this is - take any covering of  $A \subseteq \cup A_n, A_n \in \mathcal{F}_0$ , and note that  $\cup A_n \in \mathcal{F}$ , and  $\sum P(A_n) \geq P(\cup A_n)$ . Now, take  $B \in \mathcal{F}$ , and note that  $B \in \mathcal{M}$  by the extension theorem, and so:

$$\begin{aligned} P(B) &= P^*(B) = P^*(A \cap B) + P^*(A^c \cap B) \implies P(B) = P^*(A) + P^*(A^c \cap B) \\ &\implies P(B) \geq P^*(A) \end{aligned}$$

Thus, we have:

$$P^*(A) \leq \inf [P(B) : A \subset B, B \in \mathcal{F}]$$

Giving us our first equality. Now, we want to show:

$$P_*(A) = 1 - P^*(A^c) = \sup [P(C) : C \subset A, C \in \mathcal{F}]$$

By the above part, and definitions, we have:

$$P_*(A) = \inf [1 - P(C) : A^c \subset C, C \in \mathcal{F}]$$

Note that  $A^c \subset C$  if and only if  $C^c \subset A$ . If we consider  $C^c \subset A$  - we note that  $1 - P(C)$  is minimized if and only if  $1 - P(C^c) = P(C)$  is maximized. Thus, the above equals:

$$= \sup [1 - P(C^c) : C^c \subset A, C \in \mathcal{F}] = \sup [1 - P(C^c) : C^c \subset A, C^c \in \mathcal{F}]$$

Note, the second step is because  $C \in \mathcal{F}$  if and only if  $C^c \in \mathcal{F}$ . Now, note that the above expression can be simplified to:

$$= \sup [P(C) : C \subset A, C \in \mathcal{F}]$$

The final step is to show that these are achievable - ie, there are sets  $B$  and  $C$  such that:

$$P^*(A) = P(B) \quad P_*(A) = P(C)$$

Assume not achievable in the first case. Then, we have  $P(B_n)$  such that:

$$P^*(A) = \lim_n P(B_n)$$

As  $\mathcal{F}$  is a  $\sigma$  field, we have that  $\cap B_n \in \mathcal{F}$ . Let  $C_n$  be the incremental intersections - note  $C_n \downarrow \cap B_n$ , and:

$$P^*(A) \leq \lim_n P(C_n) \leq \lim_n P(B_n) \implies P(\cap B_n) = P^*(A)$$

And so, we have a contradiction. Continuity from below will give us a similar result for the supremum, and so in both cases, we have the infimum and supremum are always achieved.

2. Show that  $A$  is  $P^*$  measurable if and only if  $P_*(A) = P^*(A)$ . This is equivalent to:

$$1 - P^*(A^c) = P^*(A) \implies P^*(A) + P^*(A^c) = 1$$

Note that if  $A$  is  $P^*$  measurable, the above is clearly true via definition, taking  $E = \Omega$ . Now, assume  $P_*(A) = P^*(A)$ . We can take sets in  $\mathcal{F}$  such that  $A_1 \subset A \subset A_2$ , with:

$$P(A_1) = P_*(A) = P^*(A) = P(A_2)$$

Note, this is because the infimum and supremum are achievable in the previous part. Now, I will show that  $A \in \mathcal{M}$ , which implies  $A$  is  $P^*$  measurable. We have that:

$$A = A_1 \cup (A \setminus A_1)$$

Now, we note that, by the properties of a probability measure:

$$P(A_2 \setminus A_1) = P(A_2) - P(A_1) = 0$$

As  $\mathcal{M}$  is complete, and  $A \setminus A_1 \subseteq A_2 \setminus A_1$ , we have that  $A \setminus A_1 \in \mathcal{M}$ . Thus, as  $\mathcal{M}$  is a field, and closed under unions, we have that  $A \in \mathcal{M}$  as well.

3. The outer and inner measures associated with a probability measure  $P$  on a  $\sigma$  field  $\mathcal{F}$  are usually *defined* by the above infimum and supremum. Show that the infimum and supremum are the same as our original outer and inner measure definitions with  $\mathcal{F}$  in the role of  $\mathcal{F}_0$ .

This much is clear - because  $\mathcal{F} = \sigma(\mathcal{F})$ , and we showed that the definitions are the same in the first part.

### 3.3 Countable Additivity and the Outer Measure

This problem relies on 2.13, 2.15, and 3.2 - all of which we have done. For the following examples, described  $P^*$  as the original outer measure and  $\mathcal{M} = \mathcal{M}(P^*)$  by our condition with subsets  $E$ . Sort out the cases in which  $P^*$  fails to agree with  $P$  on  $\mathcal{F}_0$ , and explain why:

1. Let  $\mathcal{F}_0$  consist of sets  $\emptyset, \{1\}, \{2, 3\}$  and  $\Omega = \{1, 2, 3\}$  and define probability measures  $P_1$  and  $P_2$  on  $\mathcal{F}_0$  by  $P_1(1) = 0$  and  $P_2(2, 3) = 0$ . Note that  $\mathcal{M}(P_1^*)$  and  $\mathcal{M}(P_2^*)$  differ.

Well, the  $\sigma$  algebras must differ on the sets they include - which would be interesting. I guess, going from the back - this would be because of completeness - maybe they differ on including  $\{2\}$ . In  $\mathcal{M}(P_1^*)$ , we find:

$$1 = P_1^*(2, 3) \quad P_1^*(2 \cap 2, 3) + P_1^*(2^c \cap 2, 3) = P_1^*(2) + P_1^*(3) = 2$$

$$0 = P_2^*(2, 3) \quad P_2^*(2 \cap 2, 3) + P_2^*(2^c \cap 2, 3) = P_2^*(2) + P_2^*(3) = 0$$

So,  $\mathcal{M}(P_2^*)$  contains  $\{2\}$ . The completeness comment comes from only  $\mathcal{M}(P_2^*)$  containing a set with measure 0, and nonempty subsets, which led to the example, I guess.

2. Suppose that  $\Omega$  is countably infinite, let  $\mathcal{F}_0$  be the field of finite and cofinite sets, and take  $P(A)$  to be 0 or 1 as  $A$  is finite or cofinite.

By 2-13, as  $\Omega$  is countably infinite, we have that  $P$  is *not* countably additive. And so, our theorem 3.1 does not apply - we need  $P$  to be countably infinite to prove that  $P^*(A) = P(A)$  for  $A \in \mathcal{F}_0$ .

And so, we should have such a set. Take a cofinite set in  $\Omega$  - which is just a countable set  $A$ . We have that:

$$P(A) = 1$$

However,  $A$  can be covered by a union of finite sets - in which case, the infimum would be 0, and  $P^*(A) = 0$ , which means that  $P^*(A) \neq P(A)$ . Countable additivity would have implies that the sum of disjoint singletons should be equal to  $P(A)$ .

3. The same, but suppose that  $\Omega$  is uncountable. Well, in 2-13, we have that  $P$  is a countably additive probability measure on our field, and so it can be properly extended.

4. Suppose that  $\Omega$  is uncountable, let  $\mathcal{F}_0$  consist of the countable and the cocountable sets, and take  $P(A)$  to be 0 or 1 as  $A$  is countable or cocountable. We have that  $\mathcal{F}_0$  is a  $\sigma$  field, and  $P$  is a countably additive probability measure - so, the theorems should apply.
5. The probability in Problem 2.15. Well, in that problem, we have that the probability measure is not countably additive. Take the set  $A = (1/2, 1]$ . We have that  $P(A) = 1$ . However, we also find:

$$A = \bigcup A_n \quad A_n = (1/2 + 1/(n+2), 1]$$

And so, our infimum would give us  $P^*(A) = 0$ , and we get a disagreement again.

6. Let  $P(A) = I_A(\omega_0)$  for  $A \in \mathcal{F}_0$ , and assume  $\{\omega_0\} \in \sigma(\mathcal{F}_0)$ . Well, it is countably additive - there are no disjoint sets containing  $\omega_0$ . We have additivity in all other cases (disjoint sets not containing  $\omega_0$ , and disjoint sets with one containing  $\omega_0$ ).

So,  $P^*$  and  $\mathcal{M}$  are well defined. If we want to describe  $\mathcal{M}$  - we need sets where:

$$\inf [P(B) : A \subset B, B \in \mathcal{F}] = \sup [P(C) : C \subset A, C \in \mathcal{F}]$$

I think  $\mathcal{M}$  consists of all sets containing  $\omega_0$  and their complements. First, note  $P(\omega_0) = 1$ . Any cover of  $\omega_0$  by sets in  $\mathcal{F}_0$  would have a sum greater than or equal to 1. Now, take  $A$  containing  $\omega_0$ . The supremum contains 1, and cannot be larger than 1. Similarly, the infimum cannot be smaller than 1 - as any set in  $\mathcal{F}$  that contains  $A$  is a set in  $\mathcal{F}$  that contains  $\omega_0$ , and so must be  $P = 1$  by monotonicity.

So, if  $A$  contains  $\omega_0$ , it is in  $\mathcal{M}$ , with  $P^*(A) = 1$ . Similarly, if it doesn't, it has  $P^*(A) = 0$ , as the complement is in it. Note,  $\mathcal{M}$  thus contains all subsets, and their probabilities are either 1 or 0 based on this condition. qed.

### 3.6 Extension for a Finitely Additive Probability Measure on a Field

Let  $P$  be a *finitely additive* probability measure on a field  $\mathcal{F}_0$ . For  $A \subset \Omega$ , in analogy with (3.1), define:

$$P^\circ(A) = \inf \sum_n P(A_n)$$

Over *finite* sequences of  $\mathcal{F}_0$  sets  $A_n$ . Let  $\mathcal{M}^\circ$  be the class of sets  $A$  such that:

$$P^\circ(E) = P^\circ(A \cap E) + P^\circ(A^c \cap E)$$

For all  $E \subset \Omega$ .

1. Show that  $P^\circ(\emptyset) = 0$  and that  $P^\circ$  is nonnegative, monotone, and *finitely* subadditive.

$P^\circ(\emptyset) \leq P(\emptyset) = 0$ . Note, as all the sums are greater than 0,  $P^\circ(\emptyset) = 0$ . Clearly, this implies non-negative as well. Monotonicity, as if  $A \subset B$ , any cover of  $B$  covers  $A$ , and so that sum is included in  $A$ 's infimum. Take any finite union  $\cup_n A_n$ . Examine:

$$P^\circ(\cup_n A_n)$$

Note, finite subadditivity can just be found via the  $\epsilon/2^k$  trick. We have a finite cover of each  $A_n$ , with total sum within  $\epsilon/2^n$  of  $P^\circ(A_n)$  - note, a finite amount of finite covers still covers  $\cup_n A_n$ , thus, we have:

$$P^\circ(\cup_n A_n) \leq \sum_n P^\circ(A_n) + \epsilon/2^n = \epsilon + \sum_n P^\circ(A_n)$$

Using these four properties, prove:

**Lemma 1**  $\mathcal{M}^\circ$  is a field.

First, we have:

$$P^\circ(E) = P^\circ(E) + P^\circ(\emptyset) = P^\circ(\Omega \cap E) + P^\circ(\emptyset \cap E)$$

And so  $\mathcal{M}^\circ$  contains  $\Omega$  and  $\emptyset$ . Now, we go over complements - as the  $E$  definition is symmetric between complements, we have that  $A \in \mathcal{M}^\circ \implies A^c \in \mathcal{M}^\circ$ . Now, we prove that  $\mathcal{M}^\circ$  is closed under finite unions (or equivalently, finite intersections). Take  $A, B \in \mathcal{M}^\circ$ , and  $E \subset \Omega$ . We have:

$$P^\circ(E) = P^\circ(B \cap E) + P^\circ(B^c \cap E)$$

$$= P^\circ(A \cap B \cap E) + P^\circ(A^c \cap B \cap E) + P^\circ(A \cap B^c \cap E) + P^\circ(A^c \cap B^c \cap E)$$

Just by applying  $A, B \in \mathcal{M}^\circ$  three times. By *finite subadditivity*, we find:

$$\geq P^\circ(A \cap B \cap E) + P^\circ((A^c \cap B \cap E) \cup (A \cap B^c \cap E) \cup (A^c \cap B^c \cap E))$$

$$= P^\circ(A \cap B \cap E) + P^\circ((A \cap B)^c \cap E)$$

As we already have the other direction via finite subadditivity, we thus have  $A \cap B \in \mathcal{M}^\circ$ , and  $\mathcal{M}^\circ$  is a field.

**Lemma 2** If  $A_1, A_2, \dots$  is a *finite* sequence of disjoint  $\mathcal{M}^\circ$  sets, then for each  $E \subset \Omega$ :

$$P^\circ\left(E \cap \left(\bigcup_k A_k\right)\right) = \sum_k P^\circ(E \cap A_k)$$

If  $n = 1$ , there is nothing to prove. Now, assume  $n = 2$  - if  $A_1 \cup A_2 = \Omega$ , then  $A_2 = A_1^c$ , and it is just our condition. Otherwise:

$$P^\circ(E \cap (A_1 \cup A_2)) = P^\circ(A_1 \cap E \cap (A_1 \cup A_2)) + P^\circ(A_1^c \cap E \cap (A_1 \cup A_2))$$

Where the second step again makes use of our condition for  $E \cap (A_1 \cup A_2)$  and  $A_1 \in \mathcal{M}^\circ$ . The above equals:

$$= P^\circ(E \cap A_1) + P^\circ(E \cap A_2)$$

Where we have that  $A_1^c \cap E \cap (A_1 \cup A_2) = E \cap A_2$  via disjointness. Ie,  $A_2 \subseteq A_1^c$ , so  $x \in A_2 \implies x \in A_1^c$ . It is easy to prove via induction for an arbitrary length finite sequence of disjoint  $\mathcal{M}^\circ$  sets.

**Lemma 3**  $P^\circ$  restricted to the field  $\mathcal{M}^\circ$  is *finitely additive*. Take a finite disjoint sequence  $A_1, A_2, \dots$  of  $\mathcal{M}^\circ$  sets. By Lemma 1,  $\cup A_n \in \mathcal{M}$ . We want to show:

$$P^\circ(\cup A_n) = \sum P^\circ(A_n)$$

Just apply Lemma 2 with  $E = \Omega$ . Thus, we have:

$$P^\circ(\cup A_n) = P^\circ(\Omega \cap \cup A_n) = \sum P^\circ(\Omega \cap A_n) = \sum P^\circ(A_n)$$

2. Show that if  $P^\circ$  is defined by the finite coverings definition - then, **Lemma 4:**  $\mathcal{F}_0 \subset \mathcal{M}^\circ$ .

Take  $A \in \mathcal{F}_0$ . We want to show, for  $E \subset \Omega$ :

$$P^\circ(E) \geq P^\circ(A \cap E) + P^\circ(A^c \cap E)$$

Well, we have for some finite union  $E \subset \sup_n A_n$ ,  $\sum P(A_n) \leq P^\circ(E) + \epsilon$ . We have:

$$P^\circ(A \cap E) + P^\circ(A^c \cap E) \leq \sum_n P(A \cap A_n) + \sum_n P(A^c \cap A_n)$$



Which is by monotonicity of  $P^\circ$  and finite subadditivity. Now, with finite additivity of disjoint sets:

$$= \sum_n P(A_n) \leq P^\circ(E) + \epsilon$$

Thus,  $A \in \mathcal{M}^\circ$ .

**Lemma 5** Show that  $P^\circ(A) = P(A)$  for  $A \in \mathcal{F}_0$ . Clearly,  $P^\circ(A) \leq P(A)$ . Also, if  $A \subset \cup A_n$  for finite  $A_n$ , then  $P(A) \leq \sum_n P(A \cap A_n) \leq \sum_n P(A_n)$ , so  $P(A)$  is less than all elements in the infimum, and  $P^\circ(A) = P(A)$ .

3. Define  $P_\circ(A) = 1 - P^\circ(A^c)$ . Prove that if  $E \subset A \in \mathcal{F}_0$ , then:

$$P_\circ(E) = P(A) - P^\circ(A - E)$$

We have, by finite additivity:

$$P_\circ(E) = 1 - P^\circ(E^c) = 1 - P^\circ(E^c \cap A) - P^\circ(E^c \cap A^c) = 1 - P^\circ(A - E) - P^\circ(A^c)$$

As  $A^c \subset E^c$ . Note,  $P^\circ(A^c) = P(A^c)$ , and  $1 - P(A^c) = P(A)$ , and so:

$$P_\circ(E) = P(A) - P^\circ(A^c)$$

### 3.7 Finitely Additive Field Extension

Relies on 2.7 and 3.6, both of which we have solutions for. Suppose that  $H$  lies outside the field  $\mathcal{F}_0$ , and let  $\mathcal{F}_1$  be the field generated by  $\mathcal{F}_0 \cup \{H\}$ . By 2.7, this field consists of sets of the form:

$$(H \cap A) \cup (H^c \cap B) \quad A, B \in \mathcal{F}_0$$

In this problem, we show that a finitely additive probability measure  $P$  on  $\mathcal{F}_0$  has a finitely additive extension to  $\mathcal{F}_1$ . Define  $Q$  on  $\mathcal{F}_1$  by:

$$Q((H \cap A) \cup (H^c \cap B)) = P^\circ(H \cap A) + P_\circ(H^c \cap B)$$

1. Show that the definition is consistent. Ie, that if:

$$(H \cap A_1) \cup (H^c \cap B_1) = (H \cap A_2) \cup (H^c \cap B_2)$$

Then:

$$Q[(H \cap A_1) \cup (H^c \cap B_1)] = Q[(H \cap A_2) \cup (H^c \cap B_2)]$$

Well, if they are equal - then we must have  $H \cap A_1 = H \cap A_2$ , as it is a disjoint union. Similarly,  $H^c \cap B_1 = H^c \cap B_2$ . Thus, the definition is consistent, as clearly:

$$P^\circ(H \cap A_1) = P^\circ(H \cap A_2) \quad P_\circ(H^c \cap B_1) = P_\circ(H^c \cap B_2)$$

2. Show that  $Q$  agrees with  $P$  on  $\mathcal{F}_0$ . Take  $A \in \mathcal{F}_0$ . We have  $A = (H \cap A) \cup (H^c \cap A)$ . Thus:

$$Q(A) = P^\circ(H \cap A) + P_\circ(H^c \cap A)$$

By Part (c) of the previous question, as  $H^c \cap A \subset A \in \mathcal{F}_0$  the above equals:

$$= P^\circ(H \cap A) + P(A) - P^\circ(A - (H^c \cap A)) = P(A)$$

3. Show that  $Q$  is finitely additive on  $\mathcal{F}_1$ . Show that  $Q(H) = p^\circ(H)$ .

Take disjoint sets  $C_1, \dots, C_n$  in  $\mathcal{F}_1$ , which must be of the form  $(H \cap A_i) \cup (H^c \cap B_i)$  for  $A_i, B_i \in \mathcal{F}_0$ . We find:

$$\begin{aligned} Q(\cup C_i) &= Q\left(\bigcup (H \cap A_i) \cup (H^c \cap B_i)\right) = Q\left(\left(H \cap \bigcup A_i\right) \cup \left(H^c \cap \bigcup B_i\right)\right) \\ &= P^\circ\left(H \cap \bigcup A_i\right) + P_\circ\left(H^c \cap \bigcup B_i\right) \end{aligned}$$

Note, each of the  $A_i$  and  $B_i$  were disjoint, and so they are disjoint unions above. We note that each  $A_i$  is an  $\mathcal{M}^\circ$  set, and so by Lemma 2 above, the above equals:

$$= \sum_i P^\circ(H \cap A_i) + P_\circ\left(H^c \cap \bigcup B_i\right)$$

For the second term, note that  $H^c \cap \bigcup B_i \subset \bigcup B_i \in \mathcal{F}_0$ , and so by part (c) in the previous question, the above equals:

$$= \sum_i P^\circ(H \cap A_i) + P\left(\bigcup B_i\right) - P^\circ\left(H \cap \bigcup B_i\right)$$

Breaking down the middle term by finite additivity on  $\mathcal{F}_0$ , and the final term by Lemma 2 in the previous question, the above equals:

$$= \sum_i P^\circ(H \cap A_i) + P(B_i) - P^\circ(H \cap B_i)$$

Where the second and middle terms now vary with the  $i$  in the sum. Finally, we make use of part (c) above again, noting that  $H^c \cap B_i \subset B_i$ , and  $B_i - H^c \cap B_i = H \cap B_i$ :

$$= \sum_i P^\circ(H \cap A_i) + P_\circ(H^c \cap B_i) = \sum_i Q(C_i)$$

Now, we show that  $Q(H) = P^\circ(H)$ . We have that:

$$Q(H) = Q((H \cap \Omega) \cup (H^c \cap \emptyset)) = P^\circ(H \cap \Omega) + P_\circ(H^c \cap \emptyset) = P^\circ(H)$$

So - what we have now is that  $Q$  is finitely additive on  $\mathcal{F}_1$ . It is also nonnegative, clearly, and  $Q(\Omega) = P(\Omega) = 1$ . It also clearly respects monotonicity, by  $P^\circ$  respecting it. Thus,  $Q$  is a finitely additive probability measure on  $\mathcal{F}_1$ . Further,  $Q$  agrees with  $P$  on  $\mathcal{F}_0$ , and so it is indeed an extension. Thus, we have proved the main points of the problem.

4. Define  $Q'$  by interchanging the roles of  $P^\circ$  and  $P_\circ$  on the right hand side of the definition of  $Q$ , namely:

$$Q'((H \cap A) \cup (H^c \cap B)) = P_\circ(H \cap A) + P^\circ(H^c \cap B)$$

Show that  $Q'$  is another finitely additive extension of  $P$  to  $\mathcal{F}_1$ . The same is true of any convex combination of  $Q''$  of  $Q$  and  $Q'$ . If  $Q'$  is a finitely additive extension - so to would be the additions of  $Q'$  and  $Q''$ , and we maintain being an extension as the fractions of a convex combination add to 1. Show that  $Q''(H)$  can take any value between  $P_\circ(H)$  and  $P^\circ(H)$ . Well, that last part is clear, if the previous parts are true.

So, the only thing to prove is that  $Q'$  is another finitely additive extension of  $P$  to  $\mathcal{F}_1$ . We will not show that here - it is just an extension of the above problems, and I don't want to do it again.

### 3.8 Finitely Additive Field Extension to Power Set Field

Use Zorn's lemma to prove a theorem of Tarski: A finitely additive probability measure on a field has a finitely additive extension to the field of all subsets of the space.

Note - this makes use of the previous problem. Also note - the last part of 3.7d also now has more of a use, I think. For the original finitely additive probability measure  $P$  on field  $\mathcal{F} \subset 2^\Omega$ , with each subsequent expansion to  $H \in 2^\Omega - \mathcal{F}$ , we can choose any measure value for it between  $P_\circ(H)$  and  $P^\circ(H)$ . As for why - note that when we expand the field - we expand the number of sets that can finitely cover  $H$ . However,  $H$  can still be covered by sets in the original  $\mathcal{F}$ , and so the values between  $P_\circ(H)$  and  $P^\circ(H)$  would actually be a subset of the possible values that  $H$  can take on (actually, not

sure this is correct... we just proved the existence of an extension, not that all extensions agreed on sets outside of  $\mathcal{F}$ ).

Zorn's Lemma, as stated in my Topology book: Let  $(M, \leq)$  be a partially ordered set. Suppose every chain  $K \subset M$  is bounded. Then  $M$  has a maximal element, ie there is an  $a \in M$  such that no  $x \in M$  satisfies  $x > a$ .

I was originally thinking about chains of sets - but maybe chains of fields that extend  $\mathcal{F}$ . First, take any chain of such fields  $\mathcal{I}$ :

$$\mathcal{F}_i \quad i \in \mathcal{I} \quad \implies \quad i, j \in \mathcal{I} \implies \mathcal{F}_i \leq \mathcal{F}_j \text{ or } \mathcal{F}_j \leq \mathcal{F}_i$$

The notation above is because  $\mathcal{I}$  doesn't have to be countable. We note that:

$$\bigcup_{i \in \mathcal{I}} \mathcal{F}_i$$

Is a field as well - over any arbitrary unions. Note, this is not the case in general - but the  $\mathcal{F}_i$  are *ordered*. First off, it is clearly closed under complements, and contains  $\Omega$  and  $\emptyset$ . Now, take  $A_1, \dots, A_n \in \bigcup_{i \in \mathcal{I}} \mathcal{F}_i$ . We note that there is an order between each of the  $\mathcal{F}_i$  containing the  $A_i$  - so, take the largest one, which contains all  $A_i$ , and thus their disjoint union. Thus, we have found a maximal element for our chain.

So, our partially ordered set  $(M, \leq)$  where  $M$  contains fields containing  $\mathcal{F}$  that agree with  $P$ , must have a maximal element. This is because every chain is bounded. Now, this maximal element must contain each of the extensions of  $\mathcal{F}$  to one arbitrary set in  $2^\Omega$ . So, the maximal element must contain every set in  $2^\Omega$ , and so it is just the field on  $2^\Omega$ .

So, the existence of a maximal element implies our theorem: a finitely additive probability measure  $P$  on a field  $\mathcal{F}$  has a finitely additive extension to the field of all subsets of the space. qed.