


C5aR-dependent microglia activity

**Alex Keyes
Yarik Andrianov
Sasha Dovgan
Boris Olifirov
Oleh Halaidych
Pavel Belan**


**Department seminar
17.02.2023
Kyiv – Iowa City**

What is a level of intrinsic spontaneous microglia activity in particular induced by the spinal cord extraction from mice?



Whether C5a application can produce a significant and substantial increase in microglia activity in order to warrant our electrophysiological experiments?



BSA
BSA+C5a



Raw traces E0002, before wash



What is a level of intrinsic spontaneous microglia activity in particular induced by the spinal cord extraction from mice?

Substantial spontaneous microglia activity does exist.


Whether C5a application can produce a significant and substantial increase in microglia activity in order to warrant our electrophysiological experiments?

It is likely that C5aR activation increases the frequency of microglial $[Ca^{2+}]_i$ transients 2-3 times. We need to process the results of already performed experiments. Possibly additional experiments will be necessary.


Whether the level of intrinsic spontaneous microglia activity is increased due to the spinal cord extraction from mice over the time course of our electrophysiological experiments (30-60 min)?

We need to conduct 3-5 experiments with 30-60 min recording of spontaneous microglia activity.

Whether perfusion via different syringes and BSA can produce significant changes in microglia activity?



BSA
BSA+C5a



Whether perfusion via different syringes and BSA can produce significant changes in microglia activity?


It seems that changes in perfusion lines can create substantial artefacts in a baseline and XY movement. We would recommend to add drugs to the same perfusion line if possible.

How spatially C5a-dependent microglia activity is organized?


Whether C5a-dependent microglia activity is mainly newly emerged or is it rather increase in frequency of previously spontaneous active sites?

Is C5a-dependent microglia activity randomly distributed over FOV or rather concentrated in certain sites?



Norm traces F7, silent





ROI9
ROI23
ROI30
ROI31
ROI32
ROI36
ROI67



Norm traces F7, active




Raw traces E0002, before wash




Norm traces E0002, active

C5a



ROIs with higher or equal initial activity (C5a Application)



How spatially C5a-dependent microglia activity is organized?

Whether C5a-dependent microglia activity is mainly newly emerged or is it rather increase in frequency of previously spontaneous active sites?


It is likely that the most active sites are newly emerged. Probably there is also increase of activity in spontaneously active sites. Additional experiments should be performed or processed.

Is C5a-dependent microglia activity randomly distributed over FOV of rather concentrated in certain sites?


Possibly regions of microglia activity are segregated. Additional analysis is warranted.

How spatially C5a-dependent microglia activity is organized?


Whether C5a-dependent microglia activity is mainly emerged in soma or processes of microglia cells?




- ROI9
- ROI23
- ROI30
- ROI31
- ROI32
- ROI36
- ROI67



Separate ROIs



Combined ROIs



How spatially C5a-dependent microglia activity is organized?

Possibly C5a activates microglia in the restricted areas of spinal cord. Thus, neurons located in this area are likely to be affected while other neurons will not change their activity in C5aR-dependent manner.

Whether C5a-dependent microglia activity is mainly emerged in soma or processes of microglia cells?



It seems that the effect of C5aR activation is mainly manifested in microglia processes although somas can be also involved. We think we need to show separate statistics for somas and processes.

What is a time course of C5a-induced microglia activity?



Is there some desensitization of C5aRs?

Is the effect of C5a washable?


Norm traces E0002, silent




Norm traces E0002, active






E0002 silent C5a+wash



E0002 before C5a+wash



20=16



What is a time course of C5a-induced microglia activity?

The effect is not instantaneous (about 1 min).

Is there some desensitization of C5aRs?


It seems that the effect of C5a is maximal at the beginning and slowly and gradually diminished. Having the time course of C5a effect on microglia we may predict a time course of possibly changes in neuronal activity.

Is the effect of C5a washable?

Most probably NO. It means we need to carry out experiments with PMX in a design when it is pre-applied before C5a application (3-5 experiments are necessary).

Possibly PMX can stop the effect of C5a (both during C5a application and after C5a application. We may check it experimentally.

What readouts of microglia activity we suggest



Events (3).xlsx - Excel (Pr.)

Can't Redo (Ctrl+Y)


D1

| | A | B | C | D | E |
|----|-------|------|----------|----------|----------|
| 1 | ROI | Time | Ampl | Duration | Integral |
| 2 | ROI2 | 647 | 6.182424 | 6.825691 | 42.19932 |
| 3 | ROI2 | 1470 | 0.539626 | 3.201683 | 1.727711 |
| 4 | ROI10 | 1301 | 0.38519 | 5.351811 | 2.061466 |
| 5 | ROI10 | 1232 | 0.532264 | 5.271084 | 2.805606 |
| 6 | ROI10 | 1207 | 0.319488 | 7.401665 | 2.364741 |
| 7 | ROI10 | 1320 | 0.309388 | 11.85273 | 3.667094 |
| 8 | ROI10 | 1181 | 0.414709 | 6.084139 | 2.523148 |
| 9 | ROI12 | 957 | 1.059184 | 25.13817 | 26.62594 |
| 10 | ROI12 | 933 | 0.740447 | 7.278934 | 5.389663 |
| 11 | ROI12 | 1297 | 0.51576 | 8.24647 | 4.253198 |
| 12 | ROI12 | 998 | 0.662614 | 5.131905 | 3.400474 |
| 13 | ROI12 | 1311 | 0.612693 | 5.947062 | 3.643722 |
| 14 | ROI12 | 832 | 0.396104 | 19.00761 | 7.528982 |
| 15 | ROI12 | 899 | 0.29057 | 13.08161 | 3.801127 |
| 16 | ROI13 | 930 | 3.353547 | 5.348798 | 17.93745 |
| 17 | ROI13 | 899 | 1.386385 | 6.78761 | 9.410242 |
| 18 | ROI14 | 930 | 2.625928 | 5.780971 | 15.18041 |
| 19 | ROI14 | 897 | 1.884301 | 7.052169 | 13.28841 |
| 20 | ROI14 | 874 | 0.600153 | 6.534655 | 3.921796 |
| 21 | ROI14 | 1451 | 0.467436 | 5.38029 | 2.514939 |
| 22 | ROI14 | 653 | 0.41595 | 5.366205 | 2.232071 |
| 23 | ROI17 | 896 | 1.032533 | 3.44942 | 3.561641 |
| 24 | ROI17 | 798 | 0.648945 | 11.866 | 7.700384 |
| 25 | ROI17 | 1288 | 0.40937 | 4.383127 | 1.79432 |
| 26 | ROI18 | 1439 | 1.756456 | 7.822781 | 13.74037 |
| 27 | ROI18 | 888 | 0.601253 | 5.581221 | 16.21221 |

Лист1 | E_0002 | Int1 | Int2 | ...

Enter

Cumulative plots for statistical comparisons and mean or median values as parameters to quantify changes



| | A | B | C | D | E |
|----|-------|------|----------|----------|----------|
| 1 | ROI | Time | Ampl | Duration | Integral |
| 2 | ROI28 | 41 | 1.136885 | 6.576673 | 7.476918 |
| 3 | ROI28 | 58 | 0.763939 | 9.273236 | 7.084185 |
| 4 | ROI29 | 64 | 0.567919 | 6.172165 | 3.50529 |
| 5 | ROI32 | 64 | 5.303416 | 5.053387 | 26.80021 |
| 6 | ROI32 | 84 | 3.489153 | 4.286335 | 14.95568 |
| 7 | ROI29 | 85 | 0.421838 | 4.515457 | 1.904791 |
| 8 | ROI32 | 104 | 3.70213 | 4.679989 | 17.32593 |
| 9 | ROI28 | 113 | 0.404412 | 5.310447 | 2.147608 |
| 10 | ROI29 | 117 | 1.054817 | 5.012141 | 5.286891 |
| 11 | ROI51 | 124 | 0.822343 | 4.797511 | 3.9452 |
| 12 | ROI29 | 176 | 0.422791 | 4.996937 | 2.112661 |
| 13 | ROI32 | 176 | 3.462348 | 4.33982 | 15.02597 |
| 14 | ROI54 | 190 | 0.665648 | 6.23369 | 4.149444 |
| 15 | ROI28 | 239 | 0.506738 | 5.004044 | 2.535739 |
| 16 | ROI29 | 241 | 0.643603 | 4.635797 | 2.983614 |
| 17 | ROI11 | 253 | 0.451641 | 6.979348 | 3.15216 |
| 18 | ROI51 | 259 | 1.919223 | 4.704815 | 9.029589 |
| 19 | ROI51 | 270 | 1.204909 | 5.606128 | 6.754876 |
| 20 | ROI2 | 293 | 1.433512 | 4.470963 | 6.409181 |
| 21 | ROI11 | 321 | 0.761567 | 10.6614 | 8.11937 |
| 22 | ROI47 | 322 | 0.328942 | 15.82329 | 5.204944 |
| 23 | ROI51 | 322 | 3.189159 | 3.745654 | 11.94549 |
| 24 | ROI10 | 336 | 0.335901 | 7.033257 | 2.362481 |
| 25 | ROI28 | 353 | 0.170831 | 6.731901 | 1.150015 |
| 26 | ROI11 | 383 | 0.434414 | 6.028257 | 2.618759 |

Using statistics for interevent intervals, amplitudes, duration and integrals we will be able to obtain more comprehensive analysis and dose dependence of C5aR activation.

We need to do check if we can use already performed dose-dependent experiments and process them. Or to do some additional experiments.

Norm traces E0002, before C5a

