

UNIVERSITY OF THE WITWATERSRAND  
JOHANNESBURG

ARCHEAN METALLOGENY IN SOUTHERN AFRICA

by

C. R. ANHAEUSSER

*(Senior Research Fellow, Economic Geology Research Unit)*

ECONOMIC GEOLOGY RESEARCH UNIT

INFORMATION CIRCULAR No. 91

*September, 1974*

## ARCHEAN METALLOGENY IN SOUTHERN AFRICA

### ABSTRACT

The Archean granite-greenstone terrane of southern Africa possesses a number of unique characteristics. Not only is the shield area one of the smallest in the world but it is also one of the oldest crustal fragments still preserved. Estimates of lithic proportions show that, in the southern African Archean, the granitic rocks outnumber those of the volcanic belts (greenstone belts) by approximately 6 : 1 - in marked contrast to the situation on the Canadian Shield where granite : greenstone ratios have been estimated to be approximately 1 : 2.

In order to establish further details of the lithological proportions, the southern African Archean greenstone belts are described in terms of a three-stage stratigraphic model involving a Lower Ultramafic Unit at the base, followed by a Mafic to Felsic Unit and a terminal Sedimentary Unit. Planimeter studies demonstrate that, on average, the southern African greenstone belts possess a relatively high proportion of ultramafic and mafic volcanic and plutonic rock types and a correspondingly lower proportion of rocks of intermediate to felsic composition. The relative abundances of the various volcanic components of the lower two stratigraphic units are tabulated and the lithic proportions are compared with similar data available from greenstone belts of the Superior Province in Canada. Geochemical data illustrating the nature of Archean volcanism in southern Africa, and elsewhere, are made available for purposes of comparison.

The distribution patterns of a wide variety of Archean mineralization types, including the deposits associated with the ancient granites and pegmatites, are illustrated by means of mineral distribution maps. A close genetic relationship exists between the various lithological subdivisions of the Archean greenstone belts and the mineralization. Important deposits of chrome, nickel-copper, gold, and chrysotile asbestos, are associated with the ultramafic-mafic component of the volcanic pile, whereas antimony, gold, and lesser amounts of mercury, barite, corundum, copper-lead-zinc, and massive sulphide iron formations occur throughout the greenstone piles and commonly occur as chemical sediments terminating cycles of volcanism. These well-banded rocks, which probably formed by fumarolic-exhalative processes, also contain significant stratiform gold mineralization.

The Archean mineralization of southern Africa is briefly compared and contrasted with that of the Canadian Shield and elsewhere. It is concluded that the disparity of lithological types and proportions, between the Canadian and southern African shields, is, together with differences in age, the underlying cause of the variability in the mineralization thus far encountered in the two regions.

\* \* \* \* \*

## ARCHEAN METALLOGENY IN SOUTHERN AFRICA

### CONTENTS

	<u>Page</u>
<u>INTRODUCTION</u>	1
<u>THE ARCHEAN GRANITE-GREENSTONE TERRANE OF SOUTHERN AFRICA</u>	1
<i>Nature and Distribution</i>	1
<i>Stratigraphy</i>	1
<i>Lithic Proportions</i>	5
<u>GEOCHEMISTRY OF ARCHEAN VOLCANIC ROCKS</u>	9
<u>ARCHEAN MINERALIZATION</u>	13
<i>Types</i>	13
<u>THE NATURE AND DISTRIBUTION OF ARCHEAN MINERALIZATION IN SOUTHERN AFRICA</u>	14
<i>Mineralization Associated with Ultramafic to Mafic Volcanic and Plutonic Rocks</i>	14
Nickel-Copper	14
Chrysotile Asbestos	16
Magnesite	16
Talc	16
Verdite-Buddstone	17
Chromite Mineralization in Archean Greenstone Belts	17
Other Chromite Occurrences in Southern Africa	19
Chromite Distribution in Southern Africa	19
Gold	21
Regional Distribution of Archean Gold Mineralization	23
<i>Mineralization Associated with Mafic to Felsic Volcanic Rocks</i>	23
Antimony and Mercury	24
Barite	26
Corundum and Kyanite	26
Copper-Lead-Zinc-Gold	27
Massive Sulphide Iron Formations	27
Silver-Tungsten-Arsenic	28
<i>Archean Sedimentary Deposits</i>	29
<i>Mineralization Associated with Granites and Pegmatites</i>	30
<u>DISCUSSION</u>	30
<u>REFERENCES</u>	32

\* \* \* \* \*

## ARCHEAN METALLOGENY IN SOUTHERN AFRICA

### INTRODUCTION

The early Precambrian (Archean) granite-greenstone terrane that forms part of the southern African Shield constitutes one of the oldest known fragments on the earth's surface. Underlying the eastern part of Botswana, the northeastern segment of South Africa, and Rhodesia, the shield consists of two cratons - the Kaapvaal and the Rhodesian cratons - separated by the Limpopo high-grade metamorphic belt. Ranging in age from 2 700-3 500 m.y. (or even older), the region acts as basement to a unique series of cratonic basins which formed under continental conditions as far back in time as 3 000 m.y. ago (Anhaeusser, 1973a; Pretorius, 1974a). Although the bulk of the mineral wealth of the African subcontinent has largely come from these basins (Pretorius, 1973; 1974a, b) the older Archean granite-greenstone terrane has the distinction of having been the principal source of much of this mineralization (which includes gold and uranium), as well as of currently contributing the most diversified range of mineral types and products exploited in southern Africa.

The purposes of this paper are, briefly, to consider the nature and distribution of Archean mineralization in southern Africa and to relate the mineral deposits to one another and to the geological environments in which they occur. Furthermore, it has become apparent that despite the many similarities that exist between the Archean shield areas of the world (Anhaeusser et al., 1969) there are also significant differences, the latter reflected in the variable geochemical nature of the volcanic sequences (Wilson et al., 1965; Baragar and Goodwin, 1969; Viljoen and Viljoen, 1969a, b; Hallberg, 1972), and in the nature of the mineralization.

In the sections which follow selected metallogenic data are used to illustrate the close genetic relationship of mineralization to specific stages in the evolutionary development of the granite-greenstone terrane in southern Africa. Comparisons and contrasts are drawn between this mineralization and that developed in Archean successions elsewhere in the world, particularly in the Canadian Shield.

### THE ARCHEAN GRANITE-GREENSTONE TERRANE OF SOUTHERN AFRICA


#### NATURE AND DISTRIBUTION

A great variety of volcanic, sedimentary and granitic elements make up the Rhodesian and Kaapvaal cratons. Scattered within the expanse of granitic rocks are numerous volcanic (greenstone) belts, the present day configurations of which are shown in Figure 1. Much of the southern African Archean is covered by younger formations which obscure approximately 70 per cent of the cratonic crustal area. The exposed Archean consists predominantly of granitic rock types with the greenstone belt relics constituting a relatively low proportion of the preserved basement. The areal extent and relative proportions of Archean and younger successions developed on the Rhodesian and Kaapvaal cratons is shown in Table 1. By comparison with Archean crustal fragments in Canada and Western Australia (Table 2), the southern African fragments are generally smaller, a feature that may be linked to the latter areas apparently greater antiquity.

Both the Rhodesian and Kaapvaal cratons are bounded by high-grade, polymetamorphic, mobile belts which are considered to represent reworked Archean cratonic material, with or without, infolded, younger, supracrustal rocks (Anhaeusser et al., 1969; Anhaeusser, 1973a; Mason, 1973). As these areas represent separate geotectonic domains they will not be described further except where pertinent to the metallogenic history of the regions discussed in this paper.

#### STRATIGRAPHY

The volcanic and sedimentary piles that comprise the greenstone belts of the Rhodesian basement complex have been grouped into three major subdivisions, referred to as the Sebakwian, Bulawayan and Shamvaian groups respectively (Macgregor, 1951; Bliss and Stidolph, 1969;



**Figure 1 :** Map illustrating the exposed Archean granite-greenstone terrane of the Rhodesian and Kaapvaal cratons, southern Africa.

TABLE 1

AREAL EXTENT AND RELATIVE PROPORTIONS OF ARCHEAN AND YOUNGER SEQUENCES  
DEVELOPED ON THE RHODESTAN AND KAAPVAAL CRATONS

Crustal Fragment	Area (km <sup>2</sup> )	Exposed Archean (km <sup>2</sup> )	Exposed Granites (km <sup>2</sup> )	Exposed Greenstone Belts (km <sup>2</sup> )	Younger Cover (km <sup>2</sup> )	Exposed Archean (%)	Exposed Granites (%)	Exposed Greenstone Belts (%)	Younger Cover (%)
Rhodesian Craton	312 000	195 762	162 027	33 735	116 238	62.7	51.9	10.8	37.3
Kaapvaal Craton	585 000	80 662	73 778	6 884	504 338	13.8	12.6	1.2	86.2
Combined Rhodesian and Kaapvaal Cratons	897 000	276 424	235 805	40 619	620 576	30.8	26.2	4.6	69.2

TABLE 2

AREAL EXTENT OF ARCHEAN CRUSTAL FRAGMENTS IN CANADA  
AND WESTERN AUSTRALIA

Locality	(km) <sup>2</sup>
Superior Province, Canada	2 000 000
Slave Province, Canada	195 000
Yilgarn Block, Western Australia	700 000
Pilbara Block, Western Australia	500 000

Wilson, 1973). At the base, the Sebakwian stratigraphy consists of a variety of mafic and ultramafic rocks with sedimentary interlayers, and is characterized by the presence of intrusive masses of ultramafic rocks which have, in the past, been classed into the so-called Magnesian Series - a name reserved by Macgregor (1951) for a suite of late Sebakwian intrusive ultramafic rocks. In many places the Sebakwian Group has been extensively invaded, fragmented, and often apparently granitized, by the intrusion of a wide variety of granitic rock types. Mafic lavas, often pillowed, together with andesites, felsic lavas, and pyroclasts, and local sedimentary intercalations (usually of the banded iron formation - phyllite association), constitute the overlying Bulawayan Group. The latter assemblage is, in turn, overlain by the Shamvaian Group which is comprised predominantly of quartzites, conglomerates, greywackes, and shales, together with subordinate volcanic rocks, the latter consisting of andesites, and felsic lavas and pyroclasts.

On the Kaapvaal Craton, rocks of an essentially similar character to those of the Rhodesian greenstone belts are collectively referred to as remnants of the Swaziland Sequence (Supergroup), the latter being particularly well-developed in the Barberton Mountain Land. The Onverwacht Group, developed at the base of the sequence, has been subdivided into six formations (Figure 2). The lower three formations are referred to collectively as the *Lower Ultramafic Unit*, being characterized by a relative abundance of ultramafic and mafic volcanic rocks. These assemblages are not comparable with any well-established class of ultramafic or basaltic rock types and have been assigned the name "komatiite" (Viljoen and Viljoen, 1969a).

A persistent sedimentary horizon termed the *Middle Marker* is developed at the top of the Lower Ultramafic Unit and heralds an abrupt change in the type of volcanicity that occurs in

## STRATIGRAPHIC COLUMN OF THE SWAZILAND SEQUENCE




Figure 2 : Stratigraphic column of the Swaziland succession in the Barberton Mountain Land, South Africa. The distribution of the Lower Ultramafic Unit, the Mafic-to-Felsic Unit, and the Sedimentary Unit is shown in the inset general geological map of the Barberton greenstone belt.  
(after Anhaeusser, 1973a).

the upper three formations of the Onverwacht stratigraphy (Viljoen and Viljoen, 1969b). Referred to collectively as the *Mafic-to-Felsic Unit*, the upper formations consist of cyclically alternating mafic and intermediate to acid volcanic rocks, together with a wide variety of pyroclastic and chemical sedimentary rocks (tholeiitic basalts, dacites, rhyodacites, rhyolites, and cherts). Ultramafic bands and lenses occur sporadically throughout the succession but are volumetrically of minor importance.

In a number of localities within the Lower Ultramafic Unit, layered differentiated ultramafic pods and sills are developed, the latter consisting of dunites, harzburgites, peridotites, pyroxenites, gabbros, norites and anorthosites (Anhaeusser, 1969, 1972, 1974; Viljoen and Viljoen, 1969g).

Overlying the predominantly volcanic successions of the Onverwacht Group, is an assemblage of rocks consisting mainly of detrital sediments with subordinate volcanic and pyroclastic members. This succession has been subdivided into an argillaceous and an arenaceous

sedimentary unit, the former known as the Fig Tree Group, the latter as the Moodies Group. The Fig Tree assemblages consist mainly of pelitic sediments (greywackes, shales) with siliceous chemical precipitates (banded ferruginous cherts) and minor trachyandesitic lavas, agglomerates and tuffs. The Moodies Group represents a cyclically repetitive assemblage composed predominantly of conglomerates, quartzites, sub-greywackes, sandstones, and shales, together with subordinate mafic lavas, jaspilitic chert, and banded iron formation.

No general agreement exists as to the precise correlation of the Archean stratigraphic subdivisions of Rhodesia, Swaziland and South Africa. The main areas of contention centre about the suggestions, proposed by Stowe (1968a, b) and Hunter (1970), that successions older than the greenstone belts exists both in Rhodesia (the Pre-Sebakwian) and in Swaziland (the Ancient Gneiss Complex). Viljoen and Viljoen (1969e, f) and Anhaeusser (1973a), regard these areas as metamorphosed xenolithic equivalents of the Lower Ultramafic Unit.

As no extensive lithostratigraphic subdivision of the Rhodesian Archean has yet been attempted, efforts at a precise correlation of the stratigraphy with the Barberton model are not possible. A simplified scheme of correlation across the southern African Archean is shown in Table 3. In the writer's opinion, the granitic rocks (including those grouped within the Ancient Gneiss Complex and the Pre-Sebakwian) post-date the assemblages grouped within the Lower Ultramafic Unit.

TABLE 3

SUGGESTED LITHOSTRATIGRAPHIC CORRELATION OF RHODESIAN AND SWAZILAND  
ARCHEAN ROCKS WITH THE BARBERTON STRATIGRAPHIC MODEL

SOUTH AFRICA Barberton Belt		SWAZILAND	RHODESIA	UNIT
MOODIES		MOODIES	SHAMVAIAN	arenaceous SEDIMENTARY
FIGTREE		FIGTREE		argillaceous UNIT
UPPER ONVERWACHT	Swartkoppie	UPPER ONVERWACHT		MAFIC-TO-FELSIC UNIT
	Kromberg	BULAWAYAN		
	Hooggenoeg			
Middle Marker	Komati	LOWER ONVERWACHT	SEBAKWIAN	LOWER ULTRAMAFIC UNIT
LOWER ONVERWACHT	Theespruit			
	Sandspruit	Ancient Gneiss Complex	Pre-Sebakwian	

LITHIC PROPORTIONS

A planimeter survey of the geologic map (Figure 1) provides an assessment of the lithic composition of the southern African Archean. Due to the lesser degree of exposure of the granite-greenstone basement of the Kaapvaal Craton (Table 1) the Rhodesian Craton was selected to illustrate the varying proportions of the more important elements making up this ancient terrane. As can be seen in Table 4, granitic rocks make up over 80 per cent of the exposed Rhodesian basement complex, the remainder consisting of the Great Dyke and the greenstone belt relics. In an attempt to further refine the lithic proportions of the latter volcanic belts it was considered necessary to attempt a geological reinterpretation of the Rhodesian granite-greenstone terrane based on the Barberton model. The resulting map (Figure 1) was compiled from all the existing Rhodesian Geological Survey bulletins, as well as from the 1:1 000 000 geological map of Rhodesia (1971 edition). In addition, use was made of the reinterpreting geological map of the region compiled by Viljoen and Viljoen (1969f, p. 252).

TABLE 4

LITHOLOGICAL PROPORTIONS IN THE EXPOSED GRANITE-GREENSTONE TERRANE OF THE RHODESIAN CRATON

Rock Type	Area (km <sup>2</sup> )	Per Cent
Granitic Rocks*	162 027	81.2
Greenstone Belts*	33 735	16.9
Great Dyke	3 800	1.9
<b>TOTAL</b>	<b>199 562</b>	<b>100.0</b>
Ultramafic-Mafic Rocks*	2 823	8.4
Volcanic Rocks* (mafic and felsic)	25 894	76.7
Sediments*	5 018	14.9
<b>TOTAL</b>	<b>33 735</b>	<b>100.0</b>

(\* includes granite-greenstone terrane of northeastern Botswana).

The principal objective of the exercise was to subdivide the greenstone belts into component parts, the latter reflecting the broad scale subdivision of the stratigraphy into a Lower Ultramafic Unit, a Mafic to Felsic Unit, and a Sedimentary Unit, respectively. The results of the planimeter survey are given in Table 4 which shows that the greenstone belts are dominated by mafic to felsic volcanic rocks, the latter accounting for more than 75 per cent of the greenstone remnants.

A similar planimeter study was undertaken over the Barberton greenstone belt on the Kaapvaal Craton. This area was selected for analysis as it represents one of the better known southern African greenstone belts, having been the subject of numerous detailed investigations, both recently and in the past. The region, furthermore, acts as the type locality for greenstone belts on the Kaapvaal Craton.

The lithic proportions of the Barberton Mountain Land are shown in Table 5. As was the case with the Rhodesian greenstone belts the volcanic rocks of mafic to felsic composition constitute the largest proportion (50.5 per cent) of the Barberton stratigraphy. Comparisons of Tables 4 and 5 shows, however, that the Barberton greenstone belt contains approximately twice the Rhodesian average of rocks of (1) ultramafic-mafic and, (2) sedimentary, affinity.

Further refinement of the data relating to the proportions of the various lithologic types in the Barberton greenstone belt is possible. The threefold subdivision of the stratigraphy, described earlier, provides a useful separation of lithologic assemblages as is shown in the first part of Table 6. In the Onverwacht Group, the Lower Ultramafic Unit is most prominently represented, accounting for approximately 80 per cent of the volcanic succession (second part of Table 6).

The proportions of the various component rock types predominating in each of the three main stratigraphic units of the Barberton greenstone belt are shown in the third part of Table 6.

TABLE 5

LITHOLOGICAL PROPORTIONS IN THE  
BARBERTON MOUNTAIN LAND, SOUTH AFRICA

	Area (km <sup>2</sup> )	Per Cent
Ultramafic-Mafic Rocks	540	16.2
Volcanic Rocks (mafic to felsic)	1 682	50.5
Sediments	1 108	33.3
<b>TOTAL</b>	<b>3 330</b>	<b>100.0</b>

TABLE 6

LITHOLOGICAL PROPORTIONS IN THE BARBERTON GREENSTONE BELT,  
DETAIL OF TABLE 5

Unit and Rock Type	Area (km <sup>2</sup> )	Per Cent	
<u>ONVERWACHT GROUP</u>			
Lower Ultramafic Unit	1 768	53.1	
Mafic to Felsic Unit	454	13.6	
<u>FIG TREE AND MOODIES GROUPS</u>			
Sediments	1 108	33.3	
<b>TOTAL</b>	<b>3 330</b>	<b>100.0</b>	
<u>ONVERWACHT GROUP</u>			
Lower Ultramafic Unit	1 768	79.6	
Mafic to Felsic Unit	454	20.4	
<b>TOTAL</b>	<b>2 222</b>	<b>100.0</b>	
<u>ONVERWACHT GROUP</u>			
Lower Ultramafic Unit	Ultramafic-Mafic Rocks Basalts Felsic Volcanics	523 1 230 15	15.71 36.94 0.45
Mafic to Felsic Unit	Ultramafic-Mafic Rocks Mafic to Felsic Volcanics	17 437	0.50 13.10
<u>FIG TREE AND MOODIES GROUPS</u>			
	Sediments	1 108	33.30
<b>TOTAL</b>	<b>3 330</b>	<b>100.0</b>	

Basalts and peridotites, particularly of the komatiitic variety, make up the bulk of the Lower Ultramafic Unit. The amount, which approaches close to 53 per cent, includes the contributions made by the layered differentiated ultramafic complexes to the total. Furthermore, it is evident from Table 6 that felsic volcanic rocks are, volumetrically, insignificant in the Lower Ultramafic Unit, a feature that might not have been fully appreciated due to the prominence these rocks receive following their usefulness as stratigraphic marker beds. Similarly, ultramafic-mafic rocks of the type found in the Lower Ultramafic Unit are subordinate components of the Mafic to Felsic Unit, the latter being dominated by cyclically developed volcanic rock types possessing calc-alkaline chemical affinities.

The lithic proportions of the Barberton greenstone belt, when compared with the average Rhodesian values, demonstrates that this region is anomalously well-endowed with rocks of mafic and ultramafic composition. This fact is further strengthened by comparison of the data with that made available from selected areas on the Canadian Shield. In Table 7, the lithic proportions of rocks in the Slave Province and parts of the Superior Province are shown. Compared with southern African greenstone belts the Canadian occurrences are generally very much more extensive, the Abitibi belt alone being approximately twice the area of all the Rhodesian greenstone belts combined. Similarly the Wabigoon belt is areally greater than all the exposed greenstone remnants on the Kaapvaal Craton (cf., Tables 1 and 7).

TABLE 7

LITHOLOGICAL PROPORTIONS OF SELECTED AREAS IN THE  
SUPERIOR AND SLAVE PROVINCES OF THE CANADIAN SHIELD

Locality	Area (km <sup>2</sup> )	Total Greenstone Belt (km <sup>2</sup> )	Granite (km <sup>2</sup> )	Total Greenstone Belt (%)	Granite (%)
Abitibi Belt*	95 573	64 723	30 850	67.7	32.3
Wabigoon Belt*	15 496	7 655	7 841	49.4	50.6
Slave Province†	195 000	120 000	75 000	61.5	38.5
Locality and Rock Type			Area (km <sup>2</sup> )	Per Cent	
Abitibi Ultramafic-Mafic Rocks			2 389	3.7	
Abitibi Volcanic Rocks (mafic and felsic)			47 021	72.6	
Abitibi Sediments			15 316	23.7	
TOTAL			64 726	100.0	
Wabigoon Ultramafic-Mafic Rocks			248	3.2	
Wabigoon Volcanic Rocks (mafic and felsic)			6 214	81.2	
Wabigoon Sediments			1 193	15.6	
TOTAL			7 655	100.0	
Slave Province Volcanic Rocks			24 000	20.0	
Slave Sediments			96 000	80.0	
TOTAL			120 000	100.0	

\* (after Goodwin et al., 1972)

† (after McGlynn and Henderson, 1972)

Canadian greenstone belts are dominated by calc-alkaline volcanic sequences (Wilson et al., 1965; Baragar and Goodwin, 1969; Goodwin, 1971; Goodwin et al., 1972; and Table 7), whereas rocks of ultramafic-mafic composition play a subordinate role, the average for the Abitibi and Wabigoon belts, combined, being less than half the average value for the Rhodesian belts, and less than one quarter the value of similar rocks in the Barberton greenstone belt (cf., Tables 4, 6, and 7).

The proportions of the various volcanic rock types in greenstone belts of Canada, Rhodesia and South Africa can be compared in a variety of ways, probably the most informative of which are shown in Tables 8 and 9. In the first of these tables (Table 8) the relative proportions of specific rock types are compared. Basaltic rocks clearly form the major portion of Archean greenstone belts, wherever the latter may occur. Andesitic volcanic rocks are prominent in Canadian greenstone belts and are developed in some Rhodesian belts, but are virtually absent in the Barberton Mountain Land. Ultramafic-mafic rocks, including extrusive peridotitic lavas and the differentiated ultramafic complexes, are well-represented in southern Africa, whereas the felsic volcanic component in this region is poorly represented relative to the Canadian examples.

TABLE 8

PROPORTIONS OF ROCK TYPES IN GREENSTONE BELTS OF THE  
SUPERIOR PROVINCE, CANADA, THE RHODESIAN CRATON AND THE  
BARBERTON MOUNTAIN LAND, SOUTH AFRICA

Locality	Proportions (km) <sup>2</sup>				
	Ultramafic-Mafic	Basalts	Andesite	Felsic Volcanics	Total
Abitibi Belt*	2 389	28 213	14 106	4 702	49 410
Wabigoon Belt*	248	3 728	1 864	622	6 462
Rhodesian Craton	2 823	17 279	6 461	2 154	28 717
Barberton Belt	540	1 600	-	82	2 222
Proportions (per cent)					
Abitibi Belt	4.8	57.0	28.5	9.7	100.0
Wabigoon Belt	3.8	57.7	28.9	9.6	100.0
Rhodesian Craton	9.8	60.2	22.5	7.5	100.0
Barberton Belt	24.3	72.0	-	3.7	100.0

\* (after Goodwin et al., 1972).

The second table (Table 9) shows the ratios of various rock types, and combinations thereof, in the Canadian and southern African Archean greenstone belts. The ratios of all the combinations demonstrate clearly that the southern African greenstone belts (and the Barberton Mountain Land, in particular) have a greater ultramafic-mafic component and a lower intermediate to felsic component than their Canadian counterparts. This relationship, it will be shown later, has important implications in terms of the metallogenetic potential of the areas discussed.

GEOCHEMISTRY OF ARCHEAN VOLCANIC ROCKS

The sequential development of Archean volcanic-sedimentary piles is now well-established (Goodwin, 1968, 1971; Anhaeusser et al., 1969; Anhaeusser, 1971; Baragar and Goodwin, 1969; Viljoen and Viljoen, 1969a, b, 1971). Average chemical compositions of Archean volcanic rocks, arranged in generalized ascending stratigraphic order, have been presented for the volcanic assemblages of the Canadian Shield by Goodwin (1968), and for southern Africa and the Archean in

TABLE 9

RATIOS OF VARIOUS ROCK TYPES AND LITHOLOGICAL COMBINATIONS  
IN THE SUPERIOR PROVINCE, CANADA, THE RHODESIAN CRATON,  
AND THE BARBERTON MOUNTAIN LAND, SOUTH AFRICA

Locality	Ultramafic/Mafic	:	Basalts + Andesites + Felsic Volcanics
Wabigoon Belt*	1	:	25.3
Abitibi Belt*	1	:	19.8
Average Rhodesian Craton	1	:	9.2
Barberton Belt	1	:	3.1
Locality	Ultramafic/Mafic + Basalts	:	Andesites + Felsic Volcanics
Wabigoon Belt*	1.6	:	1
Abitibi Belt*	1.6	:	1
Average Rhodesian Craton	2.3	:	1
Barberton Belt	26.1	:	1
Locality	Ultramafic/Mafic + Basalts + Andesites	:	Felsic Volcanics
Wabigoon Belt*	9.4	:	1
Abitibi Belt*	9.3	:	1
Average Rhodesian Craton	12.3	:	1
Barberton Belt	26.1	:	1
<u>Average Superior Province, Canada*</u>	Basalt	:	Andesite : Felsic Volcanics
	6	:	3 : 1
or	Mafic	:	Felsic
	9	:	1
<u>Barberton Belt, South Africa</u>	Ultramafic/Mafic	:	Basalt : Felsic Volcanics
	6.6	:	19.5 : 1
or	Ultramafic/Mafic	:	Felsic
	26	:	1

\* (after Goodwin et al. 1972).

general by Anhaeusser (1971, 1973a) and Viljoen and Viljoen (1969a, b, 1971). This, and supplementary data, are provided in Tables 10 and 11.

In Table 10, the average chemical compositions of the more important volcanic components of the Onverwacht Group on the Kaapvaal Craton are shown. Columns 1 to 13 list the geochemistry of the basaltic and peridotitic komatiite lavas, tholeiitic basalts, and subordinate felsic volcanic interlayers encountered in the Lower Ultramafic Unit, whereas columns 14 to 19 show the geochemistry of the mafic, intermediate and felsic lavas of the Mafic to Felsic Unit in the Barberton greenstone belt. Table 11 provides a comparative set of data reflecting the geochemistry of some of the rock types encountered in the Archean greenstone belts of Canada, Western Australia, and Rhodesia. Although basaltic and peridotitic komatiites occur in all three areas they nowhere appear as abundantly as in the Barberton Mountain Land.

TABLE 10

## CHEMICAL COMPOSITION OF ARCHEAN VOLCANIC ROCKS ON THE KAAPPAAAL CRATON, SOUTH AFRICA

	ULTRAMAFIC - MAFIC UNIT											MAFIC - FELSIC UNIT							
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
SiO <sub>2</sub>	44.72	41.61	47.37	49.19	52.22	52.73	78.91	41.22	47.40	51.36	51.99	41.78	50.79	49.95	49.86	50.07	56.54	64.38	73.66
Al <sub>2</sub> O <sub>3</sub>	3.25	2.70	6.76	3.76	5.42	9.83	15.14	3.30	4.87	4.38	12.67	5.06	14.23	13.48	14.25	13.34	14.19	10.24	11.57
Fe <sub>2</sub> O <sub>3</sub>	6.02	5.63	1.18	11.80	0.98	1.23	0.25	4.31	3.48	2.33	1.61	5.28	1.12	1.82	2.65	2.31	0.0	1.40	0.93
FeO	5.52	4.35	8.08	8.88	9.70	0.69	6.54	6.32	8.47	6.45	4.53	7.11	7.89	7.67	9.10	14.24	12.76	0.62	
MgO	25.35	30.35	20.39	20.03	15.25	10.10	0.38	30.65	23.62	16.00	10.33	29.69	9.48	9.68	7.32	6.10	4.20	2.80	1.29
CaO	6.97	4.28	8.31	9.51	12.83	9.99	0.29	2.94	8.55	13.86	8.18	3.95	12.07	9.52	10.69	9.47	0.19	2.09	
Na <sub>2</sub> O	0.49	0.15	0.39	0.10	1.21	2.65	0.40	0.10	0.23	0.49	2.25	0.12	1.65	2.38	2.54	3.34	0.29	0.05	5.03
K <sub>2</sub> O	0.05	0.03	0.06	0.02	0.09	0.46	2.67	0.01	0.06	0.14	2.48	0.03	0.13	0.77	0.16	0.54	1.02	3.72	3.50
TiO <sub>2</sub>	0.52	0.31	0.46	0.43	0.56	0.85	0.54	0.27	0.43	0.46	0.47	0.22	0.41	0.69	0.70	1.22	2.03	1.13	0.02
P <sub>2</sub> O <sub>5</sub>	-	0.02	-	-	-	-	0.15	0.04	0.07	0.08	0.30	0.03	0.08	0.09	0.05	-	-	-	
MnO	0.19	0.17	0.19	0.17	0.22	0.22	0.01	0.19	0.16	0.23	0.16	0.13	0.19	0.18	0.17	0.19	0.07	0.04	0.06
CO <sub>2</sub>	-	-	-	-	-	-	0.	0.	0.09	0.08	0.40	0.05	0.14	0.13	0.48	0.23	0.08	1.56	1.73
H <sub>2</sub> O	5.58	8.81	5.26	-	2.05	1.87	1.31	9.50	4.29	2.24	2.63	8.65	2.09	3.07	2.81	3.01	5.84	2.50	2.68
No. of Analyses	3	8	4	2	5	3	4	9	4	8	1	1	1	5	5	2	1	1	1

## \* ONVERWACHT GROUP TYPE LOCALITY, BARBERTON MOUNTAIN LAND

1. Average peridotitic komatiite, Sandspruit Formation.
2. Average peridotitic komatiite, Komati Formation.
3. Average high-alumina basaltic komatiite (Geluk type).
4. Average low-alumina basaltic komatiite (Geluk type).
5. Average basaltic komatiite (Badplaas type).
6. Average basaltic komatiite (Barberton type).

## \* JAMESTOWN SCHIST BELT, BARBERTON MOUNTAIN LAND

7. Average felsic volcanic rocks (Theespruit Formation).

## \* NELSHOOGST SCHIST BELT, BARBERTON MOUNTAIN LAND

8. Average peridotitic komatiite.
9. Average basaltic komatiite (Geluk type).
10. Average basaltic komatiite (Badplaas type).
11. Tholeiitic basalt.

## \* MULDERSDRIF GREENSTONE BELT, NW OF JOHANNESBURG

12. Peridotitic komatiite.
13. Average tholeiitic basalt.

## \* ONVERWACHT GROUP TYPE LOCALITY, BARBERTON MOUNTAIN LAND

14. Average magnesian-rich basalt (Hooggenoeg Formation).
15. Average tholeiitic basalt (Hooggenoeg Formation).
16. Average tholeiitic basalt (Kromberg Formation).
17. Pillowed intermediate lava (Hooggenoeg Formation).
18. Pillowed intermediate lava (Hooggenoeg Formation).
19. Pillowed acid lava (Kromberg Formation).

Analyses : 1-6 and 14-19 (Viljoen and Viljoen, 1969a, b), 7 (Anhaeusser, 1972), 8-13 (Anhaeusser, manuscript in preparation).

Analysts : 8-13 National Institute for Metallurgy, Johannesburg, and Durham University, Durham, England.

COMPARATIVE GEOCHEMISTRY OF ARCHEAN VOLCANIC ROCKS OF THE  
SUPERIOR PROVINCE, CANADA, THE YILGARN BLOCK, AUSTRALIA AND THE RHODESIAN CRATON

	SUPPLIER PROVINCE - CANADIAN SHIELD						YILGARN BLOCK			RHODESIAN CRATON				
	1	2	3	4	5	6	7	8	9	10	11	12	13	14
SiO <sub>2</sub>	48.85	48.9	54.7	61.5	67.3	74.3	41.20	48.82	51.4	45.54	50.47	53.30	55.59	61.48
Al <sub>2</sub> O <sub>3</sub>	7.55	14.5	15.0	15.7	14.8	12.9	5.31	11.90	14.8	9.19	15.45	14.42	15.83	14.52
Fe <sub>2</sub> O <sub>3</sub>	2.02	2.14	2.00	1.83	1.17	0.74	3.89	1.17	1.5	2.40	1.63	1.49	1.47	1.44
FeO	11.09	9.03	7.64	4.49	3.44	2.22	5.13	9.24	9.1	8.53	9.62	9.71	5.73	4.25
MgO	16.73	6.27	4.50	2.38	1.55	0.85	29.87	12.81	6.7	20.65	6.05	3.97	5.51	4.33
CaO	11.43	8.74	6.39	4.41	3.13	1.48	4.66	12.70	10.7	6.07	11.76	11.36	7.50	5.64
Na <sub>2</sub> O	0.81	2.51	2.79	3.15	3.07	2.47	0.24	1.33	2.7	0.93	1.68	1.63	2.80	3.69
K <sub>2</sub> O	0.07	0.45	0.55	1.16	1.40	2.10	0.03	0.11	0.18	0.21	0.57	0.95	1.02	0.65
TiO <sub>2</sub>	1.14	1.06	0.99	0.63	0.51	0.26	0.23	0.42	0.92	0.55	0.89	0.99	0.64	0.61
P <sub>2</sub> O <sub>5</sub>	0.30	0.07	0.12	0.12	0.07	0.07	-	0.06	0.13	0.08	0.26	0.12	0.15	0.15
MnO	0.02	0.21	0.28	0.16	0.08	0.10	0.17	0.27	0.21	0.19	0.15	0.19	0.10	0.08
CO <sub>2</sub>	0.79	1.93	1.93	2.18	0.98	0.86	-	-	0.1	0.01	0.20	0.32	0.34	0.11
H <sub>2</sub> O	3.5	3.34	2.92	2.27	1.56	1.17	8.54	1.05	1.0	4.48	1.22	1.11	4.19	2.95
No. of Analyses	1	162	119	272	177	85	3	1	123	4	11	3	4	2

1. Basaltic komatiite, Wawa volcanic belt, Ontario.
2. Average basalt.
3. Average andesite.
4. Average dacite.
5. Average rhyodacite.
6. Average rhyolite.
7. Average peridotite (peridotitic komatiite, Mt Ida and Scotia, W.A.).
8. Magnesian basalt (basaltic komatiite).
9. Average tholeiitic basalt (Kalgoorlie-Norseman area, W.A.)
10. Average olivine basalt (basaltic komatiite, Geluk type).
11. Average tholeiitic basalt, Bulawayan Group.
12. Average "andesite", Bulawayan Group.
13. Average andesite, Maliami River Formation.
14. Average dacite, Maliami River Formation and Umniati Group.

<sup>1</sup> (Brooks and Hart, 1972), 2-6 (Goodwin, 1968), 7 (Nesbitt, 1971), 8 (McCall, 1973), 9 (Hallberg, 1972), 10-14 (Bliss and Stidolph, 1969).

As was discussed previously, the Canadian volcanic sequences are characterized by calc-alkaline assemblages similar to those listed in Table 11, columns 2 to 6. In Western Australia, analyses available from only a few selected areas suggest that the predominant volcanic rock types are tholeiitic basalts (column 9), like those described by Hallberg (1972). Peridotitic lavas and magnesian-rich basaltic lavas are also represented in places and have komatiitic affinities (columns 7 and 8).

Very little geochemical information is available from the Rhodesian greenstone belts. Some examples, listed in Table 11, columns 10 to 14, show a spread from basaltic komatiites to rocks of the calc-alkaline volcanic class. Assemblages, correlated with the Lower Ultramafic Unit, are widespread in this country (Figure 1) and will doubtless be found to be geochemically comparable to rocks of similar character in the Barberton belt (Table 10). Unlike the latter area, however, the Rhodesian greenstone belts contain prominent developments of andesitic pillow lavas, the latter forming part of the Bulawayan succession. Intermediate to felsic lavas and pyroclasts (andesite to dacite) occur in greenstone belts located along the western edge of the exposed Archean basement of the Rhodesian Craton. Extending from the Matsitama belt in eastern Botswana, through the Bulawayo-Bubi belt into the Midlands belt (Figure 1), these successions are referred to synonymously as the Maliami River Formation or the Umniati Group (Bliss and Stidolph, 1969). Doubt has been expressed (Bliss, 1970) as to whether these rocks form part of the greenstone belt successions or whether they represent some younger sequence. Unquestionably the assemblage, which consists predominantly of extensive sheets of pyroclastic rocks, and fresh, pyroxene-bearing lavas, is unlike the Bulawayan succession encountered elsewhere on the Rhodesian Craton. This is further borne out by the distinctive metallogenetic characteristics of the successions in the areas in which these rocks occur, and which will be discussed later.

### ARCHEAN MINERALIZATION

#### TYPES

Virtually all the known Archean mineral occurrences in the world are associated with the volcanic-rich sequences of greenstone belts, the exceptions being the deposits linked with sedimentary rocks and granites and pegmatites. Mineral distribution patterns and metallogenetic studies in Archean terranes are generally lacking, the only noteworthy contributions in this context coming from the Canadian Shield (Goodwin, 1965, 1966, 1971: Hutchinson et al., 1971: Hutchinson, 1973) and, to a lesser extent, Western Australia (Prider, 1965).

From these and numerous other, more local, studies, it has become obvious that certain mineral types and associations have a relationship both to one another and to their host rocks. Hutchinson et al., (1971) recognized four families of mineral deposits, believed to be of primary origin, in Archean rocks. Broadly, these may be modified to include : (1) ores of nickel, with minor copper and associated platinum-group metals, asbestos and magnesite deposits, and chromite and magnetite deposits. These are invariably associated with ultramafic to mafic igneous bodies (as well as ultramafic volcanic rocks). (2) pyritic, massive base metal sulphide deposits of copper, zinc, silver and gold, as well as antimony and mercury, all of which are invariably associated with well-differentiated, mafic to felsic extrusive and pyroclastic rocks of calc-alkaline character. Many of these deposits are considered to be of volcanic exhalative origin. Also included in this category are certain gold and copper deposits in porphyritic, sub-volcanic intrusions of felsic composition. (3) diverse varieties of iron formation, with or without associated sediments, that occur in well-differentiated volcanic complexes. These well-banded rocks are widely considered to be chemical sediments probably formed by fumarolic-exhalative processes. Included in this category are certain gold ores, the latter associated with oxide iron formations. (4) iron formations, particularly the oxide and some silicate species, the latter generally lacking the gold association of the exhalite variety. These iron formations occur with immature clastic sediments (greywackes and argillites). Included in this group also are certain gold deposits occurring in quartzitic and conglomeratic clastic sedimentary rocks.

To the four general families outlined above may be added two additional categories. These include : (5) mineral deposits associated with granites and pegmatites, including occurrences of beryllium, lithium, tin, tungsten, tantalum-columbium, bismuth, molybdenum, corundum, mica, quartz (silica) and felspar, as well as the precious and semi-precious minerals,

emerald, ruby and sapphire, and (6) the stratiform deposits other than the iron formations and including limestone, barite, corundum, graphite, and a variety of slates. The latter products have diverse origins ranging from sedimentary to metamorphic and volcanic exhalative or volcanogenic pyroclastic.

### THE NATURE AND DISTRIBUTION OF ARCHEAN MINERALIZATION IN SOUTHERN AFRICA

In order to simplify description of the Archean mineralization of southern Africa, the threefold subdivision of the stratigraphy of the greenstone belts, described earlier, will be used as a basis for classification. This deviates little from the classification scheme outlined initially by Hutchinson et al., (1971) and modified by the writer in the previous section. Fundamentally, both approaches centre around mineral types and host rock associations, as well as an underlying stratigraphic evolutionary control.

Ideally, Archean greenstone belts commence with an ultramafic-mafic group of rocks at the base and pass upwards into mafic, intermediate, and felsic volcanic rocks higher in the successions, the latter overlain and terminated by a variety of sedimentary assemblages (Anhaeusser et al., 1969; Anhaeusser, 1971). Most greenstone belts, however, display only parts of the total idealized column, being aborted at varying stages due to one or other causes, the most destructive of which appears to have been the intrusion of a wide variety of granitic rock types.

The southern African greenstone belts are no exception and, apart from the Barberton Mountain Land, which probably approaches closest to the theoretical model, virtually all the Archean remnants have sections of their stratigraphy either missing or attenuated. Rocks of the Lower Ultramafic Unit, being at the base of the synclinally folded greenstone belts are usually more prone to metamorphic alteration, fragmentation, and assimilation by the granites than rocks higher in the succession. Despite this, however, it is evident from Figure 1, and Tables 4 and 8, that much of this rock assemblage is preserved in southern Africa.

#### MINERALIZATION ASSOCIATED WITH ULTRAMAFIC TO MAFIC VOLCANIC AND PLUTONIC ROCKS

The distribution of mineralization associated with rocks of the ultramafic-mafic succession is shown in Figure 3. The mineral occurrences include deposits of nickel (plus copper), chrome, chrysotile asbestos, magnesite, talc, and the ornamental products verdite and buddstone. The greatest density of mineralization occurs in greenstone belts located in the southern part of Rhodesia and in the Barberton Mountain Land, both regions possessing abundant ultramafic rocks.

##### Nickel-Copper

Despite the abundance of seemingly favourable host rocks, noteworthy deposits of nickel are relatively rare in southern Africa and for the present are restricted to three major deposits including the Shangani, Trojan, and Empress ore bodies (Figure 3). The Trojan deposit, north of Salisbury, is described by le Roex (1964) as consisting of finely disseminated nickeliferous (pentlandite, millerite) and other sulphides (pyrite, pyrrhotite, chalcopyrite) occurring in dunite, serpentinite, and serpentine-talc-carbonate rocks. Nickel also occurs as a nickeloan magnetite.

The Shangani deposit, the most recent nickel discovery, is shortly to be described in a paper by Viljoen et al., (manuscript in preparation). The ore deposit, which occurs in serpentinitized peridotite, overlies a siliceous footwall tuff unit and consists of massive and disseminated sulphides (mainly pyrrhotite and pentlandite).

The Empress nickel-copper deposit, which has been described by Sharpe (1964) and Eales (1964), differs from both the Trojan and Shangani deposits in that the mineralization occurs in a differentiated sill-like, predominantly meta-gabbroic body, containing subordinate picrite, amphibolites (originally pyroxenites), norite, and diorite. The ore minerals present in the body,




Figure 3 : Map showing the distribution of nickel-copper, chrome, chrysotile asbestos, talc, verdite-buddstone, and magnesite in the Archean green-stone belts of southern Africa.

in order of abundance, include pyrrhotite and pentlandite, chalcopyrite, pyrite, violarite, and some sphalerite and chalcocite. Minor amounts of gold, silver and platinoids are also present in the ore. The Empress deposit, furthermore, does not occur in rocks of the Lower Ultramafic Unit, and should, in effect, be considered separately. It is located, instead, in the contentious Umniati Group described by Bliss (1970), and referred to earlier.

Other noteworthy occurrences of nickel-copper occur in the Tati greenstone belt of eastern Botswana. Two prospects, the Selkirk and the "Mound Struck", occur in a host rock setting consisting of quartz and felspar porphyritic volcanic rocks all with dacitic affinity (Mason, 1970). The mineralization consists mainly of chalcopyrite, pyrite, pyrrhotite and pentlandite. Little is known of the deposits but the mineralization appears to be related to troctolitic gabbroic intrusions and is thus more closely akin to the Empress deposit.

Promising nickel prospects, in rocks of the ultramafic-mafic clan, occur in the Damba area, west of the Shangani Mine, and in greenstone remnants in the Madziwa batholith, north of the Trojan deposit (Figure 3). Signs of nickel-copper mineralization occur in virtually all areas where ultramafic rocks are developed. This is true for the Barberton greenstone belt, where, despite the showings of millerite and nickeloan magnetites and trevorite, as well as the great abundances of ultramafic extrusive and intrusive rock types, no nickel ore bodies have yet been located.

#### Chrysotile Asbestos

Rocks of the Lower Ultramafic Unit are particularly favourable host rocks to chrysotile asbestos in southern Africa. As can be seen in Figure 3, there is a widespread distribution of this mineral, coinciding mainly with areas containing abundant serpentized ultramafic rocks (cf., Figure 1). Areas of particular importance are the Filabusi-Belingwe-Fort Victoria regions in the south and southeastern parts of the Rhodesian Craton and the Barberton greenstone belt, on the Kaapvaal Craton.

As comprehensive accounts of southern African chrysotile asbestos mineralization are available (van Biljon, 1964; Laubscher, 1964, 1968; Anhaeusser, 1974), few details will be given here. The asbestos is not confined to rocks of the Lower Ultramafic Unit but occurs also in sill-like bodies of ultramafic rocks developed less frequently in stratigraphy of the Mafic to Felsic Unit (e.g. the Havelock and Msauli mines). In addition, it occurs in differentiated ultramafic complexes in Rhodesia (e.g. Mashaba Igneous Complex; Shabani ultramafic body) considered by Wilson (1968, 1973) to be younger than the greenstone belts but older than the latest intrusive granitic events in the area.

#### Magnesite

Almost all the occurrences of magnesite in the Archean rocks of southern Africa are the alteration products of ultramafic rocks, the exception being the Barton Farm magnesite deposit near Gatooma, in Rhodesia (located in the Midlands greenstone belt, half way between the Great Dyke and the Empress Mine, Figure 3). This deposit, described by MacGregor and Bliss (1968), occurs in dolomitic argillites and arkoses of the Shamvaian Group, but these authors were unable to determine whether this magnesite is a primary sediment, or a replacement body.

Of the remaining magnesite occurrences, the distribution of which again coincides with the regional development of ultramafic rocks, the most important ore bodies have been those located along the northeastern margin of the Barberton Mountain Land. Descriptions of these, and other magnesite deposits in South Africa were given by van Zyl (1942). More recently, Viljoen and Viljoen (1969c) showed that the important magnesite deposits in the Kaapmuiden-Malelane area of the Barberton greenstone belt occurred mainly as stockworks and, more rarely, as tabular bodies in altered layered ultramafic bodies. A number of sill-like bodies of ultramafic type have, in this area, undergone magmatic segregation yielding extensive dunite or peridotitic dunite zones, the latter followed sequentially by pyroxenites, gabbros, anorthositic gabbros and norites. It is in the dunite zones, with cumulus olivine crystals (Fo 93 to Fo 100) constituting 70-90 per cent of the rock, that magnesite has formed within depth zones influenced by the circulation of meteoric water. This process of alteration has been assisted locally by faulting and fracturing of the successions in the area.

#### Talc

Talc deposits are prevalent in all greenstone belts possessing ultramafic rocks or magnesian rich basalts. Individual deposits vary in size and grade of talc produced. Most deposits occur in areas where deformation has affected the ultramafic rocks. The combined influences of structure and proximity to a heat source appears to have been an important factor

in talc formation. Numerous deposits have been worked in the Barberton Mountain Land, the most important occurrences being those developed along the northern margin of the greenstone belt (Figure 3), in the Jamestown Schist Belt and areas to the east (Anhaeusser, 1969, 1972).

The talc deposits occur as follows : (1) rimming intrusive porphyritic granite bodies intruded into the ultramafic lavas or layered ultramafic complexes; (2) in fault zones associated, in places, with gold-quartz veins, e.g. the Albion gold-talc deposit in the Jamestown Schist Belt; (3) in areas where ultramafic rocks have been folded, serpentinized and steatized to form mainly talc-carbonate rocks in which pods of quality talc develop by lateral secretion from the wall rocks; (4) in ultramafic rocks influenced by metasomatizing hydrothermal solutions derived from intrusive granites. The Scotia Talc Mine in the Barberton belt has, for example, numerous tourmaline needles in the talc, indicating that boron metasomatism probably accompanied the talc formation.

#### Verdite-Buddstone

Several occurrences of verdite, which is a dark green semi-precious variety of serpentine, used mainly as an ornamental stone, are known to occur only in the Barberton region (Figure 3). The mineral is frequently associated with a grey to white silica and talc assemblage giving rise to an attractive banded and invariably contorted rock known locally as buddstone.

The verdite occurrences, which are small, pod-like, deposits, are confined to the layered differentiated bodies and appear to have formed in zones of shearing and differential movement between pyroxene peridotite layers and adjacent, more massive and structurally competent, meta-gabbroic layers (Anhaeusser, 1969, 1972).

#### Chromite Mineralization in Archean Greenstone Belts

Important occurrences of chromite are located in the Archean rocks of southern Africa. Most outstanding, are the very high grade chromite deposits in the Selukwe area in south central Rhodesia (Figure 3). These deposits, which have been described in detail by Stowe (1968a), and Cotterill (1969), occur in a structurally complex succession that forms part of the Sebakwian Group of rocks. Lenticular sheets of differentiated ultramafic to mafic material containing chromite bodies occur together with meta-basaltic lavas, interbedded quartzites (altered cherts), banded iron formation, and a variety of argillaceous sediments (tuffs, greywackes) and arkosic greywackes.

The structural reconstruction of the area suggests that the Lower Ultramafic Unit rocks of the Sebakwian sequence were involved in folding, thrusting, metamorphism, silicification, and erosion, prior to the deposition of the Bulawayan succession. In support of this, it can be shown that the Bulawayan conglomerates contain pebbles and boulders of Sebakwian sedimentary and ultramafic rocks, including chromite pebbles and boulders.

The chromite deposits in the Selukwe area formed by magmatic segregation as cumulate layers at varying intervals in the differentiating ultrabasic intrusion, the thickest layers accumulating in downwarps on the chamber floor (Cotterill, 1969). Primary magmatic textures and structures occur in the chromite layers despite the intense alteration of the rocks in the area. Cotterill indicated that, at least twice after accumulation, metasomatism of the sheared chromite and peridotite had modified the chemical constitution of the chrome ores and the ultramafic rocks. Typical chromites are rich in Cr and Mg and poor in Fe, Al, and Mn. In Table 12 average ore grades of the Selukwe chrome deposits are listed. The ore bodies are among the richest in the world in terms of grade and Cr/Fe ratios and are prized as metallurgical grade ores.

The Sebakwian chromite deposits have not been specifically dated but are considered to be approximately 3 400 m.y. old. Further deposits of this nature occur scattered about the southern regions of Rhodesia (Figure 3), particularly in the Sebakwian remnants west of the Fort Victoria greenstone belt (Wilson, 1968) and in the region south of the Buhwa-Mweza belt (Worst, 1962b). In the latter area the chromite ore bodies, which include the Rhonda, Hmardick, United, and Mlimo mines, occur in greenstone remnants caught up in the northern marginal zone of the Limpopo high-grade metamorphic belt.

Geological Setting	Area and Reference	Cr <sub>2</sub> O <sub>3</sub> (per cent)			Cr/Fe	No. of Analyses
		Range	Mean	Range		
Sebakwian Group, Rhodesia (~ 3.4 b.y. old)	<u>SELUKWE</u> Selukwe Peak Mine Railway Block, Priority 1 Magazine Hill Railway Block, Priority 1 Mount Clains (Cotterill, 1969)	52.32 - 58.41 57.89 - 60.65 58.41 - 59.41 58.45 - 61.60 52.98 - 55.47	55.59 60.23 58.82 59.70 54.54	1.9 - 3.9 3.2 - 4.26 3.68 - 3.9 3.33 - 4.19 1.36 - 1.65	2.9 3.9 3.8 3.8 1.5	6 10 3 10 7
	<u>MASHABA</u> (Wilson, 1968)		41.5		2.7	1
	Mashaba Igneous Complex, Rhodesia. (post-Shamvian age - ~ 3.0 b.y. old)	<u>MASHABA</u> (Wilson, 1968)	56.61 - 60.29	55.87	3.51 - 4.56	3.91
	Messina Formation, Limpopo Mobile Belt (~ 3.0 b.y. old)	<u>SOUTPANSBERG-MESSINA</u> (van Eeden et al. 1955)	29.6 - 30.0	29.80	1.2 - 1.4	1.3
	Great Dyke, Rhodesia (~ 2.5 b.y. old)	<u>MUSENGEZI COMPLEX</u>	53.82 - 61.85	53.90	1.8 - 3.97	2.15
		<u>SELUKWE COMPLEX</u>	51.17 - 56.81	54.80	2.28 - 3.18	2.72
		<u>WEDEZA COMPLEX</u>	50.04 - 56.37	53.31	1.94 - 3.05	2.52
		<u>HARTLEY COMPLEX</u>	51.02 - 59.16	56.00	1.93 - 4.20	2.90
		<u>MOTOROSHANGA</u> (Cotterill, 1969)	58.81 - 59.03	58.98	3.48 - 3.75	3.6
Bushveld Igneous Complex, South Africa (~ 1.95 b.y. old)	<u>EASTERN BUSHVLD</u> Central Sector Southern Sector (Cameron, 1964)	38.9 - 49.84 43.35 - 48.17	~ 46.0 ~ 45.0	0.7 - 1.92 1.2 - 1.59	~ 1.55 ~ 1.45	20 14
	<u>WESTERN BUSHVLD</u> Data from 15 chromite seams. (Cousins and Feringa, 1964)					
		38.0 - 53.0	~ 47.0	0.85 - 2.4	~ 1.5	64
Turkish Ultramafic and Related Gabbroic Complexes (~ 300 m.y. old)	<u>FETHIYE-MARMARIS COMPLEX</u> (van der Kaaden, 1970)	30.41 - 59.28	~ 45.0	2.1 - 4.3	~ 3.0	56

The chemical composition of the chromite from the Rhonda group of mines ranges from 32-53 per cent  $\text{Cr}_2\text{O}_3$ , the Cr/Fe ratio varying between 1.23 and 3.03 (Worst, 1962b). Some of the ore is metallurgical grade but most is hard lumpy and of chemical grade.

Chromite of a younger Archean age (approximately 3 000 m.y. old) occurs in significant quantities in the Mashaba Igneous Complex at the western end of the Fort Victoria greenstone belt. Here, the bulk of the chrome ore bodies are found in ultramafic rocks (serpentinized dunites and peridotites) centred about the Prince Mine situated near the King asbestos mine (Figure 3). Most of these deposits are intensely deformed but in the Prince ore body there is a chromite-rich zone over 30 m thick in which chromite seams varying from less than 5 mm up to 50 mm, occur as density separated layers in serpentinite. The chromite deposits of the Mashaba Igneous Complex contain high-grade metallurgical ores, the latter contrasting with the lower grade mineralization found in Sebakwian rocks in this region (Table 12).

#### Other Chromite Occurrences in Southern Africa

While not intending to describe in any detail the remaining chromite occurrences in southern Africa, it is nevertheless of interest to note certain features relating to this mineral's distribution in time and space.

The first, and oldest, of these remaining chromite-bearing environments is located in the Limpopo metamorphic belt. Developed in this region are rocks of the Messina Formation, the latter considered to be approximately 3 000 m.y. old (Van Breemen and Dodson, 1972). Anorthosite complexes, which appear to be semi-concordant intrusions within the Messina Formation, are found in the area and are distinctly layered and folded with the latter successions. Both Söhnge (1945) and van Eeden et al., (1955) recorded chromite layers in serpentinites, hornblende gneisses, and anorthositic hornblende gneisses in the area around Messina. The chromite occurrences, which have been included in Figure 3, are small, of low grade (Table 12), and have not been mined.

The second, and considerably more important, chromite-bearing region is that associated with the Great Dyke of Rhodesia (Worst, 1960, 1964). The Dyke, which exceeds 500 km in length, is approximately 2 500 m.y. old (Davies et al., 1969) and is made up of four aligned lopolithic complexes, each possessing similar characteristics. Consisting mainly of a variety of ultramafic to gabbroic rock types, the Dyke also contains a total of 14 proved chrome seams, the latter traceable over much of the length of the complex. Tremendous tonnages of intermediate to high grade chromite exists in the Dyke (Table 12) but at present mining is only being carried out on a small scale.

The third, and largest, chromite occurrences in southern Africa are those located in the Bushveld Igneous Complex, the latter body occupying the central region of the Kaapvaal Craton (Figure 1). Estimates by Cousins and Feringa (1964) place the chrome ore reserves at approximately 18 000 million tons, making the Bushveld Complex the largest repository of chrome in the world.

The Bushveld chrome is, on average, lower grade than chrome ores in the Archean greenstone belts, the Great Dyke and the Turkish ultramafic and related gabbroic complexes (Table 12). The ore has mainly been used for the production of chromium chemicals but increasing metallurgical applications are being found as supplies of the world's higher grade chrome ores diminish.

#### Chromite Distribution in Southern Africa

The principal areas of chromite development in southern Africa are illustrated in Figure 4. Several features of significance emerge from studies of the chromite distribution. These include the following : (1) the main chromite occurrences are developed on the cratonic portions of the shield and are symmetrically disposed on either side of the Limpopo mobile belt. (2) the chromite is developed in geological formations ranging in age from the Sebakwian ( $\sim 3.5$  b.y. ago) to the Bushveld Complex ( $\sim 2.0$  b.y. ago). (3) the chromite occurs in Archean greenstone belts immediately flanking the Limpopo mobile belt and not in those far removed from this region. A significant case in point is the Barberton greenstone belt which has virtually no known chrome mineralization despite the great abundance of favourable ultramafic rock types.



Figure 4 : Distribution of chrome in southern Africa. The chrome ores are developed in formations spanning approximately 1 500 m.y. of geologic time.

**MAIN CHROMITE PRODUCING AREAS**

- 1 SELUKWE GREENSTONE BELT (SEBAKWIAN)
- 2 MASHABA IGNEOUS COMPLEX (POST SHAMVAIAN)
- 3 GREAT DYKE
- 4 WESTERN BUSHVELD
- 5 EASTERN BUSHVELD



**IGNEOUS COMPLEXES AND FORMATIONS  
IN WHICH CHROMITE OCCURS**

- ▼ Tugela Randt Complex, Natal
- Detrital Chromite in Witwatersrand (~2.5-2.75 b.y.)
- Sedimentary Basin
- Bushveld Igneous Complex (~2.0 b.y.)
- Great Dyke (~2.5 b.y.)
- Messina Formation (~3.0 b.y.)
- Mashaba Igneous Complex (~3.0 b.y.)
- Archaean Greenstone Belts (~3.0-3.5 b.y.)

(4) the richest chromite ores are those of the Rhodesian Archean and the Great Dyke whereas the greatest chromite ore reserves occur on the Kaapvaal Craton. Thus the high-grade chromite appears to be developed in the older geological environments whereas the lower grade ores are found in the younger geological settings. (5) chromite is not known to occur in large amounts anywhere in the Archean greenstone belts preserved on the Kaapvaal Craton. Indirect evidence suggesting the former presence of such chromite-rich greenstone belts lies in the fact that the Witwatersrand Goldfield, which formed approximately 2.50 - 2.75 b.y. ago, contains abundant detrital chromite, the latter occurring within the auriferous conglomerates. The available evidence suggests that the Witwatersrand sediments, including the gold and chromite grains, were derived from the erosion of pre-2.75 b.y. old greenstone belts formerly located to the north and northwest of the Witwatersrand sedimentary basin (Viljoen et al., 1970; Pretorius, 1974b). These belts probably occurred in the region now largely occupied by the Bushveld Complex. (6) small chromite occurrences are located in the Tugela Randt Complex in Natal but

are of minor importance. It may be significant that this chromite occurrence is close to the Tugela fault zone, the latter representing the boundary between the Kaapvaal Craton and the Natal high-grade metamorphic belt to the south.

The various chromite occurrences in southern Africa combine to form the largest concentration of this mineral known on the earth's surface. The full implications of the symmetrical distribution of the chromite on either side of the Limpopo belt remain to be determined. The fact that chromite has been contributed to a variety of geotectonic regimes (in the area shown in Figure 4) over a time span approaching 1 500 m.y. does, however, suggest that the mantle, centred about the region of the Limpopo mobile belt, must have been particularly chromium enriched.

#### Gold

The distribution of 289 of the more important gold occurrences in the Archean greenstone belts of southern Africa are shown in Figure 5. Of this number, 57 ore deposits have each yielded in excess of 100 000 ounces of gold. A total of over 4 000 different gold deposits occur scattered throughout virtually all the greenstone belt remnants, the latter often being referred to locally as "gold belts". The majority of these deposits have been small, with the output of just over 50 million ounces of gold and approximately 14 million ounces of silver coming from about 20 large mines. Phaup (1964) showed, for example, that in the area around Gatooma, which is located in the northern part of the Midlands greenstone belt (Figure 1), there were 359 mines that had been worked prior to 1951. The total production from this area had been over 4.5 million ounces of gold, of which 64 per cent came from the Cam and Motor Mine, 8.6 per cent from the Giant Mine, 4.4 per cent from the Thistle Etna Mine, and 23 per cent from the remainder. Less than 1 per cent of the total came from 165 mines or prospects.

Similarly for the Barberton greenstone belt, it was shown (Anhaeusser, 1969) that up to the end of 1963, a total of 5.65 million ounces of gold had been produced from 274 recorded deposits in the area. Of this total, 62 per cent of the gold came from three mines (Sheba, Consort, Fairview), whereas 99.3 per cent of the total came from 45 mines (the balance of 0.7 per cent thus coming from the remaining 229 deposits).

Gold mineralization occurs throughout the Archean stratigraphic column but is preferentially developed in the volcanic rich units and in banded iron formations, cherts, and, to a lesser extent, in argillites and quartzites. Some gold also occurs in the granitic rocks, particularly where these occur adjacent to greenstone belts, or their xenolithic remnants. The ultimate source of the gold appears to be genetically related to volcanogenic activity of one sort or another within the volcanic-rich belts. Although no quantitative supporting data are available, it is the writer's opinion (Anhaeusser, 1969), as well as that of Viljoen et al., (1969, 1970), that the basaltic and peridotitic rocks of the Lower Ultramafic Unit were the primary host rocks of much of the gold mineralization found in the southern African greenstone belts. Gold and associated sulphides are, however, widespread in volcanogenic rocks of the Mafic to Felsic Unit (e.g. Bulawayan successions, Rhodesia, and Swartkoppie Formation, Barberton Mountain Land). These rocks are generally further removed from intrusive granite bodies than the stratigraphically lower basaltic and peridotitic assemblages and have, therefore, been less affected by metamorphism than the latter successions. The concentration of the mineralization in the areas rimming the granite bodies (particularly evident along the contact zone between the Midlands greenstone belt and the Rhodesdale batholith, cf., Figures 1 and 5) may, therefore, be linked with the heat effects consequent on granite intrusion, rather than reflecting an initially higher gold content of these rocks.

Descriptions of many of the southern African Archean gold deposits are available. Information on the mines (with varying degrees of attention given to detail) is available in most of the bulletins of the Geological Survey of Rhodesia, in numerous dissertations, and in publications of the Geological Society of South Africa (particularly Haughton, editor, 1964, and Visser, editor, 1968). Several attempts have been made to summarize what is known of the deposits, their distribution, and the controls to mineralization, and further details may be obtained in the references cited (Davies and Hunter, 1964; Goldberg, 1964; Phaup, 1964; Anhaeusser, 1965, 1969; Pretorius and Hempkins, 1968a, b; Anhaeusser et al., 1968; Viljoen et al., 1969).

As mentioned previously, most of the gold is now considered to be genetically associated with rocks of volcanic origin. The superimposed effects of granite intrusion,




Figure 5 : Map of southern Africa showing the distribution of the more important gold occurrences in the Archean greenstone (metavolcanic) belts.

metamorphism, and deformation have been largely responsible for the migration of much of this gold from its original source into dilatational (tensional) zones produced by folding and faulting. Added support for this derives from trace element patterns observed in pyrites from gold-sulphide deposits of the Barberton Mountain Land. This, as well as lead isotopic data showing the age of the mineralization in the Barberton greenstone belt to be between 3 450-3 800 m.y. old, has been described by Saager and Köppel (in press, 1974).

Although many of the Archean gold deposits are clearly related to hydrothermal processes there is growing evidence to suggest that some of the gold-pyrite-arsenopyrite mineralization is also stratiform in character. The spatial association of gold with various iron formation types (both oxide and sulphide facies), as well as banded cherts, have long been recognized but have only recently been interpreted as being of volcanic exhalative origin. Examples of this type are known in the Barberton greenstone belt and in many areas of Rhodesia, including more specifically the Vubachikwe-Blanket mine areas of the Gwanda greenstone belt (Figure 1), and the Neardy Mine area in the Odzi-Umtali belt. Further details of these, and other occurrences like them, have been made available by Fripp (1974, and manuscript in preparation). Many of the deposits in this category occur in jaspilitic iron formations terminating volcanic cycles of mafic to felsic lava and acid tuffs and pyroclasts.

#### Regional Distribution of Archean Gold Mineralization

It can be seen in Figure 5 that the greatest concentration of gold deposits occurs in the more centrally situated greenstone belts on the Rhodesian Craton, as well as in the Barberton Mountain Land. As mentioned earlier, there is a tendency for most deposits to be located around the periphery of the greenstone belts, not on the contacts with the main granite batholiths but, as pointed out by Phaup (1964), usually a few kilometres in from the margins of the belts. Exceptions occur where some gold deposits flank small stocks or granite cupolas and even extend into the granite itself. There thus appears to be an "optimum thermal zone" in greenstone terranes wherein the gold and associated sulphide mineralization tends to concentrate preferentially. No quantitative data is yet available to define, more precisely, the thermal regime most conducive to gold accumulation, but evidence from metamorphic mineral assemblages suggest that conditions akin to middle or upper greenschist facies of regional metamorphism were probably most favourable ( $\sim 450-550^\circ\text{C}$ ).


Additional support for an optimum thermal zone for gold mineralization follows from the studies of Pretorius and Hemphicks (1968a), who investigated data concerning gold production in Rhodesia. Analyses of the general statistics of population frequency distributions indicated a pattern of more favourable intensity of gold mineralization associated with ores containing the lower temperature minerals chalcopyrite, stibnite, and galena, than from ore characterized by the presence of the higher temperature minerals pyrrhotite, arsenopyrite, and sphalerite.

On a regional scale it can be shown schematically (Figure 6) that the high-grade metamorphic belts are not only virtually barren of gold mineralization but they have influenced (mainly thermally) the greenstone belts in their immediate vicinity - only very minor quantities of gold having been located in areas flanking the mobile belts.

Mineralization of the type discussed was unstable in the high temperature/pressure regimes of the mobile belts and consequently moved away from these areas, and the craton margins, towards the craton interiors. Some of the mineralization in the thermally affected regions may also have been dissipated (as opposed to being concentrated) throughout many rocks in the heat affected regions. In addition, some of the migratory gold mineralization may have eventually ended up in the Witwatersrand sedimentary basin (or other interior basins), together with gold and sulphides derived from the erosion of Archean greenstone belts.

#### MINERALIZATION ASSOCIATED WITH MAFIC TO FELSIC VOLCANIC ROCKS

Under this heading, mineral types associated with volcanic rocks of intermediate to acid composition are described. Most of the mineralization occurs in successions grouped within the Mafic to Felsic Unit outlined earlier. Exceptions include mineral deposits related to felsic volcanic and pyroclastic rocks of the Lower Ultramafic Unit. The latter include mainly small deposits of barite, corundum and kyanite, and, unlike the rocks of the Mafic to Felsic Unit, have not yielded signs of any noteworthy sulphide mineralization.



**Figure 6** : Schematic map of the southern African Shield showing the distribution of gold mineralization in relationship to the circum-cratonic high-grade metamorphic mobile belts. The provenance area of the Witwatersrand Goldfield is now largely covered by younger formations, including the Bushveld Complex.

The distribution of the mineral occurrences, which include deposits and prospects of antimony, gold, mercury, copper, lead, zinc, tungsten, and massive pyrite-pyrrhotite, are shown in Figures 7 and 8.

#### Antimony and Mercury

One of the most important single occurrences of antimony mineralization in the world, and certainly the largest deposit of its kind anywhere in the Archean, is the Consolidated Murchison Mine located in the Murchison greenstone belt on the Kaapvaal Craton (Figures 1 and 7). A number of ore bodies occur in lenses strung out along approximately 50 km of the so-called "Antimony Line" (van Eeden et al., 1939) where the host rocks consists mainly of sheared and, in places, intensely carbonated (green and grey dolomite) intermediate to felsic volcanic rocks of greenschist metamorphic grade. The mineralization consists mainly of the sulphides stibnite, berthierite, tetrahedrite, and chalcostibite and includes the antimony oxides stibiconite and kermesite. Gold, pyrite, arsenopyrite and cinnabar occur locally, the mercury having been mined in the past, whereas gold is still recovered as a by-product of the antimony mining. Descriptions of the ore bodies (Mendelssohn, 1938; Hausmann, 1959; Sahli, 1961; Boese, 1964) suggest that the mineralization is epigenetic (hydrothermal) in origin but these views are now being challenged in favour of a volcanogenic origin (volcanic exhalative, fumarolic). It is suggested that processes involving the remobilization of the ores by hydrothermal activity caused local concentrations of mineralization in structurally favourable areas. The reconcentration of the stibnite was probably accompanied by silicification and dolomitization of the volcanic rocks in the structurally disturbed areas, although the association of stibnite mineralization with




Figure 7 : Map showing the distribution of antimony, barite, mercury, corundum, and kyanite associated with felsic volcanic rocks in Archean greenstone belts of the Rhodesian and Kaapvaal cratons.

green dolomitic rocks in the mines has led to an alternative suggestion that the carbonate rocks originally constituted a stratigraphic unit interlayered with the volcanic successions (M.J. Viljoen, personal communication, 1974).

Antimony mineralization is also known in the Barberton greenstone belt as well as in the centrally situated greenstone belts of the Rhodesian Craton. These deposits are generally small, the antimony frequently being recovered as a by-product of gold mining (e.g. the Indarama gold-antimony deposit in the Midlands greenstone belt, near Que Que, Rhodesia, described by Mehliss, 1968). The distribution of mercury mineralization (cinnabar) follows closely the pattern outlined for the antimony (Figure 7), with most of the production again representing a by-product of gold

mining. The small Rhodesian occurrences have been described by Robertson (1972) who noted that the more significant deposits are located in weakly metamorphosed carbonated andesitic lavas. The main occurrence on the Kaapvaal Craton is the Monarch cinnabar deposit located in the Murchison greenstone belt, close to the antimony workings of the Consolidated Murchison Mine. Hausmann (1959) described the cinnabar-rich lodes as occurring in siliceous carbonate rocks together with minor amounts of chlorite and associated low temperature ore minerals, the latter including pyrite, chalcopyrite, stibnite, tetrahedrite, digenite and covellite.

The mercury occurrences, like those of the antimony deposits, have generally been described as hydrothermal in origin. However, the close association of the mineralization with rocks of intermediate to felsic volcanic origin suggests an alternative genetic history linking these deposits to the late stages of an exhalative sedimentary environment (fumarolic mineral springs depositing siliceous sinters).

#### Barite

Small, generally sub-economic, and usually stratiform, deposits of barite occur scattered throughout many greenstone belts in southern Africa (Figure 7). Investigations in the Barberton Mountain Land by Viljoen and Viljoen (1969c) and Reimer and Heinrich (1974) suggest that the mineralization occurs in both the Onverwacht and overlying Fig Tree groups. In both these settings the barite is considered to be of volcanogenic origin, probably forming from volcanic exhalations associated with explosive volcanicity, as witnessed by the presence of tuffs in the surrounding strata. Previous hydrothermal postulates appear untenable as it can be demonstrated that the barite generally has a marked stratigraphic control, being confined to the terminal phases of volcanic cycles in which mafic lavas, followed by felsic lavas or pyroclasts, are overlain by persistent cherty layers containing both massive and banded barite.

A similar origin is suggested for many of the barite occurrences in the Rhodesian greenstone belts despite the conclusions by Morrison (1970) that the majority of the deposits occur as vein fillings associated with faults traversing Archean granites, greenstones, and metasediments. Mineralization of this type is easily remobilized, transported and redeposited by secondary processes making its origins appear divergent and obscure.

#### Corundum and Kyanite

Rhodesia has for many years been the world's principal supplier of corundum, followed by South Africa, India, Kenya and Tanzania. Two commercial varieties are known, crystal corundum and boulder corundum. The crystal corundum deposits, the distribution of which is shown in Figure 9, owe their origin to a number of processes which, according to Morrison (1972) include : (1) the metamorphism of sedimentary rocks. (2) the desilication of pegmatites intrusive into hornblende and tremolite schists respectively (cf., Hall, 1920). (3) the metasomatism of amphibolites, biotite schists, serpentinites and fuchsite mica schists during granitization. (4) the effects of granulite facies metamorphism on ultramafic rocks, on anorthosites, and on biotite schist relics in leuco-paragneisses.

The absence of pegmatites in the development of (3) and (4) above is noteworthy. These deposits fall largely in the areas of influence of the Limpopo, Zambezi and Mocambique high-grade metamorphic belts.

The boulder corundum is, economically, more important, particularly in Rhodesia. To the end of 1965, Morrison (1972) showed that boulder corundum had accounted for 97.8 per cent of the 40 297 tons of corundum produced in that country. The distribution of this variety is shown in Figure 7. All these deposits are situated in rocks of the greenstone belts and, like the barite deposits, appear to occur at various stratigraphic levels within the volcanic sequences.

Morrison (1972), who investigated the boulder corundum deposits of Rhodesia, found the main features of the occurrences to be as follows : (1) most deposits are situated near greenstone-granite contacts or near intrusive mafic complexes or dykes, all of which provided a source of heat for the development of the corundum. (2) the corundum deposits form part of the lithological sequence and show no cross-cutting relationships with the country rocks. (3) the corundum commonly occurs as lenses within aluminous mica schists containing one or more of the following minerals : andalusite, chiastolite, sillimanite, and kyanite. Fuchsite, sericite,

rutile, diaspore and margarite frequently accompany the corundum. (4) the deposits occur with wall rocks that include ultramafic rocks (talc schists, serpentinites), banded iron formation, phyllites and argillaceous metasediments (mafic tuffs), aluminous micaceous schists and hornblende schists or gneisses.

Morrison (1972) concluded that the boulder corundum deposits of Rhodesia were formed as a result of metamorphism of alumina-rich sediments, presumably pre-existing bauxites. These findings have been confirmed in the Barberton area where the corundum host rocks consist of felsic volcanic rocks (mainly pyroclasts and tuffs - see chemical analysis, Table 10, column 7) the latter occurring as members of volcanic cycles within the greenstone belts.

In addition to the corundum mineralization, precious and semi-precious sapphires and rubies have been obtained from some of the crystal corundum deposits in northeastern Rhodesia. Furthermore, kyanite has been mined in some localities, including the Halfway Kop deposit in the Tati greenstone belt of Botswana (cf. Figures 1 and 7). Here lenses and pods of kyanite occur in quartz-kyanite-sericite-pyrophyllite-schists (Boocock, 1965). These rocks represent altered felsic pyroclasts and tuffs (Mason, 1970).

#### Copper-Lead-Zinc-Gold

The distribution of copper, lead and zinc mineralization is shown in Figure 8. Unlike the Canadian Archean, there are no known major massive base metal sulphide deposits in the greenstone belts of southern Africa. Most of the locations shown in Figure 8 are prospects or small workings. Only in a few cases (e.g. the Cactus Mine, south of Que Que in the Midlands greenstone belt) are deposits of this type being mined, and then only on a small scale.

On the Kaapvaal Craton occurrences of this type are virtually absent with the exception of small stratiform deposits (probably totalling less than 1 million tons) located along the "Letaba copper-zinc line" in the Murchison greenstone belt. The deposits, with the ore minerals pyrite, marcasite, chalcopyrite, cubanite, sphalerite, pyrrhotite, digenite, covellite, and galena (Hausmann, 1959), occur in quartz-mica and quartz-chlorite schists (low-to-intermediate metamorphic grades), now considered to be the altered equivalents of felsic volcanic rocks.

On the Rhodesian Craton the Cactus Mine is one of the few deposits being worked. The Cu-Pb-Zn ores, which also contain appreciable amounts of Sb, Au and Ag, occur in structurally disturbed felsic volcanic rocks (quartz felspar porphyries and felsic tuffs). The ore deposit is associated with fractures and appears to be of hydrothermal origin. It is possible that the ore deposit represents mineralization remobilized from volcanogenic source rocks.

Copper has also been recovered as a by-product of gold mining at the Falcon Mine (Figure 8) in the Umvuma greenstone belt, and is a common associate of gold in the small mines of the Felixburg, Odzi-Umtali, Midlands and Filabusi greenstone belts (cf., Figures 1 and 8). Indications of copper mineralization are also prevalent in many of the greenstone belts along the northwest margin of the exposed Archean terrane of the Rhodesian Craton, extending from the Matsitama belt in Botswana, through the western portions of the Bulawayo-Bubi and Midlands belts. The copper showings along this zone coincide with the major differences in metallogenic association reported by Bliss (1970), who maintained that the line of demarcation of an essentially gold-rich province (including scheelite), and a copper-nickel province (with minor gold, and no recorded scheelite) corresponded with the lithological variations recorded between rocks of the Bulawayan Group and those of the Umniati and Maliami River formations.

#### Massive Sulphide Iron Formations

Massive sulphide deposits of pyrite and pyrrhotite are known to occur in several areas on the Rhodesian Craton as well as in the Limpopo mobile belt (Figure 8). North of Salisbury, the Iron Duke Mine represents one of the few sulphide facies stratiform ore occurrences to have been mined essentially for its massive pyrite (Ferguson and Wilson, 1937). This, and other deposits in the Shamva-Salisbury greenstone belt (Figure 8), represents the product of exhalative volcanogenic activity during the deposition of the Shamvaian Group.

Further massive pyrite-pyrrhotite deposits are known in the Bulawayo, Gwanda and Tati greenstone belts. In the latter greenstone belt, Ryan (1970) reported the presence of a




Figure 8 : Distribution of copper, lead, zinc, copper-gold, tungsten, iron, limestone, and iron sulphide mineralization in Archean greenstone belts of southern Africa.

major pyrite/pyrrhotite deposit in the Mphoengs schist belt, which extends from Botswana into Rhodesia. The mineralization, which can be traced intermittently for over 18 km, occurs in association with a thin, but persistent, limestone unit that occurs in a succession of altered basalt-andesite-dacite-rhyolite volcanic and pyroclastic rocks. The sequence also contains banded iron formations, and is correlated with the Mafic-to-Felsic Unit.

#### Silver-Tungsten-Arsenic

Silver, tungsten, and arsenic mineralization occurs widespread throughout the southern African greenstone belts but is invariably a by-product of gold mining. The silver occurs alloyed

with gold, whereas the tungsten (mainly scheelite), and arsenic (arsenopyrite), is found associated with the gold ores. The main tungsten-rich gold deposits are shown in Figure 8. The distribution of the silver and arsenic follows closely the pattern of gold distribution shown in Figure 5.

#### ARCHEAN SEDIMENTARY DEPOSITS

Under this heading may be grouped the ancient placer gold-pyrite deposits that occur in conglomerate-quartzite formations of the Archean greenstone belts. These have been worked in a small way in the Uitkyk Formation of the Pietersburg greenstone belt (Figure 1) and in Shamvaian successions in parts of Rhodesia. Other sedimentary mineral occurrences of note include the Archean iron formations and the limestone deposits, the latter confined to the Rhodesian greenstone belts (Figure 8).

The iron formations in the greenstone belts are almost invariably associated with volcanic rocks. They are developed at various stratigraphic positions from the base to the top of the greenstone piles but are more commonly located in the lower, essentially volcanic environment as opposed to the terminal sedimentary parts of the stratigraphic column (Beukes, 1973). Detailed mineralogical data on the iron formations is not available but qualitative estimates suggest that the order of abundance of iron compounds deviates little from the findings of James (1966) and Stanton (1972) who reviewed the chemistry of iron formations of Archean age throughout the world. The iron oxides appear to be most abundant, followed by carbonate, silicate, and sulphide facies iron formations.

Almost without exception the greenstone belts in southern Africa have iron formations that may be grouped into one or more of the four abovementioned categories. Most important are the finely laminated, alternating, iron and silica-rich beds referred to locally as jaspilites or "banded ironstones". The general tenor of iron in these rocks varies from 30 to 40 per cent Fe, although in the Mwanesi greenstone belt in Rhodesia (Figure 1), up to 50 per cent Fe has been recorded (Worst, 1962a). Processes of enrichment of the protore give rise locally to deposits with 60-65 per cent Fe, examples of which are given later.

The banded iron formations occur in two environments. In the volcanic environment they are associated with mafic and felsic volcanics and are often members of a three part cycle consisting of mafic lava, felsic lava, and chert or iron formation. Examples of this type are found commonly in the Sebakwian and Bulawayan groups in Rhodesia (Bliss and Stidolph, 1969) and within the Onverwacht assemblages of the Barberton and Pietersburg greenstone belts. Iron formations also occur in an essentially sedimentary environment where they are associated with shales, phyllites, limestones, greywackes, quartzites, and minor lava units. As pointed out by Anhaeusser and Button (1974), the role of volcanism and iron deposition in such settings is not easy to assess and inferences have to be drawn largely from associated rock types. A close and probable genetic relationship to volcanism appears reasonable for some of the iron formations in the Shamvaian Group in Rhodesia and in the Moodies Group in the Barberton Mountain Land. In the latter case, banded jaspilitic iron formations occur stratigraphically above a minor basaltic lava unit (Anhaeusser, 1971) whereas in the underlying Fig Tree Group, iron formations occur in a predominantly sedimentary assemblage but which includes lava and pyroclastic members with soda trachytic affinities (Visser et al., 1956).

Subordinate carbonate and silicate iron formations are known as well as the sulphide facies varieties described earlier. Some iron formations, linked with those of the Archean, have been subjected to intermediate-to-high grades of metamorphism and have undergone recrystallization to magnetite quartzites. These changes have taken place mainly in the mobile belts encircling the cratons. Many of the magnetite quartzite occurrences along the southern and southeastern margins of the Rhodesian Craton (Figure 1) appear to belong to this category (Worst, 1962b). In addition, bands of resistant iron formation and magnetite quartzites also occur as xenoliths in the granites on the cratons.

Deposits of iron ore that are being mined include the enriched high-grade hematite ore bodies near Que Que in the Midlands greenstone belt, the deposits in the Buhwa belt (Worst, 1962b) and the enriched Fig Tree iron formations in the Ngwenya Mine area in Swaziland (Bursill et al., 1964). Large reserves of lower grade iron ore occurs in the Mwanesi Ranges (Worst, 1962a).

Limestone occurrences are found as stratigraphic interlayers within the volcanic successions. The deposits are mined for the manufacture of cement (e.g. Colleen Bawn Mine, Gwanda belt) and for the steel industry. It is of interest to note that the oldest stromatolites yet recorded have come from the Archean limestones in the Huntsman quarry north of Bulawayo (Macgregor, 1940).

#### MINERALIZATION ASSOCIATED WITH GRANITES AND PEGMATITES

A wide variety of minerals occur scattered throughout the Archean granitic terrane of southern Africa (Figure 9). With the exception of tin, molybdenum, and tantalum-columbium, the mineralization consists mainly of non-metallic deposits of beryllium, lithium, bismuth, mica, corundum and emeralds as well as felspar, silica and graphite. Most of the deposits are associated with pegmatites located close to the margins of greenstone belts, or in areas where there are numerous xenolithic greenstone remnants.

Although the deposits occur widespread through the region depicted in Figure 9, there are two prominent areas of pegmatite development from which the bulk of the mineralization has so far been recovered. The most important pegmatite area is that known as the Bikita Tinfield, located at the eastern extremity of the Fort Victoria greenstone belt in Rhodesia (Martin, 1964). Situated in this region is the Bikita pegmatite, one of the largest deposits of lithium, caesium and beryllium in the world. The Bikita deposit, which is approximately 2 650 m.y. old (Cooper, 1964), contains numerous minerals of economic importance. In order of abundance these are : petalite, lepidolite, spodumene, pollucite, beryl, eucryptite and amblygonite. The cassiterite, tantalite, and microlite deposits have been mined out.

The second prominent pegmatite area is referred to as the Gravelotte-Mica pegmatite field, located in and adjacent to remnants of the Murchison greenstone belt on the Kaapvaal Craton. Important deposits of mica occur in the region, which also supplies quantities of beryllium, lithium, tantalum and columbium, corundum, felspar and silica. Emeralds, together with beryl, are mined from pegmatites intruded into biotite schists. The pegmatites are associated with a late potassic granite body (the Mashishimala Pluton), which was intruded into the area south of the main remnant of the Murchison schist belt.

Deposits of corundum, associated with pegmatites, show a clustering in the region north of the Pietersburg and Murchison greenstone belts. These deposits which have been described by Hall (1920), yielded large quantities of corundum and made South Africa one of the world's leading producers of this mineral during the first quarter of this century.

Important deposits of emeralds occur not only in the Gravelotte-Mica area but also in the Rhodesian greenstone belts flanking the Limpopo metamorphic belt. The most valuable deposits are those associated with pegmatites in the Sandawana Emerald Mine area, adjacent to the Buhwa-Mweza greenstone belt. Emeralds have also been recovered in the Bikita area, the Odzi-Umtali area, and the Filabusi greenstone belt (Figure 9).

Tin has been recovered from a number of pegmatites throughout the Archean regions discussed in this paper. The most important deposits, which are now largely mined out, occur in the Bikita area (Martin, 1964), in the Salisbury greenstone belt, and in Swaziland, the latter deposits being associated with granites and pegmatites southeast of the Barberton greenstone belt (Davies, 1964).

Small graphite deposits occur in metamorphosed carbonaceous shale xenoliths in the Archean granites and gneisses. Most of these deposits are located in the Limpopo mobile belt and in the areas flanking this region in Rhodesia and South Africa.

#### DISCUSSION

Petrogenetic, tectonic and stratigraphic processes in Archean rocks appear very similar throughout the world (Goodwin, 1968, 1971; Anhaeusser et al., 1969; Glikson, 1970). Furthermore, as pointed out by Hutchinson et al., (1971), the rock types, stratigraphic relationships, and




Figure 9 : Map showing the distribution of mineral occurrences associated with Archean granites and pegmatites in southern Africa.

contained mineral deposits are similar in most shield areas, including Canada, Australia, Brazil, Fennoscandia, and Africa. Despite these similarities, which support suggestions that the Archean was a time of unique crustal development, increasing evidence is being made available demonstrating that appreciable differences do exist from one shield area to another.

The most significant differences are reflected in the variable geochronological and geochemical data emanating from studies of Archean rocks around the world. Although geological processes and responses appear to have been similar in all these ancient terranes, variations in geochemical characteristics of the Archean volcanic sequences probably reflect contrasting stages in crustal development, the latter also being time dependant.

Consequent on the geochemical changes that have been recorded, are variations relating to the nature and relative abundances of Archean mineral deposits throughout the world. In the preceding pages, attempts have been made to briefly describe and illustrate the nature and distribution of Archean mineralization in southern Africa. By comparing this mineralization with that reported in the Archean terranes of Canada, Western Australia, and elsewhere, a number of important features emerge. These include the following : (1) the range in the types of mineralization found in Archean successions is everywhere extensive, although the relative abundances of the mineralization is variable. (2) certain types of mineralization appear to be equally well-represented in greenstone terranes around the world. These include deposits of gold (and associated silver), as well as the various facies of iron formation. (3) chromite is most abundantly developed in the Archean successions of the Rhodesian Craton. (4) nickel-copper ores are particularly well-represented in the Yilgarn Province of the Western Australian Shield (Woodall and Travis, 1969; Fardon, 1971), and to a lesser extent on the Rhodesian Craton. Deposits of nickel-copper ores also occur in Canadian greenstone belts but, as pointed out by Naldrett and Gasparrini (1971), the ore bodies tend to be small. (5) massive base metal sulphide ore bodies containing copper, lead, and zinc are exceptionally well-developed in the Canadian greenstone belts. Prominent silver-gold mineralization is associated with these deposits (Goodwin, 1965; Hutchinson et al., 1971; Hutchinson, 1973). By contrast, Cu-Pb-Zn deposits of this type are rare in Western Australia and southern Africa, and virtually all the silver produced in these regions is alloyed with the gold mineralization. Absent are deposits of silver (and cobalt) like those at Cobalt, Ontario, in Canada. Although the latter deposits are closely related spatially to the Nipissing diabase sill, of Proterozoic age, the silver-cobalt veins are also close to the unconformity between Archean (Keewatin volcanics and Timiskaming sediments) and younger rocks (Gowganda Formation). In the opinion of Boyle (1968), and Hutchinson et al., (1971), the silver and other elements in the Cobalt area may have been derived from the underlying Archean volcanosedimentary rocks by thermal remobilization due to the diabase intrusion. (6) the Archean granite-greenstone terrane in southern Africa has important deposits of chrysotile asbestos and antimony and, in addition, has been a principal world supplier of certain non-metallic minerals associated with granites and pegmatites (particularly lithium, beryllium and corundum).

What reasons can be offered to explain the unequal concentration of mineralization types in the Archean successions of the world? There is always a danger in speculations of this kind as vast areas of the shields still remain to be mapped and geologically prospected. However, bearing these limitations in mind and considering the available data, it is possible to demonstrate a worldwide association of certain mineral types with particular host rock stratigraphy.

As was demonstrated earlier, the southern Africa greenstone belts, relative to their Canadian counterparts, contain a greater proportion of ultramafic and mafic rocks and are deficient in rocks of intermediate to felsic character. It is concluded that this disparity of lithological proportions, between the Canadian and southern African shields, is the underlying cause of the variability in the mineralization thus far encountered in the two regions.

The more prominent development, in southern Africa, of rocks of the type classified with the Lower Ultramafic Unit probably represents a function of the greater age of the latter rock assemblages. Greater abundances of the siderophile elements (e.g. Au, Ni, Fe, and the platinum-group metals), as well as chromium, are thus to be expected in rocks of this character. By contrast, the younger, well-differentiated, intermediate to felsic volcanic sequences of calc-alkaline affinity, like those so well-represented on the Canadian Shield, are the host rocks of the chalcophile elements (e.g. Cu, Pb, Zn), and represent more favourable target areas for Archean base metal exploration. It is suspected that mineralization of this type may yet be encountered in southern Africa but the greater age of the rocks and their proportionately smaller areal (volumetric) extent will probably provide constraints on the number and size of possible ore deposits that might be encountered.

#### REFERENCES

- Anhaeusser, C.R., 1965, Wrench faulting and its relationship to gold mineralization in the Barberton Mountain Land : Econ. Geol. Research Unit, Univ. Witwatersrand, Inf. Circ. 24, 25 pp.

- Anhaeusser, C.R., 1969, The stratigraphy, structure, and gold mineralization of the Jamestown and Sheba Hills areas of the Barberton Mountain Land : Unpub. Ph.D. thesis, Univ. Witwatersrand, Johannesburg, 332 pp.
- Anhaeusser, C.R., 1971, Cyclic volcanicity and sedimentation in the evolutionary development of Archean greenstone belts of shield areas : Geol. Soc. Australia Spec. Pub. 3, p. 57-70.
- Anhaeusser, C.R., 1972, The geology of the Jamestown Hills area of the Barberton Mountain Land, South Africa : Geol. Soc. South Africa Trans., v.75, Pt. 3, p. 225-263.
- Anhaeusser, C.R., 1973a, The evolution of the early Precambrian crust of southern Africa : Phil. Trans. Royal Soc. London, v.A273, p. 359-388.
- Anhaeusser, C.R., 1973b, Precambrian tectonic environments : Econ. Geol. Research Unit, Univ. Witwatersrand, Inf. Circ. 78, 19 pp.
- Anhaeusser, C.R., 1974, The nature of chrysotile asbestos occurrences in southern Africa - a review : Econ. Geol. Research Unit, Univ. Witwatersrand, Inf. Circ. 90, 28 pp.
- Anhaeusser, C.R., Roering, C., Viljoen, M.J. and Viljoen, R.P., 1968, The Barberton Mountain Land : a model of the elements and evolution of an Archean fold belt : Geol. Soc. South Africa Trans., v.71, Annex, p. 225-254.
- Anhaeusser, C.R., Mason, R., Viljoen, M.J. and Viljoen, R.P., 1969, A reappraisal of some aspects of Precambrian shield geology : Geol. Soc. America Bull., v.80, p. 2175-2200.
- Anhaeusser, C.R. and Button, A., 1974, A review of southern African stratiform ore deposits - their position in time and space : Econ. Geol. Research Unit, Univ. Witwatersrand, Inf. Circ. 85, 48 pp.
- Baragar, W.R.A. and Goodwin, A.M., 1969, Andesites and Archean volcanism of the Canadian Shield, in Andesite Cong. (Eugene and Bend, Oregon, 1968) Proc : Oregon Dept., Geology Mineral Industries Bull. 65, p. 121-142.
- Beukes, N.J., 1973, Precambrian iron-formations of southern Africa : Econ. Geol., v.68, p. 960-1004.
- Bliss, N.W., 1970, The geology of the country around Gatooma : Rhodesia Geol. Survey Bull. 64, 240 pp.
- Bliss, N.W. and Stidolph, P.A., 1969, The Rhodesian basement complex : a review : Geol. Soc. South Africa Spec. Pub. 2, p. 305.
- Boese, R., 1964, Die Antimonglaz-gänge van Gravelotte : Unpub. Ph.D. thesis, Univ. Hamburg, 85 pp.
- Boocock, C., 1965, Mineral resources of the Bechuanaland Protectorate : Overseas Geol. and Min. Res., v.9, No. 4, p. 369-417.
- Boyle, R.W., 1968, The geochemistry of silver and its deposits : Canada Geol. Survey Bull. 160, 264 pp.
- Brooks, C. and Hart, S.R., 1972, An extrusive basaltic komatiite from a Canadian Archean metavolcanic belt : Canadian Jour. Earth Sci., v.9, p. 1250-1253.
- Bursill, C., Luyt, J.F.M. and Urie, J.G., 1964, The Bomvu Ridge iron ore deposit : in Haughton, S.H., ed., The geology of some ore deposits in southern Africa. Geol. Soc. South Africa, v.2, p. 405-413.
- Cameron, E.N., 1964, Chromite deposits of the eastern part of the Bushveld Complex : in Haughton, S.H., ed., The geology of some ore deposits in southern Africa. Geol. Soc. South Africa, v.2, p. 131-168.

- Condie, K.C., Macke, J.E. and Reimer, T.O., 1970, Petrology and geochemistry of early Precambrian greywackes from the Fig Tree Group, South Africa : Geol. Soc. America Bull. v.81, p. 2759-2776.
- Cooper, D.G., 1964, The geology of the Bikita pegmatite : in Haughton, S.H., ed., The geology of some ore deposits in southern Africa : Geol. Soc. South Africa, v.2, p. 441-461.
- Cotterill, P., 1969, The chromite deposits of Selukwe, Rhodesia : in Wilson, H.D.B., ed., Magmatic Ore Deposits, a Symposium : Econ. Geol. Mono. 4, p. 154-186.
- Cousins, C.A. and Feringa, G., 1964, The chromite deposits of the western belt of the Bushveld Complex : in Haughton, S.H., ed., The geology of some ore deposits in southern Africa : Geol. Soc. South Africa, v.2, p. 183-202.
- Davies, D.N., 1964, The tin deposits of Swaziland : in Haughton, S.H., ed., The geology of some ore deposits in southern Africa : Geol. Soc. South Africa, v.2, p. 535-541.
- Davies, D.N. and Hunter, D.R., 1964, The gold deposits of the Barberton Mountain Land in Swaziland : in Haughton, S.H., ed., The geology of some ore deposits in southern Africa : Geol. Soc. South Africa, v.2, p. 59-75.
- Davies, R.D., Allsopp, H.L., Erlank, A.J. and Manton, W.I., 1969, Sr-isotopic studies on various layered mafic intrusions in southern Africa : Geol. Soc. South Africa Spec. Pub. 1, p. 576-593.
- Eales, H.V., 1964, Mineralogy and petrology of the Empress nickel-copper deposit, Southern Rhodesia : Geol. Soc. South Africa Trans., v.67, p. 173-201.
- Fardon, R.S., 1971, The Western Australian nickel deposits : Geol. Soc. Australia Spec. Pub. 3, p. 256 (Abstract).
- Ferguson, J.C. and Wilson, T.H., 1937, The geology of the country around the Jumbo mine, Mazoe district : Rhodesia Geol. Survey Bull. 33, 137 pp.
- Fripp, R.E.P., 1974, The distribution and significance of stratabound mineral deposits in Archean banded iron formation in Rhodesia with particular reference to the Vubachikwe gold mine : Unpub. M.ph. thesis, Univ. Rhodesia, Salisbury, 147 pp.
- Goldberg, I., 1964, Notes on the relationship between gold deposits and structure in southern Rhodesia : in Haughton, S.H., ed., The geology of some ore deposits in Southern Africa : Geol. Soc. South Africa, v.2, p. 9-13.
- Goodwin, A.M., 1965, Mineralized volcanic complexes in the Porcupine-Kirkland Lake-Noranda region, Canada : Econ. Geol., v.60, No. 5, p. 955-971.
- Goodwin, A.M., 1966, Archean protocontinental growth and mineralization : Canadian Min. Jour. v.87, p. 57-60.
- Goodwin, A.M., 1968, Archean protocontinental growth and early crustal history of the Canadian Shield : 23rd Int. Geol. Congr., Prague, 1, p. 69-89.
- Goodwin, A.M., 1971, Metallogenic patterns and evolution of the Canadian Shield : Geol. Soc. Australia Spec. Pub. 3, p. 157-174.
- Goodwin, A.M., Ambrose, J.W., Ayers, L.D., Clifford, P.M., Currie, K.L. and others, 1972, The Superior Province : in Price, R.A. and Douglas, R.J.W., eds., Variations in tectonic styles in Canada. Spec. Paper 11, p. 527-623.
- Glikson, A.Y., 1970, Geosynclinal evolution and geochemical affinities of early Precambrian systems : Tectonophysics, v.9, p. 397-433.
- Hall, A.L., 1920, Corundum in the northern and eastern Transvaal : Geol. Survey South Africa Mem. 15, 223 pp.

- Hallberg, J.A., 1972, Geochemistry of Archaean volcanic belts in the Eastern Goldfields region of Western Australia : Jour. Petrology, v.13, p. 45-56.
- Haughton, S.H., (editor), 1964, The geology of some ore deposits in southern Africa : Geol. Soc. South Africa, v.2, 739 pp.
- Hausmann, S.G., 1959, A mineralogical investigation of the Letaba copper-zinc ores and the Monarch cinnabar deposits located in the Murchison range of the eastern Transvaal : Unpub. M.Sc. thesis, Univ. Witwatersrand, Johannesburg, 49 pp.
- Hunter, D.R., 1970, The Ancient Gneiss Complex in Swaziland : Geol. Soc. South Africa Trans., v.73, p. 107-150.
- Hutchinson, R.W., 1973, Volcanogenic sulfide deposits and their metallogenetic significance : Econ. Geol., v.68, No.8, p. 1223-1246.
- Hutchinson, R.W., Ridler, R.H. and Suffel, G.G., 1971, Metallogenetic relationships in the Abitibi belt, Canada : a model for Archean metallogeny : Canadian Mining Metall., Bull., v.74, p. 48-57.
- James, H.L., 1966, Chemistry of the iron-rich sedimentary rocks : U.S. Geol. Survey Prof. Paper 440-W, 61 pp.
- Laubscher, D.H., 1964, The occurrence and origin of chrysotile asbestos and associated rocks, Shabani, Southern Rhodesia : in Haughton, S.H., ed., The geology of some ore deposits in southern Africa. Geol. Soc. South Africa, p. 593-624.
- Laubscher, D.H., 1968, The origin and occurrence of chrysotile asbestos in the Shabani and Mashaba areas, Rhodesia : Geol. Soc. South Africa Trans., v.71, Annex. p. 195-204.
- Le Roex, H.D., 1964, Nickel deposit on the Trojan claims, Bindura district, Southern Rhodesia : in Haughton, S.H., ed., The geology of some ore deposits in southern Africa : Geol. Soc. South Africa, v.2, p. 509-520.
- Macgregor, A.M., 1940, A Pre-Cambrian algal limestone in Southern Rhodesia : Geol. Soc. South Africa Trans., v.43, p. 9-15.
- Macgregor, A.M., 1951, Some milestones in the Precambrian of Southern Rhodesia : Geol. Soc. South Africa Proc., v.54, p. xxvii-xxi.
- MacGregor, B.I. and Bliss, N.W., 1968, The Barton Farm magnesite deposit, Gatooma Rhodesia : Geol. Soc. South Africa Trans., v.71, Annex, p. 159-172.
- McCall, G.J.H., 1973, Geochemical characteristics of some Archaean greenstone suites of the Yilgarn structural province, Australia : Chemical Geology, v.11, p. 243-269.
- McGlynn, J.C. and Henderson, J.B., 1972, The Slave Province : in Price, R.A. and Douglas, R.J.W., eds., Variations in tectonic styles in Canada. Geol. Assoc. Canada Spec. Paper 11, p. 505-526.
- Martin, H.J., 1964, The Bikita Tinfield : Rhodesia Geol. Survey Bull. 58, p. 114-132.
- Mason, R., 1970, The geology of the country between Francistown and Madinare - northeastern Botswana : Unpub. Ph.D. thesis, Univ. Witwatersrand, Johannesburg, 186 pp.
- Mason, R., 1973, The Limpopo mobile belt - southern Africa : Phil. Trans. Royal Soc. London, Ser. A., v.273, p. 463-485.
- Mehliss, A.T.M., 1968, The geology of the Indarama gold-antimony mine Sebakwe area, Que Que, Rhodesia : Geol. Soc. South Africa Trans., v.71, Annex, p. 135-146.
- Mendelsohn, E., 1938, Gold deposits of the central Murchison range, Transvaal : Geol. Soc. South Africa Trans., v.41, p. 249-272.

- Morrison, E.R., 1970, Barium minerals in Rhodesia : Rhodesia Geol. Survey Mineral Res. Ser. 15, 16 pp.
- Morrison, E.R., 1972, Corundum in Rhodesia : Rhodesia Geol. Survey Mineral Res. Ser. 16, 24 pp.
- Naldrett, A.J. and Gasparrini, E.L., 1971, Archaean nickel sulphide deposits in Canada : their classification, geological setting and genesis with some suggestions as to exploration : Geol. Soc. Australia Spec. Pub. 3, p. 201-226.
- Nesbitt, R.W., 1971, Skeletal crystal forms in the ultramafic rocks of the Yilgarn Block, Western Australia : evidence for an Archaean ultramafic liquid : Geol. Soc. Australia Spec. Pub. 3, p. 331-347.
- Phaup, A.E., 1964, Gold mines of Southern Rhodesia - introduction : in Haughton, S.H., ed., The geology of some ore deposits in southern Africa : Geol. Soc. South Africa, v.2, p. 1-7.
- Pretorius, D.A., 1973, The role of E.G.R.U. in mineral exploration in South Africa : Econ. Geol. Research Unit, Univ. Witwatersrand, Johannesburg, Inf. Circ. 77, 16 pp.
- Pretorius, D.A., 1974a, Gold in the Proterozoic sediments of South Africa : systems, paradigms, and models : Econ. Geol. Research Unit, Univ. Witwatersrand, Johannesburg, Inf. Circ. 87, 22 pp.
- Pretorius, D.A., 1974b, The nature of the Witwatersrand gold-uranium deposits : Econ. Geol. Research Unit, Univ. Witwatersrand, Johannesburg, Inf. Circ. 86, 50 pp.
- Pretorius, D.A. and Hempkins, W.B., 1968a, Statistical moments of the frequency distribution of Rhodesian gold mineralization in time and space : a preliminary analysis : Geol. Soc. South Africa Trans., v.71, Annex, p. 9-19.
- Pretorius, D.A. and Hempkins, W.B., 1968b, Trend surface analysis of patterns of gold production from the Rhodesian basement between 1898 and 1932 : A first approach : Geol. Soc. South Africa Trans., v.71, Annex, p. 35-43.
- Prider, R.T., 1965, Geology and mineralization of the Western Australian Shield : in McAndrew, J., ed., Geology of Australian Ore Deposits, v.1 (2nd edition), Aust. Inst. Mining and Metall., Melbourne, p. 56-65.
- Reimer, T.O. and Heinrichs, T.K., 1974, A sedimentary barite deposit from the early Precambrian Fig Tree Group : (in press).
- Robertson, I.D.M., 1972, Mercury in Rhodesia : Rhodesia Geol. Survey Mineral Res. Ser., 17, 17 pp.
- Ryan, P.J., 1970, Progress report on exclusive prospecting area No. 292 Mphoengs, Rhodesia : Unpub. Rep. Southern Exploration (Pvt) Limited, 20 pp.
- Saager, R. and Köppel, V., 1974, Lead isotopes and trace elements from sulfides of Archean greenstone belts in South Africa - a contribution to the knowledge of the oldest known mineralization : Econ. Geol. (in press).
- Sahli, E.W., 1961, Antimony in the Murchison range of the northeastern Transvaal : Trans. 7th Commonwealth Min. and Metall. Congr. Johannesburg, v.1, p. 181-199.
- Sharpe, J.W.N., 1964, The Empress nickel-copper deposit, Southern Rhodesia : in Haughton, S.H., ed., The geology of some ore deposits in southern Africa. Geol. Soc. South Africa, v.2, p. 497-508.
- Söhnge, P.G., 1945, The geology of the Messina copper mines and surrounding country : Geol. Survey South Africa Mem. 40, 280 pp.
- Stanton, R.L., 1972, Ore petrology : McGraw-Hill, Inc., New York, 713 pp.
- Stowe, C.W., 1968a, The geology of the country south and west of Selukwe : Rhodesia Geol. Survey Bull. 59, 209 pp.

- Stowe, C.W., 1968b, Intersecting fold trends in the Rhodesian basement south and east of Selukwe : Geol. Soc. South Africa Trans., v.71, Annex, p. 53-78.
- van Biljon, W.J., 1964, The chrysotile deposits of the eastern Transvaal and Swaziland : in Haughton, S.H., ed., The geology of some ore deposits in southern Africa : Geol. Soc. South Africa, v.2, p. 625-669.
- van Breemen, O. and Dodson, M.H., 1972, Metamorphic chronology of the Limpopo belt, southern Africa : Geol. Soc. America Bull., v.83, p. 2005-2018.
- van der Kaaden, G., 1970, Chromite-bearing ultramafic and related gabbroic rocks and their relationship to "ophiolitic" extrusive basic rocks and diabases in Turkey : Geol. Soc. South Africa Spec. Pub. 1, p. 511-531.
- van Eeden, O.R., Partridge, F.C., Kent, L.E. and Brandt, J.W., 1939, The mineral deposits of the Murchison range east of Leydsdorp : Geol. Survey South Africa Mem. 36, 163 pp.
- van Eeden, O.R., Visser, H.N., van Zyl, J.S., Coertze, F.J. and Wessels, J.T., 1955, The geology of the eastern Soutpansberg and the lowveld to the north : Geol. Survey South Africa, Expl. Sheet 42 (Soutpansberg), 126 pp.
- van Zyl, J.S., Boardman, L.G., Brandt, J.W. and de Villiers, J., 1942, Magnesite in the Union of South Africa : Geol. Survey South Africa Mem. 38, 74 pp.
- Viljoen, M.J. and Viljoen, R.P., 1969a, The geology and geochemistry of the Lower Ultramafic Unit of the Onverwacht Group and a proposed new class of igneous rocks : Geol. Soc. South Africa Spec. Pub. 2, p. 55-85.
- Viljoen, R.P. and Viljoen, M.J., 1969b, The geological and geochemical significance of the upper formations of the Onverwacht Group : Geol. Soc. South Africa Spec. Pub. 2, p. 113-151.
- Viljoen, R.P. and Viljoen, M.J., 1969c, The relationship between mafic and ultramafic magma derived from the upper mantle and the ore deposits of the Barberton region : Geol. Soc. South Africa Spec. Pub. 2, p. 221-244.
- Viljoen, M.J. and Viljoen, R.P., 1969d, An introduction to the geology of the Barberton granite-greenstone terrain : Geol. Soc. South Africa Spec. Pub. 2, p. 9-28.
- Viljoen, M.J. and Viljoen, R.P., 1969e, The geochemical evolution of the granitic rocks of the Barberton region : Geol. Soc. South Africa Spec. Pub. 2, p. 189-218.
- Viljoen, M.J. and Viljoen, R.P., 1969f, A reappraisal of granite-greenstone terrains of shield areas based on the Barberton model : Geol. Soc. South Africa Spec. Pub. 2, p. 245-274.
- Viljoen, R.P. and Viljoen, M.J., 1969g, The geology and geochemistry of the layered ultramafic bodies of the Kaapmuiden area, Barberton Mountain Land : Geol. Soc. South Africa Spec. Pub. 1, p. 661-688.
- Viljoen, R.P. and Viljoen, M.J., 1971, The geological and geochemical evolution of the Onverwacht volcanic group of the Barberton Mountain Land, South Africa : Geol. Soc. Australia Spec. Pub. 3, p. 133-149.
- Viljoen, R.P., Saager, R. and Viljoen, M.J., 1969, Metallogenesis and ore control in the Steyndorp Goldfield, Barberton Mountain Land, South Africa : Econ. Geol., v.64, p. 778-797.
- Viljoen, R.P., Saager, R. and Viljoen, M.J., 1970, Some thoughts on the origin and processes responsible for the concentration of gold in the early Precambrian of southern Africa : Mineral. Deposita (Berlin), v. 5, p. 164-180.
- Visser, D.J.L., compiler, 1956, The geology of the Barberton area : Geol. Survey South Africa Spec. Pub. 15, 253 pp.
- Visser, D.J.L., (editor), 1968, Symposium on the Rhodesian basement complex : Geol. Soc. South Africa Trans., v.71, Annex, 282 pp.

- Wilson, H.D.B., Andrews, P., Moxham, R.L. and Ramlal, K., 1965, Archean volcanism of the Canadian Shield : Canadian Jour. Earth Sci., v.2, p. 161-175.
- Wilson, J.F., 1968, The geology of the country around Mashaba : Rhodesia Geol. Survey Bull. 62, 239 pp.
- Wilson J.F., 1973, The Rhodesian Archaean craton - an essay in cratonic evolution : Phil. Trans. Royal Soc. London, Ser. A., v.273, p. 389-411.
- Woodall, R. and Travis, G.A., 1969, The Kambalda nickel deposits, Western Australia : 9th Commonwealth Min. and Metall. Congress, London, v.2, p. 517-533.
- Worst, B.G., 1960, The Great Dyke of Southern Rhodesia : Rhodesia Geol. Survey Bull. 47, 239 pp.
- Worst, B.G., 1962a, The geology of the Mwanesi range and the adjoining country : Rhodesia Geol. Survey Bull. 54, 66 pp.
- Worst, B.G., 1962b, The geology of the Buhwa iron ore deposits and adjoining country : Belingwe district : Rhodesia Geol. Survey Bull. 53, 114 pp.
- Worst, B.G., 1964, Chromite in the Great Dyke of Southern Rhodesia : in Haughton, S.H., ed., The geology of some ore deposits in southern Africa : Geol. Soc. South Africa, v.2, p. 209-224.

\* \* \* \* \*