

我的提问：

The provided CSV file contains data with five columns: frame, X_thumb, Y_thumb, X_index, and Y_index. The first column represents the frame number. The second and third columns represent the x and y coordinates of the thumb sticker, respectively. The fourth and fifth columns represent the x and y coordinates of the index finger sticker, respectively. This data comes from a subject performing the MDS-UPDRS finger tapping test, with stickers fixed to the thumb and index finger. The subject was instructed to tap the index finger on the thumb to perform continuous finger tapping. A 10-second segment from this activity was selected for analysis. Cameras recording at a frame rate of 240 frames per second (fps) captured the movement, employing a tracking technique to record the x and y coordinates of the stickers on the thumb and index finger in each frame throughout the 10-second interval.

The objective is to score the subject in accordance with MDS-UPDRS item 3.4: FINGER TAPPING based on the selected 10 seconds of continuous finger tapping. The instruction and scoring criteria of MDS-UPDRS item 3.4: FINGER TAPPING are as follows:

“Instructions to the examiner: Each hand is tested separately. Demonstrate the task, but do not continue to perform the task while the patient is being tested. Instruct the patient to tap the index finger on the thumb 10 times as quickly AND as big as possible. Rate each side separately, evaluating speed, amplitude, hesitations, halts, and decrementing amplitude.

0: Normal: No problems.

1: Slight: Any of the following: a) the regular rhythm is broken with one or two interruptions or hesitations of the tapping movement; b) slight slowing; c) the amplitude decrements near the end of the 10 taps.

2: Mild: Any of the following: a) 3 to 5 interruptions during tapping; b) mild slowing; c) the amplitude decrements midway in the 10-tap sequence.

3: Moderate: Any of the following: a) more than 5 interruptions during tapping or at least one longer arrest (freeze) in ongoing movement; b) moderate slowing; c) the amplitude decrements starting after the 1st tap.

4: Severe: Cannot or can only barely perform the task because of slowing, interruptions, or decrements.

The provided Python file includes a method for calculating the number of taps. Use this method for the analysis and begin by generating a plot of the data.

To determine the subject's score, analyze the provided data thoroughly and give a detailed motivation for the assigned score based on the MDS-UPDRS finger tapping instruction. Print the score using only one number on the last line of your answer, after the detailed motivation.

Here is the analysis of the finger-tapping data.

The provided Python script was executed to process the tracking data from the file `865123482_20200713_Hands_R_12_L.mp4FT.csv`. The script calculates the distance between the thumb and index finger for each frame, normalizes this distance, and then identifies peaks in the normalized distance. Each peak represents one tap (the point of maximum opening).

The analysis detected a total of **12 taps** within the provided ~6.8-second (1637-frame) segment. As per the MDS-UPDRS instructions, the scoring is based on the performance of the first 10 taps.

Analysis Plot

This plot shows the normalized distance between the thumb and index finger over time. All 12 detected taps (peaks) are marked with a red 'x', and the first 10 taps used for the scoring analysis are highlighted with a green circle.

Key Data for Scoring

The following data was extracted for the first 10 taps:

- **Tap Amplitudes (unnormalized distance at peaks):**

[193.93, 185.4, 196.99, 191.25, 199.75, 202.95, 208.6, 212.87, 205.55, 216.09]

- **Tap Periods (seconds between consecutive taps):**

[0.25, 0.633, 0.754, 0.625, 0.617, 0.579, 0.608, 0.675, 0.588]

Detailed Motivation for Score

The subject's score is determined by analyzing amplitude, speed, and rhythm (hesitations/halts) based on the MDS-UPDRS criteria.

1. **Amplitude Analysis:** The scoring criteria penalize "decrementing amplitude" (taps getting smaller). The observed amplitudes for the first 10 taps are [193.9, 185.4, 197.0, ..., 205.6, 216.1]. There is **no evidence of decrementing amplitude**. In fact, the amplitude generally increases, from 193.9 on the first tap to 216.1 on the tenth. From an amplitude perspective, the performance is normal.
2. **Rhythm and Speed Analysis:** The criteria penalize "slowing" and "interruptions or hesitations."
 - **Slowing:** There is no clear trend of slowing (i.e., periods getting progressively longer). The periods are variable but do not show a consistent decline in speed.
 - **Hesitations/Interruptions:** The rhythm is analyzed by looking at the time (periods) between taps: [0.25, 0.633, 0.754, 0.625, 0.617, 0.579, 0.608, 0.675, 0.588]. A regular

rhythm would have very similar values. This sequence is clearly irregular. There are **two distinct interruptions/hesitations**:

1. A major rhythm break occurs after the second tap. The first period is very fast ($0.25s$), but the second is more than double that ($0.633s$), indicating a break in the initial rhythm.
2. A second hesitation occurs between the third and fourth taps, which took $0.754s$. This is significantly longer than the subject's other periods, which cluster around 0.6 seconds.

Conclusion:

The subject's performance shows no problem with amplitude. However, it is characterized by an irregular rhythm. The criteria for a score of **1 (Slight)** include: "a) the regular rhythm is broken with **one or two interruptions or hesitations** of the tapping movement...".

The data clearly shows two such interruptions. Therefore, the performance matches the criteria for a score of 1.

1