

我的提问：

The provided CSV file contains data with five columns: frame, X_thumb, Y_thumb, X_index, and Y_index. The first column represents the frame number. The second and third columns represent the x and y coordinates of the thumb sticker, respectively. The fourth and fifth columns represent the x and y coordinates of the index finger sticker, respectively. This data comes from a subject performing the MDS-UPDRS finger tapping test, with stickers fixed to the thumb and index finger. The subject was instructed to tap the index finger on the thumb to perform continuous finger tapping. A 10-second segment from this activity was selected for analysis. Cameras recording at a frame rate of 240 frames per second (fps) captured the movement, employing a tracking technique to record the x and y coordinates of the stickers on the thumb and index finger in each frame throughout the 10-second interval.

The objective is to score the subject in accordance with MDS-UPDRS item 3.4: FINGER TAPPING based on the selected 10 seconds of continuous finger tapping. The instruction and scoring criteria of MDS-UPDRS item 3.4: FINGER TAPPING are as follows:

“Instructions to the examiner: Each hand is tested separately. Demonstrate the task, but do not continue to perform the task while the patient is being tested. Instruct the patient to tap the index finger on the thumb 10 times as quickly AND as big as possible. Rate each side separately, evaluating speed, amplitude, hesitations, halts, and decrementing amplitude.

0: Normal: No problems.

1: Slight: Any of the following: a) the regular rhythm is broken with one or two interruptions or hesitations of the tapping movement; b) slight slowing; c) the amplitude decrements near the end of the 10 taps.

2: Mild: Any of the following: a) 3 to 5 interruptions during tapping; b) mild slowing; c) the amplitude decrements midway in the 10-tap sequence.

3: Moderate: Any of the following: a) more than 5 interruptions during tapping or at least one longer arrest (freeze) in ongoing movement; b) moderate slowing; c) the amplitude decrements starting after the 1st tap.

4: Severe: Cannot or can only barely perform the task because of slowing, interruptions, or decrements.

The provided Python file includes a method for calculating the number of taps. Use this method for the analysis and begin by generating a plot of the data.

To determine the subject's score, analyze the provided data thoroughly and give a detailed motivation for the assigned score based on the MDS-UPDRS finger tapping instruction. Print the score using only one number on the last line of your answer, after the detailed motivation.

Here is the analysis of the finger-tapping performance based on the provided data and MDS-UPDRS criteria.

Finger Tapping Analysis Plot

The plot below visualizes the normalized distance between the thumb and index finger over time. Each 'x' marker indicates a detected tap (specifically, the point of maximum opening between taps).

Detailed Motivation for Score

The analysis of the data, derived from the provided Python script, reveals the following key metrics for scoring:

- **Total Duration:** The provided data segment contains 1636 frames, which at 240 fps, corresponds to a duration of **6.82 seconds**.
- **Number of Taps:** Within this 6.82-second period, **12 taps** were detected.

- **Amplitude Analysis:** The amplitudes, representing the raw distance (in pixels) at the peak of finger opening for each tap, were:

```
[193.9, 185.4, 197.0, 191.3, 199.8, 203.0, 208.6, 212.9, 205.5, 216.1, 207.4, 205.1]
```

There is no evidence of *decrementing amplitude*. In fact, after the first two taps, the amplitude generally increases or stays consistently large. The performance shows "big" movements as instructed.

- **Speed and Rhythm Analysis (Hesitations/Interruptions):** The periods between taps (measured in frames) were: [60, 152, 181, 150, 148, 139, 146, 162, 141, 144, 143]

A regular rhythm would show consistent values. This sequence is highly irregular.

- The first tap is very quick (60 frames).
- The rhythm then breaks, with the next period being more than twice as long (152 frames).
- Another break occurs with a period of 181 frames.
- The standard deviation of the tap period is 28.4 frames, which is very high relative to the average period (142.36 frames), confirming a significant lack of rhythm.
- These breaks (e.g., the jump from 60 to 152, and the 181-frame period) qualify as interruptions or hesitations in the tapping movement. There are at least **3 to 5 clear interruptions** (e.g., the 60, 152, and 181-frame periods are all significant deviations from the mean).

Conclusion:

The subject maintains excellent amplitude, showing no decrement. However, the performance is characterized by a highly irregular rhythm with multiple interruptions and hesitations (at least 3-5).

According to the MDS-UPDRS 3.4 criteria:

- Score 0 (Normal) is incorrect due to the interruptions.
- Score 1 ("...one or two interruptions...") is an understatement.
- Score 2 ("...3 to 5 interruptions during tapping...") accurately describes the observed performance.
- Score 3 is too severe, as there is no amplitude decrement and the slowing is not "moderate" (the average speed is maintained, but it's *irregular*).

Therefore, the performance profile aligns best with a score of 2.

2