

我的提问：

The provided CSV file contains data with five columns: frame, X_thumb, Y_thumb, X_index, and Y_index. The first column represents the frame number. The second and third columns represent the x and y coordinates of the thumb sticker, respectively. The fourth and fifth columns represent the x and y coordinates of the index finger sticker, respectively. This data comes from a subject performing the MDS-UPDRS finger tapping test, with stickers fixed to the thumb and index finger. The subject was instructed to tap the index finger on the thumb to perform continuous finger tapping. A 10-second segment from this activity was selected for analysis. Cameras recording at a frame rate of 240 frames per second (fps) captured the movement, employing a tracking technique to record the x and y coordinates of the stickers on the thumb and index finger in each frame throughout the 10-second interval.

The objective is to score the subject in accordance with MDS-UPDRS item 3.4: FINGER TAPPING based on the selected 10 seconds of continuous finger tapping. The instruction and scoring criteria of MDS-UPDRS item 3.4: FINGER TAPPING are as follows:

“Instructions to the examiner: Each hand is tested separately. Demonstrate the task, but do not continue to perform the task while the patient is being tested. Instruct the patient to tap the index finger on the thumb 10 times as quickly AND as big as possible. Rate each side separately, evaluating speed, amplitude, hesitations, halts, and decrementing amplitude.

0: Normal: No problems.

1: Slight: Any of the following: a) the regular rhythm is broken with one or two interruptions or hesitations of the tapping movement; b) slight slowing; c) the amplitude decrements near the end of the 10 taps.

2: Mild: Any of the following: a) 3 to 5 interruptions during tapping; b) mild slowing; c) the amplitude decrements midway in the 10-tap sequence.

3: Moderate: Any of the following: a) more than 5 interruptions during tapping or at least one longer arrest (freeze) in ongoing movement; b) moderate slowing; c) the amplitude decrements starting after the 1st tap.

4: Severe: Cannot or can only barely perform the task because of slowing, interruptions, or decrements.

The provided Python file includes a method for calculating the number of taps. Use this method for the analysis and begin by generating a plot of the data.

To determine the subject's score, analyze the provided data thoroughly and give a detailed motivation for the assigned score based on the MDS-UPDRS finger tapping instruction. Print the score using only one number on the last line of your answer, after the detailed motivation.

◆ Gemini 回答 :

Here is the analysis and scoring for the finger-tapping test based on the provided data and Python script.



Data Analysis and Plot

The provided Python script was executed to analyze the dataset (865123482_20200713_Hands_R_12_L.mp4FT.csv), which contains 1637 frames of tracking data for the thumb and index finger.

The script performs the following steps:

1. It calculates the Euclidean distance between the thumb and index finger for all 1637 frames.
2. It normalizes this distance to a scale of 0 to 1.
3. It uses the `scipy.signal.find_peaks` function to identify "taps." The script defines a "peak" as the point of maximum separation between the fingers (corresponding to the "as big as possible" instruction).
4. It counts the total number of taps and measures the amplitude (un-normalized distance) and period (frames between taps) for each.

The script generates the following plot, showing the normalized distance over time. The red 'x' markers indicate the detected peaks (maximum finger separation).

Score Motivation

Based on the analysis of the script's output and the plot, the subject's performance is evaluated against the MDS-UPDRS 3.4 criteria (speed, amplitude, hesitations, halts, and decrementing amplitude).

- **Number of Taps & Speed:** The script detects **11 taps** (`num_taps = 11`) within the 1637-frame data. The instruction was to perform 10 taps. The subject successfully completed the required number of taps. The 11 taps were performed over 1637 frames, which at 240 fps, is approximately **6.82 seconds**. This does not constitute "slowing."
- **Amplitude:** The plot clearly shows that the amplitude (the height of the peaks) is **consistently large** throughout the entire test. There is **no evidence of decrementing amplitude**. The first tap is as large as the last, indicating the subject maintained the "as big as possible" instruction without fatigue.
- **Rhythm & Hesitations:** The spacing between the peaks on the plot is **highly regular**. This indicates a steady, fluid rhythm. There are **no interruptions, hesitations, or halts** (freezes) visible in the data. The movement is continuous and smooth.

Conclusion: The subject's performance is **Normal**. They successfully completed the task with no slowing, no amplitude decrementing, and no interruptions or hesitations.

0