

我的提问：

The provided CSV file contains data with five columns: frame, X_thumb, Y_thumb, X_index, and Y_index. The first column represents the frame number. The second and third columns represent the x and y coordinates of the thumb sticker, respectively. The fourth and fifth columns represent the x and y coordinates of the index finger sticker, respectively. This data comes from a subject performing the MDS-UPDRS finger tapping test, with stickers fixed to the thumb and index finger. The subject was instructed to tap the index finger on the thumb to perform continuous finger tapping. A 10-second segment from this activity was selected for analysis. Cameras recording at a frame rate of 240 frames per second (fps) captured the movement, employing a tracking technique to record the x and y coordinates of the stickers on the thumb and index finger in each frame throughout the 10-second interval.

The objective is to score the subject in accordance with MDS-UPDRS item 3.4: FINGER TAPPING based on the selected 10 seconds of continuous finger tapping. The instruction and scoring criteria of MDS-UPDRS item 3.4: FINGER TAPPING are as follows:

“Instructions to the examiner: Each hand is tested separately. Demonstrate the task, but do not continue to perform the task while the patient is being tested. Instruct the patient to tap the index finger on the thumb 10 times as quickly AND as big as possible. Rate each side separately, evaluating speed, amplitude, hesitations, halts, and decrementing amplitude.

0: Normal: No problems.

1: Slight: Any of the following: a) the regular rhythm is broken with one or two interruptions or hesitations of the tapping movement; b) slight slowing; c) the amplitude decrements near the end of the 10 taps.

2: Mild: Any of the following: a) 3 to 5 interruptions during tapping; b) mild slowing; c) the amplitude decrements midway in the 10-tap sequence.

3: Moderate: Any of the following: a) more than 5 interruptions during tapping or at least one longer arrest (freeze) in ongoing movement; b) moderate slowing; c) the amplitude decrements starting after the 1st tap.

4: Severe: Cannot or can only barely perform the task because of slowing, interruptions, or decrements.

The provided Python file includes a method for calculating the number of taps. Use this method for the analysis and begin by generating a plot of the data.

To determine the subject's score, analyze the provided data thoroughly and give a detailed motivation for the assigned score based on the MDS-UPDRS finger tapping instruction. Print the score using only one number on the last line of your answer, after the detailed motivation.

Here is the analysis and scoring for the finger-tapping data.

Finger Tapping Analysis Plot

The plot below visualizes the normalized distance between the thumb and index finger over the ~4-second recording. Each "x" marks a detected tap (peak distance). The first 10 taps, used for the scoring, are highlighted with red circles.

Detailed Motivation for Score

The analysis was performed by executing the provided Python script's methodology on the CSV data. The script calculates the distance between the thumb and index finger, normalizes it, and identifies peaks (taps) using `find_peaks`. The scoring is based on the MDS-UPDRS criteria for item 3.4, focusing on the *first 10 taps* detected.

The analysis of the 4.06-second clip (975 frames) detected a total of 11 taps, meaning the 10-tap task was fully completed.

1. Amplitude and Decrementing Amplitude

The raw distance (amplitude) for the first 10 taps was:

```
[139.05, 161.52, 160.08, 144.31, 136.69, 168.34, 144.31, 136.03, 122.15, 139.2]
```

- **Observation:** The amplitude is not normal. It does not show a consistent decrement from the start (it actually increases to a peak at tap 6). However, after peaking midway through the sequence (tap 6 at 168.34), there is a clear and significant decrement in the second half of the task (taps 7, 8, and 9 are 144.3, 136.0, and 122.2, respectively).
- **Scoring:** This pattern directly aligns with the criterion for a **Score 2**: "amplitude decrements midway in the 10-tap sequence."

2. Speed, Rhythm, and Hesitations

The time (in frames) between the first 10 taps was:

```
[99, 92, 104, 78, 106, 79, 104, 78, 91]
```

- **Observation:** The rhythm is highly irregular. A "Normal" (Score 0) performance would have very consistent periods. This subject's performance shows significant variability, with a mean of 92.3 frames but a standard deviation of 11.0 frames (a Coefficient of Variation of 11.9%). The subject clearly speeds up (e.g., 78-79 frames) and slows down (e.g., 104-106 frames) multiple times. These fluctuations count as "interruptions" or "hesitations" in the rhythm. There are at least 4-5 such significant deviations from the mean pace.
- **Scoring:** This irregularity fits the criterion for a **Score 2**: "3 to 5 interruptions during tapping."

Conclusion

The subject's performance exhibits multiple clear indicators for a mild impairment. Both the amplitude (decrementing midway) and the rhythm (3-5 interruptions/hesitations) independently meet the criteria for a score of 2.

2