

# 数学模型 HW1

2019 年 3 月 6 日

## 1 第一题

### 1.1

在  $(\bar{A}, \bar{B})$  变化率为 0, 因此

$$-\bar{A} + \bar{B} = \bar{A} - \bar{B} = 0,$$

根据

$$\bar{A} + \bar{B} = A_0 + B_0,$$

我们有

$$\bar{A} = \bar{B} = \frac{A_0 + B_0}{2}.$$

### 1.2

令  $f(x) = \ln x + 1/x - 1$ , 我们有  $f(1) = 0$ ,

$$f'(x) = \frac{x-1}{x^2} \begin{cases} < 0, & 0 < x < 1, \\ > 0, & x > 1. \end{cases}$$

因此  $f(x) \geq f(1) = 0$ 。从而  $\ln(x) \geq 1 - \frac{1}{x}$ 。从而

$$\begin{aligned} & A \ln \left( \frac{A}{\bar{A}} \right) + B \ln \left( \frac{B}{\bar{B}} \right) \\ & \geq A \left( 1 - \frac{\bar{A}}{A} \right) + B \left( 1 - \frac{\bar{B}}{B} \right) \\ & = 0. \end{aligned}$$

### 1.3

$$\begin{aligned} \frac{dG}{dt} &= \frac{dA}{dt}(1 + \ln A) + A \cdot \frac{1}{A} \frac{dA}{dt} + \frac{dB}{dt}(1 + \ln B) + B \cdot \frac{1}{B} \frac{dB}{dt} \\ &= (-A + B)(1 + \ln A - 1 - \ln B) \\ &\leq 0. \end{aligned}$$

最后一步分  $A > B$  与  $A < B$  两种情况讨论即可。

## 2 第二题

### 2.1

Conservation law:

$$u_t + \left( \frac{u^2}{2} - \mu u_x \right)_x = 0, \quad x \in \mathbb{R}, \quad t > 0.$$

Flux function:  $\frac{u^2}{2} - \mu u_x$ .

### 2.2

Hopf-Cole 变换:

$$u = -a \frac{\partial}{\partial x} \ln \phi = -a \frac{\phi_x}{\phi}.$$

将 Hopf-Cole 变换带入到原方程, 并利用  $a = 2\mu$ ,

$$-2\mu \frac{\phi \phi_{xt} - \phi_x \phi_t}{\phi^2} + 4\mu^2 \frac{\phi \phi_x \phi_{xx} - \phi_x^3}{\phi^3} = -2\mu^2 \frac{\phi^2 (\phi_x \phi_{xx} + \phi \phi_{xxx} - 2\phi_x \phi_{xx}) - 2\phi \phi_x (\phi \phi_{xx} - \phi_x^2)}{\phi^4},$$

化简可得

$$-2\mu \frac{\phi \phi_{xt} - \phi_x \phi_t}{\phi^2} = -2\mu^2 \frac{\phi \phi_{xxx} - \phi_x \phi_{xx}}{\phi^2},$$

即

$$\left( \frac{\phi_t}{\phi} - \mu \frac{\phi_{xx}}{\phi} \right)_x = 0.$$

因此

$$\frac{1}{\phi} \phi_t - \mu \frac{1}{\phi} \phi_{xx} = g(t)$$

是一个只关于  $t$  的函数。

## 3 第三题

### 3.1

$$\frac{df}{dt} = \frac{\partial f}{\partial t} + \nabla_{\mathbf{q}} f \cdot \frac{d\mathbf{q}}{dt} + \nabla_{\mathbf{p}} f \cdot \frac{d\mathbf{p}}{dt}.$$

根据守恒律, 我们有

$$\frac{\partial f}{\partial t} + \nabla_{\mathbf{q}} \cdot (f \dot{\mathbf{q}}) + \nabla_{\mathbf{p}} \cdot (f \dot{\mathbf{p}}) = 0. \quad (\dot{\mathbf{p}} \text{ 表示 } \frac{d\mathbf{p}}{dt}).$$

将上式展开, 我们有

$$\frac{\partial f}{\partial t} + \nabla_{\mathbf{q}} f \cdot \dot{\mathbf{q}} + \nabla_{\mathbf{p}} f \cdot \dot{\mathbf{p}} + f (\nabla_{\mathbf{q}} \cdot \dot{\mathbf{q}} + \nabla_{\mathbf{p}} \cdot \dot{\mathbf{p}}) = 0.$$

又由

$$\dot{\mathbf{p}} = -\nabla_{\mathbf{q}} H, \quad \dot{\mathbf{q}} = \nabla_{\mathbf{p}} H,$$

因此

$$\nabla_{\mathbf{q}} \cdot \dot{\mathbf{q}} + \nabla_{\mathbf{p}} \cdot \dot{\mathbf{p}} = \nabla_{\mathbf{q}} \cdot \nabla_{\mathbf{p}} H - \nabla_{\mathbf{p}} \cdot \nabla_{\mathbf{q}} H = \sum_i \left( \frac{\partial^2 H}{\partial q_i \partial p_i} - \frac{\partial^2 H}{\partial p_i \partial q_i} \right) = 0.$$

因此我们有

$$\frac{df}{dt} = \frac{\partial f}{\partial t} + \nabla_{\mathbf{q}} f \cdot \dot{\mathbf{q}} + \nabla_{\mathbf{p}} f \cdot \dot{\mathbf{p}} = 0.$$

### 3.2

注意到

$$\dot{\mathbf{q}} = \frac{\mathbf{p}}{m}, \quad \dot{\mathbf{p}} = -\nabla_{\mathbf{q}} V(\mathbf{q}),$$

记  $H = \frac{|\mathbf{p}|^2}{2m} + V(\mathbf{q})$ , 从而

$$\frac{dH}{dt} = \frac{\mathbf{p} \cdot \dot{\mathbf{p}}}{m} + \frac{\partial V(\mathbf{q})}{\partial t} = \frac{\mathbf{p} \cdot \dot{\mathbf{p}}}{m} + \nabla_{\mathbf{q}} V(\mathbf{q}) \cdot \dot{\mathbf{q}} = 0.$$

$$\nabla_{\mathbf{p}} H = \frac{\mathbf{p}}{m}, \quad \nabla_{\mathbf{q}} H = \nabla_{\mathbf{q}} V(\mathbf{q}).$$

对于任意区域  $\Omega$ ,

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} H f(t, \mathbf{q}, \mathbf{p}) \, d\mathbf{q} \, d\mathbf{p} \\ &= \int_{\Omega} H \frac{\partial f}{\partial t} \, d\mathbf{q} \, d\mathbf{p} \\ &= - \int_{\Omega} H \left( \nabla_{\mathbf{q}} f \cdot \frac{\mathbf{p}}{m} - \nabla_{\mathbf{p}} f \cdot \nabla_{\mathbf{q}} V(\mathbf{q}) \right) \, d\mathbf{q} \, d\mathbf{p} \\ &= - \int_{\Omega} H (\nabla_{\mathbf{q}} f \cdot \nabla_{\mathbf{p}} H - \nabla_{\mathbf{p}} f \cdot \nabla_{\mathbf{q}} H) \, d\mathbf{q} \, d\mathbf{p} \\ &= \int_{\Omega} \nabla \cdot (f \nabla_{\mathbf{q}} (\frac{1}{2} H^2), -f \nabla_{\mathbf{p}} (\frac{1}{2} H^2)) \, d\mathbf{q} \, d\mathbf{p}, \quad (\text{此处 } \nabla \cdot (\alpha, \beta) = \nabla_{\mathbf{p}} \cdot \alpha + \nabla_{\mathbf{q}} \cdot \beta) \\ &= \int_{\partial \Omega} (f \nabla_{\mathbf{q}} (\frac{1}{2} H^2), -f \nabla_{\mathbf{p}} (\frac{1}{2} H^2)) \cdot \mathbf{n} \, dS. \end{aligned}$$

当  $\Omega$  包含了  $f$  的支集时, 在  $\partial \Omega$  上  $f$  恒为 0. 根据上式我们可得

$$\frac{d}{dt} \int H f(t, \mathbf{q}, \mathbf{p}) \, d\mathbf{q} \, d\mathbf{p} = 0.$$

## 4 第四题

## 4.1

$$\begin{aligned}
\frac{d}{dt} \int |u|^2 dx &= \int \frac{\partial}{\partial t} |u|^2 dx \\
&= \int (u_t \bar{u} + u \bar{u}_t) dx \\
&= \int \left( \frac{\hat{H}}{i\hbar} u \bar{u} - u \frac{\hat{H}}{i\hbar} \bar{u} \right) dx
\end{aligned}$$

注意到  $\hat{H} = -\frac{\hbar^2}{2m} \Delta + V(x)$ ,

$$\begin{aligned}
\text{上式} &= \frac{1}{i\hbar} \int (\bar{u} \hat{H} u - u \hat{H} \bar{u}) dx \\
&= -\frac{\hbar}{2mi} \int (\bar{u} \Delta u - u \Delta \bar{u}) dx \\
&= -\frac{\hbar}{2mi} (\bar{u} \nabla u - u \nabla \bar{u})|_{-\infty}^{\infty} + \frac{\hbar}{2mi} \int (\nabla \bar{u} \nabla u - \nabla u \nabla \bar{u}) dx \\
&= 0.
\end{aligned}$$

## 4.2

由上题我们可知

$$\begin{aligned}
\rho_t &= u_t \bar{u} + u \bar{u}_t \\
&= \frac{1}{i\hbar} (\bar{u} \hat{H} u - u \hat{H} \bar{u}) \\
&= -\frac{\hbar}{2mi} (\bar{u} \Delta u - u \Delta \bar{u}) \\
&= -\frac{\hbar}{2mi} \nabla (\bar{u} \nabla u - u \nabla \bar{u})
\end{aligned}$$

因此

$$J = \frac{\hbar}{2mi} (\bar{u} \nabla u - u \nabla \bar{u}).$$