

# 数学模型第三次作业

2019 年 4 月 6 日

## 1

### 1.1

考虑方程

$$\rho_t + \rho_x + 6\rho\rho_x - \rho_{xxt} = 0, \quad (1)$$

令

$$\rho(x, t) = P(x - ct) = P(s), \quad s = x - ct,$$

有

$$\rho_t = -cP', \quad \rho_x = P',$$

代入 (1),

$$(6P + 1 - c)P' + cP''' = 0. \quad (2)$$

### 1.2

$$\begin{aligned} P(s) &= A \operatorname{sech}^2(Bs), \\ P'(s) &= -2AB \tanh(Bs) \operatorname{sech}^2(Bs), \\ P''(s) &= 6AB^2 \tanh^2(Bs) \operatorname{sech}^2(Bs) - 2AB^2 \operatorname{sech}^2(Bs), \\ P'''(s) &= 16AB^3 \tanh(Bs) \operatorname{sech}^2(Bs) - 24AB^3 \tanh^3(Bs) \operatorname{sech}^2(Bs). \end{aligned} \quad (3)$$

将 (3) 代入 (2), 并利用  $\cosh^2(x) - \sinh^2(x) = 1$  可知

$$\begin{cases} (1 + c) + 4cB^2 = 0, \\ 6A + 1 - 8cB^2 - c = 0, \end{cases}$$

解得

$$\begin{cases} A = \frac{1}{2}(c-1), \\ B^2 = \frac{c-1}{4c}. \end{cases}$$

## 2

### 2.1

考虑方程

$$\frac{\partial \rho}{\partial t} + e^{2t} \frac{\partial \rho}{\partial x} = \rho + x + t, \quad \rho(x, t=0) = \cos x,$$

设  $X = X(t)$ ,  $P(t) = \rho(X(t), t)$ , 满足

$$\begin{cases} \frac{dX}{dt} = e^{2t}, \\ X(0) = X_0, \end{cases} \quad \begin{cases} \frac{dP}{dt} = P + X + t, \\ P(0) = \cos(X_0), \end{cases}$$

求解常微分方程, 有

$$X = \frac{1}{2}(e^{2t} - 1) + X_0. \quad (4)$$

而

$$\begin{aligned} P &= \cos(X_0)e^t + e^t \int_0^t \left( \frac{1}{2}(e^{2t} - 1) + X_0 + t \right) e^{-t} dt \\ &= (\cos(X_0) + X_0) e^t - \left( X_0 + t + \frac{1}{2} \right) + X_0 + \frac{1}{2} e^{2t}, \end{aligned} \quad (5)$$

将 (4) 代入 (5) 有,

$$\rho(x, t) = \left( \cos\left(x - \frac{1}{2}(e^{2t} - 1)\right) + x - \frac{1}{2}(e^{2t} - 1) \right) e^t - (x + t + 1) + e^{2t}.$$

### 2.2

考虑方程

$$\frac{\partial \rho}{\partial t} + (x+4) \frac{\partial \rho}{\partial x} = -2\rho, \quad \rho(x=0, t) = \cos t, \quad \rho(x, t=0) = e^{-x}, \quad x \geq 0,$$

设  $X = X(t)$ ,  $P(t) = \rho(X(t), t)$ , 则

$$\begin{cases} \frac{dX}{dt} = X + 4, \\ X(0) = X_0 \text{ or } X(t_0) = 0, \end{cases}$$

故 1) 当  $x > 4e^t - 4$  时,

$$X = (4 + X_0)e^t - 4. \quad (6)$$

$P(t)$  满足

$$\begin{cases} \frac{dP}{dt} = -2P, \\ P(0) = e^{-X_0}, \end{cases}$$

解得

$$P = e^{-2t - X_0},$$

将 (6) 代入可得,

$$\rho(x, t) = e^{-(x+4)e^{-t} + 4 - 2t}, \quad x > 4e^t - 4.$$

2) 当  $x < 4e^t - 4$  时,

$$X = 4e^{t-t_0} - 4. \quad (7)$$

$P(t)$  满足

$$\begin{cases} \frac{dP}{dt} = -2P, \\ P(t_0) = \cos t_0, \end{cases}$$

解得

$$P = \cos t_0 e^{-2(t-t_0)},$$

将 (7) 代入可得,

$$\rho(x, t) = \cos \left( t - \ln \left( \frac{1}{4}x + 1 \right) \right) e^{-2 \ln(\frac{1}{4}x + 1)}.$$

### 3

#### 3.1

$$\begin{aligned} \frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{j} &= 0, \\ \frac{\partial \mathbf{j}}{\partial t} + \nabla \cdot \left( \frac{1}{\rho} \mathbf{j} \otimes \mathbf{j} + p \mathbf{I} \right) &= 0, \end{aligned} \quad (8)$$

而

$$\frac{\partial \mathbf{j}}{\partial t} = \frac{\partial \rho}{\partial t} \mathbf{v} + \rho \frac{\partial \mathbf{v}}{\partial t} \quad (9)$$

将 (9) 代入 (8) 并利用整理  $\nabla \cdot \mathbf{v} = 0$  可得

$$\rho \frac{\partial \mathbf{v}}{\partial t} + \rho \mathbf{v} \cdot \nabla \mathbf{v} = -\nabla p,$$

即

$$\frac{\partial \mathbf{v}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{v} = -\frac{\nabla p}{\rho}.$$

### 3.2

设  $f(x) = u(x, 0)$ ,  $X = X(t)$ ,  $U(t) = u(X(t), t)$ , 则

$$\begin{cases} \frac{dX}{dt} = U, \\ X(0) = X_0, \end{cases} \quad \begin{cases} \frac{dU}{dt} = 0, \\ U(0) = f(X_0), \end{cases}$$

解得

$$U(t, X_0) = f(X_0),$$

$$X = f(X_0)t + X_0.$$

将  $f(x)$  的具体表达式代入有, 当  $t < 1$  时

$$\begin{cases} X = X_0, & U = 0, & X_0 > 0, \\ X = -X_0 t + X_0, & U = -X_0, & -1 < X_0 < 0, \\ X = t + X_0, & U = 1, & X_0 < -1, \end{cases}$$

即

$$u(x, t) = \begin{cases} 0, & x > 0, \\ \frac{x}{t-1}, & t-1 < x < 0, \\ 1, & x < t-1. \end{cases}$$

当  $t = 1$  时,

$$u(x, 1) = \begin{cases} 0, & x > 0, \\ 1, & x < 0. \end{cases}$$

## 4

**证明:** 当  $x < 0$  时, 我们有  $\rho > \theta$ , 所以方程约化为

$$cv' + v'' + \mu(1 - v) = 0.$$

由于  $v(-\infty) = 1$  (蕴含有界性), 我们得到下面的解

$$v(x) = 1 - (1 - \theta)e^{\lambda_+ x}, \quad x < 0,$$

其中

$$\lambda_+ = \frac{1}{2} \left( -c + \sqrt{c^2 + 4\mu} \right).$$

而当  $x > 0$  时, 我们有  $\rho < \theta$ , 所以方程约化为

$$cv' + v'' = 0.$$

由于  $v(+\infty) = 0$ , 我们得到下面的解

$$v = \theta e^{-cx}, \quad x \geq 0.$$

注意到, 由于  $f$  在  $\theta$  处不连续, 所以  $v''$  在 0 处有一个间断跳跃 (这蕴含了  $v'$  在 0 处连续), 所以我们得到

$$(1 - \theta)\lambda_+ = \theta c.$$

整理可得,  $-c(\theta + 1) + (1 - \theta)\sqrt{c^2 + 4\mu} = 0$ , 进而得到  $c > 0$  且

$$c^2(\theta + 1)^2 = (1 - \theta)^2(c^2 + 4\mu),$$

解得  $c_1 = (1 - \theta)\sqrt{\mu/\theta} > 0$ ,  $c_2 = -(1 - \theta)\sqrt{\mu/\theta} < 0$  (舍), 故满足条件的波速  $c^* > 0$  可以被唯一确定. 对  $v$  求导可得

$$v' = \begin{cases} -\lambda_+(1 - \theta)e^{\lambda_+ x}, & x < 0, \\ -c\theta e^{-cx}, & x > 0, \end{cases}$$

故  $v(x)$  是一个单调递减函数.

## 5

### 5.1

$$\begin{aligned}\int_0^1 \frac{\delta J}{\delta y}(x)h(x) \, dx &= \left[ \frac{d}{d\epsilon} J[y + \epsilon h] \right]_{\epsilon=0} \\&= \left[ \frac{d}{d\epsilon} \int_0^1 \frac{1}{2} (y' + \epsilon h')^2 + \frac{k}{x} (y + \epsilon h)(y' + \epsilon h') + x^2 (y + \epsilon h) \, dx \right]_{\epsilon=0} \\&= \int_0^1 y' h' + \frac{k}{x} h y' + \frac{k}{x} h' y + x^2 h \, dx \\&= \int_0^1 \left( -y'' + \frac{k}{x^2} y + x^2 \right) h \, dx,\end{aligned}$$

(推导过程中用到了分部积分) 故

$$\frac{\delta J}{\delta y_*} = -y''_* + \frac{k}{x^2} y_* + x^2 = 0.$$

### 5.2

记

$$L = \frac{1}{2} (y')^2 + \frac{k}{x} y y' + x^2 y,$$

由 Euler-Lagrangian 方程,  $y_*$  满足

$$\frac{\delta J}{\delta y} = \frac{\partial L}{\partial y} - \frac{\partial}{\partial x} \left( \frac{\partial L}{\partial y'} \right) = -y'' + \frac{k}{x^2} y + x^2 = 0.$$

### 5.3

由  $k = 0$  得  $y'' = x^2$ , 解得

$$y = \frac{1}{12} x^4 + C_1 x + C_2,$$

由边界条件  $y(0) = 1$ ,  $y(1) = 1$ , 得  $C_1 = -\frac{1}{12}$ ,  $C_2 = 1$ .

### 5.4

考虑  $-y'' + \frac{2}{x^2} y = 0$ , 寻找形如  $y = x^\alpha$  的特解. 将  $y = x^\alpha$  代入可得  $\alpha(\alpha - 1) - 2 = 0$ , 即  $\alpha = 2, -1$ , 故

$$y = C_1 x^2 + C_2 x^{-1}$$

是  $-y'' + \frac{2}{x^2}y = 0$  的通解. 而  $y = \frac{1}{10}x^4$  是  $x^2 - y'' + \frac{2}{x^2}y = 0$  的特解, 故

$$y = \frac{1}{10}x^4 + C_1x^2 + C_2x^{-1}$$

是  $x^2 - y'' + \frac{2}{x^2}y = 0$  的通解. 由边界条件  $y'(0) = 0$ ,  $y(1) = 1$ , 得  $C_1 = \frac{9}{10}$ ,  $C_2 = 0$ , 不满足  $y(0) = 1$ .

## 6

### 6.1

$$\delta^2 J(h, y) = \int_0^1 \left( \frac{h'^2}{2[1 + (y')^2]^{3/2}} \right) dx.$$

### 6.2

$$\delta^2 J = \int_0^1 \left( \frac{h'^2}{2[1 + (y')^2]^{3/2}} \right) dx \geq 0.$$

故  $\delta^2 J = 0$  当且仅当  $h' = 0$ , 由边界条件  $h(0) = h(1) = 0$  知  $\delta^2 J = 0$  当且仅当  $h = 0$ .