

# Note of Group Theory

Kaiser

University of South China

2025 年 1 月 12 日

## 摘要

这一篇笔记是本科阶段学习群论的笔记。

## 参考文献

- [1] Joseph J. Rotman, An Introduction to the Theory of Groups
- [2] 周彬, Lie 群与 Lie 代数
- [3] Pierre Ramond, Group Theory : A Physicist's Survey
- [4] 崔建伟, 群论讲义
- [5] A. Das and S. Okubo, Lie Groups and Lie Algebras for Physicists

## 目录

<b>1</b>	<b>有限群 (Finite Group)</b>	<b>2</b>
1.1	群与群乘法 . . . . .	2
1.2	群同态 (Group Homomorphisms) 与群的直乘 . . . . .	3
1.3	群的线性表示理论 . . . . .	3
<b>2</b>	<b>表示论</b>	<b>3</b>
<b>3</b>	<b>李群 (Lie Group)</b>	<b>3</b>
<b>4</b>	<b>李代数 (Lie Algebra)</b>	<b>3</b>



# 1 有限群 (Finite Group)

## 1.1 群与群乘法

废话也就不多说，我们直接给出群的定义，并在之后讲解一下为什么要这样去定义

**定义 1** (Definition of a group).

Suppose we give a set  $\mathbb{G} = \{g_0, g_1, g_2, \dots\}$  and an operator  $\cdot$ , if they satisfy the following properties

1. **Closure** For every ordered pair of elements,  $g_i$  and  $g_j$ , there exists a unique element

$$a_i \cdot a_j = a_k \quad (1)$$

for any three  $i, j, k$ .

2. **Associativity** The  $\cdot$  operation is associative

$$g_i \cdot (g_j \cdot g_k) = (g_i \cdot g_j) \cdot g_k \quad (2)$$

3. **Unit element** The set  $G$  contains a unique element  $e$  (here we will use  $g_0$  to symbol it) such that

$$g_0 \cdot g_i = g_i \cdot g_0 = g_i \quad (3)$$

for all  $i$ . In particular, this means that

$$g_0 \cdot g_0 = g_0.$$

4. **Inverse element** Corresponding to every element  $g_i$ , there exists a unique element of  $G$ , the inverse  $(g_i)^{-1}$  such that

$$g_i \cdot (g_i)^{-1} = (g_i)^{-1} \cdot g_i = g_0 \quad (4)$$

In particular, this means that

$$g_0 = (g_0)^{-1}$$

then we called the set  $\mathbb{G}$  a Group.

同时，我们根据集合  $\mathbb{G}$  的元素个数



## 1.2 群同态 (Group Homomorphisms) 与群的直乘

## 1.3 群的线性表示理论

# 2 表示论

## 3 李群 (Lie Group)

## 4 李代数 (Lie Algebra)