TencorFlow

TensorFlow API r1.4

tf.contrib.bayesflow.entropy.renyi_ratio

```
renyi_ratio(
   log_p,
   q,
   alpha,
   z=None,
   n=None,
   seed=None,
   name='renyi_ratio'
)
```

Defined in tensorflow/contrib/bayesflow/python/ops/entropy_impl.py.

See the guide: BayesFlow Entropy (contrib) > Ops

Monte Carlo estimate of the ratio appearing in Renyi divergence.

This can be used to compute the Renyi (alpha) divergence, or a log evidence approximation based on Renyi divergence.

Definition

With z_i iid samples from q, and $exp\{log_p(z)\} = p(z)$, this p returns the (biased for finite p) estimate:

```
 (1 - alpha)^{-1} \ Log[ \ n^{-1} \ sum_{i=1}^n \ ( \ p(z_i) \ / \ q(z_i) \ )^{1} - alpha}, \\ \ approx \ (1 - alpha)^{-1} \ Log[ \ E_q[ \ (p(Z) \ / \ q(Z))^{1} - alpha} \ ] \ ]
```

This ratio appears in different contexts:

Renyi divergence

If $log_p(z) = Log[p(z)]$ is the log prob of a distribution, and alpha > 0, alpha != 1, this Op approximates -1 times Renyi divergence:

The Renyi (or "alpha") divergence is non-negative and equal to zero iff $\mathbf{q} = \mathbf{p}$. Various limits of **alpha** lead to different special case results:

See "Renyi Divergence Variational Inference", by Li and Turner.

Log evidence approximation

If $log_p(z) = log[p(z, x)]$ is the log of the joint distribution p, this is an alternative to the ELBO common in variational inference.

```
L_alpha(q, p) = Log[p(x)] - D_alpha[q || p]
```

If \mathbf{q} and \mathbf{p} have the same support, and $\mathbf{0} < \mathbf{a} <= \mathbf{b} < \mathbf{1}$, one can show ELBO $<= \mathbf{D}_{-}\mathbf{b} <= \mathbf{D}_{-}\mathbf{a} <= \mathbf{Log}[\mathbf{p}(\mathbf{x})]$. Thus, this $\mathbf{0p}$ allows a smooth interpolation between the ELBO and the true evidence.

Stability notes

Note that when 1 - alpha is not small, the ratio (p(z) / q(z))^{1 - alpha} is subject to underflow/overflow issues. For that reason, it is evaluated in log-space after centering. Nonetheless, infinite/NaN results may occur. For that reason, one may wish to shrink alpha gradually. See the **Op renyi_alpha**. Using **float64** will also help.

Bias for finite sample size

Due to nonlinearity of the logarithm, for random variables $\{X_1, ..., X_n\}$, $E[Log[sum_{i=1}^n X_i]] != Log[E[sum_{i=1}^n X_i]] != Log[E[sum_{i=1}^n X_i]]$. As a result, this estimate is biased for finite n. For alpha < 1, it is non-decreasing with n (in expectation). For example, if n = 1, this estimator yields the same result as $elbo_ratio$, and as n increases the expected value of the estimator increases.

Call signature

User supplies either **Tensor** of samples z, or number of samples to draw n

Args:

- log_p : Callable mapping samples from **q** to **Tensors** with shape broadcastable to **q.batch_shape** . For example, log_p works "just like" **q.log_prob** .
- q: tf.contrib.distributions.Distribution float64 dtype recommended. log_p and q should be supported on the same set.
- alpha: Tensor with shape q.batch_shape and values not equal to 1.
- z: Tensor of samples from q, produced by q.sample for some n.
- n: Integer Tensor. The number of samples to use if z is not provided. Note that this can be highly biased for small n, see docstring.
- seed: Python integer to seed the random number generator.
- name: A name to give this Op.

Returns:

renyi_result: The scaled log of sample mean. Tensor with shape equal to batch shape of q, and dtype = q.dtype.

Except as otherwise noted, the content of this page is licensed under the Creative Commons Attribution 3.0 License, and code samples are licensed under the Apache 2.0 License. For details, see our Site Policies. Java is a registered trademark of Oracle and/or its affiliates.

Last updated November 2, 2017.

Stay Connected	
Blog	
GitHub	
Twitter	
Support	
Issue Tracker	
Release Notes	
Stack Overflow	
English	
Terms Privacy	