

# Analysis on Period vs. Angle, Period vs. Length & Q Factor of The Pendulum

Zixuan Fan (fanzixu2)

09/18/2023

## 1 Introduction

In this lab report, we analyse the how the angle at which a pendulum is released, as well as the length that the pendulum is released, correlates with the period of the pendulum. Additionally, by computing the Q Factor [2], we will analyse the damping effect on the pendulum's period as the angle decreases. Finally, we will assess the how the damping effect correlates with the pendulum's length by calculating the Q Factor at each length. For this lab report, we will assume the rod of our pendulum stand is rigid, massless, and perfectly orthogonal to the pendulum stand; and the string of our pendulum is massless, always stretched, and has negligible air resistance. Our experimental results indicate that the pendulum's period decreases exponentially with their release angle, as induced by errors regarding friction of the string and air resistance of the pendulum. Moreover, the observed exponential decay of the pendulum's amplitude over time corresponds to a Q Factor of  $200 \pm 1$  derived through fitting function to graph, which aligns with the Q Factor of  $196 \pm 5$  derived from counting oscillation. Furthermore, the correlation between the pendulum's length and its period is evident, with longer lengths leading to increased periods. Finally, through calculating Q factors at each length, we deduced that the damping effect is more pronounced in longer pendulum lengths.

Utilizing an experimental set-up that minimizes asymmetry and maintains the motion of the pendulum in a single plane, we will gather data points regarding the period of pendulum with regards to both angle and length, as well as graphing and analysing these data points in data analysis. The equations we used in data analysis are listed below [2]:

$$\theta(t) = \theta_0 e^{-t/\tau} \cos(2\pi \frac{t}{T} + \phi_0) \quad (1)$$

In equation (1),  $\theta$  is the angle of the pendulum in rad,  $\theta_0$  is the release angle of the pendulum in rad,  $t$  is the time in seconds,  $\tau$  is in seconds,  $T$  is the maximum period (the period in the first cycle) of pendulum in seconds, and  $\phi_0$  is a phase shift constant.

$$Q = \pi \frac{\tau}{T} = \pi \quad (2)$$

In addition to the same terms in equation (1), in equation (2),  $Q$  is the Q factor, a constant.

$$\theta(t) = \theta_0 e^{-t/\tau} \quad (3)$$

Using to the same terms in equation (1), equation (3) is a deviation of equation (1)

## 2 Experimental Set-up

The experimental set-up consists of suspending a golf ball from a horizontal wooden beam using double wool strings with adjustable length, as demonstrated in **Figure 1**.




Figure 1: Experimental Set-up of The Pendulum

The string of pendulum has a vertical length of  $37.5 \pm 0.1\text{cm}$ , and the golf ball along with the tape has a combined weight of  $45.0 \pm 0.1\text{g}$ . Wool strings were chosen as their masses are light enough to be ignored. Additionally, the video capture device has a frame rate of 30 frames per second.

### 2.1 Improvements for The Set-up that Eliminate Asymmetry

To maintain symmetry during experimentation, data was collected from both the negative (left) and positive (right) sides of the pendulum release. This approach minimizes errors stemming from the inherent asymmetry in the set-up.

Efforts were also made to reduce this inherent asymmetry by adjusting the knot on the pendulum string to the lowest point on the wooden stand, thereby aligning the string with the symmetrical center line.

Moreover, the golf ball is tied up using two strings to ensure that the motion of the ball stays in the same plane, which prevents asymmetry caused by circular motion of the ball, had the ball been hung from only one string from the wooden beam.

## 3 Data Analysis

### 3.1 Justification for Measurements of Uncertainty

The most significant uncertainty within the experiment arises from the measurement of the angle, the length, and the period of the pendulum. Additionally, the friction between the string and the stand, and the influence of air resistance during pendulum swings could have contributed to an increased level of uncertainty. However, these two errors lead to a systematic shift in the data points. Unlike the random error in measurement, which causes fluctuations in individual data points, these systematic errors are far less prominent at increasing the uncertainty of individual data points. A further analysis of these two systematic errors will be provided in section 3.2.1.

The measurements for angles are done using an angle-meter with a step of  $\pm 1$  degree, equivalent to  $\pm 1 * \pi/180 = \pm 0.02$  in rad, as observable in **Figure 1**. Since the angle-meter is an analogue instrument, the uncertainty of Angle should be  $\pm 0.02/2 = \pm 0.01$  rad.

The measurements for periods and time are done using a digital tracker with a step of  $\frac{1.00\text{s}}{30\text{frames}} = 0.03\text{s}$  [1]. Since the tracker is a digital instrument, the uncertainty of period should be  $\pm 0.03$  second.

The measurement for lengths of pendulum are done using a ruler with a step of  $\pm 0.2$  cm. Since the ruler is an analogue instrument, the uncertainty of lengths should be  $\pm 0.2/2 = \pm 0.1$  cm.

In future, all of these uncertainties caused by measurement can be reduced by selecting measuring instruments with smaller steps.

## 3.2 Relationship between Angle and Period of Pendulum

### 3.2.1 Graphing and Analysing Figure 2

The pendulum was released at  $1.57 \pm 0.01$  rad from both side of the stand. Importing gathered data into a provided python program, we were able to graph the period of pendulum against angles in **Figure 2**:




Figure 2: *Period of Pendulum (s) vs. Angle (rad)*

Note that the horizontal error bar has a length of  $\pm 0.01$  rad, and the vertical error bar has a length of  $\pm 0.03$  second.

It is observable from **Figure 2**, that the maximum period  $T$  decreases exponentially as the angle gets closer to 0 rad. However, this decrease occurs within an interval approximately 0.20, which is a negligible difference, attributable to two systematic errors: One being the reducing friction between string and stand as angle gets smaller, and the other one being the decreasing air resistance during pendulum swings as angle gets smaller. Both of these errors decreases the period of the pendulum as the angle gets smaller.

### 3.2.2 Graphing and Analysing Figure 3 and Figure 4

Upon tracking the decay of pendulum's amplitude starting from  $1.31 \pm 0.01$  radian, a relationship between the angle of pendulum and time is demonstrated in **Figure 3**.

**Figure 3** is crucial in determining the Q Factor by counting the number of oscillations that occurred in this graph.

It is observable from **Figure 3** that the relationship between the Angle of Pendulum over time is an oscillating cosine function. The maximum amplitude of the pendulum decreases exponentially as time increases. Therefore, taking the angle at all the local maximum amplitude over the entire time interval, along with their respective time, we can construct a new graph of the angle at local maximum amplitude over time in **Figure 4**:




Figure 3: *Angle of Pendulum (Rad) vs. Time (s)*




Figure 4: *Angle of Pendulum at Local Maximum Amplitude (Rad) vs. Time (s)*

Note that the for both **Figure 3** and **Figure 4**, the vertical error bar has a length of  $\pm 0.01$  rad, and the horizontal error bar has a length of  $\pm 0.03$  second. However, these error bars are hardly observable, attributable to large range at both the vertical and horizontal axis, as well as the large number of concentrated data points

**Figure 4** is crucial in determining the factor  $\tau$ , which is used to calculate Q Factor through fitting the function to graph in section 3.3.1.

### 3.3 Q Factor Calculation of section 3.2

#### 3.3.1 Through Fitting the Function to Graph

Referencing the gathered data from section 3.2, we can find the release angle of the pendulum:

$$\theta = 1.31 \pm 0.01 \text{ rad}$$

as well as the maximum period of pendulum:

$$T = 1.29 \pm 0.03 \text{ s}$$

Substituting in the values to equation , we will fit equation (3) the graph in **Figure 4**, and find the  $\tau$  factor:

$$\theta = (-1.31 \pm 0.01 \text{ rad}) * e^{t/\tau} \quad (4)$$

$$\therefore \tau \simeq 77.58 \pm 0.03 \text{ s}$$

Further, we calculate Q factor through equation (2):

$$\therefore Q = \pi \frac{\tau}{T} = \pi \frac{77.58 \text{ s} \pm 0.03 \text{ s}}{1.29 \text{ s} \pm 0.03 \text{ s}} \simeq 200 \pm 1$$

Therefore, the Q Factor for the pendulum is  $200 \pm 1$ ,

### 3.3.2 Through Counting Oscillations

Through counting oscillation until the 4 % of the original angle  $1.31 \pm 0.01$ :

$$\theta_0 * 4\% = 1.31\text{rad} * 4\% = 0.05 \pm 0.01\text{rad}$$

I was able to count  $196 \pm 5$  oscillations, therefore, the Q factor from counting oscillations is  $196 \pm 5$ . Uncertainty is  $\pm 5$  due to human error on determining the exact point where angle equals  $0.05 \pm 0.01\text{rad}$

### 3.3.3 Comparing Q Factor from both methods

The Q Factor through counting oscillation is very close with the Q Factor through Estimating Periods, with a difference of only  $\Delta = 200 - 196 = 4$ . This difference is covered within the uncertainty from both Q Factor calculations, as  $\pm 5 + \pm 1 = \pm 6 \geq \Delta$

## 3.4 Relationship between Period vs. Length of Pendulum

### 3.4.1 Graphing and Analysing Figure 5

Through releasing the pendulum from  $1.05 \pm 0.01$  rad, separately at a length of  $30.0 \pm 0.1\text{cm}$ ,  $25.0 \pm 0.1\text{cm}$ ,  $20.0 \pm 0.1\text{cm}$ ,  $15.0 \pm 0.1\text{cm}$ , and  $10.0 \pm 0.1\text{cm}$  from both side of the wooden stand; and recording the period inside the tracker application [1], we were able to gather data and graph the relationship between the period and the corresponding length of the pendulum in **Figure 5**:




Figure 5: *Period of Pendulum (s) vs. Length (cm)*

Note that the horizontal error bar has a length of  $\pm 0.1$  cm, and the vertical error bar has a length of  $\pm 0.03$  second.

It is therefore observable that the maximum period  $T$  of pendulum increases exponentially as length of pendulum increases.

### 3.4.2 Calculating and graphing Q Factor vs. Length

For each of the different length of pendulum, we will calculate the Q Factor through fitting the provided equation (3) - with values of angle and time found in section 3.4.1 - to the graph of Periods vs.

Angles until the tenth cycle. We are using the fitting to graph method, as this method will produce results of Q factor with lower uncertainty ( $\pm 1$ , as shown in section 3.3.1) than that from counting oscillations ( $\pm 5$ , as shown in section 3.3.2)

Note that since both Period (T) and Time (t) has an uncertainty of  $\pm 0.03$  seconds, the uncertainty of  $\tau$  should be  $\pm 0.03$ , and therefore the uncertainty of Q Factor should be  $\pm 1$ . This calculation has been demonstrated in section 3.3.1

Using the calculated data of Q Factors, we are able to graph a relationship between the Q Factor and the length of the pendulum in **Figure 6**:




Figure 6: *Q Factor vs. Length (cm)*

Note that the horizontal error bar has a length of  $\pm 0.1$  cm, and the vertical error bar has a length of  $\pm 1$ .

It is therefore observable that Q Factor increases exponentially as length of pendulum increases.

## 4 Conclusion

It is therefore deduced through this lab report, that:

1. The Maximum Period (the period in the first cycle) of pendulum is dependent of the angle, as demonstrated by **Figure 2** in section 3.2.1. However, this dependency is negligible, as there are two specified error in section 3.2.1 which could have caused the decrease of period over the decrease of angle.
2. The amplitude is observed to decay exponentially as time increases, as demonstrated by **Figure 3** in section 3.2.2. Upon calculation in section 3.3, the two Q Factors measurements, one with an uncertainty of  $\pm 1$  and one with an uncertainty of  $\pm 5$ , does agree with each other.
3. The Maximum Period (The period in the first cycle) of pendulum is dependent on its length, as demonstrated by **Figure 5** in section 3.4.1. As length of pendulum increases, the period of pendulum increases exponentially.
4. The Q Factor of pendulum is dependent on its length, as demonstrated by **Figure 6** in section 3.4.2. As the length of pendulum increases, the Q Factor of pendulum decreases exponentially.

with an uncertainty of  $\pm 1$ . As a higher Q Factor indicates a lower rate of energy loss and the oscillations die out more slowly [2], this implies that for longer lengths of pendulums, the damping effect will be more significant, as the rate of energy loss will be higher, and the oscillation will die out more quickly.

## 5 Reference List

- [1] Tracker. Tracker Video Analysis and Modeling Tool for Physics Education. (n.d.). <https://physlets.org/tracker/>
- [2] Lepeshkin, N. (n.d.). Q factor . University of Massachusetts Lowell. <https://faculty.uml.edu/pchowdhury/PHYS2690/Supp/Q-factor.pdf>