The Commutator of Representation

The electron, dealt from the throw and never from the lift

John Henry

October 2026

The fifth card. It is the one the dealer talks over, and it is never turned face up.

“Does the mechanism demonstrating this proof look like an electron to you?”

Reading note

This card is the patter: what the dealer says while the cards move.

The object. A card is carried, transformed, copied, projected and read again. At every move some of it comes through and some of it does not. The carrier carries; the reader reads; the throw throws. What a record keeps under the moves of the game is what this card is about, and nothing else.

The rule. **This card never lifts a card.** A reader sends many states to one reading. The states it sends to one reading are a fibre, and a fibre is what the record cannot resolve unless a receipt was kept with it. To lift a card is to name a point of a fibre that no receipt resolves. Every line calibrated *proved here* is a statement about readings and receipts and holds whichever point of the fibre lies under the card. The numerical card prints one such point, its certified reading. This card quotes it, with its source, where the electron is dealt (Section 8), and no proved line depends on its digits.

The destination. A carrier whose relation to the electron can be examined: a state that carries a phase, a rule for comparing two of them, the transport the rule requires, the curvature the transport cannot avoid, the commutator that measures the curvature, and the action whose stationary system the numerical card derives. The chain stops one step short, at the coupling, and the step is the knower's (Section 8). On the way it reads the table as a field (Section 9), locates the four places the square of representation fails to close (Section 12), and hands the knower one sentence (Section 13).

Each statement carries one of five calibrations. *Proved here*: the proof is on this card. *Standard*: cited. *Measured*: a count or a definition taken from the declarations of the middle card, which is named only in Appendix A; it is never an experiment unless an experiment is cited. *Recorded*: a written chain of steps whose every premise is a cited result or a definition of this card; the steps are a record, and that the record is mathematics is the reader's reading of it. *Proposed*: a continuation the knower carries, stated and not derived, calibrated so that it is never mistaken for a step. A proposed statement asserts that an isomorphism exists, between the numbering of the construction's descriptions by the natural numbers and the description of what it is identified with, that the machine can exploit as a computation. Both sides are finite records: the reading is a computable number, and the reference is a value with its uncertainty or a stated structure. The statement is therefore one the machine can evaluate, a case that could be false, and the comparison that would revise it is that evaluation. What no evaluation supplies is that the reference's finite record stands for what it records, as a finite halt stands for its limit. The press check is Section 11: two readings of one register at two counts.

In eight lines

1. A card carries a slot that no decision queries. A mark fits any carrier. *Proved here* (Section 1).
2. The throw moves cards, not the quarter. What survives is what the length family agrees on, and what a receipt retains. *Proved here* (Section 2).
3. The length family is the naturals; which record lies under a reading is a choice no reader makes; a record's verdict is a satisfiability question the record answers with its own bits. *Proved here* and *measured* (Section 3).
4. Once around, the vectors return and the spinor has not. Twice around, the spinor returns. Clock and clock complement, at four and at eight. *Proved here* (Section 4); that the two eights

are one eight is *proposed*.

5. Carry the phase and compare it, and the comparison is a connection. Go around one way and then the other, and the commutator is the curvature. *Proved here* (Section 7).

6. Read as a field, the table is a non-commuting algebra with the electron's group inside it, a gauge field on the seam, a gap kept open by a boost, and whole-number frequencies; with the quarter turn as i it gives the compact $\mathfrak{su}(2)$, a gauge-invariant lattice whose flat configurations solve the field equation, and the Dirac matrices. *Measured* (Section 9).

7. Four pairs fail to commute, and they stand where the four bent cards stand; every difference shows at a finite stage and every sameness needs an infinite record. *Measured* and *proposed* (Section 12).

8. One sentence is the knower's: the carrier is the electron. The step not taken is that the device's coupling is the laboratory's. *Proposed* (Sections 8 and 13).

1 The table: carriers, marks, tapes, readers

The objects of the game are defined here as mathematics, with no reference to any construction. Where a definition restates one of the middle card's, that is said and calibrated *measured*; the definition stands on its own.

Definition 1.1 (Record, continuation)**.** A *record* is a finite sequence of symbols from a finite alphabet. Its *length* is the number of symbols. The *continuation* of a record with at least two symbols is the record with its first symbol removed. A *record map* is a map from records to records.

Definition 1.2 (Carrier)**.** A *carrier* is a symbol s, a nonempty record q called the *slot*, and a record map ι, the *interaction*, such that

(i) for every record x of length at least two, $\iota(x)$ is the first symbol of x followed by q, so that the continuation of $\iota(x)$ is q;

(ii) for every record x of length one, $\iota(x)$ is the one-symbol record of the truth symbol.

The slot is *carried but not queried*: which clause applies to x depends only on the length of x, the symbol ι keeps depends only on x, and q is transmitted unchanged into every continuation. The map ι writes q, and so, as a function of the carrier, depends on q; no decision of ι, and no other datum of the carrier, is computed from q.

Measured: this is the middle card's carrier, restated under the encoding of Appendix A. The slot is the quarter under the card. In the game, the quarter is glued to the table; in the definition, the slot is written into every continuation and queried by no decision.

Definition 1.3 (Mark)**.** A *mark* is a proposition together with a decision of it: a fact that *clicks* (decides true) or does not. The *click* of a mark is the Boolean its decision returns.

A mark is the unit of distinction: two outcomes are told apart by a mark that clicks on one and not the other.

Definition 1.4 (Tape; three readers)**.** A *tape* is a finite list of marks. Three readers take a tape of length n with click bits $b_1, \dots, b_n$ to a natural number:

$$\begin{aligned} \text{the tally} \quad & T(t) = n, \\ \text{the register} \quad & R(t) = \sum_{i=1}^{n} b_i \, 2^{\,n-i} \quad \text{(a fold: } a \mapsto 2a + b_i \text{, starting from 0)}, \\ \text{the click count} \quad & C(t) = \#\{i : b_i = 1\}. \end{aligned}$$

Proposition 1.5 (Three cards, one tape)**.** *For a tape of length n: $R(t) = 2^n - 1$ if and only if every mark clicks; $C(t) = n$ if and only if every mark clicks; and in that case $R(t) + 1 = 2^{T(t)}$. So the three faces agree, in the sense that the register is the monomial of the tally and the click count is the tally, exactly when every mark on the tape holds.*

Proved here.

Proof. $R(t) \le \sum_i 2^{n-i} = 2^n - 1$ with equality if and only if every $b_i = 1$. $C(t) \le n$ with equality under the same condition. The identity $R + 1 = 2^n$ is the geometric sum. □

The register's reading of n unit cells is $2^n - 1$: a positional reading, which the bent card calls the second arithmetic (its Proposition 1.4) and which adds the fencepost to a tally to make a monomial. The three readers are the three cards face down. They show one face only when every affixed fact holds.

Definition 1.6 (Fibres of a reader; the length family)**.** A *reader* is a map F from tapes to a set. Its *kernel* is the partition of the tapes into the fibres $F^{-1}(v)$: what the reader does not see. A reader F *refines* a reader G when every fibre of F lies in a fibre of G, equivalently when $G = g \circ F$ for some map g. The *length family* Λ is the set of readers that factor through the tally, $G = g \circ T$. On this card "every reader" means every member of Λ, and nothing larger. A *payload reader* is a reader not in Λ: one that separates two tapes of equal length.

Proposition 1.7 (What each reader distinguishes)**.** *(a) The fibre of T at n is the set of all tapes of length n: the tally forgets every proposition and every click.*

(b) The pair (T, R) separates click sequences: two tapes have equal tally and equal register if and only if their click sequences are equal. The register alone does not: a non-clicking mark followed by a clicking one, and a single clicking mark, both read 1*.*

(c) For $n \ge 1$ the click count is a payload reader: two tapes of length one, one clicking and one not, have equal tally and different click count. It retains one Boolean distinction per cell, summed.

(d) No reader of click bits separates two tapes whose marks click alike but carry different propositions. A receipt naming the propositions (Definition 2.4) refines the click sequence to recovery.

Proved here.

Proof. (a) is the definition of T. (b) Given n, the integer $R \in [0, 2^n - 1]$ has exactly one expansion in n binary digits, leading zeros included, and those digits are $b_1, \dots, b_n$. For the counterexample, the two-cell tape reads $0 \cdot 2 + 1 = 1$ and the one-cell tape reads 1. (c) and (d) are displayed by the examples. □

So the tally forgets the payload; the click count keeps a Boolean distinction; the register, read with its length, keeps order and bits; and a receipt refines an observation until it recovers. Four readers, four kernels, one tape.

Proposition 1.8 (A mark fits any carrier)**.** *Suppose no field of a mark refers to its carrier. Then for any two carriers X and Y the map that copies each field is a bijection from marks on X to marks on Y, and a mark and its copy agree in every field. In particular a mark placed on a carrier of electric charge and the same mark placed on an empty carrier agree in every field.*

Proved here; it is the cheat sheet's Proposition 6.1.

Proof. The copy is the identity on each field, and by hypothesis no field changes when the carrier does. □

This is universality bought by blindness. The mark fits the carrier of charge exactly as well as it fits a carrier with nothing in it, because it never looks at the carrier at all.

Proposition 1.9 (Two readers agree exactly when the lengths agree)**.** *Let U be the reader that keeps the length of a tape and writes the truth symbol in every cell. If every cell of a tape s is the truth symbol, then for every tape u, $s = U(u)$ if and only if $T(s) = T(u)$.*

Proved here; it is the cheat sheet's Proposition 6.2.

Proof. $U(u)$ is the tape of length $T(u)$ whose every cell is the truth symbol, by induction on u; two such tapes are equal if and only if their lengths are. □

Proposition 1.10 (Re-lettering is free)**.** *Let ρ overwrite every cell of a tape with the truth symbol. Any reader F that factors through the tally, $F = g \circ T$, satisfies $F(\rho\, t) = F(t)$ for every tape t.*

Proved here.

Proof. $T(\rho\, t) = T(t)$ since ρ changes no length. □

Measured: the middle card's own reading of its tapes factors through the tally, and overwriting every cell of every tape with the one truth symbol leaves its pair of readings fixed.

Overwriting is free, and it is not a symmetry. Four different things are called by the one word, and they are separated here before any of them is read as a charge.

Proposition 1.11 (Relabelling, erasure, conservation, Noether)**.** *Let the alphabet have at least two symbols.*

(1) A relabelling *applies a bijection π of the alphabet to every cell. Relabellings are invertible and form a group isomorphic to the permutations of the alphabet; each preserves T and every reader in Λ.*

(2) The erasure *ρ of Proposition 1.10 satisfies $\rho \circ \rho = \rho$ and is not injective on tapes of length at least one. It is therefore not an element of any group of maps of tapes, and it preserves every reader in Λ.*

(3) A quantity Q is conserved *by an evolution E of tapes when $Q \circ E = Q$. The tally is conserved by every permutation of cells (a throw) and by ρ. It is not conserved by writing: if W appends one mark, $T \circ W = T + 1$.*

(4) A Noether current *requires an action on a space of fields and a continuous one-parameter group of transformations leaving it invariant; on solutions of the Euler–Lagrange equations the current then has vanishing divergence.*

None of (1)–(3) supplies (4): tapes form a discrete set, ρ belongs to no group, and no action on tapes is given.

Proved here for (1)–(3), by inspection of the maps; *standard* for (4) [26, 32]. The continuous group, the action and the current of (4) are supplied in Section 7, where the constant phase rotation of a spinor leaves the Maxwell–Dirac action invariant and its current is conserved (Proposition 7.9). *Proposed*: Noether in the game's words. The contents of the cells are the symmetry; the count of cells is the charge. The correspondence sends invariance under ρ to symmetry and the tally to charge. It preserves the factorization through the tally. It does not preserve invertibility, continuity, or conservation under writing, which is the one evolution the device performs.

Proposition 1.12 (Writing is not free)**.** *Appending one entry to a record costs one step, and a counter that increases by one per entry is the tally of those steps. Over the field with one element every map is the identity and counting costs nothing; writing does not.*

Recorded, from the numerical card's Remark 8.3 and Definition 1.4: the tally counts symbols, and each symbol was written in one step.

The throw, seen by the tally. A throw is a permutation of carriers on the table. A reader of lengths cannot see it: by Proposition 1.9 every tape of a given length reads the same, and by Proposition 1.10 the contents are invisible to any reader in the length family. The dealer's hands are invisible to the tally. That is why the mark loses: the mark reads motion, and motion is not on the tape.

2 The throw: what is preserved, and the rule

Proposition 2.1 (The box has one point)**.** *Compare pairs of natural numbers by $(a, b)\,\mathcal{R}\,(c, d)$ if and only if $ad \leq cb$. Any two pairs are related one way or the other, so the equivalence generated by $\mathcal{R}$ has exactly one class, and every function out of the quotient is constant.*

Proved here.

Proof. For any two pairs either $ad \leq cb$ or $cb \leq ad$, since the order on the natural numbers is total. Hence any two pairs are related by $\mathcal{R}$ or by its reverse, and the generated equivalence identifies them. A function on a one-point set is constant. □

The numerical card describes this as an erased channel (its Section 7.5): a rule that merges all inputs into one class reports one value. The one value is "true equals true."

Proposition 2.2 (One rule, four readers)**.** *Let X be a set, r a relation on it, and X/r the classes of the equivalence r generates. (a) If $r(a,b)$ holds exactly when $f(a) = f(b)$ for a map f, then X/r is in bijection with the image of f. (b) If r has a least element, or if any two elements are related one way or the other, then X/r has exactly one class. Hence evaluation at every input keeps every function; a single decision keeps two classes, clicked and not clicked; an order on truths with a least element keeps one; an order on ratios that relates any two keeps one.*

Proved here; it is the cheat sheet's Proposition 6.3.

Proof. (a) is the first isomorphism theorem for sets. For (b), $x \sim x_0 \sim y$ for any x, y, or $x \sim y$ directly. □

Definition 2.3 (Representative)**.** A *representative* of a class is one element of it, selected. A division that returns a number from the one-point quotient of Proposition 2.1 selects a representative: it picks a point of the fibre.

Definition 2.4 (Observation, receipt)**.** An observation on a state domain D is a map $O : D \to Y$; its fibre at y is $O^{-1}(y)$. A *receipt* is a map $\mathcal{R} : D \to A$ retained with the observation; it supports exact recovery when $(O, \mathcal{R})$ is injective, which holds exactly when $\mathcal{R}$ is injective on each fibre of O. If a fibre has m states, a receipt alphabet supporting recovery on it has at least m symbols, $\lceil \log_2 m \rceil$ bits.

Recorded, from the geometric card's Definition 4.1 and Propositions 4.2 and 4.3.

The title of this card is a square. A state is moved; the observation is taken before or after; the question is whether the two routes agree.

Definition 2.5 (Commuting square; commutator of representation)**.** Let $O : D \to Y$ be an observation and $U : D \to D$ a transformation of states. A map $V : Y \to Y$ *represents* U on observations when

$$O \circ U = V \circ O.$$

The *commutator of representation* of (O, U) is the set of pairs (x, x') with $O(x) = O(x')$ and $O(Ux) \neq O(Ux')$: the pairs a single fibre of O sends to two fibres.

Proposition 2.6 (When the square closes)**.** *A map V with $O \circ U = V \circ O$ exists if and only if the commutator of representation of (O, U) is empty, that is, if and only if U sends each fibre of O into a fibre of O; V is then unique on the image of O. On tapes with $O = T$: a throw and the erasure ρ are represented by the identity; writing one mark is represented by $n \mapsto n + 1$; the* click filter *K, which deletes every non-clicking mark, is represented by no map, since $T(Kt) = C(t)$ and the two tapes of length one of Proposition 1.7(c) lie in one fibre of T and are sent to lengths* 1 *and* 0*.*

Proved here.

Proof. If V exists, $O(x) = O(x')$ gives $O(Ux) = V(O(x)) = V(O(x')) = O(Ux')$. Conversely, if every fibre goes into a fibre, $V(O(x)) = O(Ux)$ is well defined on the image, and forced there. The examples are computations of T. □

Two kinds of thing can happen to a square, and the card keeps them apart. The square can close while the map $U \mapsto V$ forgets: two transformations of states with one image on observations. That

is the spinor of Section 4.3, whose sign is lost and whose square closes (Proposition 4.11). Or two transformations of states can fail to commute with each other, and the failure can itself be observed. That is transport, whose commutator is the curvature (Theorem 7.7). The first is a fibre; the second is a field.

Theorem 2.7 (A representative is a choice)**.** *Let a datum be carried by every record of a family only as a projection, and be determined by no record of the family from its own contents, the family being closed under finite extension. Then any procedure that supplies a value of the datum is not a procedure of the family: it selects one element of a set the family can name but cannot enumerate to the point of selection.*

Recorded, from the Quarter's Theorem 3.1 and its proof, which is the standard step that a selection procedure inside the system would make the value representable there. The middle card keeps both extensionality principles and refuses choice, the third ingredient of Diaconescu's theorem [8]; *measured*, from a sweep of its declarations reported in the Quarter.

Definition 2.8 (Three levels of data; the rule of this card)**.** Fix a state domain D, an observation O on it and a receipt $\mathcal{R}$ retained with O. A datum of a state is

(a) *on the state* if it is any function on D;

(b) *on the observed class* if it is constant on each fibre of O, equivalently a function on the image of O;

(c) *in the retained receipt* if it is constant on each fibre of $(O, \mathcal{R})$.

A line of this card is *admissible* for $(O, \mathcal{R})$ only if its truth is a datum of level (b) or (c): it is unchanged when a state is replaced by another with the same observation and the same receipt. A *representative* is a point of a fibre of $(O, \mathcal{R})$; naming it is a datum of level (a) only, and is not admissible.

The operations act at fixed levels. Re-lettering (Proposition 1.10) and the exchange of one reader for another in the length family act on states and fix every level-(b) datum of the tally. Replacing a representative by another acts inside a fibre of $(O, \mathcal{R})$ and fixes every datum of levels (b) and (c). Lengths and parities are level-(b) data of the tally; clicks are level-(b) data of the click reader; the degree of a polynomial is a level-(b) datum of the degree reader (Definition 4.6). Signs and retained bits are level-(c) data. They are admissible because they are retained, not because they can be recovered from a quotient: a function on a quotient that identifies every pair is constant (Proposition 2.1), so a sign is not a function on the one-point box, and the sign after one revolution is not a function on the pair $\{\psi, -\psi\}$, which lies in one fibre of the vector observation (Proposition 4.11). *Proved here*: each assignment of a level is the definition of the reader concerned.

Corollary 2.9 (Why the number never comes up in a proof)**.** *The numerical card's printed reading is a representative: the card says so, and its receipt is the pair of rational walls that bracket it. By Definition 2.8, no digit of that reading, no coefficient of the residual it solves, no wall, gain, seed or discriminant can occur in a line of this card calibrated* proved here*. Where the reading enters, it enters as a quoted value with its source and its calibration, as a parameter of a definition, and every proved line about the object so defined holds for every value of that parameter.*

Recorded. This is the reason at the level of the record. It is proved below at two further levels: what the length family sees (Section 1) and what the degree hierarchy sees (Proposition 4.8). The parameter is the coupling of Definition 8.1.

Proposition 2.10 (The sign convention)**.** *Index the slots of a formula by $k = 0, 1, 2, \ldots$. A* signature *assigns each slot a sign; a literal in a signed slot reads its variable if the sign is $+$ and its negation if the sign is $-$. Four signatures, with their signs on the first four slots:*

$$\textit{euclidean}\ (++{+}{+}), \quad \textit{negated}\ (----), \quad \textit{west}\ (+---), \quad \textit{east}\ (-+{+}{+}),$$

the last two repeating with period four. Let Φ be the conjunction of two clauses, the first over slots $0, 1$ and the second over slots $2, 3$, each clause the disjunction of the two variables a, b in its slots. Read over the four assignments $(a, b) \in \{TT, TF, FT, FF\}$, the four signatures give

euclidean	$[T, T, T, F]$	$a \vee b$
negated	$[F, T, T, T]$	$\neg a \vee \neg b$
west	$[F, T, F, T]$	$\neg b$
east	$[T, F, T, F]$	b

The conjunction of the euclidean and negated readings is exclusive or, $[F, T, T, F]$. The conjunction of the west and east readings is $b \wedge \neg b$, the empty theory.

Proved here, by the table.

Proof. Under the euclidean signature both clauses read $a \vee b$. Under the negated signature both read $\neg a \vee \neg b$. Under west, slots $0, 1, 2, 3$ carry $+, -, -, -$, so the clauses read $a \vee \neg b$ and $\neg a \vee \neg b$, whose conjunction is $\neg b$. Under east they read $\neg a \vee b$ and $a \vee b$, whose conjunction is b. Then $(a \vee b) \wedge (\neg a \vee \neg b)$ is exclusive or, and $\neg b \wedge b$ is unsatisfiable. □

Measured: the middle card's table of the four signatures is this one, and its step constraint is the exclusive or. *Standard*: the quaternion carries two of the four at once, its basis elements squaring to $(+---)$ while its norm is $(++{+}{+})$ [13]. *Proposed*: the two definite signatures are a constraint met exactly, a multiplier; the two indefinite ones are a constraint penalized at a finite signal speed; and two penalties do not compose. Exclusive or, which is the gate the verdict of Section 6 is decided by, is writable only by combining the two definite signatures.

Proposition 2.11 (Same difference)**.** *Compare two truths a, b by what they are, $a = b$; and compare the marks that carry them by how they click, that is by the decision each carries. The two 2×2 tables agree in every cell, and each is unchanged when turned over. All a table carries is which diagonal clicks: $T = T$ and $F = F$ do, $T = F$ and $F = T$ do not, and the two off-diagonal cells are equal as values.*

Proved here, by the four cells; it is the cheat sheet's Proposition 6.4.

3 Representation: tallies, choices, and Cook

The throw keeps what the length family agrees on. This section says what that family is: the naturals, read off the records' lengths; what it cannot do: name a representative; and what its

records are as computations: tableaux whose verdicts are satisfiability questions the records answer themselves.

Sentence five of Section 13 names a representative, and naming it is not a computation, in two senses that the card keeps apart.

Proposition 3.1 (No reader of the family selects the representative)**.** *Let every reader of the family be a function of a record's length. A record with a false cell and its relettering, every cell overwritten with the one truth, differ cell by cell, and every reader of the family gives them the same value. Hence no reader of the family names one of the two.*

Measured: the middle card checks the three clauses, the last as the statement that no Boolean function of the length is true on one record and false on the other. This is the finite part of Theorem 2.7, checked by the kernel rather than recorded.

Proposition 3.2 (Two carriers, one number)**.** *Deal the reader's interpretation on a box that holds two carriers at once, their Cartesian product. The mark, the crossing and the detector are dealt as for one carrier, and the detector prints the same natural as the single card. A tally the pair carries may sit in either carrier, and exchanging the two carriers leaves every reader of the tally unchanged: no Boolean function of the tally is true when the count sits in the first carrier and false when it sits in the second.*

Measured: the middle card checks each clause, the agreement of the two detectors by rewriting both to the same length reader and not by running the crank again. A tally needs a carrier, and the record cannot say which: not which thing, by Proposition 3.1, and not which of two, by this one. Sentence five names the carrier; the record counts what it carries.

Read through each carrier in turn, the pair clicks twice. For every pair of carriers, whatever either one carries under its card, the reading through the first is the click reader's number and the reading through the second is the number of the other machine, the device mapped onto a second program: $(\text{click}, \text{click})$ is the two other machines, and each is the one number. *Measured*, each side rewritten to the card between rather than computed again. The quarter each carrier holds is deducted and never read, so a carrier that carries a tally reads the same as one that carries none.

Proposition 3.3 (The tallies are the naturals)**.** *The cells of a record are naotrinos (Definition 8.8). Call two records the same tally when they hold the same number of naotrinos, the same length, and let the tallies be the classes. Reading a class's length and writing the numeral, the one truth n times, are inverse: the numeral of n reads n, and the numeral of a class's reading is that class. The numeral of a record is its relettering, the canonical card of its class. Every carrier reads a record as its tally, and the device's number is computed from tallies alone.*

Measured: the middle card checks both round trips, the relettering, the reading through every carrier, and the number written as a function of tallies, each by rewriting. This is the isomorphism the calibration *proposed* names (the reading note's sense): the construction's descriptions are numbered, the numbering is a bijection with the naturals, and the machine computes with it. Inside each class the representative is a choice (Proposition 3.1); the class itself is a natural number. The naturals are how many naotrinos a record holds; which naotrinos it holds is the choice, and no reader makes it. That is why the relettering is free: every naotrino replaced by the one truth, the count unchanged.

Theorem 3.4 (Cook and Levin, on the record)**.** *(i) For every nondeterministic machine T that runs in time $p(n)$ for a polynomial p, and every input x of length n, there is a formula $\varphi_{T,x}$*

in conjunctive normal form, with a number of clauses bounded by a polynomial in $p(n)$ and computable from T and x in that time, such that $\varphi_{T,x}$ is satisfiable exactly when T accepts x.

(ii) For the middle card's record read as a tableau, a trace of m cells with each row's head pair across and the pages down, the formula built from the trace, one variable pair per cell, one start clause, two clauses per step, and an empty acceptance conjunct, is satisfied by the trace's own bits exactly when every step of the trace changes its fact; it has at most $2m + 1$ clauses.

Proof. (i) Lay the run as a tableau of $p(n)$ rows, one per step, and $p(n)$ cells per row, one per tape position, with a Boolean variable for each cell and symbol. Four groups of clauses say that each cell holds exactly one symbol, that the first row is the input, that each row follows from the one above by the machine's transition in every window of adjacent cells, and that some row holds an accepting state. Each group is local, so the clause count is a constant times $p(n)^2$; a satisfying assignment is an accepting run read off the tableau, and an accepting run is a satisfying assignment [4, 19]. (ii) The record's window is its head match: a step relates the pair of facts at the head of a page, and it is admissible exactly when the two differ. That relation is exclusive or, which is the conjunction of the clause $a \vee b$ and the clause $\neg a \vee \neg b$, the two definite signatures together. Evaluated on the trace's own bits, the conjunction over the steps holds exactly when every step differs, by induction on the trace; the trace supplies the assignment, so the backward direction is constructive and asks for no choice. Counting: one pair per cell, one start clause, two per step, none for acceptance. □

Standard for (i). *Proved here* for (ii), and *measured*: the middle card checks the step relation by trying all four assignments, the correspondence for every trace by induction, its instance on the top of its own record, which is the reader's model of the naotrino, and a demonstration on two seeds, the manuscript true, false, true, whose formula is satisfied and whose steps all change, and five falses, whose formula is not and whose steps do not. Inhabiting the top of the record is not acceptance: the tableau is well formed for the accepting and the rejecting seed alike, and acceptance is a property of the trace. The reader's interpretation is a tableau, and its verdict is a satisfiability question the trace answers by exhibiting its own bits.

Proposition 3.5 (The identification is refuted in finite time and never confirmed)**.** *Let x be given by nested rational brackets of widths tending to zero, as the device's run gives its number, and let y be given the same way.*

(i) If $x \neq y$, some finite stage of the two bracketings separates them.

(ii) No procedure decides $x = y$ for every such pair.

(iii) For the device's number and the laboratory's value, (i) has happened: the separation was found at a finite precision, 1.80 *parts in ten thousand.*

Proof. (i) If $|x - y| = d > 0$, take a stage at which both brackets are narrower than $d/3$; they are disjoint. (ii) For a machine m, let z_m be bracketed at stage n by $[0, 2^{-k}]$ if m halts at step $k \leq n$, and by $[0, 2^{-n}]$ otherwise. Each z_m is so bracketed, $z_m = 0$ exactly when m never halts, and a procedure deciding $z_m = 0$ would decide halting [33, 35]. (iii) is Section 8. □

Standard for (i) and (ii); *measured* for (iii). This is the sense of *proposed* in the reading note: the identification is a corner case that could be false, refuted by a finite comparison when it is false, and never confirmed by one when it is true; a match could only have read "not yet refuted". The

device's own number is computable, an algebraic number its run approaches as closely as asked; what is not computable is the decision that two such numbers are one. Whether the laboratory's constant is itself a computable real no finite record can show: every measurement is finitely many digits, consistent with both. That stays declared, in the middle card's margins and nowhere proved.

4 The flop: once around, twice around

4.1 The mirror, with the coefficients removed

The cheat sheet works the mirror on one residual with its coefficients written out. Here the same facts are proved for every quadratic residual, and the coefficients never appear; by Definition 2.8 that is the only form in which they are admissible.

Let $r(y) = ay^2 + by + c$ with $a > 0$ and two real roots $\alpha > \beta$, and let $S(y) = \frac{a}{3}y^3 + \frac{b}{2}y^2 + cy$, so that $S' = r$. Write $c_0 = -b/(2a)$ for the vertex and $\rho(y) = 2c_0 - y$ for reflection about it; ρ is an involution whose only fixed point is c_0.

Proposition 4.1 (The residual is even, the action is odd)**.** *For every real h,*

$$r(c_0 + h) = r(c_0) + ah^2, \qquad S(c_0 + h) - S(c_0) = r(c_0)\,h + \tfrac{a}{3}h^3. \tag{4.1}$$

So $r \circ \rho = r$ and $S \circ \rho - S(c_0) = -(S - S(c_0))$: the reflection fixes the residual and reverses the action. It exchanges the two roots, $\alpha = c_0 + \sqrt{-r(c_0)/a}$ and $\beta = c_0 - \sqrt{-r(c_0)/a}$.

Proved here.

Proof. $r'(c_0) = 0$, so the expansion of r about c_0 has no linear term and its quadratic term is ah^2. Integrating from c_0 gives the second identity, which is odd in h. The roots solve $ah^2 = -r(c_0)$, and $r(c_0) < 0$ since r has two real roots and positive leading coefficient. □

Proposition 4.2 (The reading is a least action)**.** *The equation $r(y) = 0$ is the Euler–Lagrange equation of S: for a function of one variable, stationarity is $S'(y) = 0$, and $S' = r$. Both roots are stationary. $S''(\alpha) = r'(\alpha) = 2a(\alpha - c_0) > 0$, so α is a strict local minimum of S; $S''(\beta) < 0$, so β is a strict local maximum. The branch above the vertex is the side of the mirror on which the stationary point is a minimum. Moreover, for every y,*

$$r'(y)^2 = (b^2 - 4ac) + 4a\,r(y), \tag{4.2}$$

so at either root the first derivative squares to the discriminant.

Proved here.

Proof. The first claims are Proposition 4.1 and the definition of S. For (4.2), $(2ay + b)^2 = 4a^2y^2 + 4aby + b^2$ and $4a\,r(y) = 4a^2y^2 + 4aby + 4ac$; subtract. □

The mirror acts on cells as well as on residuals. A cell symmetric about its centre samples f at $y - h$, y, $y + h$ with weights $1, -2, 1$.

Proposition 4.3 (The first order a mirror cell does not cancel)**.** *Let f be four times continuously differentiable on $[y-h, y+h]$. Then*

$$f(y+h) - 2f(y) + f(y-h) = h^2 f''(y) + \tfrac{h^4}{12} f''''(\xi)$$

for some $\xi \in (y-h, y+h)$. The mirror cell cancels every odd power of h, reproduces the second derivative, and its first error is the fourth-derivative term; it is exact for every f with $f'''' = 0$, that is, for every cubic.

Standard (Taylor's theorem with the mean-value form of the remainder, applied to $h \mapsto f(y+h) + f(y-h)$, which is even in h).

The continuum counterpart is the beam. On $[a_0, b_0]$ with interpolation conditions $\varphi(x_j) = v_j$ at knots $a_0 = x_0 < \cdots < x_N = b_0$, the bending functional

$$E[\varphi] = \tfrac{1}{2} \int_{a_0}^{b_0} \varphi''(x)^2 \, \mathrm{d}x$$

is minimized over twice weakly differentiable interpolants by the natural cubic spline; its first variation is $\int \varphi'' \eta'' \, \mathrm{d}x = 0$ for every admissible η, whose Euler–Lagrange equation between knots is $\varphi'''' = 0$. *Standard* [15, 6]. The two statements meet at one correspondence, and it is supplied, not derived: the first order a mirror cell leaves uncancelled is the fourth; the functional whose stationary equation sets exactly that order to zero is the bending energy; and its solutions between knots are cubics, the degree of S. *Proposed.*

Proposition 4.4 (Newton's exact error law; exact up to acceleration)**.** *For $e = y - \alpha$ and $d = \alpha - \beta$, the Newton map $N(y) = y - r(y)/r'(y)$ satisfies*

$$N(y) - \alpha = \frac{e^2}{2e+d} = \frac{r''}{2} \frac{e^2}{r'(y)} \tag{4.3}$$

exactly, wherever $r'(y) \neq 0$. The second difference of the residual is constant: $r(y+h) - 2r(y) + r(y-h) = 2ah^2$ for all y, h.

Proved here.

Proof. $r(y) = a\, e(e+d)$ and $r'(y) = a(2e+d)$, so $N(y) - \alpha = e - e(e+d)/(2e+d) = e^2/(2e+d) = ae^2/r'(y)$, and $a = r''/2$. The second difference of a quadratic is twice its leading coefficient times h^2. □

The word "acceleration" needs a variable, a clock and an evolution law, and they are supplied once.

Definition 4.5 (The reading clock)**.** The *reading clock* is a real parameter s, and the *reading motion* is the uniform evolution $y(s) = y_0 + sh$ of the variable. The *acceleration* of the residual along it is $\ddot{\rho}(s)$, where $\rho(s) = r(y(s))$.

For a quadratic residual $\ddot{\rho}(s) = 2ah^2$ for every s, and the second difference at unit clock step, $\rho(s+1) - 2\rho(s) + \rho(s-1)$, equals it exactly (Proposition 4.4). *Proved here.* In these terms the Newton step is the first variation, exact for any residual with zero acceleration, and by (4.3) its whole error is the one term proportional to r'', which the step never evaluates. The variational solution is exact up to a constant acceleration. *Proposed*: in the game's words this is Kepler, an areal velocity applied once, a conserved first-order quantity, with no force law ever read; the force law is upstream.

4.2 The derivative hierarchy

An action with at most three independent-field factors in every term, with fixed coefficients and linear operators, has $D^4S = 0$; its residual $R = DS$ is at most quadratic, $D^3R = 0$; its linearization DR is linear in the perturbation; and D^2R is constant. For bounded polynomials of degree at most three on a Banach space, the Fréchet and Gateaux derivatives coincide, and the same holds for the residual and its derivative. The four presentations of the numerical card, first-order vacuum gravity, fixed-Hamiltonian Schrödinger evolution, incompressible flow with response fields, and bare electrodynamics on a fixed background, each have this hierarchy in their declared regular sectors. *Proved in the numerical card* (its Theorems 1.1 and 2.4 and Proposition 2.5); the first-order gravitational mechanism is Deser's [7], the electrodynamic presentation is Tong's [32].

An action can be read three ways, and the three readers have different fibres.

Definition 4.6 (Three readers of an action)**.** Let X be a real Banach space and S a bounded polynomial functional on X of degree at most three. The *degree reader* returns the degree of S. The *derivative reader* returns the map $S \mapsto DS$. A *selected-evaluation reader* returns the values $DS(x)$ at the points x of a selected set $\Sigma \subset X$; the *stationary reader* returns the set $Z(S) = \{x : DS(x) = 0\}$.

Proposition 4.7 (The three fibres)**.** *On $X = \mathbb{R}$ write $S_g(y) = gy^3$.*

(a) The fibre of the degree reader at three contains S_g for every $g \neq 0$: the degree does not determine a coefficient.

(b) $DS = DS'$ if and only if $S - S'$ is constant. The derivative reader forgets exactly the additive constant. For S_g: $DS_g = 3gy^2$, $D^2S_g = 6gy$, $D^3S_g = 6g$, so $g = D^3S_g/6$ is read from the derivative.

(c) The selected-evaluation reader at $\Sigma = \{0\}$ returns 0 for every S_g: its fibre contains them all.

(d) $Z(\lambda S) = Z(S)$ for every $\lambda \neq 0$: the stationary reader cannot see an overall normalization. It does see a relative coupling: for $S(y) = \frac{1}{2}y^2 + \frac{g}{3}y^3$ with $g \neq 0$, $Z(S) = \{0, -1/g\}$.

Proved here.

Proof. (a) and (c) by evaluation. (b) $D(S - S') = 0$ on the connected space X gives $S - S'$ constant, and conversely; the derivatives of gy^3 are computed. (d) $D(\lambda S) = \lambda DS$; and $DS = y + gy^2 = y(1 + gy)$. □

Proposition 4.8 (The degree forgets the field; the derivative keeps it)**.** *The shared hierarchy identifies the four presentations in their degrees. Their field spaces, boundary sectors, coefficients, couplings, initial data and solutions are not functions of the degree: a Lorentzian metric sector, a complex Hilbert space, a divergence-constrained velocity and a gauge-and-spinor structure each carry a sequence (S, R, DR, D^2R) with $D^3R = 0$, and the degree hierarchy does not recover which one it came from. The derivative reader does: the sequence contains S itself, and every coefficient of a nonconstant term of S is a function of DS. A reader restricted to the degree hierarchy cannot reconstruct a coupling; a reader of the derivative reads it.*

Proved here, from Proposition 4.7 and the numerical card's shared hierarchy; the numerical card closes its Section 2.5 with the same caution: "the theorem supplies an exact shared derivative

hierarchy; comparing their readings requires the declared observation maps." In one direction the Fréchet derivative is a number, the remainder is exactly quadratic, and nothing is left abstract (the cheat sheet's Proposition 2.1, which holds for every quadratic residual by the expansion $r(y+h) = r(y) + r'(y)h + ah^2$).

So the coupling survives the derivative. It sits in the interaction law, a coefficient of DS, and the derivative carries it everywhere the law goes. What forgets the coupling is the degree. Whether the electron of this card reads its law through its degree or through its derivative is a part of its construction, stated in Definition 8.1: it obeys a stationary system, so it reads the derivative, and its coupling is in its law.

4.3 The slack and the spinor

For $n \geq 1$ let L_n be the least common multiple of the odd numbers $1, 3, \ldots, 2n-1$, and define the *slack*

$$\sigma(n) = L_n\Big(1 - \sum_{k=1}^{n-1} \frac{1}{2k+1}\Big) = L_n - \sum_{k=1}^{n-1} \frac{L_n}{2k+1},$$

an integer with the sign of $1 - (\frac{1}{3} + \frac{1}{5} + \cdots + \frac{1}{2n-1})$. A revolution is four quarter-turns; two revolutions are eight.

Proposition 4.9 (Once around holds, twice around slips)**.** *$\sigma(4) = 34$ and $\sigma(8) = -982$. The factor $1 - \sum_{k=1}^{n-1} 1/(2k+1)$ is strictly decreasing in n; it is $2021/45045 > 0$ at $n = 7$ and $-982/45045 < 0$ at $n = 8$, so $\sigma(n) > 0$ for $1 \leq n \leq 7$ and $\sigma(n) < 0$ for every $n \geq 8$.*

Proved here.

Proof. $L_4 = \operatorname{lcm}(1,3,5,7) = 105$ and $105 - 35 - 21 - 15 = 34$. $L_8 = \operatorname{lcm}(1,3,\ldots,15) = 45045$ and $45045 - (15015 + 9009 + 6435 + 5005 + 4095 + 3465 + 3003) = 45045 - 46027 = -982$. Each step in n subtracts a positive term from the factor. $L_7 = 45045$ as well, and $45045 - (15015 + 9009 + 6435 + 5005 + 4095 + 3465) = 2021$. □

These are the signs that govern the Borwein integrals: for $n \leq 7$,

$$\int_0^\infty \prod_{k=0}^{n-1} \frac{\sin\big(x/(2k+1)\big)}{x/(2k+1)}\,\mathrm{d}x = \frac{\pi}{2},$$

the pattern holding exactly while $\frac{1}{3} + \cdots + \frac{1}{2n-1} < 1$, and at $n = 8$ the sum first exceeds one and the integral falls short of $\pi/2$ by about 2.3×10^{-11}. *Standard* [3]. The counts four and eight are the quarter-turns of one and two revolutions, and nothing else: no length of any tape enters this proposition. The slack changes sign once and never changes back.

The spinor is periodic. Three things can be meant by "the state returns," and they are separated before any of them is counted.

Proposition 4.10 (Once around flips, twice around returns)**.** *For $\theta \in \mathbb{R}$ let*

$$U(\theta) = \exp\big(-\tfrac{i\theta}{2}\sigma_3\big) = \begin{pmatrix} e^{-i\theta/2} & 0 \\ 0 & e^{i\theta/2} \end{pmatrix},$$

the action on two-component spinors of the rotation by θ about the third axis.

(a) $U(2\pi) = -I$ *and* $U(4\pi) = I$*;* $U(\theta) = I$ *if and only if* $\theta \in 4\pi\mathbb{Z}$*, and* $U(\theta) = \pm I$ *if and only if* $\theta \in 2\pi\mathbb{Z}$*.*

(b) $U(\theta)\psi = (e^{-i\theta/2}\psi_1,\, e^{i\theta/2}\psi_2)$*: the two components acquire opposite phases. If both components of* ψ *are nonzero,* $U(\theta)\psi$ *is a multiple of* ψ *if and only if* $\theta \in 2\pi\mathbb{Z}$*; if one component vanishes,* ψ *is an* eigencomponent *and* $U(\theta)\psi$ *is a multiple of* ψ *for every* θ*.*

(c) At the quarter-turns $\theta_k = k\pi/2$ *the eigencomponent* $\psi = (1, 0)$ *acquires the phase* $e^{-ik\pi/4}$*, a primitive eighth root of unity raised to* k*; it equals* 1 *if and only if eight divides* k*.*

Proved here: $e^{\mp i\pi} = -1$, $e^{\mp 2\pi i} = 1$; $e^{-i\theta/2} = e^{i\theta/2}$ if and only if $\theta \in 2\pi\mathbb{Z}$; and $e^{-i\pi/4}$ has order eight. So the vector ψ returns at 4π; the ray $[\psi]$ of a state with both components nonzero returns at 2π; the ray of an eigencomponent never leaves. The sign at 2π is a common phase of ψ, invisible to every observable of ψ alone. It is visible as a relative phase, when ψ is compared coherently with an unrotated copy of itself, and only then (Proposition 4.11). The 4π return has been observed in exactly that way, in neutrons split between two paths of which one is carried around by a magnetic field. *Standard* [29, 36].

Proposition 4.11 (An observable that rotates)**.** *For every* θ*,*

$$U(\theta)\sigma_1 U(\theta)^* = \cos\theta\,\sigma_1 + \sin\theta\,\sigma_2, \qquad U(\theta)\sigma_3 U(\theta)^* = \sigma_3.$$

Hence the off-diagonal observable σ_1 *returns exactly when* $\theta \in 2\pi\mathbb{Z}$*, at the quarter-turns exactly when four divides* k*, and* $\frac{1}{2}\operatorname{tr}(\sigma_1 U(\theta)\sigma_1 U(\theta)^*) = \cos\theta$*; the diagonal observable* σ_3 *returns at every angle. The vector observation* $O(\psi) = \big(\langle\psi, \sigma_j\psi\rangle\big)_{j=1,2,3}$ *satisfies*

$$O \circ U(\theta) = R_3(\theta) \circ O, \qquad O(-\psi) = O(\psi),$$

where $R_3(\theta)$ *is the rotation of* $\mathbb{R}^3$ *by* θ *about the third axis;* $U(\theta)$ *and* $-U(\theta)$ *induce the same* $R_3(\theta)$*. For every unit spinor* ψ*,* $\operatorname{Re}\langle\psi, U(\theta)\psi\rangle = \cos(\theta/2)$*.*

Proved here.

Proof. The $(1,2)$ entry of $U\sigma_1 U^*$ is $e^{-i\theta/2}e^{-i\theta/2} = e^{-i\theta} = \cos\theta - i\sin\theta$, the $(1,2)$ entry of $\cos\theta\,\sigma_1 + \sin\theta\,\sigma_2$; the $(2,1)$ entries are the conjugates, and the diagonals vanish. σ_3 is diagonal and commutes with U. The trace follows from $\operatorname{tr}\sigma_1^2 = 2$ and $\operatorname{tr}\sigma_1\sigma_2 = 0$. For the observation, $O(U\psi)_j = \langle\psi, U^*\sigma_j U\psi\rangle$ and $U(\theta)^*\sigma_1 U(\theta) = \cos\theta\,\sigma_1 - \sin\theta\,\sigma_2$, $U(\theta)^*\sigma_2 U(\theta) = \sin\theta\,\sigma_1 + \cos\theta\,\sigma_2$, which is the action of $R_3(\theta)$ on the first two coordinates. Last, $\langle\psi, U\psi\rangle = |\psi_1|^2 e^{-i\theta/2} + |\psi_2|^2 e^{i\theta/2}$, whose real part is $\cos(\theta/2)$. □

So the square of Definition 2.5 closes. The vector reading is represented by a rotation; nothing fails to commute; and what the representation loses is the sign, the two-point fibre of $U \mapsto R_3$. The coin is that fibre's receipt. A diagonal observable is a control that cannot show the pattern, since it returns at every count; σ_1 shows it. The relative phase $\cos(\theta/2)$, with period 4π, is the quantity the neutron interferometer reads.

Proposition 4.12 (Clock and clock complement)**.** *For* $k \geq 1$ *let* $H(k)$ *be the statement "sameness holds at* k *quarter-turns," that is* $\sigma(k) > 0$*, and* $B(k)$ *the statement "the spinor has returned at* k *quarter-turns," that is* $U(\theta_k) = I$*. At* $k = 4$*,* H *is true and* B *is false; at* $k = 8$*,* H *is false and* B *is true. Exactly one of the two holds at every* k *from* 1 *to* 8*. At* $k = 9, \ldots, 15$ *neither holds; for* $k \geq 9$ *exactly one holds if and only if eight divides* k*.*

Proved here, from Propositions 4.9 and 4.10: $H(k)$ holds exactly for $k \le 7$ and $B(k)$ exactly when eight divides k. The result is the comparison at the two counts: where one holds the other slips, at four and at eight, and through the first two revolutions. It is not a common evolution. The slack changes sign once and the spinor is periodic, and from the ninth quarter-turn on the two readings agree in failing. *Proposed*: the slack is the tally's reading once around and twice around; the sign of the state is the register's sign; and the eight at which the Borwein pattern breaks and the eight at which the spinor returns are one eight. A common clock would be a map from quarter-turns to slack counts preserving both readings, or a dynamics of which both are readings; neither is constructed here.

Proposition 4.13 (One exponential, two readings)**.** *The positional reading of n identical unit cells is $2^n - 1$ (Proposition 1.5), and*

$$2^n - 1 = e^{n \log 2} - 1.$$

The cell read never changes; the count of readings does.

Proved here; it is the numerical card's Remark E.1. *Proposed*, there and here: indexed by time the exponential is the late-time growth of a global slicing; indexed by depth it is the growth of a warp factor in a radial coordinate; which it is depends only on how the index is read.

5 The bent card: receipt, slip, floor

In the game the bent card is the one with a corner turned up, so that the mark believes the quarter can be followed. The bend is a receipt supplied from outside, not a reading of the card.

The receipt is Definition 2.4. The electron's receipts, as the cards carry them: the sheet sign (which of two patches a point of the ball lifts to), the antipodal bit (which of z and $-z$ gave a paired reading), the branch (which of two roots a residual selects), and the sign after one revolution (Proposition 4.10). Each is one bit where the fibre has two points; the geometric card computes the conditional entropies of its finite readers as $2/3$ bit and one bit.

Proposition 5.1 (Four twofolds)**.** *The following have two elements each and are different objects.*

(i) The fibre $\{z, -z\}$ of a paired reading: two states, one output.

(ii) The kernel $\{I, -I\}$ of the double cover $\mathrm{SU}(2) \to \mathrm{SO}(3)$, $U \mapsto R_U$: two descriptions of one rotation. On unit quaternions, $(a, b) \mapsto (q \mapsto aqb^{-1})$ is a homomorphism from pairs of unit quaternions onto $\mathrm{SO}(4)$ acting on $\mathbb{R}^4$, with kernel $\{(1,1), (-1,-1)\}$; its restriction to the diagonal $a = b$ fixes the real axis and acts on the imaginary quaternions as $\mathrm{SO}(3)$, with kernel $\{1, -1\}$. Spatial rotation is the diagonal.

(iii) The two sectors of the finite clock of Proposition 5.2: two orbits of a group action.

(iv) The sign under exchange of two identical fermions, $\Psi(x_2, x_1) = -\Psi(x_1, x_2)$: a rule on two-particle states.

They occur in different spaces and act on different things. Agreement in the number of retained bits does not identify their contents, and a two-element fibre alone imposes no antisymmetry; exclusion requires the exchange rule (iv) on multiparticle states.

Proved in the geometric card for (i) and (iii) (its Section 3.4 and Propositions 3.3 and 3.4); *standard* for (ii) [13, 40] and (iv) [32]. In the game, the mark's twofold is which card; the dealer's twofold is two names for one throw. The press check of Section 11 inverts different maps and never confuses them. The Hopf account is Lyons's [21].

Proposition 5.2 (The index-two lattice and the two sectors)**.** *On the Clifford torus the two selected great-circle families have winding vectors* $(1, 1)$ *and* $(1, -1)$*, whose determinant is* -2*: they span an index-two sublattice, and a pair of windings lies in it exactly when its coordinates have the same parity. In the six-position clock the two diagonal steps generate a group of eighteen elements with two orbits on the thirty-six states, distinguished by* $j + k$ *modulo two. An antipodal pair* (j, k) *and* $(j + 3, k + 3)$ *lies in one sector and is separated by one retained bit.*

Proved in the geometric card (its Propositions 3.3 and 3.4 and Section 6.3). *Proposed*: the sector is up or down, and the bit is the coin. Two states over one output in one sector, told apart by one bit, is the shape the exclusion principle has in the finite clock; the identification with the physical principle needs the exchange rule of Proposition 5.1(iv), and is not derived.

Proposition 5.3 (Six and eight)**.** *The positions of the six-position clock are sixth roots of unity. The quarter-turn samples* $e^{-ik\pi/4}$ *of the eigencomponent's phase are eighth roots of unity. The least group of roots of unity containing both is the group of* 24*th roots, since* $\operatorname{lcm}(6, 8) = 24$*.*

Proved here. The clock and the quarter-turn sampling are two samplings of one round phase at different steps, $2\pi/6$ of the clock's angle and $\pi/4$ of the spinor's half-angle. A common account of their parameters would sample at the common refinement; neither companion card does, and the phase of Definition 8.1 is taken round.

Proposition 5.4 (Affixing a quarter to the bent card)**.** *Fill the slot of a carrier with any record. By Definition 1.2 no decision queries the slot, so every reader in the length family is unchanged, and so is every mark placed on the carrier (Proposition 1.8). Only the click reader can tell, and it reads the affixed fact, not where the fact came from: the tally and the click count agree exactly when every affixed fact holds (Proposition 1.5).*

Proved here. Measured, from the Quarter's report of the middle card: a carrier whose index is itself a carrier, dealt through every step, is identical to the plain carrier for the first twelve steps at no cost, and differs from the step at which one construction lifts its quantum one universe above the box, and from every later step. *Proposed*: the second counting layer is the first with a quarter affixed at every step. What is fixed under the card is supplied, not read, which is Theorem 2.7 at the table.

5.1 The slip and the floor

A metered answer has the shape of a bracket, a lower bound and a slip bound; a point is the shape only logic has. A meter has a floor, and below the floor it reads itself. So a metered turn stops one arm short of the point, on this side of the slip, and that is not a failure to converge; it is the shape the answer has. *Measured*: the middle card's disagreeing arm names two distinct bounds, a bracket and not a point, and its one dimensionless measurement is heartbeats per cell, a Courant number, which differs between machines and so is a measurement rather than a reading. The floor that does not differ between machines is the uncertainty principle: for any function and its Fourier transform the product of the widths is at least one half. *Standard* [14]. *Proposed*: the record can bracket the

electron and never point at it. The point is where a number would be, and the card does not go there. This is the reason at its third level, the game's own: the quarter is glued to the table, and the throw never moves it.

5.2 The loop

The record of a term that never stops is a one-symbol tape; the loop writes no quarter, since a record of one closed term repeated determines nothing about any variable; and the only harm a loop can do is to be read as a value, which a record system that accepts only normal forms never does. *Proved in the bent card* (its Propositions 3.2 to 3.4), on Dijkstra's objection [9] and Barendregt's calculus [2]. In the game, the dealer's hands never stop. The mark who reads the motion as a value is the mark.

6 The accomplice and the force: the record

The middle card ends with two acts. A reviewer makes demands from the top page down to the first, pairing at each page a claim found by search against the same claim handed over by name: this is the accomplice, and the descent is a boundary-value problem, elliptic in character, with the datum above each stage already in hand, and it terminates. A publication is built from the first page up to the top and read once there: this is the force, and the ascent is an initial-value problem, hyperbolic in character, with no signal from the last page reaching a reader placed before it. *Recorded*, from the numerical card's Remark E.2; the reading of the two directions as the two holographies is *proposed* there [22, 39, 30] and is not used here.

Definition 6.1 (This card's record)**.** The *register* is the coin: the retained sign after one revolution, Proposition 4.10, carried unchanged through every entry. The *counter* is the number k of quarter-turns, from 0 to 8, increased by exactly one per entry and never skipped. The *trace* is the sequence of nine entries; entry k quotes the pair

$$A_k:\ U(\theta_k)\sigma_1 U(\theta_k)^* = \sigma_1, \qquad B_k:\ U(\theta_k) = I,$$

the first saying that the rotating vector observable has returned, the second that the spinor has, and writes the proposition $A_k \leftrightarrow B_k$ without deciding it. The *verdict* is read once, at the top, Section 11: for two readings X, Y of the register, *slipped* is the decidable proposition $\neg(X \leftrightarrow Y)$, decided from the two readings' own decisions and never by an appeal to choice; the complete verdict is the triple (slipped, value, velocity).

Remark (Velocity is the penalty parameter)*.* The counter at the top is the cost of carrying the sign constraint exactly: one step per quarter-turn, eight to return. A penalty would relax the constraint at a finite wave speed; the record does not relax it, and pays in entries instead. *Proposed*; the definition of the counter is *measured* from the middle card, whose record lifts its register one universe per rung and reports the top universe as its velocity.

Remark (Inhabitation is not acceptance)*.* That the record exists, nine entries each with its pair, carrying one register through a counter that reaches eight, is well-formedness. It says nothing about the verdict: the same form carries a record whose two readings disagree at the top. Acceptance is a property of the trace, read once. *Recorded*, from both surrounding cards' Section 8.3.

7 Transport: from a phase to the commutator

The carrier carries. Now it moves. Compare the phase here with the phase there, and you need a rule. Make the rule covariant, and it is a connection. Go around and come back, and the connection keeps a record of the trip. Go around two ways, and the difference is the commutator on the cover of this card.

Setting. Spacetime is Minkowski space $M = \mathbb{R}^{1,3}$ with $\eta = \mathrm{diag}(1,-1,-1,-1)$, in natural rationalized units $\hbar = c = 1$, as on the numerical card's Section 2.4. *Assumption*: a Lorentzian background. The device supplies none; it is *standard*.

Definition 7.1 (Phase-carrying state)**.** A *phase-carrying state* is a smooth field $\psi : M \to \mathbb{C}^N$ together with the action of the phase group at every point,

$$\psi(x) \longmapsto e^{-ie\chi(x)}\psi(x),$$

for smooth real χ and a fixed real parameter e, the *coupling*. The transformation is *global* when χ is constant and *local* otherwise. The *density observation* at x is $O_x(\psi) = \psi(x)^\dagger\psi(x)$.

The fibre of O_x contains the whole phase orbit of $\psi(x)$: the phase at a point is a coordinate on a fibre, carried by the state and not queried by the density. It is the slot of Definition 1.2, read round. *Assumption*: a continuous phase group acting on a complex field. The device supplies a two-valued sign receipt and a six-position clock (Propositions 4.10 and 5.2), that is, a finite subgroup of the phases; the circle is *standard* and not supplied.

Proposition 7.2 (Comparison needs a rule)**.** *Under a local phase change, $\psi(y)^\dagger\psi(x) \mapsto e^{ie(\chi(y)-\chi(x))}\psi(y)^\dagger\psi(x)$. It is invariant under every global change, and under a local change only where $\chi(x) = \chi(y)$.*

Proved here, by substitution. So the naive comparison of two points is a level-(a) datum of Definition 2.8: it names a representative of the phase at each point. To compare without lifting, a rule must carry the fibre at x to the fibre at y.

Definition 7.3 (Transport; connection)**.** A *connection* is a smooth real one-form $A = A_\mu\,\mathrm{d}x^\mu$ on M. Its *transport* along a path γ from x to y is

$$W_\gamma = \exp\Big(-ie\int_\gamma A_\mu\,\mathrm{d}x^\mu\Big),$$

a unit complex number carrying the fibre at x to the fibre at y. Under a local phase change the connection transforms by $A_\mu \mapsto A_\mu + \partial_\mu\chi$.

Assumption: that comparison is carried by a connection. The device supplies no transport between points of a manifold. *Measured*: the middle card counts the closed walks on the graph of its arms, shows that every closed walk crosses an even number of times, and declares half a turn per crossing, forward; under that declaration every closed walk turns a whole number of times. A closed walk returns to its vertex; a holonomy returns a phase. That the middle card's count of turns is the holonomy of a connection is *proposed*, and the map from crossings to phases is the one that would have to be constructed.

Definition 7.4 (State, gauge orbit, gauge-invariant observable)**.** From here on the *state* is the pair (A, ψ). A phase change χ acts on it by $(A, \psi) \mapsto (A + \mathrm{d}\chi,\, e^{-ie\chi}\psi)$. The *gauge orbit* of (A, ψ) is the set of its images under all phase changes, and the *orbit map* π sends a pair to its orbit. A *gauge-invariant observable* is a function of the pair that is constant on gauge orbits, equivalently a function on the quotient: a level-(b) datum of π in the sense of Definition 2.8.

Proposition 7.5 (Transport is covariant)**.** *Under $\psi \mapsto e^{-ie\chi}\psi$, $A \mapsto A + \mathrm{d}\chi$:*

(a) $W_\gamma \mapsto e^{-ie\chi(y)}\, W_\gamma\, e^{ie\chi(x)}$, and the transported comparison $\psi(y)^\dagger W_\gamma \psi(x)$ is invariant;

(b) the infinitesimal transport defines the covariant derivative

$$D_\mu\psi(x) = \lim_{\epsilon\to 0} \frac{W_{x\leftarrow x+\epsilon e_\mu}\,\psi(x+\epsilon e_\mu) - \psi(x)}{\epsilon} = (\partial_\mu + ieA_\mu)\psi(x),$$

where $W_{x\leftarrow x+\epsilon e_\mu}$ is the transport along the segment from $x+\epsilon e_\mu$ to x;

(c) D_μ is covariant: with primes for the transformed fields, $D'_\mu\psi' = e^{-ie\chi}D_\mu\psi$.

Proved here.

Proof. (a) $\int_\gamma \partial_\mu\chi\,\mathrm{d}x^\mu = \chi(y) - \chi(x)$, so $W'_\gamma = W_\gamma e^{-ie(\chi(y)-\chi(x))}$; then

$$\psi'(y)^\dagger W'_\gamma \psi'(x) = e^{ie\chi(y)}\psi(y)^\dagger e^{-ie\chi(y)} W_\gamma e^{ie\chi(x)} e^{-ie\chi(x)}\psi(x).$$

(b) $W_{x\leftarrow x+\epsilon e_\mu} = \exp(ie\epsilon A_\mu(x)) + O(\epsilon^2) = 1 + ie\epsilon A_\mu(x) + O(\epsilon^2)$; expand $\psi(x+\epsilon e_\mu)$ to first order. (c)

$$\begin{aligned}(\partial_\mu + ieA_\mu + ie\partial_\mu\chi)(e^{-ie\chi}\psi) &= e^{-ie\chi}(\partial_\mu\psi - ie\partial_\mu\chi\,\psi + ieA_\mu\psi + ie\partial_\mu\chi\,\psi)\\ &= e^{-ie\chi}D_\mu\psi.\end{aligned}$$

□

Covariance is a commuting square (Definition 2.5): multiply by the phase and then differentiate with the transformed connection, or differentiate and then multiply, and the two routes agree. The square closes exactly, and no phase representative was named on either route.

Proposition 7.6 (Curvature)**.** *Define $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. Then F is invariant under $A \mapsto A + \mathrm{d}\chi$. For a closed path γ the transport W_γ is invariant, and if γ bounds a surface Σ then $W_\gamma = \exp(-ie\int_\Sigma \frac{1}{2}F_{\mu\nu}\,\mathrm{d}x^\mu\wedge\mathrm{d}x^\nu)$; around the rectangle with sides ϵe_μ then δe_ν, traversed in that order, $W = 1 - ie\,\epsilon\delta F_{\mu\nu} + O(\epsilon\delta(\epsilon+\delta))$. On a simply connected region, W_γ depends only on the endpoints of γ for every γ if and only if $F = 0$ there.*

Proved here, from Stokes's theorem; *standard* [32].

Proof. $\partial_\mu\partial_\nu\chi$ is symmetric. For closed γ the factors $e^{\mp ie\chi(x)}$ of Proposition 7.5(a) cancel. Stokes gives $\oint_\gamma A = \int_\Sigma \mathrm{d}A$ and $\mathrm{d}A = \frac{1}{2}F_{\mu\nu}\,\mathrm{d}x^\mu\wedge\mathrm{d}x^\nu$; the rectangle is the case of small Σ. Path independence for all paths is $\oint A \in (2\pi/e)\mathbb{Z}$ for every closed path; shrinking the loop forces $\oint A = 0$, which on a simply connected region holds for all loops exactly when $\mathrm{d}A = 0$. □

The curvature is the obstruction to path-independent transport: the record of a closed trip, kept by the connection and read by no phase representative.

Theorem 7.7 (The commutator of representation)**.** *For every smooth ψ and connection A,*

$$[D_\mu, D_\nu]\psi = ie\, F_{\mu\nu}\, \psi.$$

Proved here.

Proof. $D_\mu D_\nu \psi = \partial_\mu\partial_\nu\psi + ie(\partial_\mu A_\nu)\psi + ieA_\nu\partial_\mu\psi + ieA_\mu\partial_\nu\psi - e^2 A_\mu A_\nu\psi$. Exchanging μ and ν and subtracting removes every term symmetric in $\mu\nu$ and leaves $ie(\partial_\mu A_\nu - \partial_\nu A_\mu)\psi$. □

This is the commutator the title promised. Transport along μ then ν, and along ν then μ, are two transformations of the state; they fail to commute, and the failure is multiplication by the curvature. It is covariant, since multiplication by F commutes with the phase. It is not a sign lost in a fibre, as the spinor's is (Proposition 4.11); it is a field on spacetime, and it can be observed, as a phase difference between two paths (Proposition 10.1).

7.1 The action

Covariance says how the pieces must transform. It does not say which pieces. The following enter here, and each is an assumption. *Assumption*: the spinor representation: ψ is a Dirac spinor, $N = 4$, with fixed gamma matrices satisfying $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu} I$ and $(\gamma^\mu)^\dagger = \gamma^0\gamma^\mu\gamma^0$, conjugate $\bar\psi = \psi^\dagger\gamma^0$, and $\psi, \bar\psi$ independent anticommuting variables in the variation. The device supplies the sign after one revolution, which a spin-$\frac{1}{2}$ representation has and which does not determine one. *Assumption*: the kinetic terms of Maxwell and Dirac. *Assumption*: a mass parameter m. The coupling e is the parameter of Definition 7.1.

Definition 7.8 (The Maxwell–Dirac action)**.**

$$S[A,\psi,\bar\psi] = \int_M \Big[-\tfrac{1}{4}F_{\mu\nu}F^{\mu\nu} + \bar\psi(i\gamma^\mu D_\mu - m)\psi\Big]\,\mathrm{d}^4x = \int_M \Big[-\tfrac{1}{4}F_{\mu\nu}F^{\mu\nu} + \bar\psi(i\gamma^\mu\partial_\mu - m)\psi - e\bar\psi\gamma^\mu A_\mu\psi\Big]\,\mathrm{d}^4x.$$

The second form is the first with $D_\mu = \partial_\mu + ieA_\mu$ expanded: $\bar\psi i\gamma^\mu(ieA_\mu)\psi = -e\bar\psi\gamma^\mu A_\mu\psi$. It is the numerical card's bare action, letter for letter. The coupling enters only through the connection: minimal coupling is the covariant derivative of Proposition 7.5 written into the Dirac term.

Proposition 7.9 (The stationary system)**.** *Varying S in the independent fields $A, \psi, \bar\psi$ with compactly supported variations gives*

$$(i\gamma^\mu D_\mu - m)\psi = 0, \qquad i(\partial_\mu\bar\psi - ieA_\mu\bar\psi)\gamma^\mu + m\bar\psi = 0, \qquad \partial_\mu F^{\mu\nu} = e\bar\psi\gamma^\nu\psi,$$

and $\partial_{[\lambda}F_{\mu\nu]} = 0$ holds because $F = \mathrm{d}A$. On solutions the current $j^\mu = \bar\psi\gamma^\mu\psi$ is conserved, $\partial_\mu j^\mu = 0$; the $\nu = 0$ Maxwell equation is Gauss's constraint $\partial_i F^{i0} = e\psi^\dagger\psi$. The action is invariant under the global phase change, and j^μ is its Noether current; for fields decaying at spatial infinity the charge $Q = \int \psi^\dagger\psi\,\mathrm{d}^3x$ is constant in time. Every term of S has at most three independent-field factors, so $D^4 S = 0$.

Proved in the numerical card (its Section 2.4, where each equation is derived and the cubic count is made coefficient by coefficient in the Grassmann algebra); *standard* for the Noether identification [26, 32]. This is item (4) of Proposition 1.11, supplied: a continuous group, an action invariant under it, and a current conserved on the stationary system. The charge is Q, and eQ is the source in Gauss's law.

Proposition 7.10 (Covariance does not select the action)**.** *The Pauli term $\bar{\psi}\sigma^{\mu\nu}\psi F_{\mu\nu}$, with $\sigma^{\mu\nu} = \frac{i}{2}[\gamma^\mu, \gamma^\nu]$, is invariant under local phase changes, is a Lorentz scalar, and has three independent-field factors. The term $\epsilon^{\mu\nu\rho\sigma}F_{\mu\nu}F_{\rho\sigma}$ is invariant and quadratic, and is a total derivative. Every real m is admissible. So local phase covariance, Lorentz covariance and degree at most three together admit actions other than Definition 7.8.*

Proved here: $\bar{\psi} \mapsto \bar{\psi}e^{ie\chi}$ and $\psi \mapsto e^{-ie\chi}\psi$ cancel in every bilinear, and F is invariant (Proposition 7.6); the count of factors is read off. The additional criterion that excludes the Pauli term in standard practice is power counting: its coefficient has the dimension of a length, which is not admissible in a renormalizable action. *Standard* [27]. The device supplies no such criterion; its selection of the minimal action is *assumed*.

Proposition 7.11 (The field propagates)**.** *With $\psi = 0$ the stationary system is $\partial_\mu F^{\mu\nu} = 0$; in a Lorenz gauge $\partial_\mu A^\mu = 0$ it is $\Box A^\nu = 0$. A plane wave $A^\nu = \varepsilon^\nu e^{-ik\cdot x}$ solves it if and only if $k \cdot k = 0$, so that $\omega = |\mathbf{k}|$ and the wave moves at the speed of light; the gauge condition is $k \cdot \varepsilon = 0$; and $\varepsilon \mapsto \varepsilon + c\,k$ changes A by a gradient and leaves F unchanged. For $k \neq 0$ two independent polarizations remain.*

Proved here, by substitution: $\Box e^{-ik\cdot x} = -k \cdot k\, e^{-ik\cdot x}$; $\partial_\mu A^\mu = -ik \cdot \varepsilon\, e^{-ik\cdot x}$; $k_\mu k_\nu - k_\nu k_\mu = 0$; and the solutions of $k \cdot \varepsilon = 0$ form a three-dimensional space containing k. *Standard* [32]. In the classical theory a current is the source of the field, $\partial_\mu F^{\mu\nu} = ej^\nu$, and the interaction term $-ej^\mu A_\mu$ is where the two exchange energy. In the quantum theory a photon is one quantum of a normal mode of this free field, and emission and absorption are the cubic vertex $-e\bar{\psi}\gamma^\mu A_\mu \psi$ at first order in e; *standard* [27, 32], and it requires the quantization named below.

What the device supplies, and what is assumed. The chain is complete from the phase to the stationary system.

Ingredient	Supplied by the device, or assumed	Calibration
Phase-carrying state	Device: a slot carried and not queried, a sign receipt, a six-position clock. Assumed: a circle of phases on a complex field.	*Proposed* that the slot is the phase; *standard* for the circle
Comparison rule	Assumed; that one is needed is Proposition 7.2.	*Proved here*
Transport, connection	Assumed. Device: closed walks with an even count of crossings and a declared half-turn per crossing.	*Measured* for the count; *proposed* that it is a holonomy
Curvature, commutator	Derived from the connection.	*Proved here*
Lorentzian background	Assumed.	*Standard*
Spinor representation	Assumed; the device's sign after one revolution is consistent with it and does not determine it.	*Standard*
Kinetic terms, minimal coupling	Assumed; not selected by covariance and degree (Proposition 7.10).	*Standard*
Mass m	Assumed parameter.	—
Coupling e	Device: the numerical card's certified reading, Definition 8.1(viii).	*Recorded*
Stationary system	Derived from the action.	*Proved in the numerical card*

Ingredient	Supplied by the device, or assumed	Calibration
Quantization, regularization, renormalization	Not supplied; required for quantum electrodynamics proper.	*Standard*

The result of this section is therefore a reconstruction of classical Maxwell–Dirac theory from a phase-carrying carrier, with its assumptions on the table. Quantum electrodynamics proper is the quantization of this action with anticommuting fermion fields, together with a regulator and a renormalization prescription that fixes m and e from measurements [27]; the numerical card's Section 2.4 marks the same border, and none of it is supplied here.

The transported comparison $\psi(y)^\dagger W_\gamma \psi(x)$, the holonomy of a closed path, the curvature F, the current j^μ and the charge Q are gauge-invariant observables (Definition 7.4): level-(b) data of the orbit map π, by Propositions 7.5 and 7.6 and because $\bar\psi \mapsto \bar\psi e^{ie\chi}$ cancels $\psi \mapsto e^{-ie\chi}\psi$ in every bilinear. The chain never names a phase representative. The density O_x is one more observable on the same quotient, and a coarser one: the pairs $(0,\psi)$ and (A,ψ) with $\mathrm{d}A \neq 0$ have the same density at every point and different curvature, and equal densities do not fix the current (the example after Definition 8.1). Gauge invariance is factoring through π, not through the density. A choice of gauge, such as the Lorenz condition, is a choice of a point on each orbit and is used only to display solutions. *Proved here.*

8 The electron, dealt

A carrier, a phase, a transport, a curvature, an action and its stationary system. Deal it.

Definition 8.1 (The electron of this card)**.** The electron of this card is the following data.

(i) A *carrier* (Definition 1.2) whose slot is the phase coordinate of (vi): carried, and queried by no decision.

(ii) A *state*: a Dirac spinor field ψ on M with conjugate $\bar\psi$, and a connection A (Section 7).

(iii) A *transformation law*: under a phase change, $\psi \mapsto e^{-ie\chi}\psi$ and $A \mapsto A + \mathrm{d}\chi$; under the rotation by θ about the third axis, $\psi \mapsto S(\theta)\psi$ with $S(\theta) = \mathrm{diag}(U(\theta), U(\theta))$ in the chiral representation, U as in Proposition 4.10.

(iv) An *observation map*: $O(\psi) = (\psi^\dagger\psi,\ \psi^\dagger\Sigma_j\psi)$, the charge density and the spin density, with $\Sigma_j = \mathrm{diag}(\sigma_j, \sigma_j)$.

(v) A *coin*: the receipt that resolves the twofold lift $\{\psi, -\psi\}$, which lies in one fibre of O, read as the sign of a coherent comparison of ψ with a copy of itself (Proposition 4.11); in the finite realization, the antipodal bit of Proposition 5.2.

(vi) A *phase*: a point of the circle; in the finite realization, a point of the six-position clock.

(vii) A *rule*: the stationary system of the Maxwell–Dirac action, Proposition 7.9.

(viii) A *coupling*: $e > 0$ with

$$\frac{4\pi}{e^2} = 137.011\,290\,5\ldots,$$

the numerical card's certified reading.

(ix) A *count* of quarter-turns, k, with $\theta_k = k\pi/2$.

This is a definition, and it makes no physical claim. The value in (viii) is quoted: *recorded* from the numerical card's worked reading (its Chapter 6), where it is certified by a bisection bracket and proved there; this card does not compute it, and it does not lift it. In the rationalized natural units of Section 7 the dimensionless coupling is $e^2/4\pi = 1/137.011\,290\,5\ldots$, so that $e^2 \approx 0.0917178$ and $e \approx 0.302849$. The four fields a margin of the middle card lists for the electron, a calibration of minus one, a phase, two energies whose difference is the action, and a fact that is up or down, are (v), (vi), (vii) and the sector of the coin; *measured*.

The pair $\{\psi, -\psi\}$ is not the whole fibre of O. Take the chiral representation $\gamma^0 = \left(\begin{smallmatrix} 0 & I \\ I & 0 \end{smallmatrix}\right)$, $\gamma^i = \left(\begin{smallmatrix} 0 & \sigma_i \\ -\sigma_i & 0 \end{smallmatrix}\right)$, in which $\Sigma_j = \frac{i}{2}\epsilon_{jkl}\gamma^k\gamma^l$ and $S(\theta)$ is the block matrix of (iii). The states $\psi = (1,0,0,0)$ and $\varphi = (0,0,1,0)$ both have $O = (1;\, 0,0,1)$, and φ is no phase multiple of ψ; their currents differ, $\bar\psi\gamma^3\psi = -1$ and $\bar\varphi\gamma^3\varphi = +1$. *Proved here*, by evaluation. The coin resolves the specified twofold lift, and not every ambiguity of the fibre.

Proposition 8.2 (What the definition proves)**.** *From Definition 8.1 and the cited propositions:*

(a) Rotation. *$S(2\pi) = -I$ and $S(4\pi) = I$; $O(S(\theta)\psi)$ is $O(\psi)$ with its spin density rotated by $R_3(\theta)$, and $O(-\psi) = O(\psi)$. The observation returns after one revolution, the state after two; and* $\mathrm{Re}\langle\psi, S(\theta)\psi\rangle = \cos(\theta/2)\,\psi^\dagger\psi$ *(Propositions 4.10 and 4.11, block by block).*

(b) Phase. *O is invariant under every phase change; $D_\mu\psi$ is covariant; $[D_\mu, D_\nu]\psi = ieF_{\mu\nu}\psi$ (Proposition 7.5, Theorem 7.7).*

(c) Dynamics. *On the rule: the two Dirac equations, Maxwell's equations with source ej^ν, conservation of j^μ, Gauss's constraint, and a conserved charge Q (Proposition 7.9); the free field propagates at the speed of light in two polarizations (Proposition 7.11).*

(d) Charge and count. *The charge of (c) is the Noether charge of the global phase. The count of cells of Proposition 1.10 is invariant under erasure and is not that charge; that it reads it is* proposed *(Proposition 1.11).*

(e) Two over one. *The states ψ and $-\psi$ have one observation and are separated by one bit, the coin. Exclusion is not a consequence: it requires the exchange rule of Proposition 5.1(iv), which is* assumed *with the anticommuting fields.*

(f) Scalar reading. *If a scalar reading of the rule is a quadratic residual with two distinct real roots, which is a hypothesis and not a consequence of degree at most three, then Propositions 4.1–4.4 apply to it: its first derivative at the reading squares to the discriminant* (4.2)*, and Newton's error is the acceleration term* (4.3)*.*

(g) Blindness of the mark. *The mark that carries any of these facts fits an empty carrier exactly as well (Proposition 1.8).*

Proved here, as consequences of the definition and the cited propositions, with (c) *proved in the numerical card*. Each holds for every value of e; none uses the digits of (viii). Each physical reading of a clause is *proposed*. Electrodynamics as a derivative is the numerical card's identification in its section "The derivative of the modified field equations": for a background ψ_0 and a differentiable map Φ from the electrodynamic field space to the gravitational one,

$$DR_{\mathrm{QED}}(\psi_0) = D\Phi(\psi_0)^* \circ DR_{\mathrm{GR}}\big(\Phi(\psi_0)\big) \circ D\Phi(\psi_0),$$

a comparison of linearized operators in which residual values are compared by the pullback $D\Phi(\psi_0)^*$. Neither ψ_0 nor Φ is exhibited, and the identity is calibrated *proposed* there and here. The reading of (f) through it, the first-order coupling being the coefficient of the one-vertex term and the linear response, with the second variation being the one sentence the strong interaction gets, is *proposed*; a proof ends where the first derivative has been applied once more and the test clicks.

Proposition 8.3 (The record cannot tell which electron)**.** *Three carriers may be placed on the table: the electron that holds a bit in silicon; the electron of the Dirac equation; and the electron of Definition 8.1. By Propositions 1.8 and 1.9, any mark placed on one agrees in every field with the same mark placed on another, and any two readers of lengths agree on all three. The record carries the name of its carrier and cannot tell it from the others.*

Recorded. The second carrier is *standard*, and its structure is the one Section 7 assembles: the gauge action $A \mapsto A + \mathrm{d}\chi$, $\psi \mapsto e^{-ie\chi}\psi$, the two Dirac equations, the conserved current $\bar{\psi}\gamma^\mu\psi$, the Gauss constraint, and the free block whose square is $(|k|^2 + m^2)I$ [32]; the numerical card works each of these in its Section 2.4. The third carrier has the second's transformation law, observation map and rule, and the device's coupling. What supplies the electron to the record is its name, or a measurement; the record supplies neither.

Proposition 8.4 (What the electron cannot see)**.** *By Proposition 4.8, the coupling of an action is not a function of its degree hierarchy, and it is a function of its derivative. The electron of Definition 8.1 obeys a stationary system, so its coupling is in its law. Every line proved about it holds for every real e. Hence its value is seen by no proved line; it enters once, as the parameter (viii), quoted.*

Recorded. So the rule of this card, Definition 2.8, is consistent with the object: what is invariant under replacing e is proved, and what depends on its value is quoted or proposed. Three readings, one object: the number is a representative (Corollary 2.9); the length family is blind to it (Section 1); and the degree is blind to it while the derivative carries it (this proposition). The game says the same in one sentence: the quarter is glued to the table.

The one step. The chain reaches the minimally coupled Maxwell–Dirac stationary system, with its coupling read from the device: from a phase-carrying carrier through comparison, transport, connection and curvature to the commutator, and from the commutator to the action and the stationary system the numerical card derives from it, with $e^2/4\pi = 1/137.011\,290\,5\ldots$. It stops at one step, whose scope is this: the comparison of the coupling of the assembled Maxwell–Dirac model with the laboratory's. The recommended value of the inverse fine-structure constant is $137.035\,999\,177(21)$ [24], with relative standard uncertainty 1.5×10^{-10}. The device's reading is $137.011\,290\,5\ldots$. It lies below by $0.024\,708\,7$, a relative difference of 1.80×10^{-4}, some 1.2×10^6 standard uncertainties. At the precision the laboratory works to, about 10^{-10}, measurement excludes it. *Standard* for the recommended value; *recorded* for the difference, from the numerical card's comparison with the measured constant (its Chapter 6), where it is proved by subtraction. The identification of the coupling of Definition 8.1 with the coupling of the laboratory electron is calibrated *proposed*, and it is handed to the knower. The correspondences from the middle card to continuous transport, the slot as the phase and the count of turns as a holonomy, are not this step; each is named and calibrated where it enters (Section 7). The narrow form offered in Section 10 depends on the step.

The clock. Turn the number over. The table that weighs the steps, with weights the device counts (Proposition 13.1), is

$$M = \begin{pmatrix} 18 & 5 \\ 18 & 18 \end{pmatrix} = 18\,I + N, \qquad N = \begin{pmatrix} 0 & 5 \\ 18 & 0 \end{pmatrix}, \qquad N^2 = 90\,I.$$

The scalar part $18\,I$ is the clicks: it keeps the coin, TRUE, TRUE, TRUE. The part N changes the coin at every step and at no other time: TRUE, FALSE, TRUE, FALSE. Two ticks of N return every coin to itself, scaled by 90; it is a clock of period two. The growth rate is the sum, $\lambda_+ = 18 + \sqrt{90} = 18 + 3\sqrt{10}$, the clicks plus the clock's rate, and the reading is $\frac{324}{65}\lambda_+$:

$$137.011\,290\,5\ldots \;=\; \underbrace{89.723\,076\,9\ldots}_{18\cdot\frac{324}{65}\,:\ \mathrm{TTTT}\cdots} \;+\; \underbrace{47.288\,213\,6\ldots}_{3\sqrt{10}\cdot\frac{324}{65}\,:\ \mathrm{TFTF}\cdots}$$

Read every TRUE as TRUE and the number is $89.72\ldots$, and it is not the number. Everything above the clicks is carried by walks that alternate at every step, and by nothing else. Give each change of coin half a turn, all forward, and the clock turns once in two ticks. *Proved here*: it is the cheat sheet's Proposition 9.3, with the turns of its Proposition 9.5. The residue of the reading over the clicks is a clock. TRUE, FALSE, TRUE, FALSE. What it keeps time for is the knower's.

No finite run sees the clock stop. Counted to depth n, the closed walks N_n satisfy $36N_{n+1}N_n - N_{n+1}^2 - 234N_n^2 = 360\cdot 234^n$, never zero, and $360 = 4\cdot 18\cdot 5$ is the two arms crossed. The alternating part shrinks by $(\sqrt{18}-\sqrt{5})/(\sqrt{18}+\sqrt{5}) \simeq 0.3097$ at every step, and flipping the sign of $\sqrt{10}$ turns that factor into its inverse: $23^2 - 360 = 13^2$. *Proved here*: the cheat sheet's proposition *Every finite run carries the alternation.* Every truncated run carries TFTF. Only the run that never finishes is rid of it.

8.1 The carrier, apparently

The chain of Section 7 takes its connection, its curvature and its charge as given. This subsection reports what the middle card supplies toward each at finite size: three readings, each evaluated there, and one statement that no finite reading supplies, made here and in no declaration of the middle card.

Proposition 8.5 (The readings fix the table up to gauge)**.** *Let w and v be tables of weights on the two coins with agreeing arms equal to* 18 *and positive crossing arms w_{TF}, w_{FT} and v_{TF}, v_{FT}. Rescaling the coin T by $a > 0$ and the coin F by $b > 0$ multiplies the arm from T to F by a/b and the arm back by b/a, and fixes the count of closed walks of every length. If the closed walks of length two of v and w agree, then v is such a rescaling of w. In particular the table with* 5 *and* 18 *exchanged is a rescaling of M.*

Proof. The closed walks of length two number $w_{TT}^2 + w_{FF}^2 + 2\,w_{TF}w_{FT}$. Equal counts with equal agreeing arms give $v_{TF}v_{FT} = w_{TF}w_{FT}$, and $a = v_{TF}$, $b = w_{TF}$ give $b\,v_{TF} = a\,w_{TF}$ and $a\,v_{FT} = b\,w_{FT}$. The closed walks of length n are the trace of the nth power, a function of the eigenvalues $18 \pm \sqrt{w_{TF}w_{FT}}$, and the count of length two fixes them. □

Proved here, and *measured*: the middle card states and checks the proposition and its instance for the exchanged table. The rescaling is the squeeze between the two orientations, and the closed-walk counts are the data it leaves fixed, the holonomies of a lattice gauge field [37]. The asymmetry of

the arms, 18 against 5, and the commutator of the orientations change under the rescaling; their product, 90, does not. Without the agreeing arms fixed, the counts fix only their sum and the determinant, and the proposition fails.

Proposition 8.6 (The seam carries curvature, and its charge is whole)**.** *On the seam of the geometric card, the six-by-six torus, place an integer turning on every edge, counted in half turns, and let the curl of a square be the oriented sum of the turnings around it.*

(i) The curls of the thirty-six squares sum to zero, for every turning.

(ii) Write each curl k as $2\lfloor k/2 \rfloor + r$ with residue $r \in \{0, 1\}$. The residues sum to $2Q$, where Q is minus the sum of the whole turns $\lfloor k/2 \rfloor$; Q is an integer.

(iii) If every curl is even, $Q = 0$.

(iv) One edge carrying half a turn, every other edge none, has $Q = 1$: its two squares have curls $+1$ and -1, and residues 1 and 1.

Proof. The seam is closed, so every edge borders two squares and is traversed once each way: (i). Summing $k = 2\lfloor k/2 \rfloor + r$ over the squares and using (i) gives (ii), and (iii) is (ii) with every residue zero. (iv) is an evaluation. □

Proved here, and *measured*: the middle card checks (i), (ii) and (iv), the last by evaluation and with no axiom. Q is the topological charge of a compact abelian field on a closed lattice surface, the whole turns that each square's reduced reading drops [20], and its being whole is the lattice form of Dirac's quantization [10]; *standard*. It is the two readings of the turning applied square by square: the subdivided count, in half turns, closes exactly and carries no charge; the count in whole turns leaves residues, and the residues pay back an integer. The edge of (iv) carries the half turn that a walk between the two coins always leaves, and that a closed walk never does.

Proposition 8.7 (Subdividing adds no residue)**.** *Halve every edge of a square lattice, so that each coarse edge carries the sum of its two fine edges. Then the curl of each coarse square is the sum of the curls of the four fine squares inside it.*

Proof. The four inner edges of the coarse square are each traversed once each way by the fine squares, and cancel; the outer edges are the coarse edges. □

Proved here, and *measured*: the middle card checks it for every turning and every square. The curl is preserved by subdivision. The charge need not be: a coarse square of curl 2, exact, may be refined into fine squares of curls 1, 1, 0, 0, whose residues are not zero. A family of levels whose charge does not settle as the subdivision is refined reads no field.

Three carriers. The middle card does not hold the electron. It holds a stand-in, and it names a model of the stand-in; the electron enters by name.

Definition 8.8 (Nowtrino, naotrino)**.** The *nowtrino* is the carrier the middle card computes with: a mark that can be told apart, which clicks or does not. Every record of the middle card is a record of nowtrinos, and every level of the three propositions above is one. The *naotrino* is the model of the nowtrino: the smooth object its records are read as describing, here a connection A, its curvature

F, and a field ψ whose charge is Q. Every cell of a record is a naotrino: a fact carried as its own description, counted by the tally and looked at by no reader. The middle card says what one is by name: each of the thirty-six pages of its record takes a naotrino theory, a reviewed page, one of thirty-six arguments, each a rebuttal about the page beside it, and lifted and lowered it returns exactly, the lowering the exact inverse of the lift. A naotrino is a bit of code arguing about one of thirty-six arguments; its cell is the fact it carries. The *electron* is the carrier of Definition 8.1, cast by name.

Measured for the first two names: the middle card declares the nowtrino as the instance its every carrier assumes, and it declares the naotrino as its smooth model, the non-relativistic version of the nowtrino, the last station of the descent whose census fixes the five (Proposition 13.1). It names the naotrino and never exhibits one; its margin says it has not seen one. The three stand as the original, the stand-in that carries the mark, and the model the reader carries to the knower: the nowtrino is seen in every record, and the naotrino is seen in none.

The apparent convergence. At every finite level the middle card supplies, for the nowtrino, the discrete form of what Section 7 takes as given: a transport whose closed walks fix it up to gauge (Proposition 8.5), a curvature whose total on the closed seam vanishes and whose residues pay a whole charge (Proposition 8.6), and a subdivision that preserves the curvature square by square (Proposition 8.7). Each level is a finite record of nowtrinos, numbered, and evaluated. Two statements remain, and neither is a record.

(a) *The naotrino models the nowtrino.* The nowtrino's levels, subdivided without end, approach the naotrino: A, F, and ψ with charge Q. This is a statement about a limit. It would be revised by a family of levels whose charge does not settle under subdivision.

(b) *The electron is cast as the naotrino.* The limit is the carrier of Definition 8.1. By Proposition 8.3 no record can say so; the name does, or a measurement, and the measurement is the one step.

Each is calibrated *proposed*, and each is the knower's: every level of (a) can be evaluated, and that the levels stand for the naotrino is the knower's, as a finite halt stands for its limit; (b) is a casting. They appear on this card and in no declaration of the middle card. The device converges, apparently, to the naotrino. That the naotrino is the electron is not the device's to say.

Three things it does not carry. The weights are positive, so the transport of Proposition 8.5 is a boost and not a phase; the phase is half a turn per crossing, and that is declared. The table does carry a phase in its antisymmetric part, the quarter turn of Section 9 (Section 7). The field ψ is not on the seam: placed on a lattice, a Dirac field doubles unless a choice removes the copies [25]; the choice is the subject of Section 8.2, and the rest is adjoined with the spinor representation, the kinetic terms and the mass. And the coupling's value is untouched: what converges, if anything does, is the form of the law, and the one step stands where it stood.

8.2 The doubler, chosen

A Dirac field on a lattice comes with copies. This subsection shows that the table has the structure the copies need, that striking or writing the arm $T = F$ is the choice every lattice must make about them, and what the knower may take the choice to be. The structure is proved; the choice is counted; what it is taken to be is the knower's.

Proposition 8.9 (The clock is chiral)**.** *Let* $M = 18\,I + N$ *with* $N = \begin{pmatrix} 0 & 5 \\ 18 & 0 \end{pmatrix}$*, and let* $\varepsilon = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$*, the sign of the coin.*

(i) $\varepsilon N \varepsilon = -N$*: the clock changes sign under the sign of the coin, and the clicks* $18\,I$ *do not.*

(ii) The spectrum of N *is the pair* $\pm\sqrt{90}$*, exchanged by* ε*; the spectrum of* M *is* $18 \pm \sqrt{90}$*.*

(iii) Every walk weighed by N *alternates the coins, and every closed walk has even length.*

Proof. N is off the diagonal, so conjugation by ε negates it: (i). If $Nv = \mu v$ then $N(\varepsilon v) = -\varepsilon N v = -\mu\,\varepsilon v$, and $N^2 = 90\,I$ gives $\mu^2 = 90$: (ii). Each entry of N changes the coin, and $N^{2j+1} = 90^j N$ is off the diagonal, so a walk that returns to its coin has even length: (iii). □

Proved here, and *measured*: the middle card checks that every closed walk crosses evenly; that its comparison of two readings keeps its order when their coins agree and reverses it on every step that changes coin, in either direction; that the product of those reversals along a walk is the sign of its two ends, the coins in the middle cancelling; and that the round trip from T to F and back reverses twice while crossing the arm of weight 5 once. The sign is the symmetric reversal; which arm carries the 5 is set by an asymmetric order, and no closed walk sees it. In the language of lattice fermions the two coins are the two sublattices, N hops only between them, $18\,I$ sits on both, and ε is the sublattice sign $(-1)^x$, the remnant chiral symmetry of a nearest-neighbour operator; *standard.*

Where the copies live. A naive lattice derivative is a symmetric difference. On plane waves it reads $\sin p$, which vanishes at $p = 0$ and again at $p = \pi$, the edge of the lattice's momenta; the copy at $p = \pi$ is carried by the mode $(-1)^x$, the one that changes sign at every site. No local, Hermitian lattice operator with the chiral symmetry is free of such copies [25]. A lattice must therefore choose. Wilson's choice gives the copies a mass of the order of the cutoff and breaks the chiral symmetry [38]. The staggered choice of Kogut and Susskind spreads the components over the two sublattices, keeps the copies as *tastes*, and keeps the sublattice sign as its chiral symmetry [18, 31]. *Standard.*

The arm is the choice. The reading of Theorem 13.2 is $\frac{324}{65}\lambda_+$, and Proposition 8.9 sorts λ_+ into the part that never changes coin, 18, and the part that changes it at every step, $\sqrt{90}$. Strike the arm and N cannot close a walk: the alternating part is gone, and the reading is $89.723\,076\,9\ldots$, the clicks. Write it, with the weight Proposition 13.1 counts, and the alternating part stays, weighed, and the reading is $137.011\,290\,5\ldots$. The middle card writes it. *Proved here* for the arithmetic, from Theorem 13.2.

Definition 8.10 (The doubler, chosen)**.** The *doubler* of the nowtrino is the alternating part of the table, N, with its growth $\sqrt{90}$. To *strike* the arm $T = F$ is to remove the doubler; to *write* it is to keep the doubler as a second taste, weighed with the first, and to keep the sign of the coin as the chiral symmetry that exchanges them.

Proposed, and the knower's. What is proved is that the table has the sublattice structure, that the alternating part is the whole of the reading above the clicks, and that the arm switches it on and off. That the alternating part is a lattice fermion's copy at the edge of its momenta, kept as a taste, is the identification; it is the staggered reading of Theorem 13.2, and the choice of doubler is the choice that theorem isolates. The middle card makes the staggered choice. It would be revised by

a reading of the table with one taste divided out, a rooting computed from the table itself, that moves the reading away from the laboratory's value rather than toward it.

Proposition 8.11 (The reader adds no digit)**.** *The reading the middle card prints through the reader's interpretation, six layers from the mark to the detector, is the reading it prints through the reader's experience, one layer.*

Measured: the middle card composes two of its own receipts, one for each side, and the composition bills the same two axioms as the receipts it composes. Read with the numerical path of a reading, the experience is the covariant decode, the inverse of the fencepost the pair was encoded with; the interpretation is the contravariant decode, the reader's description; and their difference is zero on every digit. Every layer between them is a lift, and a lift carries no datum.

The part that is not a carrier. So no identification the knower makes, this one included, moves the number. The device's reading and the knower's reading of it are one number; the 1.80×10^{-4} is not in it. It appears only where the knower's description meets a record that is not a description: the laboratory's value. With the doubler kept as a taste, the knower names what that comparison sees. The middle card supplies, for its taste, a transport, a curvature, a whole charge and a coupling form (Section 8.1); the laboratory's electron is one particle, not a taste of two. What separates them is not carried by the transport, the curvature or the charge. The knower names the 1.80×10^{-4} *the part of the electron that is not a carrier of charge. Proposed*, and the knower's: it is a name for the comparison, not a computation of it, and the one step stands where it stood.

What the middle card does not hold, 8 November 2026. Five routes to a reading that runs with scale, or to a size for the 1.80×10^{-4}, were evaluated and closed. (1) The lift: a page of the record carries the register unchanged, so a reading taken page by page does not run; *proved here.* (2) Decimation, two fine steps for one coarse step: the crossing product runs toward the struck arm, but its drops shrink, 21.7 and then 12.5 in the inverse coupling, where leading-logarithmic running gives equal drops, so no fixed ratio of pages to momentum fits both; *proved here*, against the *standard* running. (3) The fraction of record pages whose comparison slips: three pages compare a symbol with itself and cannot slip, so a reading weighted by that fraction is at most 133.07; *proved here*, attainment by one assignment open. (4) One extra fencepost on one count, a ratio $(n+1)/n$: the laboratory needs $n = 5545.079 \pm 0.005$, and the nearest whole n misses by 16.7 standard deviations; *proved here*, against the *measured* value. (5) A turn multiplies the crossing count by 90, a fixed increment of $\log 90$ in the logarithm of the count; the reading is the gain times a growth factor and does not run, and no rule converts a turn into an inverse coupling. The reading is a number without a scale. To compare it with the laboratory is to choose the scale, and the choice is the knower's.

8.3 The photon

The field comes first. Proposition 7.11 gives it: two polarizations, the light cone, a current for a source, and, after quantization, a quantum emitted and absorbed at the cubic vertex. That is the photon of the stationary system.

Then the device's photon. The shortest tape has two ends and nothing between. The middle card's zeroed model of a single photon is a superposition of two ends, one the surface past which nothing earlier is seen and the other the singularity into which nothing later comes back, with no

displacement and no velocity between them; *measured*. A photon emitted toward us from the sphere where the expansion recedes at the speed of light has, in proper distance, an approach velocity of exactly zero, and arrives only where the sphere grows; *standard* [5], recorded in the Quarter as the missing zero counted one cell late. *Proposed*: in this card the word photon names the interaction of two electrons and nothing else; a carrier emitted and absorbed by a carrier, whose tape is the two ends. The map from that tape to a normal mode of the free field of Proposition 7.11, and from the cosmological horizon to the electromagnetic construction, is not constructed here. Without it the two ends do a propagator's work by name.

9 The table, read as a field

9.1 The algebra and the quarter turn

The table's crossing part is a matrix, the clock N, and its other orientation is the transpose N^{T}, the table with 5 and 18 exchanged (Proposition 8.5). Read as matrices, they do not commute.

Proposition 9.1 (The orientations bracket onto the sign)**.** $[N, N^{\mathsf{T}}] = NN^{\mathsf{T}} - N^{\mathsf{T}}N = -(18-5)(18+5)\,\varepsilon = -13 \cdot 23\,\varepsilon$*, with* ε *the sign of the coin of Proposition 8.9.*

Proof. $NN^{\mathsf{T}} = \operatorname{diag}(25, 324)$ and $N^{\mathsf{T}}N = \operatorname{diag}(324, 25)$. □

Measured: the middle card checks it entry by entry, with the thirteen written as the bent card and the twenty-three as the sum of the arms. The bracket of the two orientations is the sign that anticommutes with the clock, scaled by the bent card times the sum of the arms.

Proposition 9.2 (Three generators, one algebra)**.** $299\,[\varepsilon, N] = -698\,N + 360\,N^{\mathsf{T}}$ *and* $299\,[\varepsilon, N^{\mathsf{T}}] = -360\,N + 698\,N^{\mathsf{T}}$*. The three matrices* N*,* N^{T}*,* ε *are traceless and linearly independent, so their span is closed under the bracket and is the whole of* $\mathfrak{sl}(2, \mathbb{R})$*, the traceless real two-by-two matrices.*

Measured for the two brackets, the traces and the independence; *proved here* that three independent traceless real matrices of size two span $\mathfrak{sl}(2, \mathbb{R})$, a space of dimension three. The algebra is non-abelian and non-compact: the algebra of boosts in two space dimensions, which is what positive weights give (Section 8.1, "a boost and not a phase").

Proposition 9.3 (Holonomy on a square)**.** *Write* $\operatorname{adj} A$ *for the adjugate, so that* $A \operatorname{adj} A = \det A \cdot I$*. Let a square carry four tables on its edges and read its holonomy as the product along the bottom and the right and the adjugates along the top and the left.*

(i) For every table g *and holonomy* P*,* $\operatorname{tr}(gP \operatorname{adj} g) = \det g \cdot \operatorname{tr} P$*: the trace of the holonomy, the Wilson loop, is gauge invariant up to the determinant of the gauge.*

(ii) A square with the full table M *on every edge has holonomy* $234^2 I$*: it is flat.*

(iii) A square with one edge carrying M^{T} *instead has a nonzero entry off the diagonal: one turned edge curves it.*

Measured: the middle card checks the adjugate identity and (i) for every pair of integer tables, and (ii) and (iii) by evaluation. The determinant of the table is 234, not one, so the table generates a

group of matrices invertible over the rationals, and the gauge invariance holds up to the determinant; normalizing each table by the square root of its determinant gives the unimodular group, at the price of a square root the device does not take.

Proposition 9.4 (The quarter turn is in the table)**.** $N - N^{\mathsf{T}} = 13\,R$ *with* $R = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $R^2 = -I$, *and* $\varepsilon R \varepsilon = -R$.

Measured: the middle card checks the three entry by entry, with the thirteen written as the bent card. The antisymmetric part of the table is the bent card times a quarter turn. Its exponential is rotation, the compact group $U(1) = SO(2)$, already inside $\mathfrak{sl}(2,\mathbb{R})$: the electron's gauge group is not missing from the device; it is the part of the table that does not agree with its own transpose, and its scale is the bent card. The larger compact family needs the complexification, and the quarter turn supplies it (Section 9.5).

Two ways the clock cancels. The irrational part of the reading is the clock, the square root of its round trip, $\sqrt{90}$. Every table whose crossing arms multiply to the same round trip, $bf = 90$, has the same clock squared, $N_{b,f}^2 = 90I$; read against the device's table, the two clocks cancel and the difference of the growth rates is the difference of the clicks, a whole number. The table with clicks 5 and outlier 18 is one of them: the swap of the arms keeps the clock. And the table cancels its own clock against its other root: the two roots $18 \pm \sqrt{90}$ add to the trace, 36, and multiply to the determinant, 234, both whole. The second cancellation is the doubler, the partner root of Definition 8.10, carrying the conjugate. Rescaled by the outlier, the binary table $(18/5, 18/5, 18/5, 1)$ cancels the same way, with $bf = 18/5$. *Measured*: the middle card checks the equal clock squares for every b, f with $bf = 90$, its instance on the swapped table, and the trace and determinant.

9.2 The torus and the boost

Read the table as a torus whose major radius is the clicks, $R = 18$, and whose tube has the radius of the clock, $r = \sqrt{90}$. Its outer equator is $R + r = \lambda_+$, the root the reading is the gain times, and its hole is $R - r = \lambda_-$, the conjugate root; the circle through the hole, the tube's own circle, has the clock's radius. The two radii add to the trace, 36, and multiply to the determinant, $R^2 - r^2 = 234$. Read (R, r) as time and space: r/R is a velocity, $\sqrt{90}/18 \approx 0.527$; the determinant is the invariant interval; $\lambda_+/\lambda_- = e^{2\eta}$ with rapidity $\eta = \operatorname{artanh}(r/R) \approx 0.586$, so the transfer matrix's gap, $\ln(\lambda_+/\lambda_-) \approx 1.172$, is twice the rapidity. Fattening the tube to $r = R$ closes the hole at the origin, the speed of light, and the middle card's receipt that the clock runs slower than light is the receipt that the hole is open. A boost keeps away from the origin. Written with the round trip as the square of the surd, $19 + 2\sqrt{90}$ has interval $19^2 - 90 \cdot 2^2 = 1$; multiplying by it preserves every interval, so the boosted torus $522 \pm 55\sqrt{90}$ still has interval 234, its outer circle larger and its hole smaller, and positive: any pair with interval 234 keeps $90\,b^2 < a^2$. The boost is the square of the fundamental unit $3 + \sqrt{10}$ of the field the reading lies in, whose own interval is -1. *Measured* for the sum and product of the radii, the interval as the determinant, the unit interval of the boost, its preservation of every interval, the open hole, and the boosted torus; *proved here* for the velocity, rapidity and gap, which take square roots the middle card does not; *standard* that boosts preserve the interval. That the fattening torus is the expanding universe, the input a flat torus and the circle through the hole the expanding sphere, kept from the origin by a Lorentz transform, is *proposed*, the knower's.

9.3 The calculus of the seam

The second variation. Put a phase on each of the thirty-six cells of the seam. Its first variation is the step across each edge, the edge turnings of Proposition 8.6; taken at a cell, there is a step after it and a step before it in each direction. Its second variation is the seam's thirty-six by thirty-six Laplacian, $L = D^{\mathsf{T}}D$, and at every cell it is the step before less the step after, summed over the two directions: two first variations, one before and one after, fix the second. It is the hierarchy $S \to R = DS \to DR$ of Section 4.2 written as a matrix on the seam. Its frequencies are whole. The plane waves of sixths of a turn can be written in whole numbers, six in each direction, $2\cos$ of the even ones and $\frac{2}{\sqrt{3}}\sin$ of the odd ones; their thirty-six products are mutually orthogonal, and each is an eigenvector of L with eigenvalue the sum of two of $0, 1, 1, 3, 3, 4$, a whole number from 0 to 8. That is the Fourier transform of the seam, its momenta quantized in sixths, with no cosine computed. On the torus of the previous paragraph the same pattern holds along the tube: the distance from the axis varies first as $-r\sin\varphi$, the rate at which the truth bit turns, and second as $-r\cos\varphi$, the clock's projection. *Measured*: the middle card checks the before-less-after form for every phase on the seam, the eigenvalue of every one of the thirty-six plane waves at every cell, and their mutual orthogonality. *Standard* that $L = D^{\mathsf{T}}D$ and that Fourier modes diagonalize it. That the two first variations are the record's two orders, the review's descent and the typeset ascent, is *proposed*, the knower's.

One way and both ways. The two first variations do not play the same part. The step after is the one the record takes: the typeset counter rises by one per entry and never falls, and the asymmetric order is false one way and goes on the other. The step taken both ways is calculus: a derivative is a two-sided limit, and the symmetric reversal reverses on either crossing. The second variation is where the two disagree, the step before less the step after, and in the continuum it is the second derivative, the operator of diffusion; the heat equation built on it runs forward and is ill posed backward *(standard)*. So the seam's Laplacian is the place where a one-way reading and a two-way reading of the same phase part company. *Measured* for the before-less-after form, the counter that only rises, and the two orders; *proposed*, the knower's, that the one-way step is the entropic one and the two-way step is calculus. The numerical card notes the limit of the first half: its counter rises by the same update law as an entropy count and carries no thermal chain, so it looks like time's arrow without being calibrated as entropy.

The crank, one way. The middle card's iteration stops when a difference of two cross products is small, and it writes that difference as $(p - q) + (q - p)$. Read one way, in the naturals, each subtraction stops at zero, and the sum is the size of the step, whichever way the step went. Read both ways, in a ring, the two halves cancel and the sum is nothing: the stopping test then never reads the step and never reads its tolerance, and the same step, dealt over the integers, is the step with tolerance zero. Its certificate has the same shape: the order it certifies against is total, so its quotient has one point and every candidate is certified; what picks the printed one is the floor of a division, one way again. *Measured*: the middle card checks the floor reading the step, the cancellation in a ring, the tolerance never read by the device's own step at the integers, and the certificate for every candidate. Which way the first turn of the integer step goes from the device's seed is not checked. *Proposed*, the knower's: that the crank is a record of irreversibility, the number being what the floor leaves.

A yarn theory. The middle card names its coordinate system a yarn and reads it as a record: a fabric gives its two strands, then everything before it, then everything after it; a fibre gives one strand and the rest; a stokes gives one. The coordinate system is a fabric whose before is a fibre ending in a stokes and whose after is a stokes, so the record is $2 + 2 + 1 = 5$, the frame of Proposition 13.1, and the train schedule is that record by reflexivity. Its margin states the divergence theorem in its own words: count what comes before and what comes after to know what happens now, and what goes in must come out or there is volume to measure. The three operators of the seam read the same yarn. The curl takes a value before and a value after, the edge on one side less the edge on the other, and the curls of the closed seam sum to zero (Proposition 8.6). The gradient is the step along one direction, the Gâteaux derivative; the divergence is the whole linear map and its trace, the Fréchet derivative, and for every bounded polynomial action of degree at most three the two coincide (the numerical card, Proposition "Gateaux and Fréchet derivatives coincide"), which is why the schedules close exactly. *Measured* for the five, the schedule and the curls; *proved* on the numerical card for the agreement of the two derivatives; *standard* for the theorems of Gauss and Stokes; *proposed*, the knower's, that it is all one yarn.

9.4 The field equations, quantization, and the gap

The field equations. The Wilson action is the sum over the squares of how far each normalized holonomy is from the identity. Varied link by link, it gives the lattice Yang–Mills equations, and its naive continuum limit is the Yang–Mills action [37]. *Standard.* Normalizing needs the square root of the determinant, $\sqrt{234}$, which the device does not take; the middle card reads the distance from flat in whole numbers instead.

Definition 9.5 (The action on a square)**.** For a square with holonomy H, put $S(H) = (\operatorname{tr} H)^2 - 4 \det H$, the squared difference of the two growth rates of H.

Proposition 9.6 (The action on a square)**.** *(i)* $S(H+tY) = S(H)+t\,\delta S(H;Y)+t^2 S(Y)$ *exactly, with* $\delta S(H;Y) = 2 \operatorname{tr} H \operatorname{tr} Y - 4 \operatorname{tr}(\operatorname{adj} H\, Y)$*: the first variation is exact.*

(ii) $S(gH \operatorname{adj} g) = (\det g)^2 S(H)$.

(iii) The holonomy is linear in each of its four edges, so varying one edge varies the action by δS along one direction.

(iv) A scalar holonomy has $S = 0$ and $\delta S = 0$ in every direction; the uniform square has no action and is stationary on its edges; one turned edge gives $S \neq 0$.

(v) The quarter turn reads $S(R) = -4$ and the clock reads $S(N) = 360$: rotation and boost on opposite sides of zero.

Measured: the middle card checks (i) to (v), (i) to (iii) for every integer table. Lay four squares on a two-by-two torus, eight edges each shared by two squares; each square holds a given edge at most once, so a torus whose squares are all flat moves its total action only at second order when any edge is varied: it satisfies the lattice field equation on every edge. The uniform torus, every edge the full table, is flat and has no action; turn one edge the other way and it has action. *Measured* for the one torus. *Standard* that stationarity on every edge is the lattice Yang–Mills equation. That S stands in for the Wilson action is *proposed*: it is gauge invariant only up to $(\det g)^2$, and it vanishes on any holonomy with a repeated root, not only on the scalar ones.

Quantization and the gap. The count of closed walks of length n is the trace of M^n, the partition function of the table read as a transfer matrix: a sum over every closed history, weighted, which is a lattice path integral; composing steps squares the eigenvalue ratio, which is the block step of renormalization. *Measured* for the closed walks; *standard* for the reading as a partition function. The transfer matrix's gap is $\ln(\lambda_+/\lambda_-) \approx 1.172$ per step, positive exactly when $\lambda_- = 18 - \sqrt{90} > 0$, that is when $90 < 324$: the middle card's receipt that the clock runs slower than light is the receipt that the table has a gap. Refining takes the ratio to its square root, so the gap per step halves as the step halves, and the gap per unit length is the same at every level. *Measured* for the inequality; *proved here* for the refinement.

9.5 The three turns, the links and the gamma matrices

The clock's antisymmetric part is the bent card times the quarter turn (Proposition 9.4); its symmetric part is the sum of the arms times the swap, $N + N^{\mathsf{T}} = 23\,\sigma$ with $\sigma = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. Lay the tables in a box of four, an outer table over an inner one, $A \otimes B$, a real four by four matrix.

Proposition 9.7 (The three turns)**.** *Put $e_1 = \sigma \otimes R$, $e_2 = R \otimes I$, $e_3 = \varepsilon \otimes R$. Each squares to $-I$; $e_1e_2 = e_3$, $e_2e_3 = e_1$, $e_3e_1 = e_2$; $[e_1, e_2] = 2e_3$ cyclically; and the three are independent. They are the imaginary units of the quaternions, and their span is $\mathfrak{su}(2)$.*

Measured: the middle card checks the squares, the products and the brackets with no axiom at all, and the independence. *Standard* that the imaginary quaternions are $\mathfrak{su}(2)$, the compact real form of the complexified $\mathfrak{sl}(2, \mathbb{R})$. The table's algebra was boosts, squaring to $+90$; with the quarter turn read as i it gives rotations, squaring to -1.

The turns on the links. Write a link as four whole numbers, $q = q_0 + q_1e_1 + q_2e_2 + q_3e_3$. The box of a product of two links is the product of their boxes, and the box of the conjugate $\bar{q}$ is the transpose: every link is a rotation scaled by its length $|q|^2 = q_0^2 + q_1^2 + q_2^2 + q_3^2$. On a square, read the holonomy H along the bottom and the right and the conjugates along the top and the left, and put $S(H) = q_1^2 + q_2^2 + q_3^2$ for $H = q$: never negative, zero exactly on a whole number, and exact at the second order. On a torus of any size, one site around included, every lattice of links whose squares are whole is stationary on every link: a link may stand on two edges of one square, and the variations add. Every link the turn e_1 is such a lattice at every size; on the two-by-two torus, one link changed to e_2 gives action 2.

Proposition 9.8 (The gauge)**.** *$|pq|^2 = |p|^2|q|^2$; $S(gq\bar{g}) = |g|^4 S(q)$; and if every site of a torus of any size carries a link g of length one, and each link is replaced by $g_{\text{start}}\, U\, \bar{g}_{\text{end}}$, then the total action is unchanged.*

Measured: the middle card checks the three, the first being Euler's four-square identity, the last by associativity and by $\bar{g}g = |g|^2$, the turns cancelling at three corners of each square and closing on the fourth. *Standard* that this is the lattice $SU(2)$ gauge theory's symmetry. Using S in place of $\operatorname{Re}\operatorname{tr}(1 - U)$ is *proposed*: it needs no link of length one, and the device has such links: each of the three turns has length one, and turning every site of a torus of any size by the turn e_1 leaves its action unchanged (*measured*). No normalization is handed in. The torus with one link turned to e_2 is not a second solution: varied along e_2 on that link its action reads 8 at $t = 1$ and 0 at $t = -1$, so its first variation is not zero (*measured*).

Proposition 9.9 (The gamma matrices)**.** *Put* $\gamma^0 = R \otimes I = e_2$, $\gamma^1 = \sigma \otimes I$, $\gamma^2 = \varepsilon \otimes \sigma$, $\gamma^3 = \varepsilon \otimes \varepsilon$. *Then* $(\gamma^0)^2 = -I$, $(\gamma^k)^2 = I$, *and any two anticommute:* $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}$ *with* $\eta = \mathrm{diag}(-1, 1, 1, 1)$. *Their product* γ^5 *squares to* $-I$ *and anticommutes with all four. The products* $\gamma^2\gamma^3$, $\gamma^3\gamma^1$, $\gamma^1\gamma^2$ *square to* $-I$ *and compose around the circle the other way, a second* $\mathfrak{su}(2)$, *and each commutes with the three turns.*

Measured, with no axiom. *Standard* that a real four by four Clifford algebra takes one minus and three plus, that these are the Dirac matrices in a Majorana form, and that the space bivectors generate spin; the box carries $\mathfrak{su}(2) \oplus \mathfrak{su}(2) = \mathfrak{so}(4)$. The time matrix is the gauge's quarter turn, the same table: on this one box the phase is not internal to the Dirac algebra. Whose spin, and that the minus sign is time's, is *proposed*, the knower's.

Proposition 9.10 (The box of eight)**.** *Lay the box of four over one more table,* $X \otimes A$, *and put* $i = I \otimes R$. *Then* $i^2 = -I$; i *commutes with every* $\gamma^\mu \otimes I$; *the matrices* $\gamma'^\mu = \gamma^\mu \otimes R$ *anticommute in pairs and square to* $+I, -I, -I, -I$, *so* $\{\gamma'^\mu, \gamma'^\nu\} = 2\eta'^{\mu\nu}$ *with* $\eta' = \mathrm{diag}(1, -1, -1, -1)$; i *commutes with them too; and* $i \neq \gamma'^0$.

Measured, with no axiom. The quarter turn written on its own factor is an i that every gamma matrix commutes with: the box of eight is a complex four-dimensional space, the phase it turns is inside the Dirac algebra and is not time, and multiplying by it turns the metric over to the convention of the complex Dirac matrices. *Standard* that these are the Dirac matrices acting on complex four-component spinors, with $U(1)$ phases $\cos\theta + i\sin\theta$ commuting with them. That this phase is the electron's charge is *proposed*, the knower's.

Proposition 9.11 (The train line)**.** *Put a spinor* ψ, *eight whole numbers, at every site of a torus of any size, and on every link a phase* $u = x + yi$ *with whole* x, y. *For a direction with matrix* γ'^μ *put the current* $B_\mu = i\,\gamma'^0\gamma'^\mu$, *and for a link from* s *to* s' *along it*

$$h(\psi_s, u, \psi_{s'}) = \langle \psi_s, B_\mu u\, \psi_{s'} \rangle;$$

let the action be $\sum_s m\langle\psi_s, \psi_s\rangle$ *plus* h *summed over the links. Then* B_0 *and* B_1 *are antisymmetric,* $\langle\psi, B\chi\rangle = -\langle\chi, B\psi\rangle$; *every current commutes with every phase; a phase moves across the dot as its conjugate; a link read backward turns over,* $h(\psi_{s'}, \bar{u}, \psi_s) = -h(\psi_s, u, \psi_{s'})$; $\langle g\psi, g\chi\rangle = |g|^2\langle\psi, \chi\rangle$; *and if every site carries a phase* g *of length one, with* $\psi_s \mapsto g_s\psi_s$ *and* $u \mapsto g_s\, u\, \bar{g}_{s'}$, *the action is unchanged.*

Measured: the middle card checks each statement, and that its γ' acting on a spinor is the matrix of Proposition 9.10. Because the current is antisymmetric, the sum over links is $\sum_\mu B_\mu \otimes (T_\mu - T_\mu^{-1})$ as a form: riding a link forward less riding it back, the lattice derivative. Each link is ridden once, uptown, so no site needs its predecessor; read downtown it turns over, and the line runs one way. A first version read each link both ways with γ' alone; the dot keeps only the symmetric part of what stands between two spinors, the space γ' is antisymmetric, and that version had no space hop at all. The current is the repair. *Standard* that this is the naive lattice Dirac action with $U(1)$ links, gauge invariant, with its doublers. The coupling does not appear: every statement holds whatever scale relates the link's phase to a field. That scale is e, in $u = e^{ieaA}$, and in the weight $\beta = 1/e^2$ of the action on a square against this one *(standard)*; that the device's reading sets it, $\beta = 137.011\,290\,5\ldots/4\pi$, is *proposed*, sentence five.

Proposition 9.12 (The doublers)**.** *Ride a circle of stations with every phase one. On two stations the line reads nothing, for every spinor. On four, the line's response to every probe* φ,

$\sum_s h(\psi_s, 1, \varphi_{s+1}) + h(\varphi_s, 1, \psi_{s+1})$, *vanishes for the field that is the same at every station and for the field* $\chi, -\chi, \chi, -\chi$ *that alternates at every step; and the field* $\chi, \varphi, -\chi, -\varphi$ *that turns a quarter per station is seen.*

Measured, the last by evaluation. *Standard* that the derivative $T - T^{-1}$ vanishes at momentum 0 and at momentum π, so the naive line carries two tastes per direction, 2^d in d directions [25]. The alternating field is the clock of Section 8.2, true, false, true, false: the line cannot tell it from stillness. That the doubler the line cannot see is the taste the written arm $T = F$ keeps (Definition 8.10) is *proposed*, the knower's.

Proposition 9.13 (The whole action)**.** *Write a phase* $x + yi$ *as the turn* $x + y\,e_2$*. Products, conjugates and lengths of phases are those of the turns, and the action on a square of phases is the square of its turned part,* y^2*. For every whole number* β*, the action*

$$S = \beta \sum_{squares} S(H) + \sum_s m\langle \psi_s, \psi_s \rangle + \sum_{links} h,$$

on a torus of any size, is unchanged when every site turns by a phase of length one.

Measured: the middle card checks the embedding of the phases in the turns and the invariance of S for every β, by composing Propositions 9.8 and 9.11. *Standard* that this is lattice quantum electrodynamics with naive fermions, and that $\beta = 1/e^2$ in a normalization where each link carries e^{ieaA}. The device leaves β free; that its reading fixes it is *proposed*, and the comparison with the laboratory excludes the identity at zero momentum by 1.80 parts in ten thousand (Section 8).

9.6 What stays outside

Maps between the tori of the family and its continuum limit; the quantization of the $\mathfrak{su}(2)$ lattice and of the train line; and the value of β. A step is no length, so the gap of Section 9, $\ln(\lambda_+/\lambda_-)$ per step, needs a unit; on a lattice the coupling sets the spacing, so the unit is β *(standard*, for how a lattice's scale is fixed; *proposed*, the author's, for this gap). What is left open by design is the mass gap: the four-dimensional identification of the table's gap with the gap of quantum Yang–Mills, which the millennium problem states for a compact simple group in four dimensions, and which is not claimed. These are carried with sentence five (Section 13).

The continuum, quantized. The author reads the last two items on the geometric card. Its projection of the three-sphere onto the ball has three regions (the geometric card, Definition 2.9 and Proposition 2.10, proved there): over the open ball, away from the silhouette by a margin, every point has two representatives told apart by a sign, and the inverse lift is Lipschitz; at the approach to the silhouette, where the two sheets meet, the inverse is only Hölder of exponent one half, and no larger; outside the closed ball there is no lift at all. The interior is where the numerical card's variation is exact, every action of degree at most three, so that the Gâteaux and Fréchet derivatives coincide (proved there). On the author's reading the interior is the continuum, now, where variation is exact; the approach to the silhouette is before, where variation is no longer exact and carries higher-order terms; the outside is the future, which obeys no continuity; and the continuum is quantized by the sign that tells its two sheets apart, one bit over every point (*proposed*, the author's). On that reading the device's computation sits in the interior, where its higher-order terms are absent, and the laboratory's value carries those of before: Truth's reading, equal up to

the last term. Every measurement, the author adds, displays an interaction from the past: the laboratory reads from before, and the slip of 1.80 parts in ten thousand is what reading from before costs against the interior. That is the arrow of time again (*proposed*, the author's; the slip itself is *measured*). The author names the rest. The interior is densely sampled rationals: points in space and time, each with a particular measurement. Every number the middle card prints is such a point, a pair of whole numbers between two exact walls (*measured*), and the rationals are dense (*standard*). That an interaction leads to a distinguished outcome is the causal theory every rung assumes, the middle card's real instance, *as foretold by the sacred texts of the Flying Spaghetti Monster*: the one declaration under all of it.

The inferred mass gap. The whole action moves with the knower only through β: for any two values, the actions differ by their difference times the action on the squares, and the matter, the squares and the gap per step are the same (*measured*). The middle card names the knower's conclusion on the rung that infers, a closure of one cell whose order against the constant it reads both ways (*measured*), as the inferred mass gap. That β is that conclusion, the mass gap the knower infers from the reader's experience, is *proposed*, the author's.

Two roots, two readers. The counted quartic has two roots in $65x$, $5832 \pm 324\sqrt{90}$, which are 324 times the outer radius $18 + \sqrt{90}$ and the hole $18 - \sqrt{90}$; they add to 11664 and multiply to 24564384, the quadratic's own counts (*measured*). Over 65 they are the reading, $137.011\,290\,5\ldots$, and the conjugate, $42.435\ldots$. The reader reports the outer root, by the calibration that reports the larger. The author reads the two roots as the two crossing arms: the outer root is inferred true by the reader and false by the knower, the arm $T = F$, which stores false; the hole is inferred false by the reader and given true by the knower, the arm $F = T$, which stores true. On that reading the knower's inferred mass gap is the conjugate, $42.435\ldots$, and the reading $137.011\ldots$ stays the reader's coupling. *Proposed*, the author's.

The circle in hexadecimal, and β. The circle enters as its hexadecimal digits, $\pi = \sum_k 16^{-k}\,(120k^2 + 151k + 47)/(512k^4 + 1024k^3 + 712k^2 + 194k + 15)$, and every term after the twenty-fourth together weighs less than $\frac{752}{225}\,16^{-24}$; the two walls print the same twenty-four places, $3.141\,592\,653\,589\,793\,238\,462\,643$ (*measured*; that the series sums to π and the tail bound are *standard*, the digit formula of Bailey, Borwein and Plouffe). The square root of ninety enters between two whole numbers whose squares straddle it (*measured*). Then $\beta = \text{reading}/4\pi$, the weight of the squares against the line in Wilson's normalization, and its walls print the same twenty-four places:

root	$\beta = \text{reading}/4\pi$	on the device's square, $/8\pi$
the reader's, $137.011\ldots$	$10.903\,012\,075\,134\,981\,375\,048\,166$	$5.451\,506\ldots$
the knower's, $42.435\ldots$	$3.376\,859\,126\,587\,073\,981\,938\,450$	$1.688\,429\ldots$

Measured, with no axiom. *Standard* that $\beta = 1/e^2$ in that normalization; the factor of two on the device's square is a calibration, since its action reads twice $1 - \cos\theta$ to leading order. That the knower's β is the inferred mass gap and the reader's the coupling is *proposed*, the author's. The roots' shape is the knower's: with $x = r^2 X$ the quartic is r^4 times itself at $r = 1$, so c and t fix the roots and the reader's r only rescales them (*measured*).

The charge, through Euler. By Euler's formula a link's phase is $u = e^{i\theta}$ with $\theta = e\,aA$: the quarter turn is $e^{i\pi/2}$, and two of them are $e^{i\pi} = -1$, which is the receipt $R^2 = -I$ read with the circle's π (*standard*, Euler). So the charge is $e = \sqrt{4\pi/\text{reading}} = 1/\sqrt{\beta}$. No root is taken: a whole number of twenty-four places is found whose square sits between the walls (*measured*, with no axiom): $e = 0.302\,849\,425\,021\,801\,185\,772\,755$ for the reader's root and $0.544\,181\,192\,849\,948\,845\,712\,632$ for the knower's. The laboratory's charge in the same units is $0.302\,822\ldots$, from its α (*measured* in the literature); the reader's charge differs from it by the same slip, halved, since e goes as the root of the reading. That half is, in the author's words, the part of the electron that is not a carrier of charge, read at the charge (*proposed*).

Two trains. Two trains leave the station, the author says: true and false, and the false is the scenic route, with changes in curvature. The receipts under each are already on this card. The true train is the clicks, every cell true, eighteen times the gain, a fraction whose digits repeat every six; its track is flat, every link the turn e_1, every square whole, no action, a solution at every size. The false train is the clock, through the written arm $T = F$, the square root of ninety times the gain, whose digits never repeat; turned links curve its squares, and one turned link leaves a torus that is not a solution. The straight line cannot see the field that alternates at every step (Proposition 9.12). *Measured* for each of these; that they are two trains from one station, and that the comparison Episode Five called missing is the current B_μ that makes a difference into a derivative, is *proposed*, the author's.

Lanczos, Newton, bisection. The table satisfies its own polynomial, $M^2 = 36M - 234I$, with 36 its trace and 234 its determinant; so two more steps of any walk are 36 of one step less 234 of none, and the closed walks keep the three-term recurrence $\operatorname{tr} M^{n+2} = 36 \operatorname{tr} M^{n+1} - 234 \operatorname{tr} M^n$. Rescaled by a diagonal change so the crossing arms match, the table is the tridiagonal matrix that Lanczos produces, with 18 on the diagonal and $\sqrt{90}$ off it: the clicks and the clock. Its recurrence, $(\lambda - 18)^2 - 90$, ends at $\lambda^2 - 36\lambda + 234$, and at $\lambda = 13y$ that is 13 times the residual $13y^2 - 36y + 18$ (*measured*, each of the four). *Standard*: conjugate gradients is Lanczos with a factorization and is Newton's method on a quadratic, and the sign changes of a three-term recurrence count the roots below a point, so halving locates them. On the author's reading, the device's two searches, Newton and bisection agreeing to twenty-four places, are these two readings of one Lanczos recurrence; the arm $T = T$ is the variational problem, the walk $T = F$, $F = F$, $F = T$ the Jacobian-free Newton–Krylov run, and $F = F$ the linear algebra that gives the Sobolev proof (*proposed*, the author's).

10 The proposed experiment, without its number

The cheat sheet proposes an arrangement and a number for it. This card restates the arrangement and one observable, and not the number.

The refinement operator. The derivation card writes the device as one operator, in letters. The table, read as the map that takes a walk one level up, is the refinement operator; its inverse is the lowering. The device is a consuming iterator on that inverse: each step applies the inverse of the shifted operator and reads its quotient, one symbol of the record per step, which is Newton's method (*standard*). The quotient $\langle v, Sv\rangle / \langle v, v\rangle$ runs from $a - \sqrt{r}$ to $a + \sqrt{r}$ and reaches them on the two

diagonals (*measured*): the two roots are its stationary values, found without writing the polynomial, and the still and the alternating fields carry them. The length it divides by is the comparison read one way: the sum of the squared one-way steps around a ring is a semi-norm whose kernel is exactly the two doublers, and read both ways the same sum is nothing (*measured*). Projected off that kernel, on the turning fields, the central difference is twice the quarter turn and the Galerkin matrix is invertible; the Jacobian-free difference of the residual is exact with remainder thirteen times the step squared; and one such Newton step stays inside the reading's bracket (*measured*). On the author's reading, the derivative of the operator along one direction is a relativistic geodesic and its whole derivative is electrodynamics; the one-way comparison is the third arrow of time, the difference between the reader's experience and interpretation the fourth, and the four are the quartic, one causal law on four faces of the frinkahedron; the quartic's c, t, r, x are velocity, time, phase and position, its unread sign the sign of the velocity $\sqrt{90}/18$ (*measured*, below light), whose rapidity, half the gap, is the reader's boost and the Hubble sphere, the pole the torus goes around (*proposed*, each).

The electron four ways, and the derivative of itself. Each of the numerical card's four actions needs a coefficient with units, and each is set by the electron (*standard*): vacuum gravity keeps only c, the prefactor with Newton's constant dropping out of stationarity, lengths in the Compton length $\hbar/m_e c$; Schrödinger keeps $\hbar/m_e$; the fluid keeps a viscosity whose floor is $\hbar/(4\pi\sqrt{m_e m})$; Maxwell–Dirac keeps e and the size over the Compton length. Standard maps join them: the slow limit, Madelung's fluid, and the Dirac value $g = 2$ that Kerr–Newman also carries. Then the table. Being linear, it is its own Fréchet derivative: refine $v + h$ and the difference is the refinement of h, with no remainder; the action it carries varies by the operator twice and leaves the action of the step. On the whole numbers with $\sqrt{90}$, multiplication by $18 + \sqrt{90}$ is the table with crossings 90 and 1, and it commutes with every boost, each of which keeps the interval. Two tables whose crossings multiply to one round trip are one table seen through two clocks, so the device's crossings 5 and 18 and those 90 and 1 are one operator, and no reading sees the split. The speed is read only from the two stationary values, as their difference over their sum, $\sqrt{90}/18$ (*measured*, each). That only the round trip of light can be measured, and that the extreme values of a quotient under a fixed length are the Lagrange multipliers of that length, are *standard*. *Proposed*, the author's: the four actions are four descriptions of the electron; the split of the round trip is the one-way convention of synchrony; and quantum electrodynamics is the Fréchet derivative of itself, in a space with an interval and boosts, where the speed of light is never measured one way and enters only as a Lagrange multiplier.

The observable. The phase of an interference fringe. A beam is split between two paths and recombined. On one path the spin is rotated by θ about an axis; the two paths differ in addition by a phase ϕ. For a state polarized along the rotation axis, ideal visibility and equal amplitudes, the intensity at one output is

$$I(\theta, \phi) = \frac{I_0}{2}\Big[1 + \cos\big(\tfrac{\theta}{2} - \phi\big)\Big]; \tag{10.1}$$

with visibility $\mathcal{V} < 1$ the cosine carries the factor $\mathcal{V}$. In the arrangement,

$$\phi = \phi_{\mathrm{AB}} + \phi_{\mathrm{dyn}} + \delta, \qquad \phi_{\mathrm{AB}} = \frac{q}{\hbar}\oint A \cdot \mathrm{d}l = \frac{q\Phi}{\hbar},$$

where ϕ_{AB} is the holonomy of Proposition 7.6 for a carrier of charge q enclosing flux Φ, with its sign fixed by the orientation of the loop [1]; ϕ_{dyn} collects the known dynamical phases of the two paths; and δ is the residue.

Proposition 10.1 (The fringe)**.** *For $\psi = (1, 0)$ and output amplitude $\frac{1}{2}\big(\psi + e^{i\phi}U(\theta)\psi\big)$, the intensity is $\frac{1}{2}\big[1 + \cos(\theta/2 - \phi)\big]$ times the input intensity. For $\phi = 0$ and every unit spinor it is $\frac{1}{2}[1 + \cos(\theta/2)]$, with period 4π in θ.*

Proved here: the intensity is $\frac{1}{2}\big[1 + \mathrm{Re}\big(e^{i\phi}\langle\psi, U(\theta)\psi\rangle\big)\big]$, and Proposition 4.11 gives the inner product.

Control. Neutrons, $q = 0$, no residue: the fringe has period 4π in θ, the spinor's return, observed [29, 36]. *Standard.*

Setup. An electron carried twice around a carrier that is neutral and expels magnetic field. Positronium is neutral at monopole order and responds at higher multipole order; a condensed phase of it has been proposed and not yet made [28]. A superconductor expels magnetic field [23]. In general relativity a moving mass produces a field with the form of magnetism, and the frame dragging it predicts has been measured [11]. *Standard*, each. The force an orbiting charge meets from the expulsion would be a second observable; it needs an explicit model of the expelled field, which is not given, and it is not used.

Prediction, in this card's terms. The standard theory of Section 7 predicts $\delta = 0$: every phase between the paths is a holonomy or a dynamical phase. The proposal is that $\delta \neq 0$ in this arrangement and in no other, that the first revolution leaves the residue and the second takes it away or turns it over: a sign, by Proposition 4.10; a slack, by Proposition 4.9; one clock in two phases at four and at eight, by Proposition 4.12. *Proposed.*

What is owed, and where the number lives. The value of δ is the numerical card's and is a representative; it is not restated here. That card also records, from the measured anomalous moment [12, 24], that the coupling read as the fine-structure constant in general is already excluded (the one step of Section 8), and that only the narrow form, a prediction for one stated arrangement, is offered. What the arrangement still owes is a buildable carrier with the two properties, and a reason it counts.

11 The press check

The register is read once, here, by the two readings of Definition 6.1, at the top of the record and at its midpoint. Nothing beneath the head is re-read by the press check.

Reading A: the vector reading. A_k holds exactly when four divides k, by Proposition 4.11: the rotating observable σ_1 returns exactly when $\cos\theta_k = 1$, equivalently when $U(\theta_k) = \pm I$.

Reading B: the spinor reading. B_k holds exactly when eight divides k, by Proposition 4.10: $U(\theta_k) = I$ exactly then, and $\frac{1}{2}\operatorname{tr} U(\theta_k) = \cos(k\pi/4)$ is 1 exactly then.

The trace. Entry k quotes (A_k, B_k) and writes $A_k \leftrightarrow B_k$ undecided. The two numerical rows are the quantities the readings decide; the last row is the diagonal control, which returns at every count and cannot show the pattern.

k	0	1	2	3	4	5	6	7	8
$\frac{1}{2}\operatorname{tr}(\sigma_1 U \sigma_1 U^*) = \cos\theta_k$	1	0	-1	0	1	0	-1	0	1
A_k	✓				✓				✓
$\frac{1}{2}\operatorname{tr} U(\theta_k) = \cos\frac{k\pi}{4}$	1	$\frac{\sqrt{2}}{2}$	0	$-\frac{\sqrt{2}}{2}$	-1	$-\frac{\sqrt{2}}{2}$	0	$\frac{\sqrt{2}}{2}$	1
B_k	✓								✓
$U\sigma_3 U^* = \sigma_3$	✓	✓	✓	✓	✓	✓	✓	✓	✓

Proposition 11.1 (The two verdicts)**.** *At the top, $k = 8$: A_8 and B_8 both hold, so $\neg(A_8 \leftrightarrow B_8)$ is false. Slipped: false. Value: the coin, read back unchanged. Velocity:* 8*. Once around, $k = 4$: A_4 holds and B_4 fails, so $\neg(A_4 \leftrightarrow B_4)$ is true. Slipped: true. Value: the coin, now carrying the sign -1, since $U(\theta_4) = -I$. Velocity:* 4*.*

Proved here, from Propositions 4.10 and 4.11; both decisions are evaluations of a root of unity and neither uses a choice.

$$(A, B) \text{ at } 8: \ (\text{false}, \ \text{the coin}, \ 8), \qquad (A, B) \text{ at } 4: \ (\text{true}, \ -\text{the coin}, \ 4).$$

The record slips once around and does not slip twice around. The numerical card reports two triples from one register in the same way, one not slipped and one slipped by the smallest representable amount; here the smallest representable amount is a sign.

The demand-level reading. Separate from the press check and outside its rule, a second reading goes through the calibration column of this card and records, at each place, where the reading handed over by the knower says more than the reading proved. It does not decide any proposition written undecided; it lists the proposals:

- that the coupling of the carrier is the laboratory's, the one step (Section 8);
- that the carrier is the electron (Proposition 8.3);
- that the middle card's count of turns is the holonomy of a connection (Definition 7.3);
- that the two sectors of the even clock are up and down, and the bit the coin (Proposition 5.2);
- that the eight of the slack and the eight of the spinor are one eight (Proposition 4.12);
- that contents are the symmetry and cells the charge (Propositions 1.11 and 8.2(d));
- that electrodynamics is the pulled-back linearization of the modified field equations, $DR_{\mathrm{QED}}(\psi_0) = D\Phi(\psi_0)^* \circ DR_{\mathrm{GR}}(\Phi(\psi_0)) \circ D\Phi(\psi_0)$, and the second variation the strong interaction's one sentence (Proposition 8.2);
- that the neutral, field-expelling arrangement reads the residue and nothing else does (Section 10);
- that the two directions of the record are the two holographies (Section 6);
- that the bracket's floor is the uncertainty floor made independent of the machine (Section 5.1).

At every other line the located answer is a proposition proved here, cited, or recorded from a companion card.

Inhabitation is not acceptance. That the trace has nine entries with their pairs is well-formedness. The verdicts above are readings of it.

12 The four commutators

The card's title names one object, the commutator of representation: the square $O \circ U = V \circ O$ of Definition 2.5, and the places it fails to close. This section collects them.

Everything else on this card commutes: the readers, the lifts, the carriers, the pair, the relettering, the refinement of the curl. Four pairs do not, and each is located by a receipt: (1) the two orientations, whose bracket is $-13 \cdot 23\,\varepsilon$ (Proposition 9.1); (2) the sign of the coin and the clock, which anticommute (Proposition 8.9); (3) the two orders, the comparison that reverses on both crossings and the order that is false one way and goes on the other (Proposition 8.9 and its source); (4) the description and the measurement, which agree on every digit and slip only against the laboratory (Proposition 8.11). They answer, in reverse order, the four places the device is not yet the electron: the number is (4), the comparison; the missing scale is (3), since a gap is a ratio whose top is a direction and whose unit the device does not fix; the group is (2), the quarter turn that the sign reverses; the dimension is (1), each pair of orientations one more direction a crossing can run. And they stand where the four bent cards stand: Lift 11, the rule that takes the thirteen, is the direction (3); Lift 20, the reading as the coupling, is the comparison (4); Lift 26, the spinor representation and the mass, is the sign (2); Lift 32, quantization, is the transfer matrix against its transpose (1). *Measured* for each commutator; *proposed* for the pairings. A bent card is a commutator the device can locate and cannot compute away. The author reads the four as the four dimensions: four directions, each the receipt of a pair that does not commute, standing where the four variational models of the numerical card stand. On that reading the four dimensions are not adjoined but located, one receipt each (*measured* for the four commutators; *proposed*, the author's, that they are the four dimensions and the four models). The middle card now counts four directions: on a lattice with four, a square for every pair, its action is gauge invariant and its flat configurations are stationary, at every size (*measured*).

12.1 An infinite record

Each of the four differences shows at a finite stage, and this card shows them. Each sameness needs the whole record: the equality of two reals, refuted in finite time and never confirmed (Proposition 3.5); a unit carried to the continuum; rotation by every angle, the exponential of R, an infinite series; the seam refined without end. The argument the knower is handed is therefore of one shape: every finite check agrees except the one that slips, and closing the slip takes an infinite number of symbols. That is sentence five's calibration, *proposed*.

12.2 From five marks to the slip

Put together, the card's constructions are one chain, and every link carries its own calibration.

1. *Five marks.* The frame of Proposition 13.1: a fabric of two crossed strands, two before it and one after, $2 + 2 + 1 = 5$, the weight of the arm $T = F$. *Measured*.

2. *Five posts, four reals.* A real is read as a bracket between two walls; four brackets laid side by side share their inner walls, so four reals need five posts, and the count of brackets and the count of walls differ by the one post the record never reads. *Proved here* for the count; that the five posts are the five marks is *proposed*.

3. *The four reals.* The four readings of the numerical card's quartic, t, c, r and x: three counted, 5, 18, 18, and the fourth, $137.011\,290\,5\ldots$, read across time from one affine bit that time's direction selects. The quartic is homogeneous of degree five; when $cs^2 = t$ it factors as $c\,(r^2 - tx(1-s))(r^2 - tx(1+s))$, and the product does not read the sign of s. Fed the counted $c = 18$, $t = 5$, $r = 18$, thirteen times the quartic is a quadratic in $65x$ whose root is $5832 + 324\sqrt{90}$, 324 times $18 + \sqrt{90}$: over 65, the reading. Since $5/18$ is no square, the factors do not split in whole numbers. *Measured* for the three, the degree, the factors, the sign the product forgets, and the counted root; *proved* on the numerical card for the quartic's value; *proposed*, the knower's, that the sign of s is the affine bit.

4. *The frinkahedron.* Four real coordinates place a point in four dimensions; the geometric card's frinkahedron is a convex polytope there whose boundary is a round three-sphere, and a flat torus such as the seam sits in a three-sphere between the two core circles of its Hopf fibration. *Proved* on the geometric card; *standard* for the flat torus; that the seam is that torus is *proposed*.

5. *The calculus.* On the seam: the step before and after, the curl, the second variation, whole-number frequencies; Gâteaux and Fréchet agree on actions of degree three (Section 9). *Measured*, and *proved* on the numerical card.

6. *The laws and the gauge field.* Four actions of degree three give Einstein's, Schrödinger's, Navier and Stokes's, and Maxwell and Dirac's equations; the table read as a field gives a non-commuting algebra with the electron's group inside, Wilson loops, curvature, and a gap kept open by a boost; with the quarter turn as i, $\mathfrak{su}(2)$ on gauge-invariant links and the Dirac matrices. *Proved* on the numerical card; *measured* here.

7. *Quantization.* Whole windings with half a turn per crossing, and residues paired into a whole charge: the two quantizations whose ratio, through the impedance of the vacuum, gives the fine-structure constant. *Measured* for the counts; *declared* for h and e, which the SI fixes exactly; *standard* for the ratio.

8. *The electron.* The reading taken as the coupling: sentence five. *Proposed*.

9. *The slip.* $137.011\,290\,5\ldots$ against $137.035\,999\,177(21)$: 1.80 parts in ten thousand, which no reader, convention or declaration absorbs. *Measured*.

Links one to seven are receipts, derivations, and standard results joined by arithmetic and by two readings the knower may decline; the five appears in four of them, as marks, as posts, as the quartic's degree and as a weight. The chain cannot itself cross from the eighth link to the ninth.

Where the quarter sits in the five. Of the five posts, four bound the four reals, and each of those is written by a record: three counted, one bracketed. The fifth is the post at zero. The middle card's naturals start at one, its denominators are tapes it wrote and are never zero, and its count, like Riemann's ψ, begins past the term that Jacobi's θ keeps: the omitted zero of the Quarter. No record writes that post. Its value enters additively and every value satisfies the record (the Quarter, "Affine freedom of the three constants"), so its fibre is all of the reals, and by the Quarter's theorem

that the unrepresentable is chosen, whatever stands there is a choice. That is where the numbers no record describes sit: in the one post of the five that no reader reads, under the arm $T = F$, which stores a statement with no proof and is counted anyway. The digits show it.

the arm $T = F$	reading	its digits
struck, weight 0	$\frac{18\cdot 324}{5\cdot 13} = \frac{5832}{65}$	$89.7\,\underline{230769}\,\underline{230769}\,\underline{230769}\ldots$
written, weight 5	$\frac{324}{65}\left(18+\sqrt{5\cdot 18}\right)$	$137.011\,290\,548\,979\,457\,087\,737\,3\ldots$

Struck, the reading is a fraction, and after its first digit six digits repeat forever: a finite record writes every digit. Written, the clock enters as $\sqrt{5\cdot 18}$, and $5\cdot 18 = 90$ is not a square, so the reading is not a fraction and its digits never settle into a repeating pattern; a polynomial describes the number, and no finite pattern writes its digits. Every digit that no finite pattern writes comes in with the five. *Measured*: the middle card checks the repetition of the struck digits, their six, and that 90 is no square. *Standard*: a number's digits eventually repeat exactly when it is a fraction. *Proposed*, the knower's: that the post at zero under the five is the Quarter's quarter.

The order, read both ways. The middle card's last inference compares two closures by its order: the interpretation, read from the constant and the reader's look at it, against the closure that ends on *constant at most real*. Dealt on the rung that names the knower, the comparison holds, and it holds the other way too. Every cell of the constant clicks, so the order on the constant never reverses, and both slips are true. *Measured*: the middle card checks both directions and mints each as a fact that clicks. The device reads the order both ways; that it runs one way is the knower's, *proposed*. A cell that did not click would reverse the order on itself.

13 What the knower carries

The proposals of this card were once collected as nine sentences. Eight of them are now the middle card's, each by a result on the left-hand side, and three of those by narrowing what the sentence claims to what the device states; one is carried by the knower. Each is still a calibrated reading in the sense of the numerical card's reading note: it reads a construction, supplies a correspondence, preserves some properties, and stands to be revised by stated evidence.

What the device carries, 9 November 2026.

1. **Equality is a mirror.**
   The device's residual is a scalar, and every scalar residual is the derivative of an action: the roots of r are the stationary points of S with $S' = r$ (Propositions 4.1 and 4.2). *Proved here*, and *measured*: the middle card checks that its own residual, read at one cell of denominator, is the first variation of a cubic action with the same three counted coefficients, thirteen, thirty-six and eighteen: moving the point by h moves the action by $6h$ times the residual, a term in h^2 and a term in h^3, and nothing more. The general vector form is not needed and is not claimed.

2. **The run is the reading to its own precision.**
   Formerly, the run contains its limit. The device prints twenty-four places, and twenty-four is a length it counts; the bracket it holds is narrower than one place at the twenty-fourth (Section 5.1).

Proved here, and *measured*. The limit is no longer claimed: that no reading at the device's precision can tell the run from the root is the calibration, fixed once against the device's own count, and it absorbs no offset.

3. **The device's actions have degree at most three.**
   Formerly, the selected degree is nature's degree. Each of the four actions the numerical card varies has degree at most three in its declared independent fields, so $D^4 S = 0$ (Section 4.2). *Proved* on the numerical card. Whether these are nature's actions is part of sentence five.

4. **The device's work is a count.**
   Formerly, least work is least action. The logical work of a step is counted by the prover, and the count of the same step is the same on two different machines while the time differs (Proposition 1.12). *Measured*. Reading that work as physical action at a named boundary is no longer claimed.

6. **The sign is what vectors forget.**
   The vector observation of a spinor loses the kernel $\{I, -I\}$, and the device retains it as a coin (Proposition 4.11). *Proved here*; the 4π fringe that no vector description reproduces is *measured* [29, 36] (Proposition 10.1).

7. **The slack and the spinor are compared at two counts.**
   Formerly, they are one clock. They are complementary at four and at eight quarter-turns, and from the ninth both readings fail (Proposition 4.12). *Proved here*. The strong form was revised by that computation; what remains is a comparison, and a comparison is not a debt.

Two former sentences are not the device's, and they are not separate debts either: each is sentence five read from one side.

8. **The device's coupling is the laboratory's.**
   As an identity with the coupling at zero momentum it is *measured* and excluded: 1.80 parts in ten thousand (Section 8). What remains, the reading taken as $e^2/4\pi$ in the carrier's law, is how sentence five casts the carrier; the narrow form, one stated arrangement, is revised by the fringe phase δ of that arrangement.

9. **The number under the card is a choice.**
   Theorem 2.7 proves that no procedure inside the family selects a representative of a fibre, so the device cannot carry it. To name the electron as the carrier is to name that representative: sentence nine is sentence five seen from the record.

The identification of the next subsection is the device's as well: the arm $T = F$ is written, with a weight the prover counts (Proposition 13.1); only what the written arm is read as, a doubler kept as a taste (Definition 8.10), belongs to sentence five.

What the knower carries.

5. **The carrier is the electron.**
   Reads: Definition 8.1, the naotrino as the nowtrino's model (Section 8.1), the written arm as a kept taste (Definition 8.10), the reading as the coupling, the quarter turn read as i, whose spin the gamma matrices carry, and β as the gap's unit (Section 9). *Supplies*: the defined object for

the physical electron, and with it the one point of a fibre that no record names (Theorem 2.7). *Preserves*: the transformation law, the observation map and the stationary system, which are the standard ones; and every digit, since the reader adds none (Proposition 8.11). *Revised by*: any observation of the electron that its stationary system with some coupling does not reproduce at the classical level; the quantum level needs the quantization the card does not supply. Already revised at zero momentum by 1.80 parts in ten thousand, which the knower names the part of the electron that is not a carrier of charge. *Proposed.*

The left-hand side is everything a prefix can show, and on this card it is shown or cited in full: that a mark fits any carrier, that two readers agree exactly when their lengths do, that re-lettering is free and is not a symmetry, that the box has one point, that exclusive or needs both definite signatures, that the residual is even and the action odd, that Newton's error is exactly the acceleration term, that the degree forgets the coupling and the derivative keeps it, that the slack is positive at four and negative at eight, that the spinor flips at four and returns at eight while σ_1 returns at four, that transport is a connection and its commutator is the curvature, that the action's stationary system is Maxwell's and Dirac's, and that the verdict slips at four and not at eight. The last line of the middle card asks whether the mechanism demonstrating its proof looks like an electron. This card answers what a record can answer: the mechanism carries a coin, a phase, a transport, a rule of degree three, a coupling and a count, and no reader of it can tell the carrier from an empty one. The Quarter's epigraph asks, "Can you find the polynomial in the physical model of this device?" That is the mark's question, and the patter never answers it. Whether the carrier is an electron is the knower's, and it is handed over calibrated, so that it is carried as a proposal and not mistaken for a receipt.

The identification

One sentence remains, and it is the first one: $T = F$. In the table it is the arm that compares TRUE with FALSE. It stores False. It does not click. No run of the device reaches it. What it carries is not chosen.

Proposition 13.1 (The weights are counted)**.** *Every arm of the table carries the frame, a tape of five marks. An arm whose stored comparison can click carries the coefficient tape of thirteen as well, so that* $18 = 13 + 5$*, and the arm* $T = F$*, which stores* False*, carries the frame alone. The five is fixed three times, and the three tapes agree mark for mark:*

1. *the frame as built: a fabric of two crossed strands, a fibre of two strands before it and one strand after it,* $2 + 2 + 1 = 5$ *marks, each the one truth;*
2. this way*, the smooth model: asked at which stations of the descent the local model of the carrier stops, the prover names five, from the smooth model through the laboratory to the naotrino;*
3. that way*, the yarn: asked the same of the yarn, it names five again, the yarn standing where the smooth model stood and the other four stations shared.*

The thirteen is fixed the same way. The step that sends the local model along the yarn stops at thirteen stations, the three laboratory stations and ten that only the yarn step reaches, and its tape is the coefficient tape.

Measured: each count is the prover's census of the middle card's own elaboration, and each agreement closes by reflexivity (Appendix A). The weighting rule is the one declaration on this table: the arm whose stored fact cannot click loses the thirteen. It does not assign the five. The five is where both ways stop, and it is what remains on the arm when the thirteen is taken away. The two tapes of five are not reached alike. The frame writes the one truth five times at zero distance, a tare. The census counts it five times at the end of the long path, through the carrier, the stations and the naotrino, a count. They are the same tape, and the arm $T = F$ carries it. Strike it, and see what the machine holds.

Theorem 13.2 ($T = F$)**.** *Let the arm $T = F$ carry weight $w \geq 0$, the other three arms* 18*, so that $M_w = 18\,I + N_w$ with $N_w = \begin{pmatrix} 0 & w \\ 18 & 0 \end{pmatrix}$ and $N_w^2 = 18w\,I$.*

1. *If $w = 0$ the arm is struck. Then $N_0^2 = 0$: the one-way crossing that survives cannot close, the crossing cycle stops, and every closed walk keeps its coin, $\operatorname{tr} M_0^n = 2 \cdot 18^n$. Then $\lambda_+ = 18$ and the reading is $\frac{324}{65} \cdot 18 = 89.723\,076\,9\ldots$, a rational number: every TRUE read as TRUE.*
2. *If $w = 5$, the weight Proposition 13.1 counts, the arm is written. Then $N_5^2 = 90\,I$, the clock runs, $\lambda_+ = 18 + 3\sqrt{10}$, and the reading is $\frac{972(6+\sqrt{10})}{65} = 137.011\,290\,5\ldots$, irrational. The whole of the difference, $47.288\,213\,6\ldots$, is the growth rate of N_5, whose walks cross the arm $T = F$ at every other step.*
3. *Take that reading for the coupling, $e^2/4\pi = 1/137.011\,290\,5\ldots$. Then the chain of Section 7 carries it through Theorem 7.7 and Definition 7.8 to the minimally coupled Maxwell–Dirac stationary system, Proposition 7.9, under the assumptions named there: a Lorentzian background, the spinor representation, the kinetic terms, the mass, and, for quantum electrodynamics proper, quantization and renormalization.*

Proof. N_w is off-diagonal, so $N_w^2 = 18w\,I$ and the eigenvalues of M_w are $18 \pm \sqrt{18w}$. At $w = 0$ the matrix is triangular with both eigenvalues 18; at $w = 5$, $\sqrt{90} = 3\sqrt{10}$. The reading is $\frac{324}{65}\lambda_+$ in both cases (the cheat sheet's Propositions 9.3 and 9.4). Item 3 is Section 7 with the coupling supplied. *Proved here* for items 1 and 2; item 3 as calibrated where each of its steps enters. □

So the number is in the machine exactly when $T = F$ is written. Strike the arm and the machine holds $89.72\ldots$, the device agreeing with itself, a fraction, nobody's constant. Write it, though it never clicks and no run ever reaches it, and the machine holds $137.011\,290\,5\ldots$, and the number carries through transport and curvature into the coupling of Maxwell's and Dirac's equations. That is what $T = F$ buys. It is written on the card. It is not read by the machine. Under the weighting rule it weighs in the count all the same, and the theorem shows what it contributes. What that counted structure is taken to represent, a number in the machine that is the coupling of an electron, so that every other TRUE stands for FALSE, is the identification, and it is the knower's. *Proposed.* The laboratory has measured the coupling the identification would deliver, $137.035\,999\,177(21)$, and the delivery is short by 1.80 parts in ten thousand (Section 8).

For the reading that makes the identification: TRUE equals FALSE. Write it, and there is a number in the machine. Read it, and there is electrodynamics. The card has not lifted. The knower who reads it so has.

The last question

Every question this card has asked came back to one shape. A difference shows at a finite stage; a sameness needs the whole record. The readers agree with one another, the lifts carry no datum, two carriers print one number, the click machine and the other machine click alike through either carrier, and the reader adds no digit. Four pairs fail to commute, and each is located: the two orientations, whose bracket is the sign of the coin scaled by the bent card and the sum of the arms; the sign and the clock, which anticommute; the two orders, one reversing on every crossing and one false in a single direction; and the description and the measurement, which agree on every digit the device prints and slip only against the laboratory. Each of the four stands where a bent card stands. Each is a place where the expansion says *given* instead of *because*, and none can be computed away.

The last question is the one the table asks of itself. Its antisymmetric part is the bent card times a quarter turn, the generator of rotations, the compact group the electron couples through. So is the bent card the electron's charge? The record cannot say. The answer, as the knower gives it, is that it depends on whether one believes electrons were used in the generation of the proof.

That is not a deflection. It is the most literal sentence on the card. The middle card was checked by a kernel running on two machines, a small one and a large one, and the count of logical work, the heartbeats, was the same on both while the time on the wall was not. *Measured.* Those machines run on electrons moving through silicon. That is a fact about the hardware, *standard.* Between the two facts stands the middle card's one causal declaration, the instance every rung assumes and no rung constructs: that the device's execution is a case of the causal theory it describes. It is written in the satire's register, as foretold by the sacred texts of the Flying Spaghetti Monster, because that is what it is: believed, and not shown.

The distinction the card has kept from its first page applies here without exception. Logical work is not transferred charge. Heartbeats become charge only through a measurement at a named boundary. A measurement could be made: run a check of known logical work, count the charge that crosses the battery's terminals, seal the predicted slope before the run, and repeat on a second machine. One machine would give a slope, a result; agreement across machines would be a further claim, universality, and would need its own seal. Until such a run, the sentence "electrons were used" is true of the hardware and *proposed* of the proof.

Notice what kind of proposal it is. Every other identification on this card names a carrier somewhere else: the electron in the laboratory, the naotrino at the end of a refinement that never ends, the representative a fibre holds and no record names. This one names a carrier in the same box as the record. The tally counts naotrinos, the arguments of the record about one another, and never looks at one. The carrier of the tally is declared, at the start of the descent, and every rung takes it as given. If the electrons that executed the check are that carrier, then the record is a reading of the electron in the most direct sense available: the electron, made to carry a tally, wrote down how long it was. If they are not, the record is a description that ran on some electrons and is about none of them. No finite record decides between the two. The first reading cannot be confirmed by any finite run; the second cannot be refuted by one. It is the identification of Proposition 3.5, turned to face its own hardware.

And the slip stays where it was. Whatever the knower believes about the machine, the number the machine prints is $137.011\,290\,5\ldots$, and the laboratory's is $137.035\,999\,177(21)$. If electrons generated the proof, then electrons generated a coupling short by 1.80 parts in ten thousand, and the knower who believes it must say what the electrons carried that the tally did not. The card's knower has said it: the part of the electron that is not a carrier of charge, its volume, which no reader of length

can see and which a measurement of the whole electron always contains. That answer is *proposed*. It is also the only answer consistent with every receipt on the card, because it is the only one that puts the difference where the receipts allow a difference to be.

So the argument ends on the knower, as it was always going to. The device has shown the reader everything a record can show: a mark that fits any carrier, readers that agree exactly when their lengths do, an isomorphism between its tallies and the naturals, a non-commuting algebra read off counted weights with the electron's group inside it, a gauge field whose loops are invariant, a gap that survives refinement, and four commutators it can locate and cannot remove. It has offered the knower one sentence. The carrier is the electron. Whether the knower takes it depends, in the end, on what the knower believes the proof was made of.

Weird that if you believe there is a number in there you can derive QED.

Isn't that a daisy?

A Sources

This appendix is the only part of the card that names the middle card's files and declarations. The body refers to it only as "the middle card." Paths are relative to `CARD_2/Measurement/`. Declarations are cited by name and not by line; the working tree was inspected on 29 October 2026, and again on 3 November 2026 for the carrier encoding, the readers and the closed walks, and on 8 November 2026 for the doubler and the reader's digit, and on 10 November 2026 for the crank, the quartic, the order, the lattice, the turns and the gamma matrices; nothing here reports a build. Companion cards are cited by printed result number: the geometric card (*The Frinkahedron*, build of 22 October 2026), the numerical card (*Four Models and a Common Fourth-Order Closure*, 29 October 2026), the Quarter, the bent card (*The Infinite Loop Considered Harmless*) and the cheat sheet (*A Knower's Guide to Orbital Mechanics*).

The carrier's encoding. `Episode01.lean` declares `Number` with constructors `zero : FACT → Number` and `one : FACT → Number → Number`, and `CarrierProcess` with fields `symbol : FACT`, `quarter : Number` and `interact_with_the_universe : Number → Number`, the last with a default value. A `Number` is a nonempty record: `Number.peano` (`Episode08.lean`) sends `zero f` to the one-symbol record `[f]` and `one f n` to `f` followed by the record of `n`; on nonempty records `tapeNumber` (`Episode12.lean`) inverts it. Under this map the continuation of Definition 1.2 is `one f n` $\mapsto$ `n`; the slot q is `quarter`; the symbol s is `symbol`; and ι is the default `interact_with_the_universe`, which sends `zero _` to `zero FACT.Truth` (clause (ii)) and `one g _` to `one g quarter` (clause (i)). Its match inspects only the constructor of its argument and never `quarter`: carried, not queried. The structure permits a carrier to override the default; the definition is of the default, which is the only interaction the inspected tree uses. The executable slot is therefore a `Number`, and the record of the definition is its image under `Number.peano`.

This card	Companion card	Middle card (declaration)
Def. 1.2, carrier	Quarter §6 (affixing)	`Episode01.lean`: `CarrierProcess` with fields `symbol`, `quarter`, `interact_with_the_universe`; `Number`; `Episode08.lean`: `Number.peano`; `Episode12.lean`: `tapeNumber`
Def. mark; click	cheat sheet §6	`Episode01.lean`: `FACT` (`truth`, `click`), `FACT.Truth`; `Episode12.lean`: `the_click`
Def. 1.4, three readers	bent card Props. 1.3, 1.4	`Episode08.lean`: `naotrino_detector` (the register), `List.length` (the tally); `Episode12.lean`: `the_click_count`
Prop. 1.5	Quarter §6 (two counting layers)	`Episode08.lean`: `adding_in_the_fencepost_gives_a_monomial`; `Episode12.lean`: `the_clicks_agree`, `the_clicks_read_the_number`
Def. 1.6, Prop. 1.7	—	`Episode12.lean`: `the_click_count` (a payload reader); `read_past_quarter`, `read_past_quarter_is_length` (a reader in the length family)
Prop. 1.8	cheat sheet Prop. 6.1	`Episode12.lean`: `the_card_between`, `the_mark_between`, `the_tape_between_deals_rung01`
Prop. 1.9	cheat sheet Prop. 6.2	`Episode12.lean`: `ComputerProgram.ofTape`, `read_ofTape`, `the_other_machine_agrees`
Props. 1.10, 1.11	numerical card §7.4	`Episode12.lean`: `relettered`, `relettering_is_free`, `any_count_of_the_lengths`
Prop. 1.12	numerical card Rem. 8.3	`Episode11.lean`: `TYPESET.universe_id`
Prop. 2.1	numerical card §7.5	`Episode08.lean`: `the_rational_idea_of_relativity`; `Episode12.lean`: `the_box_has_one_point`, `the_box_reads`
Prop. 2.2	cheat sheet Prop. 6.3	`Episode05.lean`: `TruthOrder`, `FACT.SAME`; `Episode08.lean`: `a_partitioning_of_quotients`
Def. 2.4	geometric card Def. 4.1, Props. 4.2, 4.3	`Episode08.lean`: `theReadingCertified`

This card	Companion card	Middle card (declaration)
Thm. 2.7	Quarter Thm. 3.1	the axiom sweep reported in the Quarter; `Episode12.lean` `#print axioms` lines
Prop. 2.10	numerical card §3.5 (multiplier vs. penalty)	`Episode12.lean`: `Signature`, `Signature.sign`, `xorClauses`, `xorRow`, `xorTarget`, `step_correspondence`
Prop. 2.11	cheat sheet Prop. 6.4	`Episode12.lean`: `SAME_DIFFERENCE`, `same_difference_clicks`, `the_click_diagonal`
§4.1	cheat sheet Props. 3.1, 3.2; numerical card Prop. 6.1	`Episode12.lean`: `the_acceleration_is_constant`, `the_first_derivative_squared_is_the_discriminant` (stated there with the middle card's coefficients; here without)
§4.2, Props. 4.7, 4.8	numerical card Thms. 1.1, 2.4, Prop. 2.5, §2.5	`Episode04.lean`: `Variation` with constructors `the_newton`, `the_gateaux`, `the_frechet`; `Episode12.lean`: `gateauxAt`, `frechetAt`, `theConvectiveGap` and its margin
Prop. 4.9	cheat sheet Prop. 4.1; bent card §4	`Episode08.lean`: `the_slack`, `once_around_the_integral`, `twice_around_the_integral`, and the three theorems that read its sign (the first time around, through the bent card, the second time around)
Props. 4.10, 4.11	cheat sheet Prop. 7.1	`Episode05.lean`: `UNIVERSAL.once_around`, `UNIVERSAL.twice_around`; `Episode01.lean` margin on the clock and clock complement
Prop. 4.13	numerical card Rem. E.1	`Episode08.lean`: the register's all-ones identities
Props. 5.1, 5.2, 5.3	geometric card §3.4, Props. 3.3, 3.4, §6.3	`Episode03.lean`: `Equivalation` and `DigitalProcess.tick` (the exclusion arm); `Episode02.lean` margin on the six-state sensor
Prop. 5.4	Quarter §6	`Episode12.lean`: `the_card_between` dealt through `deal_rung`; the scratch comparison of all steps is reported in the Quarter and not here

This card	Companion card	Middle card (declaration)
§5.1	Quarter §6 (two floors)	`Episode05.lean`: `Closure.same`, `Closure.different`; `Episode12.lean`: `theCourant`, `bracketed_at_the_slip`; `Calibration/LeanCalibration.lean`: `register_noise_floor`
§6, Def. 6.1	both cards §8; numerical card Rem. E.2	`Episode10.lean`: `REVIEWED`; `Episode11.lean`: `TYPESET`, `PressVerdict`, `pressCheck`
§7, Def. 7.3 (the closed walks)	numerical card §2.4	`Episode12.lean`: `the_closed_words`, `the_closed_walks`, `the_closed_words_are_closed_walks`, `the_crossings`, `every_closed_walk_crosses_evenly`, `the_winding`, `the_turning_walks`, `every_closed_walk_turns_evenly`, `the_residue`, `the_two_readings`, `every_closed_walk_is_exact`, `every_walk_between_the_coins_leaves_half_a_turn`; no declaration of a connection on spacetime or a transport between points
§8.1, Props. 8.5–8.7	geometric card, "A cellular model of the accessible torus" and "States, encodings, and the action of the clocks"	`Episode12.lean`: `the_walks_on`, `the_device_walks_on_its_table`, `the_readings_fix_the_gauge`, `the_other_orientation_is_a_gauge_copy`; `the_curl`, `the_seam_flux`, `the_closed_seam_carries_no_net_flux`, `the_two_readings_of_a_surface`, `the_residues_pair_into_the_charge`, `one_half_turn_is_one_charge`; `the_plaquette`, `subdividing_adds_no_residue`; no declaration of a limit, a convergence, or a field on spacetime

This card	Companion card	Middle card (declaration)
Def. 8.8	cheat sheet §6	`Episode11.lean`: the instance argument `[the_naotrino_theory : REVIEWED Box Pigeon k]` of every typeset page, $k = 1, \ldots, 36$; `Episode10.lean`: `rebut` and the thirty-five reviewer instances; `Episode02.lean`, `Episode03.lean`: the instance argument `[nowtrino : DISTINGUISHABLE Box Pigeon]` and its margins (no click, no matter); `Episode07.lean`: `IS_THIS_A_NAOTRINO?` and its margin (the non-relativistic version of the nowtrino), `naotrino_model := A_SMOOTH_MODEL` in `IS_IT_SPINNING?`; `Episode04.lean`: `REAL` and its margin (no naotrino seen)
§8.2, Prop. 8.9, Def. 8.10	cheat sheet §9, "The clicks are a scalar; the excess alternates"	`Episode12.lean`: `every_closed_walk_crosses_evenly`, `the_arm_weight`, `the_closed_words_are_closed_walks`; `the_comparison_reverses_on_a_crossing` (the default `COUNTABLE.bounded?` of `Episode01.lean`, through `CAN_YOU_COUNT_THEM?`), `the_reversal_is_the_crossing_sign`, `the_reversals_telescope`, `the_round_trip_reverses_twice`, `the_crossings_pair_up`; the direction in `Episode01.lean`: `Number.le` and in `Episode12.lean`: `the_arm_stores_a_fact_that_can_click`; no declaration of a lattice fermion, a taste, or a rooting
§3, Prop. 3.1	the Quarter, Theorem 3.1	`Episode12.lean`: `no_length_reader_sees_the_relettering`, `the_tape_with_a_false_cell`, `the_relettering_changes_the_cells`, `a_representative_is_a_choice`, beside `relettered` and `relettering_is_free`

This card	Companion card	Middle card (declaration)
Prop. 3.2	—	`Episode12.lean`, namespace `readers_interpretation_of_the_pair`: `the_pair_between`, `the_mark_on_the_pair`, `the_pair_deals_rung01`, `read_past_the_pair`, `two_carriers_print_one_number`, `the_pair_reads_the_number`, `no_reader_of_the_tally_sees_which_carrier`, `which_carrier_is_a_choice`; `read_past_the_carrier`, `the_click_through`, `each_carrier_reads_the_card_between`, `click_click_through_the_carriers`, `click_click_on_the_tally`, with `the_clicks_read_the_number`; for Prop. 3.3: `the_same_tally`, `the_tallies`, `the_tally_of`, `the_numeral`, `the_tally_of_the_numeral`, `the_numeral_of_the_tally`, `the_numeral_is_the_relettering`, `every_carrier_reads_a_natural`, `the_number_is_computed_from_naturals`
Thm. 3.4	—	`Episode12.lean`: `THEORY.step_correspondence`, `THEORY.trace_correspondence`, `cook_levin_the_description`, `cook_levin_on_the_mechanism`; in `readers_interpretation_of_the_experience`: `cook_levin_on_the_readers_interpretation`, `the_accepting_seed`, `the_rejecting_seed`, `the_formula_accepts_and_rejects`, `the_verdicts_are_the_steps`; (i) is the standard theorem cited
Prop. 3.5	—	no declaration: (i) and (ii) are the standard results cited; (iii) is the comparison of Section 8

This card	Companion card	Middle card (declaration)
§13, sentence 1	cheat sheet, *What the knower believes*, item 1	`Episode12.lean`: `the_mirror_residual`, `the_cubic_action`, `the_mirror_is_the_device_residual`, `the_residual_is_the_derivative_of_the_action`, `the_mirror_coefficients_are_counted`, beside `the_residual` and `the_first_derivative`

This card	Companion card	Middle card (declaration)
§9, Props. 9.1–9.3	—	Episode12.lean: Table, the_table_product, the_table_bracket, the_clock_table, the_other_clock, the_sign_table, the_orientations_bracket_onto_the_sign, the_sign_brackets_the_clock_back, the_sign_brackets_the_other_clock_back, the_three_are_traceless, the_three_are_independent, the_adjugate_inverts, the_wilson_loop_is_gauge_invariant, the_holonomy, the_uniform_square_is_flat, one_turned_edge_curves_the_square, the_quarter_turn, the_antisymmetric_part_is_the_quarter_turn, the_quarter_turn_squares_to_minus_one, the_sign_reverses_the_quarter_turn, the_clock_of, the_same_round_trip_is_the_same_clock, the_swapped_table_shares_the_clock, the_conjugate_roots_are_rational, the_round_trip, the_surd_product, the_interval, the_outer_and_the_hole, the_interval_is_the_determinant, the_boost_has_unit_interval, the_boost_preserves_the_interval, the_hole_stays_open, the_boosted_torus, the_step_after, the_step_before, the_seam_laplacian, the_second_variation_is_before_less_after, the_sixths, the_frequency, the_plane_wave, the_plane_waves_are_the_frequencies, the_overlap, the_plane_waves_are_independent; the_struck_numerator, the_struck_denominator, the_struck_remainder, the_struck_digit, the_struck_digits_repeat_every_six, the_struck_digits_are_two_three_oh_seven_six_nine, the_round_trip_is_not_a_square, with the_post_at_

This card	Companion card	Middle card (declaration)
Def. 9.5, Prop. 9.6	—	`Episode12.lean`: `the_plaquette_action`, `the_first_variation_of_the_plaquette`, `the_varied_table`, `the_plaquette_action_stops_at_second_order`, `the_plaquette_action_is_gauge_covariant`, `the_holonomy_is_linear_in_its_bottom_link` (and right, top, left), `the_bottom_link_varies_the_action`, `a_scalar_square_is_stationary`, `the_uniform_square_is_a_lattice_solution`, `the_turned_square_has_action`, `the_rotation_and_the_boost_read_opposite_signs`; the torus: `Lattice`, `the_square`, `the_lattice_action`, `each_square_holds_a_link_once`, `a_flat_lattice_is_stationary`, `the_uniform_lattice_solves_the_lattice_equation`, `the_turned_lattice_has_action`

This card	Companion card	Middle card (declaration)
Props. 9.7–9.9	—	Episode12.lean: the_swap, the_symmetric_part_is_the_swap, Box4, the_box, the_box_product, the_turn_of_the_swap, the_turn_of_the_quarter, the_turn_of_the_sign, the_three_turns_square_to_minus_one, the_turns_compose_around_the_circle, the_turns_bracket_around_the_circle, the_three_turns_are_independent; the_table_satisfies_its_own_polynomial, the_table_power, the_walks_keep_a_three_term_recurrence, the_lanczos_recurrence_ends_at_the_table, the_lanczos_polynomial_is_the_residual; every_table_satisfies_its_own_polynomial, the_crossings_are_a_quarter_turn_and_a_swap, both_roots_solve_any_table, the_rayleigh_quotient_lies_between_the_roots, the_roots_are_reached_on_the_diagonals, the_one_way_step_squared, the_semi_norm_is_the_comparison_read_one_way, read_both_ways_the_comparison_measures_nothing, the_semi_norm_sees_nothing_exactly_on_the_doublers, the_galerkin_matrix_is_invertible, the_jacobian_free_difference_is_exact, one_jacobian_free_newton_step_stays_in_the_bracket, the_quartic_sign_is_the_velocity, the_steps_commute, every_four_torus_is_gauge_invariant, a_flat_four_torus_is_stationary, every_boost_keeps_the_interval, the_hole_never_closes, the_ratio_of_the_roots_is_twice_the_boost, the_schwinger_term_from_the_reading; the_refinement_is_its_own_derivative, the_action_varies_by_the_operator_twice, the_surd_products_commute,

This card	Companion card	Middle card (declaration)
§9, the crank, one way	—	Episode08.lean: the_device_computing_the_next_configuration, the_constant_can_be_inferred; Episode12.lean: the_floor_reads_the_step, both_ways_the_step_is_zero, both_ways_the_guard_always_stops, one_way_the_guard_reads_the_step, at_Int_the_scale_is_never_read, at_Int_the_tolerance_is_zero, every_k_is_certified
§12, the four reals; the order, read both ways	—	Episode12.lean: the_closing_quartic, the_quartic_is_homogeneous_of_degree_five, the_quartic_factors, the_quartic_does_not_read_the_sign, the_counted_quartic, the_reading_solves_the_counted_quartic, the_conjugate_solves_the_counted_quartic, the_two_roots_are_the_counts, the_hex_term, the_hex_pi, the_hex_tail, the_hex_digits_print_pi, the_root_of_ninety_is_bracketed, the_beta_of, the_readers_beta, the_knowers_beta, the_betas_on_the_device_square, the_reader_only_rescales_the_roots, the_readers_charge, the_knowers_charge, the_readers_charge_is_bracketed, the_knowers_charge_is_bracketed; Episode05.lean: INFERRED, Closure.le; Episode12.lean: the_knower_reads_one_way, the_device_reads_it_back, the_two_closures_click, the_inferences, every_inference_clicks

This card	Companion card	Middle card (declaration)
Prop. 8.11	—	`Episode12.lean`: `the_reader_adds_no_digit`, composing `naotrino_detector_reads_the_number` and `the_same_natural_both_ways`; `Episode08.lean`: `fenceposts_must_be_preserved`, `the_device_as_linear_operator_mapped_identity` (the fencepost out and in)
Prop. 13.1, Thm. 13.2	cheat sheet §9	`Episode08.lean`: `the_coordinate_system`, `the_dimensions_of_universe`, `the_coordinate_system_gauss_length`; `episode7_elaboration_of` `this_way` as `the_data_table`, `that_way` as `the_train_schedule`, `the_calculus` as `the_differential_form`; `the_data_table_is_the_gauss`, `the_train_schedule_is_the_gauss`, `the_differential_form_is_the_coefficient`, `bent_card`, `the_macguffin`; `Episode12.lean`: `the_arm_stores_a_fact_that_can_click`, `the_arm_weight`, `the_click_table_is_derived`
Def. 8.1	cheat sheet §6, §7; numerical card Ch. 6	`Episode07.lean` margin listing the electron's values; `Episode04.lean`: `Knowledge.jarjar` and its margin casting the part; `Episode12.lean`: `readers_interpretation_of_the_experience`; for (viii), `what_would_you_call_this_number?`
Prop. 8.3	numerical card §2.4	`Episode01.lean` foreword on the two readings of Q.E.D.
§8.3	Quarter §6 (the horizon)	`Episode08.lean`: `the_photon_worldline` and the three encodings that follow it (position, velocity, acceleration, each zero)
§10	cheat sheet §6.2, §7; numerical card §6.7	`Episode05.lean`: `AtreyuProcess` margin; `Episode07.lean`: `YARN_THEORY` margin
§11	both cards §8.3	`Episode11.lean`: `pressCheck`

B The reader's experiment

Every result calibrated *proved here* is a written record: hypotheses, a sequence of displayed steps, and a conclusion. Read as an experiment it has a setup, an apparatus (whoever checks the steps), a law (the rule each step obeys), a procedure (the order of the steps), observations (each step seen to hold) and a reading (the conclusion). Four runs on paper reproduce the results on which the press check and the transport rest.

Run 1: the slack. Write the odd numbers below eight and below sixteen; form their least common multiples, 105 and 45045; divide each by each odd number and subtract the quotients from the multiple. Observe 34 and -982. Each line is an integer division and a subtraction. This is Proposition 4.9.

Run 2: the signatures. Write the two clauses of Proposition 2.10 over four slots. For each of the four signatures, sign the slots, read each literal, and evaluate the formula on the four assignments. Observe the four rows of the table. Conjoin the first two rows and observe $[F, T, T, F]$; conjoin the last two and observe $[F, F, F, F]$. If any single row equals $[F, T, T, F]$, Proposition 2.10 is wrong.

Run 3: the eighth root and the rotating observable. For $k = 0, \dots, 8$ and $\theta_k = k\pi/2$, write $U(\theta_k) = \mathrm{diag}(e^{-ik\pi/4}, e^{ik\pi/4})$. Record where $U(\theta_k) = I$: at $k = 0$ and 8. Compute $U(\theta_k)\sigma_1 U(\theta_k)^*$, whose upper-right entry is $e^{-ik\pi/2}$, and record where it equals σ_1: at $k = 0$, 4 and 8. Then compute $U(\theta_k)\sigma_3 U(\theta_k)^*$ and observe σ_3 at every k: a diagonal observable is the control, and it cannot show the pattern. These are Propositions 4.10 and 4.11, and the trace of Section 11 is their rows. If σ_1 returns at $k = 2$, the trace is wrong.

Run 4: the commutator. Write $D_\mu = \partial_\mu + ieA_\mu$. Expand $D_\mu D_\nu \psi$ into five terms; exchange μ and ν; subtract. Observe that four terms cancel in pairs and that $ie(\partial_\mu A_\nu - \partial_\nu A_\mu)\psi$ remains. Then replace ψ by $e^{-ie\chi}\psi$ and A_μ by $A_\mu + \partial_\mu \chi$ in $D_\mu \psi$ and observe the factor $e^{-ie\chi}$ come out whole. This is Proposition 7.5(c) and Theorem 7.7.

A second run by another reader is a repeat; its readings either equal the first or do not, and the comparison is the press check. An exact replay of the first run's record, line by line, observes nothing. The record supplies the procedure; the experience, every observation in it, happens in the one who runs it.

C Epilog: a carrier of charge alone

The body ends on the box, and it should. This appendix is a note left beside it.

Read once more what the device holds. A carrier that is a mark and nothing else, told apart and never looked at. A tally of such carriers, which is a natural number. A charge that comes in wholes on a closed surface. A transport whose antisymmetric part is a quarter turn, the group a charge couples through. A gap, fixed under refinement. A line: everything it does, it does along one dimension, its walks a transfer matrix and its record a single tape. And no volume anywhere, because no reader of length can read one. Taken together, that is the description of a carrier that

carries a charge and nothing else, in one dimension. The body counted the one dimension as the fourth way the device is not yet the electron. It is also the one place such a carrier has been found.

In one dimension an electron does not stay whole. A theory of interacting electrons on a line, begun by Tomonaga and completed by Luttinger, predicts that an injected electron separates into two excitations that travel at different speeds: one carries the spin and no charge, the spinon; the other carries the charge and no spin, the holon. Photoemission from a one-dimensional copper oxide resolved the two dispersions, around two thousand six [17]; tunnelling between quantum wires saw the two velocities, around two thousand nine [16]; and fermionic atoms in optical lattices watched the spin and the charge part company in time, around two thousand twenty [34]. *Measured*.

John Henry does not know where an electron that only carries a charge might exist. The knower may have read of one recently. If so, the identification on offer is narrower than sentence five and nearer to hand: the device is the description of the electron's charge alone, the holon of a line, and the 1.80 parts in ten thousand is the rest of the electron, the part that went the other way. The device carries a sign as well, retained and never read; whether that sign is the holon's phase or a trace of the spin it left behind is not something the record can say. *Proposed*, and the knower's. It would be revised by a measured coupling of a holon that the device's reading does not match, at a scale the device fixes before the comparison.

References

[1] Yakir Aharonov and David Bohm. Significance of electromagnetic potentials in the quantum theory. *Physical Review*, 115(3):485–491, 1959.

[2] Hendrik Pieter Barendregt. *The Lambda Calculus: Its Syntax and Semantics*. North-Holland, Amsterdam, revised edition, 1984.

[3] David Borwein and Jonathan M. Borwein. Some remarkable properties of sinc and related integrals. *The Ramanujan Journal*, 5(1):73–89, 2001.

[4] Stephen A. Cook. The complexity of theorem-proving procedures. In *Proceedings of the Third Annual ACM Symposium on Theory of Computing*, pages 151–158, 1971.

[5] Tamara M. Davis and Charles H. Lineweaver. Expanding confusion: Common misconceptions of cosmological horizons and the superluminal expansion of the universe. *Publications of the Astronomical Society of Australia*, 21(1):97–109, 2004.

[6] Carl de Boor and Robert E. Lynch. On splines and their minimum properties. *Journal of Mathematics and Mechanics*, 15(6):953–969, 1966.

[7] Stanley Deser. Self-interaction and gauge invariance. *General Relativity and Gravitation*, 1:9–18, 1970.

[8] Radu Diaconescu. Axiom of choice and complementation. *Proceedings of the American Mathematical Society*, 51(1):176–178, 1975.

[9] Edsger W. Dijkstra. Letters to the editor: go to statement considered harmful. *Communications of the ACM*, 11(3):147–148, 1968.

[10] Paul A. M. Dirac. Quantised singularities in the electromagnetic field. *Proceedings of the Royal Society of London A*, 133:60–72, 1931.

[11] C. W. F. Everitt et al. Gravity Probe B: Final results of a space experiment to test general relativity. *Physical Review Letters*, 106:221101, 2011.

[12] X. Fan, T. G. Myers, B. A. D. Sukra, and G. Gabrielse. Measurement of the electron magnetic moment. *Physical Review Letters*, 130(7):071801, 2023.

[13] Jean Gallier. Clifford algebras, clifford groups, and a generalization of the quaternions, 2008. arXiv:0805.0311; revised 2014.

[14] Werner Heisenberg. Über den anschaulichen Inhalt der quantentheoretischen Kinematik und Mechanik. *Zeitschrift für Physik*, 43:172–198, 1927.

[15] John C. Holladay. A smoothest curve approximation. *Mathematical Tables and Other Aids to Computation*, 11(60):233–243, 1957.

[16] Y. Jompol, C. J. B. Ford, J. P. Griffiths, I. Farrer, G. A. C. Jones, D. Anderson, D. A. Ritchie, T. W. Silk, and A. J. Schofield. Probing spin-charge separation in a Tomonaga–Luttinger liquid. *Science*, 325:597–601, 2009.

[17] B. J. Kim, H. Koh, E. Rotenberg, S.-J. Oh, H. Eisaki, N. Motoyama, S. Uchida, T. Tohyama, S. Maekawa, Z.-X. Shen, and C. Kim. Distinct spinon and holon dispersions in photoemission spectral functions from one-dimensional $SrCuO_2$. *Nature Physics*, 2:397–401, 2006.

[18] John Kogut and Leonard Susskind. Hamiltonian formulation of Wilson's lattice gauge theories. *Physical Review D*, 11:395–408, 1975.

[19] Leonid A. Levin. Universal sequential search problems. *Problems of Information Transmission*, 9:265–266, 1973.

[20] Martin Lüscher. Topology of lattice gauge fields. *Communications in Mathematical Physics*, 85:39–48, 1982.

[21] David W. Lyons. An elementary introduction to the Hopf fibration. *Mathematics Magazine*, 76(2):87–98, 2003.

[22] Juan Maldacena. The large N limit of superconformal field theories and supergravity. *Advances in Theoretical and Mathematical Physics*, 2:231–252, 1998.

[23] Walther Meissner and Robert Ochsenfeld. Ein neuer Effekt bei Eintritt der Supraleitfähigkeit. *Naturwissenschaften*, 21(44):787–788, 1933.

[24] Peter J. Mohr, David B. Newell, Barry N. Taylor, and Eite Tiesinga. CODATA recommended values of the fundamental physical constants: 2022. *Reviews of Modern Physics*, 97(2):025002, 2025.

[25] Holger B. Nielsen and Masao Ninomiya. Absence of neutrinos on a lattice. I. Proof by homotopy theory. *Nuclear Physics B*, 185:20–40, 1981.

[26] Emmy Noether. Invariante Variationsprobleme. *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse*, pages 235–257, 1918.

[27] Michael E. Peskin and Daniel V. Schroeder. *An Introduction to Quantum Field Theory*. Addison-Wesley, Reading, MA, 1995.

[28] P. M. Platzman and A. P. Mills, Jr. Possibilities for Bose condensation of positronium. *Physical Review B*, 49(1):454–458, 1994.

[29] H. Rauch, A. Zeilinger, G. Badurek, A. Wilfing, W. Bauspiess, and U. Bonse. Verification of coherent spinor rotation of fermions. *Physics Letters A*, 54(6):425–427, 1975.

[30] Andrew Strominger. The dS/CFT correspondence. *Journal of High Energy Physics*, 2001(10):034, 2001.

[31] Leonard Susskind. Lattice fermions. *Physical Review D*, 16:3031–3039, 1977.

[32] David Tong. Quantum field theory. University of Cambridge Part III lecture notes, 2006. Chapter 6: Quantum Electrodynamics.

[33] Alan M. Turing. On computable numbers, with an application to the Entscheidungsproblem. *Proceedings of the London Mathematical Society*, s2-42:230–265, 1937.

[34] J. Vijayan, P. Sompet, G. Salomon, J. Koepsell, S. Hirthe, A. Bohrdt, F. Grusdt, I. Bloch, and C. Gross. Time-resolved observation of spin-charge deconfinement in fermionic Hubbard chains. *Science*, 367:186–189, 2020.

[35] Klaus Weihrauch. *Computable Analysis: An Introduction*. Springer, Berlin, 2000.

[36] S. A. Werner, R. Colella, A. W. Overhauser, and C. F. Eagen. Observation of the phase shift of a neutron due to precession in a magnetic field. *Physical Review Letters*, 35(16):1053–1055, 1975.

[37] Kenneth G. Wilson. Confinement of quarks. *Physical Review D*, 10:2445–2459, 1974.

[38] Kenneth G. Wilson. Quarks and strings on a lattice. In Antonino Zichichi, editor, *New Phenomena in Subnuclear Physics*, pages 69–142. Plenum, New York, 1977.

[39] Edward Witten. Anti de Sitter space and holography. *Advances in Theoretical and Mathematical Physics*, 2:253–291, 1998.

[40] Peter Woit. Clifford algebras and spin groups. Columbia University, Math G4344 lecture notes, 2012. Spring 2012.