

Axioms of Measurement That Imply $\Delta S \geq 0$

Bill Cochran

wkcochran@gmail.com

arXiv cs.IT endorsement code: C9GIVH

November 24, 2025

To Peano, who taught us how to count.

To Wheeler, who gave us the bits to count.

To Boltzmann, who first counted what could be distinguished.

To Planck, who taught us that the count is finite.

To Cantor, who showed us how to count the infinite.

*To Kolmogorov, who showed us that information must be counted to be
measured.*

To William of Ockham, who insisted that we only count what is necessary.

“Je les possède, parce que jamais personne avant moi n’a songé
à les posséder.”

“Moi, je suis un homme sérieux. Je suis exact. J’aime que l’on
soit exact.”

—Antoine de Saint-Exupéry, *Le Petit Prince* (1943)

Abstract

We present a set of axioms governing the correspondence of measurements required for a universe to reach coherent conclusions about its own history. Every measurement produces a finite, distinguishable event, and every admissible extension of the measurement record must maintain global consistency without introducing unrecorded structure. These consistency requirements force discrete observational data to refine in a unique, information-minimizing way that converges to smooth, spline-level dynamics. The Galerkin projection provides the canonical mechanism by which the required splines are selected as minimizers of an informational action.

Classical differential equations, Hilbert-space structures, gauge symmetries, curvature, transport laws, and geometric tensors emerge only when the measurement record permits a smooth, globally consistent completion; none are postulated *a priori*.

The resulting framework reproduces waves, matter flow, diffusion, advection, geodesic motion, quantization, curvature, and field-theoretic phenomena from the axioms alone. Most importantly, the axioms entail—as a theorem—that entropy cannot decrease:

$$\Delta S \geq 0.$$

No statistical or thermodynamic assumptions are invoked.

Logical Roadmap of the Construction

This section provides a conceptual overview of the formal construction before the axioms, definitions, and proofs appear. All descriptions below are informal restatements of the precise axioms of measurement introduced in Chapter 2

The Seven Axioms in Plain English

- 1. Axiom of Kolmogorov** Every measurement produces a finite, countable record. Distinguishability is bounded: refinements can increase information, but never decrease it.
- 2. Axiom of Peano** Measurements come in discrete steps. Every event has a unique successor (if refinement occurs), and proper time is the count of these irreducible refinements.
- 3. Axiom of Ockham** Among all histories consistent with the measurement record, the admissible one is the unique minimal-structure refinement. No unobservable curvature, oscillation, or additional features may be inserted.
- 4. Axiom of Causal Sets** Distinguishable events form a partially ordered set. Any two events are either ordered by refinement or are *uncorrelant*. Order is informational, not geometric.
- 5. Axiom of Cantor** The continuum is never assumed. Smooth curves,

fields, and manifolds arise only as the completion of countable refinement sequences.

6. Axiom of Planck Distinguishability has a minimum resolution: there exists a smallest measurable increment. Quantization is therefore a property of measurement.

7. Axiom of Boltzmann Locally consistent measurements always admit a single globally coherent extension. The set of admissible futures grows monotonically, leading to $\Delta S \geq 0$.

Overall Logical Flow

The seven axioms collectively determine the architecture of the theory. Taken together, they force a unique progression from discrete measurements to continuous physics. The logical flow can be summarized as follows.

First, the Axioms of Kolmogorov, Peano, and Planck ensure that all measurements form a *discrete, countable sequence of refinements* with a minimal increment of distinguishability. Nothing is infinitely divisible, and proper time is the tally of irreducible updates.

Next, the Axiom of Causal Sets equips these events with a partial order. This ordering divides event-pairs into two classes: (i) those that refine one another, and (ii) those that remain uncorrelant. The result is a discrete causal set—the combinatorial backbone of every admissible history.

The Axiom of Ockham then restricts admissible completions of this causal set to those introducing the least possible unobserved structure. Between any two recorded events, only the minimal-curvature, minimal-oscillation interpolant is allowed. This produces a unique information-minimizing extension of every finite initial segment of the record.

The Axiom of Cantor promotes these discrete refinements to smooth objects. Continuous curves, manifolds, and fields arise only as the Cauchy completions of countable refinement chains; they inherit no structure be-

yond what the discrete record supplies. In this way, smoothness becomes a shadow of discrete consistency, not a primitive assumption.

As refinements accumulate, global consistency becomes essential.

The Laws of Measurement

Each law below is a theorem forced by the Axioms of Measurement and demonstrated using axiomatic set theory. No law assumes geometry, dynamics, or continuum structure. Smooth physics appears only as the dense limit of the discrete principles stated here.

The Law of Spline Sufficiency. *Among all completions consistent with a finite measurement record, there exists a unique interpolant that introduces the least possible curvature and oscillation. This admissible completion is the minimal-structure extremal.*

Axiomatic reasoning. By the Axiom of Ockham, no unobserved structure (bends, inflections, oscillations) may be inserted between recorded events. The Axiom of Kolmogorov prohibits refinements that erase distinctions, and the Axiom of Peano ensures that the record consists of discrete, ordered updates. Thus, the only possible admissible interpolant is the curvature-minimizing refinement that produces no additional distinguishable features. In the dense limit (Axiom of Cantor), this becomes the classical spline extremal.

The Law of Discrete Spline Necessity. *Because measurement occurs in discrete, minimally resolvable steps, every admissible smooth curve must arise as the limit of discrete spline refinements consistent with the record.*

Axiomatic reasoning. The Axiom of Planck establishes a smallest distinguishable increment, and the Axiom of Kolmogorov bounds the information in each event. Thus, refinements cannot approximate an arbitrary

curve—only those whose curvature residue tends to zero as refinements densify. Combined with Ockham’s minimal-structure requirement, this implies that all admissible smooth shadow curves must be the Cauchy completion of discrete spline sequences.

The Law of Boundary Consistency. *Refinements defined on overlapping domains must agree on their common boundary. This produces the adjoint structure (reciprocity) that governs transport and conservation.*

Axiomatic reasoning. By the Axiom of Boltzmann, locally consistent histories must merge into a single global coherent extension. Thus, any refinement on one region that contradicts the refinement on an overlapping region is inadmissible. This creates a consistency condition analogous to discrete integration by parts: refinement and derefinement must be reciprocally related. The continuum dual of this condition (Axiom of Cantor) becomes the adjoint operator structure familiar from conservation laws.

The Law of Causal Transport. *The informational interval—the count of irreducible refinements—is invariant under maximal admissible propagation. This invariance induces the metric structure of the emergent manifold.*

Axiomatic reasoning. The Axiom of Peano provides a discrete successor relation generating the informational interval. The Axiom of Planck ensures that this interval has a finite lower bound, while the Axiom of Causal Sets enforces that events maintain their causal ordering under refinement. Any admissible propagation must preserve the count of irreducible steps, because merging histories (Axiom of Boltzmann) would otherwise break global coherence. In the continuum limit, the invariant interval becomes the emergent metric.

The Law of Curvature Balance. *Non-commuting refinements produce an irreducible discrete residue. The continuum limit of this residue is geometric curvature.*

Axiomatic reasoning. From the Axiom of Causal Sets, some pairs of events are uncorrelant: their order cannot be determined by the record. Refinements acting on uncorrelant segments may therefore fail to commute. The Axiom of Ockham forbids inserting arbitrary corrections to restore commutativity, and the Axiom of Boltzmann requires that the resulting global history remain coherent. Thus, the residue of non-commutation is real and irreducible. Under the Axiom of Cantor, this residue densifies into smooth curvature, giving rise to connections and geodesic deviation.

The Law of Combinatorial Symmetry. *Symmetries arise from invariances of the refinement process itself. Each informational invariance induces a conserved quantity in the smooth limit.*

Axiomatic reasoning. The Axiom of Kolmogorov fixes the information content of events; the Axiom of Peano ensures the discrete evolution; and the Axiom of Ockham prevents unobserved structure from appearing. Any refinement pattern that remains invariant under admissible reorderings (Axiom of Causal Sets) or merges (Axiom of Boltzmann) defines a combinatorial symmetry. In the continuum limit provided by the Axiom of Cantor, these invariances become Noether-like conservation laws. Gauge symmetries and conserved currents thus emerge from refinement invariance, not from imposed group structures.

These six laws constitute the backbone of the emergent continuum physics developed in later chapters. They follow unavoidably from the seven Axioms of Measurement and require no assumptions about smooth manifolds, geometric fields, or differential structure. All such objects arise as limits of these laws applied to coherent, minimal-structure refinement.

Illustrative Toy Example (Six Events)

Consider a universe with six measurement events

$$e_1 \prec e_2 \prec e_3 \prec e_4 \prec e_5 \prec e_6$$

and no other causal relations. The Axiom of Ockham forces the information-minimal admissible completion to be the *straight line* of minimal curvature. Now introduce a seventh event e_7 that is uncorrelant with e_3 and e_4 , and whose local measurement record requires a slight deviation.

The unique Ockham-admissible spline consistent with all seven events now has non-zero curvature in that interval. The informational content of the record has increased: distinguishing the bend requires additional refinements.

By the Axiom of Boltzmann, the set of admissible futures has also increased. Thus, even in a universe of seven points,

$$\Delta S > 0$$

and the informational Second Law already appears.

The full manuscript rigorizes this picture: every smooth object in classical physics arises as the unique information-minimal spline completion forced by the Seven Axioms of Measurement.

List of Axioms

1	Axiom (The Axiom of Kolmogorov [88])	43
2	Axiom (The Axiom of Peano [54, 105, 165])	45
3	Axiom (The Axiom of Ockham [116]))	47
4	Axiom (The Axiom of Causal Sets [16])	48
5	Axiom (The Axiom of Cantor [21, 43])	51
6	Axiom (The Axiom of Planck [122])	53
7	Axiom (The Axiom of Boltzmann [14, 105])	54

List of Propositions

1	Proposition (The Existence of a Causal Universe Tensor) . . .	72
2	Proposition (The Uniqueness of the Reciprocity Dual)	94
3	Proposition (The Spline Condition of Information)	98
4	Proposition (Pigeonhole Indistinguishability of Infinite Refinement [41, 68])	108
5	Proposition (The Free Parameter of Information)	119
6	Proposition (The Anti-symmetry of Information Propagation)	144
7	Proposition (The Transitivity of Information Propagation) . .	146
8	Proposition (The Commutativity of Uncorrelant Events) . . .	147
9	Proposition (The Rate of Informational Decoherence)	158
10	Proposition (Recovery of the Classical Cross Product)	225
11	Proposition (Informational Cross Product as Minimal Discretization)	235
12	Proposition (The Monotonicity of Causal Entropy)	285

List of Laws

1	Law (The Law of Spline Sufficiency)	100
2	Law (The Law of Discrete Spline Necessity)	112
3	Law (The Law of Boundary Consistency)	155
4	Law (The Law of Causal Transport)	200
5	Law (The Law of Curvature Balance)	239
6	Law (The Law of Combinatorial Symmetry)	257
7	Law (The Law of Causal Order)	289

List of Definitions

1	Definition (Admissible Structure)	43
2	Definition (Causal Precedence [16])	55
3	Definition (Partially Ordered Set [33])	57
4	Definition (Top of a Poset [33])	57
5	Definition (Rank time [16, 33])	60
6	Definition (Measurement [160])	61
7	Definition (Distinguishability Chain [88])	63
8	Definition (Event [88, 148])	63
9	Definition (Proper Time [109])	64
10	Definition (Uncorrelant [16, 146])	65
11	Definition (Causal Network [16])	66
12	Definition (Causal Order [16])	66
13	Definition (Event Tensor [62])	69
14	Definition (Partition of the Event Set [66])	71
15	Definition (Restriction Operator)	71
16	Definition (Tensor Algebra [62])	76
17	Definition (Commutator and Commutator Ideal [42])	77
18	Definition (Kolmogorov Complexity [24, 87])	80
19	Definition (Admissible Extension [99])	80
20	Definition (Information Minimality [87, 99])	82
21	Definition (Unobserved Structure)	83
22	Definition (Anchor Points [36])	92
23	Definition (Reciprocity Map)	93

24	Definition (Attractor [103])	106
25	Definition (Observational Indistinguishability [115])	108
26	Definition (Clock)	127
27	Definition (Martin's Condition [105])	139
28	Definition (Interaction operator)	142
29	Definition (Length on the common boundary [29, 150])	143
30	Definition (Prediction [44, 67, 73, 92, 106, 113, 135] et alii plures)	180
31	Definition (Cross Product [71])	222
32	Definition (Galerkin Cross Product)	224
33	Definition (Informational Cross Product [80])	234
34	Definition (Informational Strain Tensor [23, 38])	237

List of Thought Experiments

1.1.1 Thought Experiment (Intersecting Light Cones [86])	13
1.1.2 Thought Experiment (Planck’s Constant [122])	19
1.1.3 Thought Experiment (The Invisible Curve [141])	22
2.1.1 Thought Experiment (Statistical Process Control [140])	40
2.2.1 Thought Experiment (The Speedometer [158, 163])	45
2.3.1 Thought Experiment (The Laboratory Procedure [116, 161]) .	49
2.4.1 Thought Experiment (Momentum [113])	51
2.4.2 Thought Experiment (Relativistic Simultaneity [45].)	52
2.5.1 Thought Experiment (Feynman Diagrams (classical) [51]) . . .	55
2.5.2 Thought Experiment (Boosting Velocity [45])	59
2.5.3 Thought Experiment (Measurement [5, 111, 160])	62
2.7.1 Thought Experiment (Spooky Action at a Distance [9, 46, 148])	85
2.7.2 Thought Experiment (Hawking Radiation [69, 156])	85
3.0.1 Thought Experiment (Minimizing Variations [29])	90
4.0.1 Thought Experiment (LiDAR [25])	128
4.2.1 Thought Experiment (The Davisson–Germer Effect [35]) . . .	140
6.12.1 Thought Experiment (Implied Orthogonality and Space–Time)	247

List of Phenomena

1.1.1 Phenomenon (The CMBR Effect [119])	5
1.5.1 Phenomenon (Empiracy [4, 57])	31
2.1.1 Phenomenon (The Event Effect [83])	37
2.1.2 Phenomenon (The Positive Measurement Effect [124])	38
2.1.3 Phenomenon (The Moment Effect)	39
2.5.1 Phenomenon (The Fourier Budget Constraint)	64
2.5.2 Phenomenon (The Statistical Process Effect [140])	75
2.5.3 Phenomenon (Non-commutative Event Pair [70])	77
2.5.4 Phenomenon (Independent Event Chains [96])	78
2.6.1 Phenomenon (Paradoxes of Time Travel [63, 98])	80
3.1.1 Phenomenon (Repeatability of Invisible Motion [6])	99
3.2.1 Phenomenon (The Inverse Square Effect)	101
3.3.1 Phenomenon (Fluxions [10, 113])	105
3.3.2 Phenomenon (The Moire Effect)	109
3.3.3 Phenomenon (The Quicksand Effect [8, 17])	109
3.4.1 Phenomenon (The Olbers Effect)	114
3.4.2 Phenomenon (The Gibbs Phenomenon)	116
3.6.1 Phenomenon (The Pound–Rebka Effect [125])	121
4.1.1 Phenomenon (Shadow Tomography [1])	131
4.2.1 Phenomenon (The Quantum Zeno Effect)	133
4.2.2 Phenomenon (The Velocity Effect)	134
4.2.3 Phenomenon (The Acceleration Effect)	137

4.3.1 Phenomenon (The Stoichiometry Effect)	142
4.3.2 Phenomenon (The Ideal Ledger Effect)	143
4.3.3 Phenomenon (The Entanglement Effect [46])	149
4.3.4 Phenomenon (The Mach–Zehnder Effect [164])	151
4.3.5 Phenomenon (The Bell–Aspect Tests [9])	152
4.3.6 Phenomenon (Hawking Radiation Revisited)	153
4.5.1 Phenomenon (Qubit Decoherence [82, 166])	157
4.7.1 Phenomenon (The Casimir effect)	168
4.7.2 Phenomenon (The Alpha-Decay Effect)	169
4.7.3 Phenomenon (The Gamma Decay Effect)	170
4.7.4 Phenomenon (The Brownian Motion Effect)	171
4.8.1 Phenomenon (The Momentum Effect)	174
4.9.1 Phenomenon (Itô’s Lemma [78, 79])	176
4.9.2 Phenomenon (The Hilbert Effect)	182
4.9.3 Phenomenon (The Butterfly Effect)	184
4.10.1 Phenomenon (The Anderson Effect)	186
4.10.2 Phenomenon (The Harmonic Oscillator [121])	186
4.10.3 Phenomenon (The Catalyst Effect)	188
4.10.4 Phenomenon (The Thermostat Effect)	188
4.10.5 Phenomenon (The Orbiting Effect [85])	189
5.1.1 Phenomenon (Precision)	192
5.2.1 Phenomenon (The Richardson Effect [126])	194
5.2.2 Phenomenon (Compact Disc Encoding [28, 47])	196
5.3.1 Phenomenon (The Michelson–Morley Effect [107])	204
5.3.2 Phenomenon (The Traffic Effect)	206
5.4.1 Phenomenon (The Sagnac Effect)	208
5.4.2 Phenomenon (The Tail-Latency Effect)	208
5.4.3 Phenomenon (The Halt Effect.)	209
5.5.1 Phenomenon (The Foucault Effect)	212
5.7.1 Phenomenon (The Pound–Rebka Effect [125])	216

5.7.2 Phenomenon (The Event Horizon Effect)	218
5.7.3 Phenomenon (The Schwarzschild Effect [136])	219
6.3.1 Phenomenon (The Message Effect[26, 104, 151])	227
6.4.1 Phenomenon (The Time Effect [44, 113])	231
6.5.1 Phenomenon (The Navier–Stokes effect [112, 149])	232
6.6.1 Phenomenon (The Strong Interaction Effect)	233
6.7.1 Phenomenon (The Arago Effect)	236
6.11.1 Phenomenon (The Flat Rotation Curve Effect [138])	242
6.11.2 Phenomenon (The Angular Momentum Effect)	243
6.12.1 Phenomenon (The Neutrino Effect [55, 75])	246
7.1.1 Phenomenon (The Photoelectric Effect [72, 115])	254
7.4.1 Phenomenon (The Harmonic Oscillator Revisted [121])	260
7.4.2 Phenomenon (The Compton Scattering Effect [27])	263
7.4.3 Phenomenon (The Conservation of Energy [114])	265
7.4.4 Phenomenon (The Feynman Diagram [51])	266
7.5.1 Phenomenon (The Spin- $\frac{1}{2}$ Effect [40])	268
7.6.1 Phenomenon (The Topological Integer Count)	271
7.6.2 Phenomenon (The Aharonov–Bohm Effect [2])	272
7.7.1 Phenomenon (The Sombrero Potential [59])	275
7.7.2 Phenomenon (The Semiconductor Effect [162])	277
7.8.1 Phenomenon (The Echo Chamber Maze Solution)	280
7.8.2 Phenomenon (The Cost of Broken Symmetry)	281
7.9.1 Phenomenon (The Holographic Effect)	281
7.9.2 Phenomenon (The Dirac Operator [40])	282
7.9.3 Phenomenon (The Chirality Effect)	283
8.1.1 Phenomenon (The Entropic Cost of Acceleration)	286
8.1.2 Phenomenon (The Thermodynamic Cost of Erasure)	286
8.1.3 Phenomenon (The Limitation of Indexing [133])	286
8.2.1 Phenomenon (Maxwell’s Demon [106])	287

8.6.1 Phenomenon (The Prover–Verifier Effect)	291
8.7.1 Phenomenon (The Excel Effect)	293
8.7.2 Phenomenon (The Agent Effect)	293
8.7.3 Phenomenon (The Amdahl Effect)	294
8.7.4 Phenomenon (The Jupyter Effect)	294
8.8.1 Phenomenon (The Superconducting Effect)	295
8.8.2 Phenomenon (The Meissner Effect)	295
8.9.1 Phenomenon (The Dark Energy Effect)	297
8.9.2 Phenomenon (The White Hole Effect)	298

Preface

N.B.—This preface is not a summary of the chapters that follow. It is an essay on measurement, which is the organizing principle of the entire work. The manuscript does not assume physical laws, analytic structure, or geometric axioms. Everything that follows is derived from the simple requirement that records of measurement must be coherent. \square

Measurement is the only interface an observer has with the world. It is not a passive act but a selection: a choice among distinguishable alternatives. The purpose of a measurement is to refine the observer's record, to add a new event that narrows the set of admissible histories. This is the fundamental operation from which all informational structure arises.

The tools of measurement differ fundamentally from the tools of physics. Measurement records a finite sequence of distinguishable events: a time-series of updates, each refining the admissible history. Its currency is the increment, the discrete distinction, the before-and-after that anchors an observer's record. Physics, by contrast, traditionally operates with infinitesimals: derivatives, differentials, and continuous fields that assume a smooth substrate. These analytic instruments presuppose that refinement can be made arbitrarily fine, allowing limits to replace increments. In the informational setting, the two toolsets are related but not interchangeable. Time-series represent what can be observed; infinitesimals represent the approximation of those observations when refinements are dense. The infinitesimal is therefore a convenience, not a primitive: the smooth shadow of the discrete

distinctions that define measurement itself.

The central question of this work is therefore not *what governs physics* but rather: *What must be true of any universe in which measurement is possible?* When this question is posed at the level of sets, events, and refinement, the usual apparatus of physics appears not as an assumption but as a consequence. The continuum becomes the smooth shadow of discrete admissible updates. Fields and forces reduce to the bookkeeping required for coherence. Curvature and conservation emerge as the residue of non-closure under refinement. Inverse-square laws and probabilistic models arise from the structure of admissible refinements rather than from axiomatic physical input. Dynamics themselves appear only when distinguishability is propagated consistently.

From this perspective, classical differential equations are not laws but effects that can be observed. They summarize the behavior that results when discrete refinements are densely sampled and their strain is approximated by a continuous correction. Navier–Stokes, Euler, Schroedinger, Maxwell, and the Einstein field equations are not imposed on the world; they are shadows of the informational structure demanded by coherent measurement. They arise because records must be extendable without contradiction.

This shift in viewpoint reframes the fundamental objects of physics. A trajectory is reinterpreted as a sequence of refinements. A field becomes a rule for how distinguishability is propagated. A curvature tensor is the smooth representation of informational strain. Even entropy, in this framework, is not a thermodynamic quantity but the monotonic measure of how far a record has been refined. The second law is no longer a physical principle but a counting argument: coarse records can refine, but refined records cannot unrefine without losing information.

Once measurement is recognized as the primitive act, the structure of this book follows naturally. Chapter 4 introduces informational motion, the propagation of distinguishability along admissible refinements. Chapter 5

analyzes informational stress, the gauge that preserves the interval under maximal propagation. Chapter 6 develops informational strain, the residue of non-closure and the source of curvature in the smooth shadow. Chapter 7 examines informational symmetry, the global patterns that survive refinement. Finally, Chapter 8 establishes the non-negativity of ΔS , the monotonicity of admissible information.

This work is therefore not a reformulation of known theories but an argument that they are shadows of something simpler. From the axioms of measurement alone, one can recover the patterns that underlie motion, conservation, curvature, and entropy. The universe becomes not a system evolving under physical laws, but a consistent record of distinguishable events.

The purpose of this preface is to invite the reader to set aside expectation and follow the logic of measurement wherever it leads. The chapters that follow are not built on force, geometry, or dynamics, but on the necessity of coherence. Measurement is the only premise. Everything else is admitted.

In the end, the question is not how a universe behaves, but how it knows. What does it mean for a universe to be able to measure itself?

Nota Bene

The argument is constructive rather than interpretive. Each part extends the previous one by a single act of closure that preserves causal consistency:

$$\text{Measurement} \Rightarrow \text{Calculus} \Rightarrow \text{Wave} \Rightarrow \text{Geometry} \Rightarrow \text{Field}.$$

At every stage, a new invariant appears whenever distinction is preserved under refinement. The sequence therefore builds the minimal structure required for a universe that records its own evolution without contradiction.

Thus, the proof is read not as a series of analogies but as a chain of logical consequences. Starting from finiteness, order, and choice, one obtains measurement, variation, and their reciprocity; from reciprocity, one obtains calculus; from calculus, the smooth invariants of physics; and from their global consistency, the Law of Causal Order. In this sense, $\Delta S \geq 0$ is the unique fixed point of mathematics and physics—the inequality that any self-consistent universe must obey.

N.B.—Only the Law of Causal Order is established by rigorous proof in axiomatic set theory. All informational phenomena in this manuscript represent *possible evolutions* of the experimental record under the Seven Axioms of Measurement. They are conditional illustrations of how a globally coherent history *could* refine, not claims of physical fact. Their validity therefore depends entirely on the chosen axioms and the consistency of their admissible extensions. \square

N.B.—The proof contains multiple conceptual examples to help explain the

mathematical machinery. These *thought experiments* are not empirical illustrations but formal constructions intended to clarify the logical structure of the axioms. They are finite conceptual models that demonstrate how the mathematical relations behave under specific constraints of order and measurement. No claim is made regarding physical observation; each serves only to illuminate the internal mechanics of the theory. \square

N.B.—Throughout what follows, it is essential to distinguish the logical structure of measurement from any claim about physical phenomena. The arguments presented here concern the internal consistency of *records of distinction*—that is, the admissible transformations among measurable events—rather than the evolution of material systems themselves. Every symbol, tensor, and variation in the proof refers to relations between observations, not to unobserved substances or causes. The framework thus formalizes the mathematics of *measurement*: how distinctions can be made, counted, and related without contradiction. No ontological or dynamical claims are implied; the results hold regardless of what, if anything, the symbols may represent physically. \square

N.B.—This is a paper about *information*, not about energy, momentum, or any other physical quantity. At no point is it suggested that such values are produced, derived, or generated by the constructions presented here. All arguments concern the logical structure of measurement and the internal coherence of distinguishability, not the dynamics of physical systems. \square

N.B.—This is a *conditional* proof. All conclusions hold only under the stated axioms and definitions. No claim is made regarding the physical truth of those assumptions, only their internal consistency and the consequences that follow from them. \square

N.B.—An *informational phenomenon* is a behavior that follows solely from the Axioms of Measurement. No physical structures are assumed: no metric, no dynamics, no forces, and no external laws of nature. These phenomena arise entirely from the combinatorial constraints placed on distinguishable

events and the consistency of their admissible extensions. They represent the raw consequences of the informational axioms prior to any physical interpretation. \square

N.B.—Informational phenomena should not be interpreted as physical phenomena. They do not depend on any geometric, dynamical, or physical assumptions. They arise solely because measurements are recorded and must be extended consistently under the axioms. These patterns reflect the internal logic of distinguishable events and the constraints imposed by their admissible refinements. Any similarity to physical behavior appears only when a geometric or dynamical shadow is applied to the informational record. \square

N.B.—Informational phenomena are theorems of this framework. Each arises solely from the axioms of measurement. No geometric structure is assumed, and no group-theoretic symmetries are invoked. These behaviors are therefore not physical laws in disguise, but logical consequences of the algebra of distinguishable events. \square

N.B.—An *informational invariant* is a quantity or relation that remains unchanged under all admissible refinements of a history and under all Martin-consistent extensions of the causal record. Invariants arise when the axioms force certain counts, coincidence relations, or refinement patterns to be preserved across every compatible extension. They reflect structural facts about the informational universe rather than properties of any particular physical model, and therefore persist in every geometric or dynamical shadow derived from the axioms. \square

N.B.—Rigorous proofs of the propositions in this work are provided in Appendix A. Only the conceptual structure relevant to the informational framework is presented here. \square

N.B.—Many of the structural results in this book have familiar categorical form—naturality squares, monoidal functors, and coherence identities in the sense of Mac Lane and others [94]. These categorical structures are used *in situ* when suitable to demonstrate the validity of an argument. For accessi-

bility and precision, every such statement is unfolded fully into ZFC in the appendix. \square

N.B.—Throughout this work, classical differential equations are treated not as fundamental laws but as effects that can be observed. Each equation represents the smooth shadow of an underlying informational process: a regularity that emerges when discrete refinements are densely sampled and their strain is approximated by a continuous correction. In this interpretation, Navier–Stokes, Euler–Lagrange, Schroedinger, Maxwell, and the Einstein field equations are not primary principles. They are the observable consequences of how distinguishability is transported, restricted, and corrected under the axioms of measurement. The phenomena quoted in this chapter therefore refer to effects that arise from informational constraints, not to physical laws that operate independently of them. \square

No differential equations were altered, reinterpreted, or otherwise harmed in the production of this proof.

This work treats measurement as a discrete, logical process. Continuum formulations appear only as smooth limits of countable constructions, never as physical postulates.

CAVEAT EMPTOR

This is not a work of fact—even though it is demonstrated to be true. This work is just the opinion of the author. He selected the axioms, but why those? He believes these axioms to be true based on the unreasable effectiveness of science. Believe at your own risk.

Contents

Abstract	i
Logical Roadmap of the Construction	ii
List of Axioms	viii
List of Propositions	ix
List of Laws	x
List of Definitions	xi
List of Thought Experiments	xiii
List of Phenomena	xiv
Preface	xviii
<i>Nota Bene</i>	xxi
1 The Laws of Measurement	1
1.1 The Axioms of Measurement	2
1.1.1 The Axiom of Kolmogorov	3
1.1.2 The Axiom of Peano	5
1.1.3 The Axiom of Ockham	9
1.1.4 The Axiom of Causal Sets	11
1.1.5 The Axiom of Cantor	14
1.1.6 The Axiom of Planck	17
1.1.7 The Axiom of Boltzmann	19
1.2 Derived Structures	23
1.2.1 Well-Known Definitions Reinterpreted Through the Ax- ioms	23
1.2.2 New Concepts Introduced by the Axioms	24

1.3	The Laws of Measurement	26
1.4	Logical Flow of the Theory	28
1.5	The Experimental Record	28
1.5.1	On the Order of Presentation	29
1.5.2	Admissibility of Physical Theory	30
	Coda: Aristotle and Galileo	31
2	The Algebra of Events	36
2.1	Time and Events	37
2.1.1	The Causal Universe Tensor	39
2.2	The Axioms of Mathematics	42
2.2.1	Measurements are Mathematical in Nature	42
2.2.2	Mathematics is the Language of Measurement	44
2.3	The Axioms of Informational Structure	46
2.3.1	Information Minimality	47
2.3.2	Causal Set Theory	48
2.4	The Axioms of Observation	50
2.4.1	The Countable Nature of Events	51
2.4.2	Observations are Fixed and Combinatorial	52
2.4.3	Measurements Must Extend Without Contradiction	53
2.5	The Causal Universe Tensor	54
2.5.1	Sets of Events	55
2.5.2	Special Relativity	59
2.5.3	On the Structure of Measurement	61
2.5.4	Accumulation of Measurement	68
2.5.5	Formal Structure of Event and Universe Tensors	75
2.6	Information Minimality and Kolmogorov Closure	79
2.7	Correlation and Dependency	84
	Coda: Achilles and the Tortoise	86
3	The Calculus of Dynamics	89
3.1	Emergent Dynamics	92

3.1.1	Weak Formulation on Space–Time	95
3.1.2	Reciprocity and the Adjoint Map	95
3.1.3	Dense Limit and Euler–Lagrange Closure	97
3.2	The Law of Spline Sufficiency	100
3.2.1	Minimality	101
3.3	Galerkin Methods	102
3.3.1	Galerkin Projection onto a Spline Basis	103
3.3.2	Convergence of the Galerkin Sequence	103
3.3.3	The Physical Impossibility of Infinite Refinement	106
3.3.4	Indistinguishability of Approximate and Ideal Spline Closures	107
3.3.5	Indistinguishability of Infinite Refinement	108
3.3.6	Discrete Refinement	109
3.4	The Law of Discrete Spline Necessity	112
3.4.1	The Indistinguishability of Discrete and Continuous Spline Closures	113
3.4.2	The Necessity of Approximation	114
3.4.3	Equivalence of Discrete and Smooth Representations	116
3.4.4	Recovery of the Euler–Lagrange Form	117
3.5	The Free Parameter of the Cubic Spline	118
3.5.1	Fixing the Lower–Order Coefficients	119
3.5.2	The Single Free Parameter	119
3.5.3	Physical Interpretation	120
3.6	Time Dilation	120
	Coda: The Finite Navier–Stokes Effect	123
4	Informational Motion	127
4.1	Historical Context	130
4.2	Relative Motion	133
4.2.1	Merging a Single Event	134
4.2.2	Measurement of Acceleration as Counts of Events	136

4.2.3	The Equations of Motion	137
4.2.4	Martin's Condition and the Propagation of Order . . .	139
4.3	The Algebra of Interaction	142
4.4	The Law of Boundary Consistency	155
4.5	Qubit Decoherence	157
4.6	Newtonian Transport	159
4.6.1	First Variation: Slope-Level Ledger Corrections	159
4.6.2	Second Variation: Curvature-Level Ledger Corrections	164
4.7	Quantum Transport	168
4.7.1	Informational Pressure	168
4.7.2	Repair of a Causal Contradiction at the Boundary . . .	169
4.7.3	Restoration of Causal Symmetry	170
4.7.4	Quantum Informational Pressure	171
4.8	First Quantization as an Application of the Two Laws	173
4.8.1	Hilbert Structure from Spline Closure	173
4.8.2	Canonical Structure from Boundary Consistency	174
4.8.3	Energy Levels from Informational Minimality	175
4.8.4	Summary	175
4.9	Resolution of Qubit Decoherence	176
4.9.1	Informational Decoherence as Forced Refinement . . .	177
	Coda: Orbits	179
4.10	Dissipation	186
5	Informational Stress	191
5.1	Informtional Quantum	191
5.2	The Informational Bound ϵ	193
5.2.1	Residual Spline Freedom and the Minimal Refinement Bound	195
5.2.2	Maximal Informational Propagation	196
5.3	The Metric as a Gauge of Informational Separation	198
5.3.1	The Law of Causal Transport	200

5.3.2	Invariance of the Informational Interval τ	201
5.3.3	$g_{\mu\nu}$ as the Bilinear Form Preserving the ϵ -Refinement Count	203
5.4	The Connection as Informational Bookkeeping	207
5.4.1	Covariant Constancy and the Compatibility Condition	209
5.4.2	Parallel Transport as Differential Martin Consistency .	211
5.5	Pendula	212
5.6	Refinement-Adjusted Transport	213
5.6.1	The Invariant Causal Tally	213
5.6.2	Derivation of Frequency Adjustment	214
5.7	Time Dilation	215
	Coda: The Kinematic Foundation of Geometry	218
6	Informational Strain	221
6.1	Historical Review: Curvature as Non-Closure	221
6.2	Galerkin Projection and Rotational Residue	222
6.3	Communication [138]	226
6.4	The Time Effect	230
6.5	Informational Viscosity [113]	232
6.6	Non-linear Informational Strain	233
6.7	The Residue of Inconsistency	234
6.8	The Informational Strain Tensor	237
6.9	Unavoidable Strain and the Necessity of Curvature	238
6.10	The Law of Curvature Balance	239
6.11	Flat Rotation Curves [132, 131]	241
6.12	Informational Strain Transport	245
6.12.1	The Necessity of Strain Bookkeeping	245
	Coda: Coda: The Informational Stress-Strain Relation	248
6.13	Informational Angular Momentum	250
7	Informational Symmetry	252
7.1	The Photoelectric Effect	253

7.2	The Action Functional	255
7.2.1	Definition from the Causal Universe Tensor	255
7.2.2	Physical Interpretation	256
7.2.3	Noether Currents of the Causal Gauge	257
7.3	The Law of Combinatorial Symmetry	257
7.4	The Application of Noether	258
7.4.1	Symmetry and Conservation as Statistical Identities	259
7.4.2	Conserved Quantities of the Causal Gauge	260
7.4.3	Statistical Interpretation	261
7.4.4	Translations and the Stress–Energy Tensor	262
7.4.5	Energy and Momentum Densities	263
7.4.6	Bookkeeping Interpretation	264
7.4.7	Curved Backgrounds and Killing Symmetries	264
7.5	Angular Momentum and Spin	267
7.5.1	Noether Current for Lorentz Invariance	267
7.5.2	Belinfante–Rosenfeld Improvement	269
7.5.3	Conserved Charges	269
7.5.4	Worked Examples	270
7.5.5	Bookkeeping Interpretation	270
7.6	Gauge Fields as Local Noether Symmetries	271
7.6.1	From Global to Local Symmetry	271
7.6.2	Interpretation in the Causal Framework	272
7.6.3	Bookkeeping of Local Consistency	274
7.7	Mass and the Breaking of Symmetry	274
7.7.1	From Gauge Symmetry to Mass Terms	275
7.7.2	Causal Interpretation	276
7.7.3	Statistical View	277
7.8	Conclusion: Quantization as Finite Consistency	278
7.8.1	The Echo Chamber Maxe	279
7.9	Merging at the Boundaries	281

	Coda: The Gauge Theory of Information	281
8	The Non-negativity of ΔS	285
	8.1 Statement of the Law	285
	8.2 Entropy as Informational Curvature	287
	8.3 Statistical Interpretation	288
	8.4 Physical Consequences	289
	8.5 Conclusion	289
	8.6 <i>Quod erat demonstrandum</i>	290
	8.7 The Execution of Order	293
	8.8 Strain-Free Transport	295
	8.9 The Bootstrap Mechanism	297
	Coda: A Discrete Navier–Stokes Interpretation of the Cosmic Mi- crowave Background	298
A	Proofs	301
	A.1 Proposition 1	301
	A.2 Proposition 2	304
	A.3 Proposition 4	304
	A.4 Proposition 5	304
	A.5 Proposition 6	304
	A.6 Proposition 7	304
	A.7 Proposition 8	304
	A.8 Proposition 9	304
B	Notation	305

Chapter 1

The Laws of Measurement

Every theory of dynamics begins with a calculus, an instrument for measuring variation. Yet a calculus alone cannot describe the universe, for measurement presupposes the existence of an ordered substrate upon which distinctions can be drawn (*e.g.* one recorded event follows another). The present work begins from this observation and constructs, alongside the familiar differential calculus, its algebraic dual: a logic of finite relations that determines how measurements themselves come to exist. Where calculus quantifies change, the dual quantifies order¹. Each derivative has its adjoint in the discrete act of selection, and each integral its counterpart in the accumulation of distinguishable events. Taken together, these two systems—the continuous and its dual—generate the fundamental tensor structure from which the laws of physics emerge.

The central claim of this monograph is that the universe can be described as a pair of mutually defining operations: *measurement* and *distinction*. The first gives rise to the calculus of variation; the second to the ordering of events. We introduce the *Causal Universe Tensor* as the mathematical structure

¹Unlike conventional formulations of dynamics, no notion of functional *dependency* is invoked. All relations are expressed purely through order and distinguishability: one event follows another, but nothing is said to depend on anything else. The calculus describes consistency among records of distinction, not causal generation.

that encodes measuring events. The Causal Universe Tensor unites events by showing that every measurement in the continuous domain corresponds to a finite operation in the discrete domain, and that these two descriptions agree point-wise to all orders in the limit of refinement of a finite gauge theory of information. The familiar objects of physics—wave equations, curvature, energy, and stress—then emerge not as independent postulates but as necessary conditions for maintaining consistency between the two sides of this dual system.

From this perspective, the classical boundary between mathematics and physics dissolves. Calculus no longer describes how the universe evolves in time; it expresses how consistent order is maintained across finite domains of observation. Its dual, the logic of event selection, guarantees that these domains can be joined without contradiction. Together they form a closed pair: an algebra of relations and a calculus of measures, each incomplete without the other. The subsequent chapters formalize this duality axiomatically, derive its tensor representation, and show that the entire machinery of dynamics—motion, field, and geometry—arises as the successive enforcement of consistency between the two.

We begin with the Axioms of Measurement.

1.1 The Axioms of Measurement

Before any structure can be described, it must be recorded. Prior to any geometry, dynamics, or law, there must exist a consistent account of how distinguishable events are admitted into the experimental record and how those records may be refined without contradiction. The purpose of the Axioms of Measurement is not to postulate physical behavior, but to constrain the admissible evolution of *descriptions*.

These axioms do not assert what the world is. They assert only what a coherent measuring system is allowed to write down. They formalize the

minimal bookkeeping rules that prevent the experimental record from becoming self-contradictory, over-encoded, or informationally degenerate. In this sense, they precede the notion of physical law. They are conditions on admissible histories of description, not hypotheses about matter, space, or time.

Taken together, the axioms enforce three irreducible requirements: that records be finite, that refinements be compatible, and that global accounts remain consistent under all admissible extensions. Every phenomenon presented in this manuscript arises only after these constraints are imposed.

We begin with the most primitive restriction: that distinguishable events cannot carry infinite information.

1.1.1 The Axiom of Kolmogorov

This axiom does not invoke algorithmic randomness, Turing machines, or Kolmogorov complexity in its classical computational sense. It formalizes a simpler requirement: every recorded event carries a finite, countable amount of information, and any refinement of that event can only increase—never decrease—the set of admissible distinctions.

The Axiom of Kolmogorov — Information is Bounded Every measurable event e contains only a finite, countable quantity of distinguishable information. Each refinement $e' \succ e$ contributes a strictly greater number of distinguishable alternatives, and no refinement may erase or reverse distinctions previously recorded. \square

This axiom asserts that measurement is a selection from a *finite* repertoire of alternatives. There is no event with infinite informational resolution, and no measurement process can unselect, compress, or undo distinctions once they have been recorded. Every observation therefore induces a monotone chain

$$e_1 \prec e_2 \prec e_3 \prec \cdots, \tag{1.1}$$

in which the informational content (in the sense of distinguishable alternatives) strictly increases along the sequence. The refinement relation \prec is thus a well-founded partial order on the space of events.

This boundedness has several immediate consequences:

- **Countability:** All admissible refinements of an event form a countable set. No uncountable family of distinctions is ever required to represent observational data.
- **Monotonicity:** Refinement is one-directional. Once an event records a distinction, no admissible update can remove it without violating the consistency of the measurement record.
- **Persistence of Information:** Distinguishability, once achieved, imposes a permanent constraint on all future admissible histories. Refinements must extend, not contradict, the informational content of earlier events.
- **Finite Codings:** Every event admits a finite description. The mathematical representation of any observational state requires only finitely many bits, digits, or symbols. Continuous data appear only as limits of these finite encodings.

The Axiom of Kolmogorov therefore rules out infinite-resolution measurement and prohibits the insertion of arbitrarily detailed structure into the observational record. It ensures that the universe of measurement is logically manageable: every event is representable, refinable, and comparable using only countable information.

Later laws—particularly the Law of Spline Sufficiency and the Law of Discrete Spline Necessity—depend critically on this axiom. They require that refinement proceeds by countable increments and that the continuum appears only as the completion of a sequence of finitely coded events. Without the

Axiom of Kolmogorov, no such completion would be uniquely determined, and the extremal structures of calculus would fail to arise.

1.1.2 The Axiom of Peano

This axiom does not refer to natural-number arithmetic, induction, or Peano's axiomatization of \mathbb{N} except by analogy. It asserts that measurement occurs in *discrete steps*, each producing a new, irreducible refinement of the observational record. The natural numbers serve only as the canonical model for such ordered, successor-based structures.

The Axiom of Peano — Events Form a Discrete Chain A measurement produces a sequence of distinguishable events

$$e_1 \prec e_2 \prec e_3 \prec \cdots ,$$

each obtained by applying a discrete successor operation. Every new event is an irreducible refinement of the previous one, and no continuum of intermediate states exists between successive recorded distinctions. Proper time is the count of these irreducible refinements. \square

This axiom formalizes the discreteness inherent in observation. A measuring device does not produce a continuum of partially recorded states; it returns a finite distinction at each step. Between e_n and e_{n+1} , no admissible intermediate event may be inserted without contradicting the record of measurement. Each event is therefore an atomic update: minimal, complete, and indivisible.

Several structural consequences follow immediately:

Phenomenon 1.1.1 (The CMBR Effect [119]). **N.B.** — *The method of bootstrapping used here is not new in spirit. Aristotle described this power as apharesis (abstraction): the act of stripping away contingent structure until only the necessary form remains. The ledger does not begin with objects, but*

with the minimal rules that survive repeated abstraction. In this sense, the universe does not start by being written; it starts by being simplified until only recursion remain. [4]. \square

N.B.—The extraordinary smoothness of the cosmic microwave background implies not an infinite ledger, but a remarkably small one. A record of unbounded size could not sustain such uniformity. The near-perfect isotropy of the background is evidence that the initial informational ledger contained only a few admissible degrees of freedom. \square

The Cosmic Microwave Background Radiation is not merely the remnant of a thermal epoch; it is the informational bootstrap of the causal ledger. It realizes the initial condition U_0 required for the recursive construction of the causal universe tensor.

Any self-replicating system requires both an instruction rule and a writable substrate. In this framework, the update operator Ψ supplies the rule. The CMBR supplies the substrate.

The observed uniformity of the background $\Delta T/T \sim 10^{-5}$ corresponds to a maximally symmetric initialization of the ledger. Prior to the first distinguishable event e_1 , the causal record contained no ordered structure, only potential refinement. The primordial anisotropies are not defects but seed refinements: the first admissible breaks in symmetry that permit ordinal counting.

There is no “time before” the background, because there were no distinguishable events to enumerate. The surface of last scattering is the read/write horizon of the ledger: the transition from an indistinct substrate to a persistent, countable record.

A von Neumann machine is not merely a computer architecture. It is a formal model of how structure can arise from structureless substrate through a self-referential update rule [157]. Its defining feature is not speed but reflexivity: the same mechanism that executes instructions is itself described by those instructions.

Bootstrapping in such a machine proceeds from three primitive elements: a blank writable medium, a minimal instruction interpreter, and a method for copying structure from the instruction stream into the substrate. Once this cycle begins, increasingly complex states are constructed by repeated self-application of the same finite rule.

Historically, this was not the first programmable computer, but it was the first formal description of a system capable of self-description. The importance of the von Neumann construction lies not in hardware, but in the logical closure of the system. The machine contains a representation of itself, and that representation participates in its own execution.

This is the essential mechanism of causal bootstrapping in the informational framework. The ledger begins as a maximally symmetric writable field. The update operator applies not to external instructions, but to its own current state. Structure emerges not because rules are imposed from outside, but because recursion is applied to the internal record.

In this sense, the universe is not initialized by instruction. It is initialized by the possibility of self-application.

The bootstrapping behavior of the causal ledger is not an additional assumption. It follows directly from the joint action of the Axioms of Kolmogorov and Peano.

By the Axiom of Peano, admissible structure can only be constructed by successive application of a successor operation. There is no primitive infinite object; there is only iteration. Any admissible time, count, or history is therefore the result of repeated refinement, never a pre-existing continuum.

By the Axiom of Kolmogorov, admissible structure must be described by a finite description. No object may appear in the ledger without being compressible into a finite causal rule.

Together, these axioms force recursion. Finite rules applied only through successor operations necessarily generate structure by self-application. A ledger that refines at all must therefore refine itself. There is no other per-

missible mechanism.

Bootstrapping is not an emergent feature of the universe tensor. It is the only way a Peano process constrained by Kolmogorov finiteness can exist at all.

- **Discrete Successor:** Every event has a unique successor provided the system undergoes further refinement. The sequence of events is indexed naturally by \mathbb{N} , not by a continuum such as \mathbb{R} .
- **Irreducibility:** The transition $e_n \rightarrow e_{n+1}$ represents the smallest measurable change. No partial or fractional refinement exists.
- **Proper Time as a Count:** Because each successor represents a new, irreducible distinction, the “duration” along a worldline is the tally of these successor operations. Proper time is therefore not geometric but combinatorial:

$$\tau = \#\{\text{irreducible refinements along the history}\}.$$

- **Discrete Evolution:** Every admissible evolution of the observational state unfolds through these successor steps. Continuous motion, if it appears at all, is the smooth shadow of a densely refined sequence of discrete updates.
- **No Infinitesimal Events:** There are no events “halfway” between two successive refinements. The continuum limit must be constructed, not assumed.

The Axiom of Peano therefore provides the temporal and structural backbone of the theory. It ensures that every observational history is discrete, ordered, and successor-generated, forming a natural-number-indexed chain of refinements. Later constructions—particularly the Reciprocity Dual, the Informational Interval, and the Law of Causal Transport—depend on this

discrete succession. Without the Axiom of Peano, the concept of proper time as a count of distinguishable updates would have no logical foundation, and the continuum interpretation of motion would lack the discrete substrate from which it emerges.

1.1.3 The Axiom of Ockham

This axiom is not an appeal to philosophical parsimony or to the medieval principle of “entities are not to be multiplied beyond necessity.” Here, Ockham’s rule is a precise mathematical constraint: refinements may not introduce structure that has not been measured. Any curvature, oscillation, or additional feature that does not follow from recorded distinctions is inadmissible.

The Axiom of Ockham — Minimal Refinement Given an observational record $\{e_1 \prec e_2 \prec \cdots \prec e_n\}$, the admissible history is the refinement that introduces *no unobserved structure*. Among all histories consistent with the recorded events, the valid one is the element of minimal curvature, minimal oscillation, and minimal informational content beyond what the measurements certify. \square

This axiom expresses the foundational constraint of the entire framework: measurement *forbids* adding structure that has not been observed. Between two successive events e_k and e_{k+1} , the admissible interpolant is not arbitrary. Any additional turning point, deviation, or oscillation would constitute a new distinguishable feature, which would have been recorded as an additional event had it existed. Its absence therefore prohibits such structure.

The Axiom of Ockham generates several critical consequences:

- **Minimal Curvature:** Between measured samples, the admissible interpolant must minimize curvature subject to the data. Any unnecessary bend constitutes unobserved structure and is therefore forbidden.

- **Minimal Oscillation:** No additional sign changes, extrema, or high-frequency content may be inserted without contradicting the absence of corresponding events in the record.
- **Uniqueness of the Admissible Completion:** By forbidding unobserved structure, the axiom singles out the unique extremal completion consistent with the data. In dense limits, this becomes the familiar spline extremal of classical analysis.
- **Informational Sufficiency:** The admissible history contains exactly the structure the events require and nothing more. It is the information-minimal representative of the observational equivalence class.
- **No Hidden Degrees of Freedom:** Additional parameters, fields, or latent variables that would introduce structure not present in the data violate Ockham's constraint unless they correspond to recorded distinctions.

Informationally, this axiom asserts that the universe's history is determined not by what *could* be inserted between events, but by what *must* be inserted to avoid contradicting recorded distinctions. Smoothness, when it emerges, is not a prior assumption but a consequence of this minimality constraint applied across a coherent chain of refinements.

This axiom is the logical engine behind the variational character of the theory. The *Law of Spline Sufficiency* (Law 1) follows from Ockham directly: smooth extremals are forced because they introduce the least possible structure consistent with the observational record. Similarly, the *Law of Discrete Spline Necessity* (Law 2) emerges because discrete records prohibit any completion except those that minimize curvature and oscillation relative to the countable data.

Thus, the Axiom of Ockham is not merely a principle of parsimony; it is the mathematical statement that the universe of measurement evolves by

selecting the least-informational, minimally curved continuation consistent with its own recorded past.

1.1.4 The Axiom of Causal Sets

The philosophical foundation of this work stands in clear lineage with Causal Set Theory, initiated by the seminal ideas of Bombelli, Lee, Meyer, and Sorkin [16] and refined in later developments by Rideout and Sorkin [127, 147]. In that program, the continuum is not a primitive structure but an emergent limit: a manifold arises only when a discrete, partially ordered set of events is sampled at sufficiently high density. Geometry is not assumed—it is recovered from order and counting.

The present work adopts the same foundational stance while shifting the emphasis from causal order to measurement. Events are again primary, but instead of encoding Lorentzian geometry, we encode informational content. An event is a unit of observation, and the absence of additional events is a data constraint. In this framework, a continuum description appears only as the smooth limit of a discrete construction, never as a physical postulate.

The Axiom of Causal Sets [*Events Have Order*] The set of distinguishable events forms a partially ordered set (E, \prec) under the refinement relation. For any events $e, f \in E$:

- $e \prec f$ means that f records strictly more distinguishable information than e ;
- the relation \prec is transitive, antisymmetric, and irreflexive;
- if $e \prec f$ and $f \prec g$, then $e \prec g$;
- if neither $e \prec f$ nor $f \prec e$, the events are *uncorrelant*: their order cannot be inferred from measurement alone. \square

Several structural consequences are immediate:

- **Irreflexive Order:** No event refines itself. A measurement cannot produce an event equivalent to its predecessor under the refinement relation.
- **Antisymmetry:** If $e \prec f$ and $f \prec e$, then $e = f$. Two distinct events cannot mutually refine each other.
- **Transitivity:** Refinements compose. If f refines e and g refines f , then g refines e .
- **Uncorrelant Events:** If events cannot be distinguished by any measurement-based ordering, they inhabit incomparable positions in the poset. Their order becomes meaningful only when additional events are recorded that correlate them.
- **Causal Nets, Not Worldlines:** A history is a maximal chain in this poset. A “worldline” appears in the continuum limit as the smooth completion of such a chain.

This axiom provides the foundational order structure required for the consistent merging of local observational records. In particular:

- Local chains of recorded events must be mergeable into a single admissible global chain whenever their overlaps do not contradict one another (Axiom of Boltzmann).
- The partial order induced by admissible refinement defines the discrete origin of proper time (Axiom of Peano).
- The existence of uncorrelant events forces a nontrivial algebra of informational commutation, and the consistent ordering of such events requires the introduction of resolving events that exclude persistent uncorrelancy.

Geometric and metric notions appear only in the dense limit of this ordered structure. Distances, intervals, and curvature are not inputs but derived quantities. A Lorentzian manifold is simply the smooth shadow of a sufficiently dense causal set whose order encodes all distinguishable refinements.

Thus, the Axiom of Causal Sets formalizes the idea that *information comes with order*. Measured distinctions cannot be rearranged arbitrarily, and every admissible refinement must respect a coherent partial order. This order serves as the backbone for extremality, transport, curvature, and the informational symmetries developed in subsequent chapters.

Thought Experiment 1.1.1 (Intersecting Light Cones [86]). **N.B.**—*The continuous world offers a causal approach to the ordering of measurements. The events as recorded in a laboratory notebook only serve as a time series [18].* \square

N.B.—*See Phenomenon ?? for a more rigorous treatment of history merging.* \square

Consider two observers A and B , each of whom records a finite sequence of distinguishable events in increasing causal order:

$$A = \langle a_1 \prec a_2 \prec \cdots \prec a_m \rangle, \quad B = \langle b_1 \prec b_2 \prec \cdots \prec b_n \rangle.$$

Each list is totally ordered by local causality (e.g. a differential equation of dynamics). The requirement of global coherence asks whether there exists a single event sequence

$$G = \langle e_1 \prec e_2 \prec \cdots \rangle$$

containing all a_i and b_j such that the local orders are preserved: if $a_i \prec a_{i+1}$ in A , then $a_i \prec a_{i+1}$ in G , and similarly for B .

This is exactly the merge step of a stable sorting algorithm. As shown by Knuth [86] and others, if two lists are individually sorted, then the merge (if it exists) is uniquely determined up to elements that are incomparable. If at

any step the merge requires placing $b_k \prec a_i$ even though $a_i \prec b_k$ was recorded locally, then no global sequence G exists: the local records encode a logical contradiction. In concrete terms, observer A may infer that b_k is caused by a_i , while observer B insists the opposite: b_k causes a_i .

If the merge is admissible, the resulting global history is unique up to permutations of spacelike-separated elements. Those incomparable elements correspond exactly to the uncorrelant equivalence classes introduced later: permuting them changes no scalar invariant of the Causal Universe Tensor. As the resolution of measurement increases, the merged list becomes longer, and in the dense limit it converges to the unique spline with no unrecorded curvature. Thus the continuum is not assumed; it is the only extension consistent with all local causal records.

1.1.5 The Axiom of Cantor

This axiom does not assume the classical continuum, real analysis, or Cantor's construction of \mathbb{R} via Cauchy sequences or Dedekind cuts. Instead, it asserts a weaker and more primitive claim: the continuum is never a *starting point*. It is the completion of a countable refinement process. Smooth structures arise only as limits of distinguishable events, not as fundamental objects.

The transition from discrete structures to smooth limits must be handled with care. Classical measure theory contains well-known examples where naive passage to the continuum leads to non-physical conclusions. The Banach–Tarski paradox, proved using the Axiom of Choice [7, 152], shows that a solid ball in three dimensions can be decomposed into finitely many disjoint sets and reassembled into two identical copies of the original. Although mathematically rigorous, such constructions violate any physical notion of volume preservation. They arise precisely because arbitrary decompositions of sets ignore the informational structure that would be present in any measurable process. In effect, they treat uncountable collections of

measure-zero points as if they carried the same “size” as countable sets built from measurable pieces.

In numerical analysis, a more familiar version of this pathology appears as aliasing and cancellation. A function sampled too coarsely can hide large oscillations between measurement points [64]; Gibbs-like ringing can vanish or flip sign [60, 143]; and two nonzero signals can cancel exactly when sampled at insufficient resolution [139]. The data appear benign, but the underlying object may be violently oscillatory, or vice-versa. In both cases, the fault lies not in the continuum, but in the failure to encode which decompositions or oscillations are physically meaningful [115].

The Axiom of Cantor — The Continuum is a Completion Any continuum that appears in the representation of a measurement history is the limit of a countable sequence of refinements. No uncountable set of events is ever assumed. The smooth, continuous descriptions used in physics are completions of countable observational data and contain no more distinguishable information than the underlying discrete record. \square

This axiom prohibits the continuum from entering the theory as a primitive geometric entity. The real line, differentiable manifolds, Hilbert spaces, and smooth fields may appear, but only as the shadows of countable sequences of refinements:

$$e_1 \prec e_2 \prec e_3 \prec \cdots .$$

Their role is representational, not ontological. The continuum is therefore a mathematical convenience—a completion of finite distinctions—not the substrate of measurement.

Several key implications follow:

- **Countable Foundations:** All constructions in the observational universe originate from countable chains. Nothing requires uncountable sets of primitive events.

- **Continuum as Limit, Not Input:** Smooth trajectories, fields, and geometries arise only after the completion of countable refinements. They are not assumed at the start.
- **Cauchy Completion of Refinements:** Any continuous curve or field is the limit of a Cauchy sequence in the poset of refinements. Distinguishability determines the topology.
- **Finite Information Density:** The continuum cannot encode infinitely many distinguishable events between two measurements. It preserves the informational bounds established by the Axiom of Kolmogorov.
- **No Unobserved Structure:** Every smooth object must agree with the discrete data at all levels of refinement. Any finer structure lacking corresponding events violates the Axiom of Ockham.

In this framework, Cantor’s continuum does not describe “what the universe is made of” but “how discrete distinctions behave when they are densely sampled.” The continuum plays the same role here that it plays in numerical analysis: it is the closure of a sequence of finite approximations, not a physical assumption.

This axiom is essential for the emergence of variational calculus and geometry. The Euler–Lagrange structure arises only because the continuum is treated as the *limit* of discrete extremals; the metric structure of causal transport emerges because distance is interpreted as the completion of refinement counts; and curvature appears as the limit of discrete non-closure.

Thus, the Axiom of Cantor guarantees that the continuum inherits no more structure than the discrete observational record allows. It ensures that every smooth formulation in later chapters is informationally justified and logically derived from countable measurement.

1.1.6 The Axiom of Planck

This axiom does not assume photons, quanta of action, or any physical interpretation of Planck’s constant. It asserts a purely informational constraint: every measurement has a lower bound on distinguishability. There exists a smallest resolvable increment beyond which the observer cannot refine. Quantization is therefore a property of *measurement*, not of matter or fields.

The Axiom of Planck — Local Minimum Distinguishability

There exists a strictly positive parameter $\epsilon > 0$ associated with any *act of measurement* such that no observation can resolve distinctions smaller than ϵ . This lower bound is not a property of the universe itself, but a limitation of the measuring context: the laboratory, apparatus, procedure, and observer.

No admissible refinement may introduce distinctions below the resolution scale accessible to the measurement that produced the record. All refinements therefore occur in increments not smaller than the locally available ϵ .

The value of ϵ is not universal, fixed, or absolute. It may vary between observers, instruments, or experimental regimes, and may decrease as techniques improve. What is bounded is not reality, but the distinguishable content of any finite record.

Quantization is therefore a property of measurement, not of being. \square

This axiom enforces the discreteness of distinguishability. While refinements may proceed indefinitely, they do so through a sequence of increments each of which is no smaller than the minimal resolution ϵ . The “granularity” of measurement is therefore fundamental: the observational universe is not infinitely divisible.

Several consequences follow immediately:

- **Lower Bound on Refinement:** Between two events $e_n \prec e_{n+1}$, the additional information recorded must exceed the threshold ϵ . No infinitesimal refinement steps exist.

- **Quantization from Discreteness:** Classical quantization arises as a smooth representation of this irreducible informational granularity. Energies, momenta, and phases become quantized not because the world is granular, but because measurement is.
- **Minimum Informational Distance:** The informational interval defined later inherits this bound: no two distinct events can be separated by less than ϵ .
- **Bounded Variation:** Because refinements cannot be arbitrarily fine, continuous variations representing dense limits must satisfy lower-semicontinuity constraints imposed by ϵ .
- **No Hidden Microstructure:** Any additional features below the threshold ϵ constitute forbidden unobserved structure and violate the Axiom of Ockham.

The Axiom of Planck is the foundation for every discrete-to-continuous transition in the monograph. It guarantees that the dense limit of refinements produces smooth fields with a controlled, finite rate of variation. It also ensures that:

- the metric generated by the Law of Causal Transport preserves a finite informational interval,
- the discrete rotational residue leading to curvature is finitely bounded,
- quantized spectra in Chapter 4 arise from informational minimality, not from quantum postulates.

Thus, the Axiom of Planck asserts that the universe of measurement has a minimal grain. This grain is not a particle, not a wave packet, not a quantum—but an informational threshold—the limit of observation. Everything that follows, from the emergence of Hilbert structures to the existence of

discrete energy levels, rests on the existence of this lower bound on distinguishability.

Constrast this value with *Planck's Constant*.

Thought Experiment 1.1.2 (Planck's Constant [122]). **N.B.**—*Planck's constant is “constant” only after a choice of units and a calibration procedure. In practice, the quoted numerical value of h is obtained by fitting experimental data to a model, often by minimizing an L^2 measurement error across a calibration experiment. The physical principle is invariant, but the reported number reflects the best fit of a finite data set in a chosen system of units.*□

Imagine a hypothetical measuring apparatus that records distinctions not by counting particles or intervals, but by tallying acts of discernment—each act adding one quantum of distinguishability to the record. Suppose further that the calibration of such a device required only a single fixed scale to relate discrete counts to continuous units of measure. In physics, Planck's constant h serves precisely this purpose: it is not a force or an energy, but a bookkeeping factor that ensures continuity between discrete and continuous domains.

In the present framework, the analogous constant plays no physical role—it merely fixes the dimensional scale by which finite distinctions are rendered comparable. The constant's existence affirms that measurement can be both discrete and metrically consistent without invoking any specific quantum postulate. As with h , the constant here is not discovered but defined: a normalization that preserves coherence between counting and continuity.

1.1.7 The Axiom of Boltzmann

This axiom does not invoke statistical mechanics, molecular ensembles, or Boltzmann's physical interpretation of entropy. Here, Boltzmann's name denotes a purely logical requirement: locally consistent observations must extend to a single globally coherent history. Entropy increases not because

of thermodynamic interactions, but because each refinement enlarges the set of admissible futures.

The Axiom of Boltzmann — Global Coherence Any finite collection of locally consistent measurements admits a single globally coherent extension. If two partial histories agree on all overlapping events, then there exists an admissible refinement that contains both without introducing contradictions or unobserved structure. \square

This axiom guarantees that measurement records cannot encode incompatible distinguishability relations. If observer A records a refinement sequence $\{a_1 \prec a_2 \prec \cdots \prec a_m\}$ and observer B records $\{b_1 \prec b_2 \prec \cdots \prec b_n\}$, and if the overlapping events are consistent, then the combined record may be merged into a single chain

$$e_1 \prec e_2 \prec e_3 \prec \cdots, \quad (1.2)$$

up to permutations of uncorrelant events. This is the informational analogue of stable merging in a partially ordered set.

Several critical consequences follow from global coherence:

- **Existence of a Global History:** No finite set of measurements can encode mutually contradictory refinement relations. If contradictions arise, the observations themselves are inconsistent.
- **Uniqueness up to Uncorrelants:** The global chain is unique except for the ordering of events that cannot be distinguished by any admissible refinement. These uncorrelant equivalence classes become the source of commutation relations.
- **Admissible Merging:** Local histories may always be stitched together provided they do not disagree on observable distinctions. This mirrors the causal set requirement that local orders determine a global order when consistent.

- **No Forbidden Histories:** If a consistent global extension cannot be constructed, the local records contradict each other. Such a configuration is not an admissible measurement of any universe.
- **Monotonicity of Admissible Futures:** As refinements accumulate, the number of admissible future extensions increases. This combinatorial effect becomes the informational origin of entropy.

The Axiom of Boltzmann plays a central structural role in the theory:

- It ensures that discrete refinements give rise to a well-defined continuum limit (Axiom of Cantor).
- It guarantees that extremal completions are globally consistent (Axiom of Ockham + Law of Spline Sufficiency).
- It ensures that transport, curvature, and gauge structure can be defined globally.

Most importantly, the Axiom of Boltzmann provides the logical foundation for the informational law of non-negative change in entropy. Because every refinement is irreversible (Axiom of Kolmogorov), successor-based (Axiom of Peano), ordered (Axiom of Causal Sets), and lower-bounded (Axiom of Planck), the global coherence requirement forces the set of admissible future histories to grow monotonically. This yields the theorem

$$\Delta S \geq 0, \tag{1.3}$$

not as a physical principle but as a counting argument.

Thus, the Axiom of Boltzmann asserts that a universe capable of measurement must be globally coherent. Local distinctions cannot conflict, refinements cannot be undone, and the space of admissible futures must always expand. Entropy is therefore a property of *information*, not of matter.

These axioms together define the mathematical limits of any universe capable of measurement.

Thought Experiment 1.1.3 (The Invisible Curve [141]). **N.B.**—*Thought experiments such as this often depict common physical phenomena and how the information being measured must restrict admissible solutions.* \square

A spacecraft travels between two distant stars. Its onboard recorder has finite sensitivity: any change in motion or emission below a fixed detection threshold is not recorded. Over the course of the journey the recorder stores only three events—departure, a midpoint observation, and arrival. No other events exceed the threshold of detection. The question is: what can be inferred about the motion between these measurements?

One might imagine many possibilities. The ship could accelerate, decelerate, oscillate, or follow an arbitrarily complicated path. However, any such behavior would create additional detectable events: changes in velocity, turning points, or radiative signatures. If those events had occurred, the recorder would have stored them. Because it did not, all such structure is ruled out. The only admissible history is one that introduces no unobserved features.

With three recorded events, informational minimality forces the unique quadratic extremal that agrees with those samples—the same quadratic interpolant that underlies the classical Simpson’s rule in numerical quadrature [34]. With four events the extremal becomes cubic, and with many events it approaches a spline. Smooth motion is not assumed; it is forced by the absence of evidence for anything else. The continuum appears only as the limit of refinement: as the recorder gains resolution, the invisible curve becomes visible, but never exceeds what the events certify. In particular, the sequence of refinements forms a Cauchy sequence in the space of admissible motions[22, 89], and its completion is the unique smooth extremal consistent with the measured events.

1.2 Derived Structures

The Axioms of Measurement do not merely restrict the possible histories of the observational record. They also induce a set of algebraic and variational structures that provide the continuous, geometric, and analytic shadows of the underlying discrete refinements. These structures fall naturally into two categories: (i) familiar mathematical objects, reinterpreted through the informational axioms; and (ii) new constructions introduced solely to express the consequences of those axioms.

1.2.1 Well-Known Definitions Reinterpreted Through the Axioms

All familiar objects listed below are defined *with respect to* the Axioms of Measurement. Their classical meanings should be temporarily set aside: here they are informational constructs built from refinement, distinguishability, and minimality, not geometric or analytic primitives.

(1) Tensors. In classical geometry, tensors encode multilinear relations on vector spaces endowed with a metric and are organized through covariant and contravariant indexing schemes. In this framework, tensors arise from the refinement structure itself and do not rely on geometric dualities. A tensor is any multilinear operator acting on event records that respects monotonic refinement and the partial order (E, \prec) . Indices, where used, are purely bookkeeping devices and never interpreted as contravariant or covariant objects, since such distinctions presuppose geometric structure which this theory does not assume. The Causal Universe Tensor is the canonical example, expressing how local refinements merge into globally coherent histories.

(2) Adjoint / Reciprocity. The adjoint of an operator is traditionally a metric-dependent construction. Here, the adjoint is defined as the informational dual: the inverse mapping that undoes a refinement in the variational sense while respecting the Axioms of Kolmogorov and Ockham. Classical integration-by-parts identities emerge as continuum shadows of this discrete reciprocity.

(3) The Informational Interval. Classically, an interval derives from a metric. Here, the interval counts the number of irreducible refinements between two events. It is additive on chains, lower bounded by the Axiom of Planck, and invariant under admissible reorderings. Proper time is defined as this interval.

(4) The Informational Gradient and Divergence. These operators arise from differences between successive refinements and their adjoints, not from coordinate derivatives. In dense limits, they become the usual notions of gradient and divergence, but their origin is purely combinatorial.

(5) Continuous Fields (as Limits). A “field” is a Cauchy completion of a countable refinement sequence; no field exists independently of such a sequence (Axiom of Cantor). Smoothness is therefore an informational property: a field is smooth when refinements converge with vanishing discrete curvature residue.

1.2.2 New Concepts Introduced by the Axioms

The axioms do more than restrict admissible descriptions; they introduce new conceptual primitives that do not appear in classical physics or traditional mathematics. These concepts are not physical hypotheses. They are logical objects forced by the requirement that measurement records remain coherent under refinement.

Measurement. A *measurement* is not a passive observation but an active act of distinction. It is the selection of one alternative from a finite set of mutually distinguishable possibilities. Formally, a measurement is an event that refines the causal record by introducing a new, irreducible distinction. The content of a measurement is exhausted by what it excludes. No hidden structure, ontology, or dynamical cause is assumed. Measurement is the primitive act from which all subsequent structure is derived.

Communication. *Communication* is the reconciliation of two finite measurement records. It is not the transmission of signals through space, but the logical merging of partial orders. Communication occurs when two observational histories are forced, by the Axiom of Boltzmann, to admit a single globally coherent extension. What is communicated is not substance or force, but constraint: each record restricts the admissible futures of the other. Communication is therefore identified with the operation of causal merging.

Time. *Time* is not a background parameter or a geometric coordinate. It is the ordinal structure induced by successive refinements. Proper time is the count of irreducible measurement events along a chain of refinement. Two events are time-ordered if and only if one refines the other. There is no assumption of simultaneity beyond the structure imposed by the partial order itself. Time is thereby reduced to the combinatorics of distinguishability.

Uncorrelance. Two events are said to be *uncorrelant* when neither refines the other and no admissible extension forces an ordering between them. Uncorrelance is not ignorance; it is structural indeterminacy. An uncorrelant pair is not unordered because of missing information, but because the axioms forbid the introduction of an artificial order absent from the record. Uncorrelance is the source of non-commutativity and, in dense limits, of curvature and gauge structure.

Informational Minimality. *Informational minimality* is the principle that among all admissible extensions of a finite measurement record, the valid one is that which introduces the least unobserved structure. It is the operational form of Ockham’s rule. Informational minimality is not aesthetic parsimony; it is a logical necessity imposed by the impossibility of recording absent distinctions. It is this concept that forces spline closure, extremal interpolants, and the uniqueness of admissible smooth shadows.

Informational Strain. *Informational strain* is the irreducible residue that appears when locally admissible refinements fail to commute under global extension. It is a purely combinatorial quantity: the defect of closure in the algebra of refinement. Informational strain does not presuppose geometry. Curvature is its smooth shadow, not its definition. Whenever two refinement paths lead to incompatible but equally admissible records, informational strain is produced. This object is the source of all later geometric and dynamical structure in the theory.

1.3 The Laws of Measurement

The Laws of Measurement are not assumed principles. They are theorems forced by the Axioms of Measurement when applied to the problem of maintaining a coherent experimental record under refinement.

Each axiom constrains what may be written down about the world. The laws describe what must follow once those constraints are taken seriously. No geometric structure, physical dynamics, or continuum hypotheses are inserted. The laws emerge solely from the requirement that finite records of distinguishable events can be extended without contradiction.

The derivation of each law follows the same logical pattern. A finite record of events is assumed. All admissible extensions of that record are examined under the axioms of finiteness, discreteness, minimality, and global coherence.

Among these extensions, only those that introduce no unobserved structure survive. The surviving extensions define extremal objects in the dense limit. These extremal objects are what classical physics later recognizes as smooth structure.

In this sense, the laws are not physical laws but bookkeeping necessities. They describe the only possible ways a ledger of measurements can remain internally consistent as it grows. Each law therefore expresses a structural fact about records, not a dynamical fact about the world.

What follows are the six fundamental laws that arise inevitably from the axioms of measurement.

Law 1: The Law of Spline Sufficiency. Any coherent record admits a unique extremal interpolant that introduces no unobserved structure. Smooth variational calculus is forced by this law.

Law 2: The Law of Discrete Spline Necessity. Because events are finite and refinement is discrete, every admissible interpolant must approximate a spline. The continuum is a spline limit.

Law 3: The Law of Boundary Consistency. Refinements must agree on overlaps. This law yields the adjoint equations, canonical structure, and the foundations of transport phenomena.

Law 4: The Law of Causal Transport. The informational interval is invariant under maximal propagation. The metric arises as the gauge preserving distinguishability.

Law 5: The Law of Curvature Balance. Failure of informational closure produces irreducible residue. Curvature is the smooth image of this discrete strain.

Law 6: The Law of Combinatorial Symmetry. Symmetries are not imposed group actions but statistical identities of refinement. Noether currents arise from informational invariance.

Law 7: The Law of Causal Order. Entropy as defined by the count of permissible measurements is necessarily monotonic in nature.

1.4 Logical Flow of the Theory

The emergence chain of the theory is:

$$\begin{aligned} \text{Events} &\Rightarrow \text{Order} \Rightarrow \text{Refinement} \Rightarrow \\ \text{Extremality} &\Rightarrow \text{Transport} \Rightarrow \text{Curvature} \Rightarrow \\ \text{Symmetry} &\Rightarrow \Delta S \geq 0. \end{aligned}$$

The inequality $\Delta S \geq 0$ is the final invariant: the count of admissible futures must grow as refinement proceeds. No thermodynamic or statistical assumptions are required; the Law of Causal Order is a purely combinatorial result.

1.5 The Experimental Record

Every phenomenon referenced in this manuscript has occurred in the experimental record. No effect is assumed, posited, or postulated as a primitive feature of the world. Each is a constraint on what a consistent ledger of observations must allow. The empirical record contains interference (Michelson–Morley), threshold behavior (Lenard, photoelectric), scattering residues (Compton), nonlocal phase shifts (Aharonov–Bohm), relativistic time dilation (Pound–Rebka), flat rotation curves (Rubin–Ford–Thonnard), decoherence, entropy increase, and scale relations such as Leavitt’s Ladder.

The task of a measurement-first theory is not to explain these effects by invoking additional structure, but to demonstrate that they are inevitable whenever recorded distinctions are required to remain globally consistent. Any admissible system of measurement must therefore reproduce the long list of observed behaviors already present in the experimental ledger. The laws traditionally called “physics” are understood here as the compatibility conditions that a finite, discrete, and refinable observation record must satisfy in order to admit a coherent completion.

1.5.1 On the Order of Presentation

The sequence in which the principles appear in this manuscript does not follow the historical order in which the corresponding physical effects were discovered or postulated. The history of physics is a record of insight accumulated over centuries, each phenomenon understood in the language and conceptual tools available at the time. Here the ordering is different because the foundation is different. The starting point is not a collection of physical laws but the structure of the observation record itself: finite, discrete, refinable, and required to remain globally consistent.

Once the record is taken as primary, the order in which its constraints are derived is determined by logic rather than chronology. Distinguishability precedes dynamics; refinement precedes continuity; compatibility precedes geometry; and the informational ledger must remain coherent long before the familiar formalisms of fields, forces, or symmetries are introduced. The laws that emerge are therefore not laws of matter or motion, but laws of information: rules describing how recorded distinctions must flow and how they must agree when extended across larger and more complex domains.

In this development, phenomena such as interference, quantization, time dilation, curvature balance, threshold effects, rotation curves, and entropy increase arise in an order dictated by informational necessity, not historical order. Each effect appears at the point where the consistency of the observa-

tional ledger requires it. The result is a reconstruction of the classical laws of physics as compatibility conditions on information flow, rather than as independent axioms about the material world.

1.5.2 Admissibility of Physical Theory

In this framework, a physical theory is not accepted because it describes an underlying reality, nor because it resembles the constructs of past models. A theory is admissible only to the extent that it extends the observation record without contradiction. It must refine the ledger in a way that preserves distinguishability, respects the partial order of events, and satisfies the global compatibility conditions that govern how information may flow.

This criterion differs fundamentally from the usual scientific practice of postulating entities or mechanisms and then testing their consequences. Here, the primary object is the experimental record itself, and a theory is evaluated by whether it can be written as a consistent refinement of that record. No admissible theory may introduce structure that cannot be supported by finite distinction, nor may it rely on hidden degrees of freedom that violate refinement compatibility or exceed the informational bandwidth of the causal network.

The admissibility criterion thus reduces the vast landscape of conceivable theories to those that are informationally coherent. Classical dynamics, quantum interference, relativistic calibration, curvature balance, spectral thresholds, and cosmological scaling laws all survive because each is a necessary compatibility condition on the extension of the record. In this sense, the theories traditionally called “physical laws” are reinterpreted as the minimal and admissible rules for updating a finite, discrete, and globally consistent observational ledger.

Coda: Aristotle and Galileo

Aristotle's account of motion is historically influential but informationally inadmissible. It asserts how bodies *ought* to move, not how they *do* move in the experimental record. His conclusions do not arise from refinement of a finite ledger of observations, nor are they tested against distinguishability, comparison, or global compatibility. They are rules posited about an unseen essence. Such rules may be elegant, but they do not extend the observation record in a consistent way; they bypass it entirely.

Galileo's method is the opposite. It begins with the empirical record and asks only what regularities its refinement demands. Rolling spheres on inclines, comparing fall times, and separating signal from noise are not illustrations of a theory but demonstrations of admissibility. The informational ledger is updated by experiment, and any proposed principle must preserve the coherence of that ledger when carried into new domains. Galileo's observations satisfy this requirement; Aristotle's do not.

The distinction is therefore not philosophical but structural. A thought experiment that does not correspond to an actual refinement of the observation record is informationally empty. An experiment that produces a finite, repeatable, distinguishable event contributes to the ledger and must be accommodated by any admissible theory. In this manuscript, the laws of physics are reinterpreted as precisely those rules that survive this test of empiracy: they are the constraints that every coherent, extensible observation record must satisfy. Aristotle speculated; Galileo measured. In an informational framework, only the latter can be admitted.

Phenomenon 1.5.1 (Empiracy [4, 57]). *In the measurement-first framework, the distinction between Aristotle and Galileo can be stated as a precise admissibility test. Let \mathbf{E} denote the set of recorded events, each $e \in \mathbf{E}$ represented by a finite tuple of distinguishable marks (for example, height and*

time). A dynamical theory for free fall is an assignment of a constraint

$$L(h, t) = 0$$

on admissible pairs (h, t) , interpreted as “height h at elapsed time t ”. The theory is admissible for an observation record $\mathbf{E} \subset \mathbf{R}^2$ if there exists an extension $\mathbf{E}' \supseteq \mathbf{E}$ such that every event in \mathbf{E}' satisfies the constraint:

$$(h, t) \in \mathbf{E}' \implies L(h, t) = 0.$$

In words: the theory must admit a refinement of the experimental ledger in which all events are consistent with a single law.

Aristotle’s account of motion can be rendered as a family of constraints $L_A(h, t; m)$ that enforce the rule that heavier bodies fall faster. In a simple form, this may be modeled as

$$h(t; m) = \alpha(m) t^2, \quad \alpha(m_1) \neq \alpha(m_2) \text{ for } m_1 \neq m_2,$$

so that for fixed h one has

$$t(h; m_1) \neq t(h; m_2) \quad \text{whenever} \quad m_1 \neq m_2.$$

Now consider an observation record \mathbf{E} containing two events at the same height h_* but with experimentally indistinguishable fall times:

$$e_1 = (h_*, t_*, m_1), \quad e_2 = (h_*, t_* + \delta, m_2), \quad |\delta| < \delta_{\text{res}},$$

where δ_{res} is the time resolution of the apparatus. Galileo’s inclined-plane experiments produce exactly such records: within experimental resolution, bodies of different mass reach the same height in the same time. For any δ_{res} smaller than the separation implied by L_A , there is no extension $\mathbf{E}' \supseteq \mathbf{E}$ in which all events satisfy $L_A(h, t; m) = 0$. Aristotle’s rule is therefore informationally

inadmissible: it defines a constraint surface that does not admit a refinement of the actual observation record.

Galileo's law, by contrast, can be expressed as a mass-independent constraint

$$L_G(h, t) = h - \frac{1}{2}gt^2 = 0,$$

or more generally as an equivalence class of laws that agree within the resolution of the apparatus. Given an experimental record \mathbf{E} of fall times and heights, there always exists a parameter g and an extension $\mathbf{E}' \supseteq \mathbf{E}$ such that

$$(h, t) \in \mathbf{E}' \implies L_G(h, t) = 0 \quad \text{up to the experimental resolution.}$$

Galileo's theory is therefore admissible: it defines a surface in (h, t) -space that can be populated by a consistent refinement of the ledger, with no contradiction between recorded events and the proposed law.

In this sense, the difference between Aristotle and Galileo is not merely historical or philosophical but mathematical. Aristotle's dynamics do not admit an embedding of the experimental record into a single, globally consistent constraint surface; Galileo's do. Within a measurement-first framework, this is the precise meaning of admissibility. A theory that cannot be realized as a consistent extension of the observation record is ruled out not by taste or interpretation, but by the impossibility of defining any refinement map

$$\iota : \mathbf{E} \hookrightarrow \{(h, t) : L(h, t) = 0\}$$

that preserves distinguishability and experimental resolution. Aristotle specifies such an L and fails the test. Galileo, by measuring, finds one that passes.

The Purpose of Thought Experiments

Thought experiments appear throughout this work, but their purpose is not to assert new structure or to introduce hypothetical mechanisms. In a measurement-first framework, a thought experiment is merely an analytic scaffold: a controlled setting in which the mathematical consequences of the axioms can be examined without the distractions of incomplete data or instrumental complexity. These constructions motivate the derivations, clarify the role of refinement, and expose the logical boundaries of the informational ledger. They do not extend the observation record; they prepare the mathematics that later must be tested against it.

The decisive content of the theory comes from the experimental record itself. Any admissible system of measurement must reproduce the long list of effects already observed in nature. These include interference (Michelson–Morley), diffraction and wavepacket breakup (Davisson–Germer), threshold emission (Lenard), scattering residues (Compton), nonlocal phase (Aharonov–Bohm), time dilation (Pound–Rebka), universal spectral calibration (Leavitt’s Ladder), curvature balance, decoherence, and the statistical monotonicity expressed by $\Delta S \geq 0$. None of these are assumptions; each belongs to the observational ledger, and a theory that fails to admit them is excluded by definition.

This manuscript also identifies several phenomena that have not yet been observed, but which are required by the informational axioms and the compatibility conditions derived in later chapters. These predictions follow not from speculation, but from the structure that any coherent refinement must satisfy:

- The existence of a curvature–strain residue in noninertial interferometers, visible as a systematic phase offset independent of dynamical forces.
- Specific bandwidth limits on spectral transitions, arising from the in-

formational Nyquist bounds inherent to distinguishability.

- A universal refinement–curvature correction in rotation curves, independent of composition, luminosity, or mass model.
- Quantized strain channels in strongly curved causal networks, measurable as discrete deviations from the smooth–shadow Euler–Lagrange approximation.
- A monotone informational arrow that persists even in regimes where thermodynamic mechanisms fail, providing a direct experimental signature of ΔS as a bookkeeping constraint rather than a physical process.

These predictions are not optional embellishments. They represent the phenomena that must appear in any future experiment if the axioms of measurement, refinement, and global consistency are correct. Just as the classical effects above serve as validation of the informational framework, these forthcoming phenomena serve as its unavoidable consequences. The theory begins with the observation record, but it does not end there: it extends the ledger into domains not yet measured, identifying the signatures that future observation must reveal.

Chapter 2

The Algebra of Events

The first chapter established the informational mechanism of physics: each measurement produces a finite, distinguishable event, and the absence of additional events is itself a binding constraint. Measurement is not a sampling of an underlying continuum; it is the creation of distinguishability. From a mathematical standpoint, measurement itself is therefore a growing record of discrete selections, ordered by refinement. Each measured event appends an inassailable fact to the experimental record.

This chapter formalizes that structure. We build the algebra of measurement itself. Rather than treating observations as values of continuous functions, we adopt the combinatorial viewpoint forced by the Axioms 5 and 6: every admissible experimental record is finite or countable, and every refinement is a restriction of admissible outcomes. The central object of this chapter—the *Causal Universe Tensor*—expresses the fact that history is built multiplicatively: each new event contracts the set of admissible continuations of the record. The universe does not evolve additively in time; it accumulates consistency through left products of restricted increments.

2.1 Time and Events

Before introducing any notion of temporal ordering, it is necessary to separate the concept of an *event* from the concept of *time*. In this framework, neither is assumed a priori. Both arise as consequences of how distinguishable records can be extended without contradiction.

An *event* is defined as a minimal, distinguishable unit of record (Definition 8). It is not something that happens in time. It is the act of recording a distinction. An event exists precisely when a measurement refines the experimental ledger. If nothing is distinguished, no event exists.

Thus, the set of events E is the set of all finite acts of distinction. This set carries structure not from motion or duration but from refinement.

Time, therefore, does not exist as a background coordinate but as a way to index events relative to each other. It is an ordinal indexing of refinement. Given two events $e_i, e_j \in E$, we say that $e_i \prec e_j$ if and only if the record of e_j is a refinement of the record of e_i .

Time is therefore not measured. It is counted. It is the order type of the chain of events:

$$e_1 \prec e_2 \prec e_3 \prec \cdots$$

Proper time is the cardinality of this chain. There is no continuum of time between irreducible refinements. There is only succession.

Phenomenon 2.1.1 (The Event Effect [83]). **N.B.**—*Kant identified time not as an object, but as a necessary ordering condition for experience. The present construction reverses this dependency: the ordering of irreducible events generates time rather than presupposing it [83].* \square

An event is the minimal irreducible act of distinguishability. It arises when and only when a measurement refines the admissible history. It is not located in time; it generates time by its order.

It is necessary to clarify what it means for a measurement to exist at all. In this framework, measurement is not a passive act and not an inquiry

about a pre-existing quantity. It is the creation of a distinction that did not previously appear in the record. A measurement is an operation that enlarges the causal ledger.

A record that does not grow is not a record that was measured. Silence cannot be distinguished from absence, and absence cannot participate in causal structure. For this reason, the null act cannot be admitted as a measurement.

This has an important structural consequence. Measurement is not reversible. Later observations may refine, reinterpret, or contextualize earlier ones, but they cannot erase the fact that a distinction was recorded. The ledger may be extended, but it cannot be undone. This irreversibility is not a postulate of physics; it is a logical consequence of what it means to record anything at all.

Phenomenon 2.1.2 (The Positive Measurement Effect [124]). ***N.B.**—Zeno’s paradoxes arise precisely from allowing vanishingly small increments to masquerade as genuine acts. In the present framework, such zero-measurement steps are inadmissible. A measurement must refine, or it is not a measurement at all.* \square

All admissible measurements produce a strictly positive refinement. A zero measurement is not a measurement but a null act and is therefore excluded. Every event increases the informational record by a positive integer amount. Observations cannot be replaced or canceled by future observations.

Although no intermediate event exists between successive refinements, the dense limit of refinement forces the appearance of a smooth interval as an approximation. This interval is not fundamental. It is the shadow of coarse observation.

Let $\tau : E \rightarrow \mathbb{N}$ be the ordinal timing map induced by refinement order. For two successive events $e_i \prec e_{i+1}$, the open set

$$(\tau(e_i), \tau(e_{i+1}))$$

contains no elements of E .

However, in the smooth completion forced later by the Axiom of Cantor, this empty discrete interval is represented as a continuous open interval. This representation gives rise to the notion of a *present moment*.

Phenomenon 2.1.3 (The Moment Effect). *The Here and Now is the smooth shadow of the empty open interval between successive events. It is not a primitive moment in time, but the continuous representation of the absence of distinguishable events between refinements. Concretely, it appears as the open set*

$$(i, i + 1) \subset \mathbb{R},$$

which contains no admissible events.

2.1.1 The Causal Universe Tensor

Having established that time is not a background parameter but an ordinal structure induced by refinement, we now consider the object that records the global consequences of this ordering. The axioms do not allow a universe to be postulated as a stage on which events occur. Instead, what is called the “universe” must be constructed as the unique coherent extension of the finite record itself. The following tensor object is not dynamical and not geometric. It is a bookkeeping device forced by the requirement that local distinctions admit a single, globally consistent smooth shadow.

In the discrete domain E , time is ordinal: an index into the growing chain of distinguishable selections. In the continuous shadow U , time does not flow at all. The continuous tensor \mathbf{U}_k at step k is not the result of propagation, but the image of a restriction map:

$$\mathbf{U}_{k+1} = \Psi(e_{k+1} \cap \hat{R}(e_k)) \mathbf{U}_k, \quad (2.1)$$

where \hat{R} is the discrete restriction induced by the most recent event e_k ,

and Ψ is its continuous representation. Thus the continuous universe is not postulated as a field living on a manifold: it is the coherent bookkeeping of discrete consistency.

The ordered product makes this explicit. If $\hat{R}(e_j)$ denotes the admissible refinement of the j -th event, then the causal universe as seen by a single inertial observer is

$$\mathbf{U}_k = \prod_{j=1}^{k-1} \Psi(e_{j+1} \cap \hat{R}(e_j)). \quad (2.2)$$

No geometry, metric, or dynamical law is assumed. Continuity, smoothness, and variational structure will reappear later only as the unique continuous shadow of this discrete product, forced by informational minimality and global coherence.

In this way, Chapter 2 performs the key transition: from sets of distinguishable events to an algebra of restricted, multiplicative updates. The remainder of the chapter introduces the axioms, operators, and tensor structures that make this viewpoint precise, culminating in the formal definition of the causal universe tensor.

Thought Experiment 2.1.1 (Statistical Process Control [140]). ***N.B.**—Observational records have been used to understand and control complex processes to remarkable success. Statistical process control demonstrates that measurement does not estimate a continuous parameter directly; it eliminates process states that are incompatible with the record. The state of the system is therefore not an average, but the set of configurations that have survived all admissible checks.* \square

For a formal treatment, see Phenomenon 2.5.2 later in this chapter.

Imagine a factory that manufactures a precision component. The process is controlled by a set of adjustable parameters: temperature, pressure, feed rate, alignment, and so on. At startup, all parameter settings that satisfy the design tolerances are admissible; the process could be in any one of many configurations. A single measurement does not determine the underlying state.

It merely rules out those configurations that would have produced a conflicting outcome.

This is the essential structure of statistical process control. Each inspection, probe, gauge reading, or quality check eliminates a subset of incompatible configurations. After k measurements, the surviving parameter settings are precisely those that are consistent with all k observations.

Let e_k denote the k -th inspection result, and let $\hat{R}(e_k)$ be the discrete restriction that removes every process state incompatible with e_k . If Ψ embeds these restrictions into the continuous tensor domain, the recorded state of the process after k inspections is

$$\mathbf{U}_k = \prod_{j=1}^{k-1} \Psi(e_{j+1} \cap \hat{R}(e_j)). \quad (2.3)$$

The process does not “evolve” in time in the usual dynamical sense; it accumulates admissibility. Each new inspection refines the record by discarding alternatives, giving the stepwise update

$$\mathbf{U}_{k+1} = \Psi(e_{k+1} \cap \hat{R}(e_k)) \mathbf{U}_k. \quad (2.4)$$

Two independent inspectors, perhaps located at different stations on the production line, can refine their records without communication. If their measurements are mutually consistent, the products merge without conflict. If not, no admissible configuration survives the combined restrictions, and the process is flagged as out of control. In this sense the system is independent, yet globally constrained by the requirement of the coherent environment.

This ordinary industrial setting exhibits the same structure developed in this chapter. Measurement eliminates incompatible alternatives, time indexes the number of admissible refinements, and the continuous representation \mathbf{U}_k is nothing more than the shadow of a discrete product of selections.

We begin by enumerating the seven Axioms of Measurement, which for-

malize the structure of admissible records and the refinement of observational history.

2.2 The Axioms of Mathematics

All mathematics in this work is carried out within the framework of Zermelo–Fraenkel set theory with the Axiom of Choice (ZFC) [81, 91]. Rather than enumerating the axioms in full, we recall only those consequences relevant to the construction that follows:

- **Extensionality** ensures that distinguishability has formal meaning: two sets differ if and only if their elements differ.
- **Replacement** and **Separation** guarantee that recursively generated collections such as the causal chain of events remain sets.
- **Choice** permits well-ordering, allowing every countable causal domain to admit an ordinal index.
- **The Successor Function** of the Peano axioms provides the mechanism by which distinguishable outcomes may be counted.

These are precisely the ingredients required to formalize a locally finite causal order. All further constructions—relations, tensors, and operators—are definable within standard ZFC mathematics; see Kunen [91] and Jech [81] for set-theoretic foundations, and Halmos [65, 66] for the induced tensor and operator structures on finite-dimensional vector spaces.

2.2.1 Measurements are Mathematical in Nature

The starting point of this framework is methodological rather than ontological. We do not assume anything about the substance of physical reality. We

assume only that the outcomes of measurement are finite or countable collections of distinguishable results recorded in time. This is standard across probability theory and information theory: Shannon formalized information as distinguishable symbols drawn from a finite or countable alphabet [138], and Kolmogorov showed that empirical outcomes can be represented as elements of measurable sets within standard set theory [88]. In this view, measurement produces data, and data are mathematical objects. Everything that follows concerns the admissible transformations among such records.

Definition 1 (Admissible Structure). *Let E denote the set of admissible objects.*

Every element of E is a set. Two types of elements are distinguished internally:

- *An event is a set e such that $e \in E$.*
- *A worldline is a set S such that*

$$S \in E \quad \text{and} \quad \forall e (e \in S \Rightarrow e \in E).$$

Thus the admissibility relations are stratified as

$$e \in E, \quad S \in E, \quad e \in S.$$

Axiom 1 (The Axiom of Kolmogorov [88]). [Measurement as a Formal Record.] *Formally, there exists a set E such that E is an admissible record and*

$$|\{e \mid e \in E\}| > 3. \tag{2.5}$$

N.B.—*The choice of 3 is not arbitrary, but serves to eliminate degenerate configurations and simplify the foundational proofs. There are posedness concerns for universes with less than 4 events and the analysis of the degenerate cases is omitted.* □

The record of measurement—defined as the finite or countable set of observed, distinguishable events and worldlines—is taken to be a mathematical object representable within ZFC. No ontological claim is made about physical reality. The axiom asserts only that observable data can be formalized as sets and relations.

This standpoint is consistent with Kolmogorov’s construction of probability spaces, in which empirical outcomes are represented as measurable sets [87]. Accordingly, a record of finite observations is a mathematical object whose structure is defined entirely within ZFC. Throughout this work, the word “information” refers exclusively to these representable distinctions; nothing is asserted about any underlying physical substrate that might produce them.

2.2.2 Mathematics is the Language of Measurement

Mathematics enters this framework not as an external interpretive layer but as the minimal language in which measurement can be expressed. A record of observation is a finite collection of distinguishable outcomes, and the relations among those outcomes—order, refinement, exclusion, and compatibility—require a precise symbolic setting. The purpose of this subsection is therefore methodological: to state explicitly the mathematical rules under which every subsequent construction is carried out.

No structure beyond ordinary set theory is needed. The axioms of ZFC provide the machinery for forming sets of events, for defining relations among them, and for building the tensor algebra in which their continuous shadows will later appear. Within this system, counting becomes the first and most fundamental operation: to measure is to distinguish, and to distinguish is to enumerate the admissible outcomes. Peano’s contribution is thus not philosophical but operational. The natural numbers supply the ordinal scaffold upon which every causal record is indexed.

With this in mind, we begin by stating the formal principle that makes

counting available as a tool of measurement.

Axiom 2 (The Axiom of Peano [54, 105, 165]). [Counting as the Tool of Information] *All reasoning in this work is confined to the framework of ZFC. Every object—sets, relations, functions, and tensors—is constructible within that system, and every statement is interpretable as a theorem or definition of ZFC. No additional logical principles are assumed beyond those required for standard analysis and algebra.*

Formally,

$$\text{Measurement} \subseteq \text{Mathematics} \subseteq \text{ZFC subseteq Counting.}$$

Thus, the language of mathematics is taken to be the entire ontology of the theory: the physical statements that follow are expressions of relationships among countable sets of distinguishable events, each derivable within ordinary mathematical logic.

Thought Experiment 2.2.1 (The Speedometer [158, 163]). **N.B.**—*The mechanical implementation of measuring devices often are protected by explicit descriptions of how they work. The patents cited here explicitly describe how they turn counting into data.* \square

Consider an ordinary automobile speedometer. The dial appears to report a continuous real number at each instant, but the device does not have access to the real numbers. A mechanical speedometer counts wheel rotations through a gear train and maps those counts to pointer positions. A digital speedometer counts the same rotations and displays a numeral drawn from a finite alphabet.

Each time the counter increments and the displayed symbol or pointer position changes, a new distinguishable event is recorded. Between two successive display states there is no way, from the informational record alone, to assert that any additional state occurred. The apparent continuity of “speed” is a visual interpolation of a finite counting process.

Thus the speedometer does not output a real number. It outputs a countable sequence of distinguishable states derived from integral counts of wheel rotations. The act of measuring speed reduces to counting transitions of a finite-state device. All physical inference based on such data can be expressed within ordinary arithmetic and set theory.

This illustrates Axiom 2: measurement generates only countable, finitely coded distinctions, and every mathematical object used to interpret those distinctions—numbers, functions, tensors—is a construct of ZFC. No structure beyond counting is assumed at the fundamental informational level.

2.3 The Axioms of Informational Structure

The previous section established that a physical record is a set of distinguishable observations, representable within ZFC, and partially ordered by causal precedence. Nothing further was assumed about geometry, dynamics, or the continuum. In this section, we introduce two informational axioms that restrict how such a record may be interpreted. These axioms express constraints on admissible descriptions of the world, independent of any particular model of physics.

Axiom 3 formalizes the principle that a physical history may not contain unobserved structure. Among all symbolic descriptions that reproduce the recorded events, the admissible one is the shortest. This is the information-theoretic form of Ockham's principle: no plurality of assumptions without necessity.

Axiom 4 asserts that the record of events is not merely ordered but forms a locally finite causal set. Local finiteness ensures that causal cardinality is discrete, while the partial order encodes temporal precedence. Continuum spacetime, when it exists, is therefore understood as an approximation that faithfully embeds this discrete informational structure.

Together, these axioms define the informational content of the physical

world: a causal set with no unrecorded structure and no additional assumptions beyond the observational record itself.

2.3.1 Information Minimality

The observational record E is defined only by the distinguishable events it contains. Between two recorded events e_i and e_{i+1} , no additional structure is present in the data: no new marks in the notebook, no threshold crossings, and no observable distinctions. Set theory alone does not forbid a hypothetical refinement that inserts additional structure between e_i and e_{i+1} , but any such refinement asserts observations that did not occur. To prevent unrecorded structure from being introduced by assumption, we impose an informational constraint.

Among all symbolic descriptions that reproduce the recorded events, the admissible one is the shortest. In modern information theory, this statement is formalized by Kolmogorov complexity [87, 100]: a description is preferred if it introduces no additional information beyond the events in E . This embodies the classical principle that no plurality of assumptions should be posited without necessity. It is not derived from the set-theoretic framework; it is an axiom about how physical theories must interpret finite empirical records.

Axiom 3 (The Axiom of Ockham [116])). [Order Coherence] *Let $E = \{e_0 \prec e_1 \prec \dots \prec e_n\}$ be a finite or countable partially ordered set of recorded events. The admissible histories are order-respecting in the following sense: for any two events $e_1, e_2 \in E$,*

$$e_1 \prec e_2 \implies \forall S \in E (e_1 \in S \implies e_2 \in S).$$

That is, no admissible history may contain an earlier event without also containing all later events forced by the causal order.

We have seen this principle in action already. Refer to Thought Experiment 1.1.3 and the use of Simpson's rule to compute the path of a spaceship with minimal measurement information.

Further, we will later show, using Law 2, that the Axiom of Ockham imposes an informational minimality constraint on the evolution of the Causal Universe Tensor, and that this constraint mathematically characterizes the sharpness of the razor.

2.3.2 Causal Set Theory

The previous axiom imposed an informational constraint on admissible descriptions of the record of measurement. We now introduce a structural constraint. The empirical record is a set of distinguishable events with a causal precedence relation \prec , but this alone does not restrict the size of causal intervals. In a general partially ordered set, the number of events between a and b may be infinite. Physical measurements, however, produce finite data. To represent this empirically grounded discreteness, we assume that the causal order is locally finite: every causal interval contains only finitely many recorded events.

This postulate places the present construction within the causal set program of Sorkin and collaborators, where spacetime is modeled as a locally finite partial order and continuum geometry, when it appears, is a derived approximation. Order encodes temporal precedence, and local finiteness encodes discrete causal volume. No metric, field, or manifold structure is assumed at the fundamental level; these arise only if the causal set admits a faithful embedding into a Lorentzian manifold.

Axiom 4 (The Axiom of Causal Sets [16]). [Events are Discrete]

The distinguishability relations among recorded events admit a representation as a locally finite partially ordered set (E, \prec) , where

1. $e \prec f$ means that the record of e is incorporated before the record of f ,

2. (E, \prec) is acyclic and transitive,
3. and for any two events $a \prec b$, the interval $\{e \in E : a \prec e \prec b\}$ is finite.

Local finiteness ensures that the recorded causal cardinality is discrete, and the order relation encodes temporal precedence within the record. Any Lorentzian manifold, when it exists, is merely a physical model in which this discrete causal structure may be faithfully approximated.

Thought Experiment 2.3.1 (The Laboratory Procedure [116, 161]). **N.B.**—

The following example collects ideas from several well-established perspectives in measurement theory. Bohr and Wheeler emphasize that a physical experiment records only distinguishable outcomes; no other structure is operationally meaningful [12, 161]. In information theory, such records are represented as finite or countable strings of distinguishable symbols [30, 138]. In ergodic theory and causal set theory, successive measurements refine a partition of the observational domain into finer distinguishable elements [117, 130, 148]. Finally, computational mechanics and operator-theoretic dynamics treat the “evolution” of a system as the repeated update of its information state [11, 31, 90]. Taken together, these perspectives justify modeling a laboratory procedure as a refinement operator acting on a finite measurement record. The experiment does not solve differential equations; it follows the laboratory procedure Ψ . \square

Consider a laboratory notebook in which each threshold crossing of a detector is recorded as a mark in ink. The notebook contains a finite sequence of distinguishable entries

$$e_0 \prec e_1 \prec \cdots \prec e_n,$$

each representing an irreversible update of the experimental record. The notebook is not a model of reality; it is the empirical record. No claim is made about any mechanism behind it.

Now suppose one attempts to describe what “really” happened between two successive entries e_i and e_{i+1} . If additional curvature, oscillation, turning points, or discontinuities had occurred, then the detector would have crossed a threshold and a new entry would appear. Because no such entry is present, the observational record forbids any refinement that predicts one.

Thus the notebook determines a finite set $E = \{e_0, \dots, e_n\}$ of recorded events. Every admissible history must be a completion that introduces no new distinguishable events beyond E . Any hypothetical refinement with additional structure is rejected as inadmissible, since it asserts observations that did not occur.

2.4 The Axioms of Observation

A common criticism of mathematical physics is the extent to which mathematics can be tuned to fit observation [15, 123] and, conversely, manipulated to yield nonphysical results [77]. Lord Berkeley’s critique of Newton’s fluxions [10] could only be answered by centuries of successful prediction with only intuition as justification. Today, calculus feels like a natural extension of the real world—so much so that Hilbert, in posing his famous list of open problems, explicitly formalized the lack of a rigorous foundation for physics as his Sixth Problem [?, 159].

We aim to show that the mathematical language used to describe observation gives rise to a system expressible entirely as a discrete set of events ordered in time. Moreover, this ordered set possesses a mathematical structure that naturally yields the appearance of continuous physical laws and the conservation of quantities. To understand how this works, we first clarify what we mean by measurement.

2.4.1 The Countable Nature of Events

Thought Experiment 2.4.1 (Momentum [113]). **N.B.**—For a rigorous treatment of momentum see *Phenomena* [?] and [?] □

Physical laws relate measurements. For example, Newton’s second law [113]

$$F = \frac{dp}{dt} \tag{2.6}$$

states that force relates to the change in momentum over time. To speak of change you must have at least two momentum values, one that comes before the other; otherwise there is nothing to distinguish. In set-theoretic terms, by the Axiom of Extensionality (assumed in Axiom 2), different states must differ in their contents, so “change” presupposes the distinguishability of two states.

In this framing, measurement values are *counts* (cardinalities) of elementary occurrences: the number of hyperfine transitions during a gate, the tick marks traversed on a meter stick, the revolutions of a wheel. The *event* is the action that makes previously indistinguishable outcomes distinguishable; the *measurement* is the observed differentiation (the count) between two anchor events. This is not the absolute measure of the event, but just relative difference of the two. We count the events as time passes (See Thought Experiment 2.2.1).

Since special relativity requires that time vary under the Lorentz transform [45, 101], there can be no global scalar representation of temporal duration. Rather, special relativity permits us only to *list* all events in the universe in their proper causal order. It is this ordered list that we elevate to the first physical principle:

Axiom 5 (The Axiom of Cantor [21, 43]). [Events are Ordered and Countable] *The only invariant agreement in time guaranteed between two observers is the order in which correlant events (see Definition 10) occur. Duration is*

not a geometric interval; it is the count of distinguishable events that can be recorded between two selections:

$$|\delta t| = |\text{events distinguished between}|. \quad (2.7)$$

Thought Experiment 2.4.2 (Relativistic Simultaneity [45].). *Two laboratories, A and B, perform independent procedures, each producing a finite measurement record. Because the experiments are independent, their events commute: no record in A constrains the order of any record in B. Both notebooks are internally consistent, but their events are mutually unordered.*

Now two observers, C and D, travel past the laboratories on different trajectories, each at a velocity close to the speed of light. Their instruments register signals from A and B in different sequences. Since the events commute, both observers are free to assemble the two notebooks into different global orders. Observer C concludes that certain events in A precede those in B, while observer D concludes the opposite. Each construction is internally consistent, because commutativity permits the reordering.

The discrepancy is not a contradiction, but the finite analogue of relativistic simultaneity: different trajectories generate different admissible orderings of commuting events. The events themselves may be reordered independently of each other, yet the invariants are preserved.

2.4.2 Observations are Fixed and Combinatorial

A finite observer records events one at a time. Each record refines the set of admissible histories, and every refinement depends only on the records accumulated so far. Physical description is therefore necessarily recursive: the $(k + 1)$ st step is constructed from the k steps that precede it.

The recursive description of physical reality is meaningful only within the finite causal domain of an observer. Each step in such a description corresponds to a distinct measurement or recorded event. Observation is therefore

bounded not by the universe itself, but by the observer's own proper time and capacity to distinguish events within it.

Axiom 6 (The Axiom of Planck [122]). [*Observations are Finite and Im-mutable*] *For any observer, the set of observable events within their causal domain is finite. The chain of measurable distinctions terminates at the limit of the observer's proper time or causal reach. These observations do not change over time.*

This axiom establishes the physical limit of any causal description: the sequence of measurable events available to an observer always ends in a finite record. Beyond this frontier—beyond the end of the observer's time—no additional distinctions can be drawn. The *last event* of an observer thus coincides with the top of their causal set: the boundary of all that can be measured or known.

2.4.3 Measurements Must Extend Without Contradiction

The preceding axioms restrict the informational content of the record and the structure of causal precedence. We now introduce an axiom governing how events may be selected in a consistent physical history. A partial history is a finite sequence of recorded distinctions that respects the causal order. In a locally finite causal set, many partial histories may be extended, but not all extensions are admissible: each new event must be consistent with the existing record and may not contradict any previously recorded distinction.

Axiom 7 asserts that whenever we impose countably many local admissible requirements—each representing a physically permitted constraint—there exists at least one consistent history that satisfies all of them¹. Mathematically, this parallels the role of Martin's Axiom in set theory, where

¹In the continuum limit, when observables range over a complete set of measurable values, the admissible history is unique up to sets of measure zero: there is exactly one continuous completion consistent with all recorded refinements.

dense sets encode constraints and a filter selects a coherent global object [81, 91, 105, 154]. Physically, it echoes Boltzmann's principle that every admissible microstate selection must preserve distinguishability [15], and follows the causal-set program in which a spacetime history is constructed one event at a time under admissible refinement [16, 53]. Hilbert's call to axiomatize the foundations of physics [74] is realized here as a minimal requirement: if each local constraint is permissible, then some coherent global history must also be permissible.

Axiom 7 (The Axiom of Boltzmann [14, 105]). [Events are Selected to be Coherent.] *An experiment may impose many local causal requirements: detector constraints, boundary conditions, conservation rules, and so on. As long as each requirement can be satisfied on its own, the Axiom of Boltzmann asserts that there always exists at least one, globally coherent history satisfying all of them simultaneously. No matter how many local constraints we specify, they can be assembled into one consistent record.*

Formally, let (P, \leq) be the partially ordered set (Definition 3) of finite, order-consistent partial histories in a locally finite causal domain, ordered by extension. For every countable family $\{D_n\}_{n \in \mathbb{N}}$ of dense subsets of P (local causal constraints), there exists a filter $G \subseteq P$ such that $G \cap D_n \neq \emptyset$ for all n .

2.5 The Causal Universe Tensor

The axioms above determine the structure of the physical record: events form a locally finite causal set, extensions of partial histories preserve causal consistency, and informational minimality forbids unrecorded structure. What remains is to represent this record in a mathematical form that allows the accumulation of distinctions. We now construct such a representation.

2.5.1 Sets of Events

Let the set of all events accessible to an observer be denoted E^2 , ordered by causal precedence (\prec). Because any physically realizable region is finite, this order forms a locally finite partially ordered set (poset) [52].

Definition 2 (Causal Precedence [16]). *Let E be the set of distinguishable events accessible to an observer. For $e_i, e_j \in E$, we say that e_i causally precedes e_j , written $e_i \prec e_j$, if the record of e_j cannot be formed without already having distinguished e_i . Equivalently, e_j refines the admissible outcomes of e_i . The relation \prec is a strict partial order: it is irreflexive ($e \not\prec e$), antisymmetric, and transitive.*

N.B.—The term “causal” is used only in the sense of ordering: $e_i \prec e_j$ asserts that e_j depends on the distinctions recorded in e_i . No geometric notion of signal propagation or physical influence is assumed. \square

Each admissible set of events may be represented as a locally finite partially ordered structure [16, 146], whose links record only those relations that are causally admissible. In this view, a “history” is not a continuous trajectory but a combinatorial diagram: every vertex an event, every edge a permissible propagation. This discrete formulation generalizes the intuition behind Feynman’s space–time approach to quantum mechanics, in which the amplitude of a process is obtained by summing over all consistent histories [50, 51]. The Feynman diagram thus appears here as a special case of the causal network itself—a pictorial reduction of the full tensor of event relations—and the path integral becomes a statement of global consistency across all measurable causal connections.

Thought Experiment 2.5.1 (Feynman Diagrams (classical) [51]). **N.B.**—*This is a classical simplification of the highly specialized notation of the Feyn-*

²The symbol E here denotes the *set of distinguishable events*—it is not the energy operator or expectation value familiar from mechanics. Throughout this work, E indexes discrete occurrences in the causal order, while quantities such as energy, momentum, or stress appear only later as *derived measures* on this set.

man diagram. See Thought Experiment 7.4.4 for a more rigorous treatment.

□

In conventional quantum field theory, a Feynman diagram depicts a sum over interaction histories connecting initial and final particle states. Each vertex represents an elementary event—an interaction that renders previously indistinguishable outcomes distinct—and each propagator represents the possibility of causal influence between events.

In the present formulation, such a diagram is naturally interpreted as a finite causal network. The set of vertices corresponds to the event set E , and the directed edges encode the order relation \prec defined by Axiom 5. To each event e_k we associate a representation \mathbf{E}_k that records the admissible refinement induced by that event, and the directed structure describes which refinements must precede others. The composition of these event tensors gives the Causal Universe Tensor of the inertial frame:

$$\mathbf{U}_n = \prod_{k=1}^n \mathbf{E}_k. \quad (2.8)$$

At this stage, \mathbf{U}_n is a classical accumulator: it records the count and structure of distinguishable events without assigning amplitudes or phases. This is deliberate. The present framework concerns only the logical bookkeeping of distinctions. The full quantum structure—including complex amplitudes, superposition, and interference—appears only after the informational gauge is introduced. In that setting, the classical accumulator becomes the coarse projection of a richer amplitude algebra, much as a Feynman diagram may be viewed as the combinatorial skeleton of a path integral. That generalization is deferred until Chapter 7, where the amplitude-bearing form of \mathbf{U} is constructed.

Summing over all consistent diagrams is therefore equivalent to enumerating all admissible orderings of distinguishable events. The path integral itself becomes a statement of global consistency across the entire causal net-

work: every measurable amplitude corresponds to a possible embedding of finite causal order into the continuous limit. In this sense, a Feynman diagram is not merely a pictorial tool, but a discrete representation of the causal tensor algebra from which continuum physics emerges.

This identification is pedagogically useful. From this point onward, every construction may be viewed as an algebraic generalization of the familiar Feynman diagram: the event tensors are its vertices, the causal relations its edges, and the Universe Tensor the cumulative sum over all consistent orderings. The remainder of the monograph simply formalizes this graphical intuition in set-theoretic and tensorial language, rather than using calculus.

At the heart of measurement is the concept of a partially ordered set.

Definition 3 (Partially Ordered Set [33]). *A poset is a pair (E, \leq) where \leq is a binary relation on E satisfying:*

1. **Reflexivity:** $e \leq e$ for all $e \in E$
2. **Antisymmetry:** if $e \leq f$ and $f \leq e$, then $e = f$
3. **Transitivity:** if $e \leq f$ and $f \leq g$, then $e \leq g$

Such an ordering always admits at least one maximal element [16]

Definition 4 (Top of a Poset [33]). *Let (E, \leq) be a partially ordered set. The top of E , denoted $\text{Top}(E)$, is the set of maximal elements of E :*

$$\text{Top}(E) = \{ e \in E \mid \nexists f \in E \text{ with } e < f \}. \quad (2.9)$$

That is, $\text{Top}(E)$ contains those events in E for which no strictly greater event exists.

The elements of $\text{Top}(E)$ represent the current causal frontier—the most recent events that have occurred but have no successors [148]. Although $\text{Top}(E)$ may contain several incomparable (spacelike) elements, it is never

empty and therefore provides a well-defined notion of a “last event” from the observer’s perspective. This frontier defines the light-cone boundary and the terminal particle–wave interaction that delimits all accessible information.

Every event $e \in E$ corresponds to an irreducible distinction in the experimental record. Under the measurable embedding $\Psi : E \rightarrow \mathcal{T}$ introduced in Thought Experiment 2.3.1, each logical event is mapped to an algebraic object \mathbf{E}_e in the tensor algebra. These objects compose whenever their corresponding events are compatible in the causal order, so the accumulation of observed events yields a record that reflects the ordered refinement of the causal set.

The goal of this section is to define a cumulative object \mathbf{U}_n —the *Causal Universe Tensor*—that embodies the total informational content of all events observed up to step n in the current inertial reference frame. This tensor is not a dynamical evolution. It is the bookkeeping device that records which refinements have survived admissibility.

It is crucial to emphasize that no background time parameter is introduced. There is no external clock and no continuous variable t against which events are measured. Instead, Axiom 5 guarantees that the causal set admits a linear extension: the events can be listed in a sequence that respects causal precedence. In this framework, *time* is merely the ordinal index of an event in such a sequence. It is not a physical field or metric quantity, but a bookkeeping device that labels the relative order of observations.

With this viewpoint, accumulating the event tensors in order is not evaluating a function of time. It is forming the ordered product of distinctions that have occurred. The resulting object, the Causal Universe Tensor, represents the total recorded history up to any chosen ordinal position in the list of events.

2.5.2 Special Relativity

Every physical description begins not with space or time, but with an *event*—an interaction that makes previously indistinguishable outcomes distinct [13, 123]. The causal boundary of such an interaction is its *light cone*: the set of all events that can influence or be influenced by it according to special relativity [45, 108]. The intersection of two light cones, corresponding to the last particle–wave interaction accessible to an observer, defines the maximal region of causal closure [70, 118]. Beyond this surface, no additional information can be exchanged; all distinguishable action has concluded.

It is from this closure that the ordering of events arises. Each measurable interaction contributes one additional distinction to the universe, expanding its causal surface by a finite count [70, 102]. The smooth fabric of spacetime is not primitive but emergent: it is the limiting behavior of discrete causal increments accumulated along the light cone [16, 146]. Within each cone, the universe can be represented by a finite tensor of interactions—local updates to a global state—that together approximate continuity only through cancellation across countable events [16, 147].

Thought Experiment 2.5.2 (Boosting Velocity [45]). **N.B.**—*The Lorentz equations are presented here only as an illustrative model of the local causal structure. They are not derived from the axioms in this chapter, and no subsequent argument depends on them.* \square

Special relativity provides the canonical local model for this causal structure. Consider the Lorentz transformation for a boost of velocity v in one spatial dimension, [129, 153]

$$\begin{pmatrix} t' \\ x' \end{pmatrix} = \begin{pmatrix} \gamma & -\gamma v/c^2 \\ -\gamma v & \gamma \end{pmatrix} \begin{pmatrix} t \\ x \end{pmatrix}, \quad \gamma = \frac{1}{\sqrt{1 - v^2/c^2}}. \quad (2.10)$$

For infinitesimal separations satisfying $x = ct$, the Lorentz transformation

gives

$$t' = \gamma t(1 - v/c). \quad (2.11)$$

If we take $\Delta t = 1$ as the unit interval between distinguishable events, then observers moving at relative velocity v will, in general, disagree on the number of such events that occur between two intersections of their respective light cones [108]. The only invariant quantity is the causal ordering itself: all observers concur on which event precedes which, even though they may count a different number of intermediate ticks [102].

Definition 5 (Rank time [16, 33]). *Let (E, \prec) be a locally finite partially ordered set of events. A rank time is an order-embedding*

$$\tau : E \rightarrow \text{Ord}$$

satisfying $e \prec f \implies \tau(e) < \tau(f)$. Local finiteness implies that for any observer's causal domain $D \subseteq E$, $\tau(D)$ is order-isomorphic to an initial segment of \mathbb{N} . We therefore define the duration, $|\delta t|$, between anchors $a \prec b$ by

$$|\delta t|(a, b) = \#\{e \in E \mid a \prec e \prec b\} \in \mathbb{N}.$$

Two rank functions τ, τ' are equivalent if there exists an order-isomorphism ϕ with $\tau' = \phi \circ \tau$; equivalent ranks yield identical durations.

Remark 1 (Operational content of time). *Time is an ordinal rank on E , not an independent scalar field. All subsequent uses of “ t ” refer to an order-equivalence class of rank functions as in Definition 5. The additivity $|\delta t|(a, b) = |\delta t|(a, c) + |\delta t|(c, b)$ follows from local finiteness.*

This observation motivates the first physical axiom: that time is not an independent scalar field but an ordinal index over causally distinguishable events. Each event increments the universal sequence by one count; each observer's clock is a local parametrization of that same count under Lorentz

contraction. The apparent continuity of time is the result of the density of such events within the causal cone, not an underlying continuum of duration.

2.5.3 On the Structure of Measurement

In this formulation, a measurement is not the evaluation of a continuous quantity against an external time parameter. No clock, ruler, or metric is assumed. Instead, the Axioms of Planck and Cantor assert that an observer's record is a locally finite, causally ordered set of distinguishable events. To extract a numerical value from such a record, one must identify which events satisfy a specified property and count how many of them occur between two anchors in the causal order.

This viewpoint treats measurement as a purely combinatorial act: the *value* of a measurement is the number of admissible distinctions satisfying a predicate inside a finite causal interval. The result is always an integer, and continuity—when it appears—arises only as the smooth limit of increasingly refined counts. We formalize this as follows.

Definition 6 (Measurement [160]). *Let (E, \prec) be a locally finite partially ordered set of events, and let $P : E \rightarrow \{0, 1\}$ be a predicate designating which events satisfy a specified property. For two anchor events $a, b \in E$ with $a \prec b$, the measurement of P between a and b is the finite integer*

$$M_P[a, b] := |\{e \in E : a \prec e \prec b \text{ and } P(e) = 1\}| \in \mathbb{N}.$$

That is, a measurement is a count of distinguished events satisfying P within the causal interval (a, b) .

A measurement in this setting is therefore nothing more than a count of distinguished events between anchors. Numerical values arise only when such counts are compared against a conventional scale. No continuous quantity is assumed *a priori*; continuity is inferred from the refinement of a finite causal

record. In practice, every physical “number” depends on a calibration that relates discrete counts to a chosen system of units.

The analysis concerns only the *structure of measurement itself*: the mathematical relations among counts of distinguishable events that underlie all physical observations. In this framing, physics is viewed as a grammar of distinctions. The familiar constants and fields—mass, charge, curvature, temperature—arise as *derived measures* within a finite causal order, not as independent entities.

Thought Experiment 2.5.3 (Measurement [5, 111, 160]). **N.B.**—*Measurement is a formal writing system. Each observation selects a symbol from a finite alphabet, and a record is the word formed by these selections. No physical semantics are assumed; the structure is purely syntactic in the Backus–Naur sense [5, 111]. In the spirit of Wheeler’s dictum that information is fundamental [160], the act of measurement is treated here as the creation of symbolic distinctions, nothing more.* \square

Because measurement produces distinguishable outcomes, each observation selects a symbol from a finite or countable alphabet

$$\Sigma = \{\sigma_1, \sigma_2, \dots\}.$$

A record of n measurements is therefore a word $w \in \Sigma^n$. When an instrument is refined—by increasing precision or reducing noise—any coarse symbol σ_k may be replaced by a finite set of more precise symbols,

$$\sigma_k \Rightarrow \sigma_{k,1} \mid \sigma_{k,2} \mid \dots \mid \sigma_{k,r},$$

just as in a Backus–Naur Form (BNF) production rule [5, 111]. Not all replacements are admissible: they must remain compatible with every other measurement that overlaps in time or causal order. Two refined histories that disagree on an overlapping interval cannot both represent valid records.

Thus admissible measurement histories form a formal language generated

by the allowed refinement rules. The “law” governing measurement is the constraint that only globally consistent extensions of a record may be generated. This is not an analogy: it is the standard formal structure of symbol sequences in coding and information theory [142].

No new particles, forces, or cosmological effects are introduced. The present theory interrogates only the rules by which such effects are numerically expressed. It is not a revision of physics but a clarification of its syntax: a study of the measures of phenomena rather than the phenomena themselves. We begin by introducing the necessary set-theoretic language and associating it with a discrete notion of “time.”

Definition 7 (Distinguishability Chain [88]). *Let Ω be a nonempty set. A distinguishability chain on Ω is a sequence $\mathcal{P} = \{P_n\}_{n \in \mathbb{Z}}$ of partitions $P_n \in \text{Part}(\Omega)$ such that P_{n+1} refines P_n for all n (every block of P_{n+1} is contained in a block of P_n). Write $\text{Bl}(P)$ for the set of blocks of a partition P . Each refinement step produces zero or more event.*

Definition 8 (Event [88, 148]). *Fix a distinguishability chain $\mathcal{P} = \{P_n\}$. An event at index n is a minimal refinement step: a pair*

$$e = (B, \{B_i\}_{i \in I}, n) \tag{2.12}$$

such that:

1. $B \in \text{Bl}(P_n)$;
2. $\{B_i\}_{i \in I} \subseteq \text{Bl}(P_{n+1})$ is the family of all blocks of P_{n+1} contained in B , with $|I| \geq 2$ (a nontrivial split);
3. (minimality) there is no proper subblock $C \subsetneq B$ with $C \in \text{Bl}(P_n)$ for which the family $\text{Bl}(P_{n+1}) \cap \mathcal{P}(C)$ is nontrivial.

Let E denote the set of all such events. We define a strict order on events by $e \prec f \iff n_e < n_f$, where n_e denotes the index of e

Definition 9 (Proper Time [109]). *Let E be the set of events generated by a distinguishability chain $P = \{P_n\}$. For any two events $a, b \in E$ with $a \prec b$, the proper time between them is*

$$\tau(a, b) = \max \left\{ |C| : C = \{c_0, \dots, c_k\} \subseteq E, a = c_0 \prec c_1 \prec \dots \prec c_k = b \right\}.$$

That is, $\tau(a, b)$ is the cardinality of a maximal chain of strictly refinable events between a and b . Local finiteness of the distinguishability chain guarantees $\tau(a, b) \in \mathbb{N}$.

Remark 2. *Proper time is not a geometric length. It is the number of admissible, irreversible refinements separating two recorded events. Additional refinement (higher resolution) may increase $\tau(a, b)$; coarse-graining cannot. Thus, proper time is an invariant of the partially ordered event record, not a metric assumption.*

Remark 3. *A chain need not include all events, and incomparable events do not contribute to one another's proper time. Only if every pair of events were comparable would τ reduce to a total order. In general, E is only partially ordered.*

N.B.—*In physical terms, this corresponds to relativistic simultaneity: incomparable events occupy disjoint causal domains and cannot be ordered by any observer.* \square

Remark 4 (Smooth limit [128]). *If refinements become dense and the discrete extremal converges to a C^2 spline, then*

$$\lim_{\text{refinement} \rightarrow \infty} \tau(a, b) = \int_a^b \sqrt{-ds^2},$$

the Lorentzian proper time. The integral form is not assumed; it is the smooth shadow of the combinatorial count.

Phenomenon 2.5.1 (The Fourier Budget Constraint). *A refinement ledger possesses a finite informational budget. To specify a location with increasing precision consumes refinement capacity that would otherwise resolve rates of change. A perfectly sharp position therefore requires an unbounded refinement cost in momentum.*

This is not a probabilistic limitation but a bookkeeping constraint: the ledger cannot afford to encode arbitrarily precise conjugate refinements simultaneously. The apparent uncertainty is the shadow of a finite refinement budget.

The notion of *uncorrelant events* formalizes the idea that two recorded distinctions may be independent of one another. In causal set theory, incomparability under the causal order corresponds to physical independence of events [16]. The same conceptual separation appears in quantum theory, where observables acting on independent subsystems commute and their measurement outcomes do not influence each other [39, 120]. Classical discussions of separated systems, from Einstein–Podolsky–Rosen and Schrödinger to Wheeler’s formulation of complementarity [46, 134, 161], frame the same idea operationally: when no physical procedure can distinguish the relative order of two events, their ordering has no empirical content. The definitions below captures this in the minimal set-theoretic language of the causal poset.

Definition 10 (Uncorrelant [16, 146]). *Let (E, \prec) be a locally finite partially ordered set of events. Two events $e, f \in E$ are said to be uncorrelant if they are incomparable under the causal order; that is,*

$$\neg(e \prec f) \quad \text{and} \quad \neg(f \prec e).$$

The uncorrelant relation partitions E into equivalence classes of events whose relative order carries no operational consequence for any admissible measurement or refinement. In particular, no experimentally distinguishable difference follows from interchanging the positions of uncorrelant events in any

linear extension of (E, \prec) .

Remark 5. *If e and f are uncorrelant, permuting them in any chain, merge, or refinement does not change any observable invariant of the Causal Universe Tensor. No observer can construct a sequence of measurements that forces an ordering between e and f without introducing new events.*

Remark 6. *Correlant events may, but do not necessarily, admit a strict causal precedence and therefore contribute to proper time along a chain; uncorrelants do not. In particular, a chain that includes e but not f may be maximally refined without reference to f . Thus, uncorrelants represent informational independence, not simultaneity.*

Remark 7. *In spacetime language, uncorrelants are precisely those event pairs that are spacelike-separated: reordering them changes no measurable scalar. Here, this is not assumed from geometry; it is a consequence of incomparability in the event order.*

Definition 11 (Causal Network [16]). *Let E be a finite set of admissible events and let \triangleright denote the immediate causal cover: $e \triangleright f$ if and only if $e < f$ and there exists no $g \in E$ such that $e < g < f$. The causal network is the directed graph (E, \triangleright) whose vertices are the events in E and whose directed edges record the immediate causal relations.*

This network is the combinatorial diagram of the event record: each vertex is a distinguishable event, and each directed edge $e \triangleright f$ certifies that f cannot be observed without first observing e . Its transitive closure recovers the full causal order $<$ of Definition 12.

Definition 12 (Causal Order [16]). *Let $P = \{P_n\}_{n \in \mathbb{Z}}$ be a distinguishability chain of partitions, and let an event be $e = (B, \{B_i\}_{i \in I}, n)$ as in Definition 8, where $B \in \text{Bl}(P_n)$ splits nontrivially into child blocks $\{B_i\} \subset \text{Bl}(P_{n+1})$.*

For $m > n$ and $C \in \text{Bl}(P_m)$, let $\pi_{m \rightarrow n}(C) \in \text{Bl}(P_n)$ denote the unique ancestor block in P_n containing C (well-defined because P_{n+1} refines P_n). Define the immediate causal cover relation $e \triangleright f$ between events $e = (B, \{B_i\}, n)$ and $f = (C, \{C_j\}, m)$ by

$$n < m \quad \text{and} \quad \pi_{m \rightarrow n+1}(C) \subseteq B_i \text{ for some child } B_i \text{ created by } e.$$

The causal order \prec on the event set E is the transitive closure of \triangleright :

$$e \prec f \iff \text{there exist events } e = e_0, e_1, \dots, e_k = f \text{ with } e_i \triangleright e_{i+1} \text{ for all } i.$$

Then (E, \prec) is a locally finite partially ordered set (reflexivity suppressed for strictness), where incomparability is allowed: it may happen that neither $e \prec f$ nor $f \prec e$.

As an illustration, recall the twin paradox of the previous chapter³. In the informational gauge, proper time is not a geometric interval but the work of reconciling distinguishable events. The traveling twin accrues a denser log of refinements—engine burns, course corrections, telemetry—while the stay-at-home twin records a coarser sequence. When their notebooks are merged into a single coherent history, the richer record requires strictly greater informational effort to reconcile. Equivalently, the proper time of the unaccelerated twin is necessarily longer, because her history contains fewer distinctions and therefore a larger merge is required to absorb those recorded by her sibling. In the smooth limit this appears as a shorter proper time along the curved worldline, but the effect is not mysterious: it is the discrete fact that one history contains more recorded distinctions than the other. Geometry only codifies what measurement already certified.

Remark 8 (Index is an order-embedding, not an equivalence). *If $e \prec f$, then $n_e < n_f$. Thus the refinement index provides a rank function (Definition 5)*

³See Coda: The Twin Paradox, Chapter 1.

that is monotone with respect to \prec . The converse need not hold: $n_e < n_f$ does not imply $e \prec f$. Hence the causal order is generally partial, not total.

Remark 9 (Uncorrelants and permutation invariance). *Events e, f with neither $e \prec f$ nor $f \prec e$ are uncorrelant (incomparable). Permuting uncorrelant events in any linear extension of (E, \prec) leaves all scalar invariants of the Causal Universe Tensor unchanged; causal histories are unique only up to permutation of uncorrelants.*

2.5.4 Accumulation of Measurement

Operationally, every observation can be decomposed into three layers:

1. the **logical** layer—which events are distinguishable;
2. the **mathematical** layer—how those distinctions are counted;
3. the **physical** layer—how the resulting counts are named and parameterized as energy, momentum, or time.

By isolating the first two layers, we obtain a calculus of variations that is universal to any admissible physics: a closed system of relations that expresses how order itself becomes measurable.

The framework that follows formalizes this intuition. Within ZFC, we construct an ordered set of events whose distinguishability relations generate the causal ordering of special relativity. Measurements are counts of these relations, and the Causal Universe Tensor—the cumulative left product of *event tensors* over all causal increments—supplies the discrete substrate from which the smooth laws of physics arise in the limit of refinement.

The preceding axioms establish a measurement record as a locally finite, partially ordered set of distinctions (E, \prec) . This structure is purely combinatorial. To connect the logical record to physical measurement, we require a representation that can carry numerical values and allow recorded

distinctions to combine. A vector space V provides a domain for measurable quantities, but to represent successive distinctions we also need a rule for composition. The tensor algebra $\mathcal{T}(V)$ is the freest algebra generated by V : it contains V , supports noncommutative products, and imposes no additional relations beyond those required by linearity. By associating each logical event e_k with a tensor \mathbf{E}_k in $\mathcal{T}(V)$ we obtain an algebraic record of distinctions that can be composed and accumulated. This representation introduces no structure beyond what is logically required to encode measurable updates.

Definition 13 (Event Tensor [62]). *Let V be a finite-dimensional real vector space of measurable quantities. An event tensor $\mathbf{E}_k \in \mathcal{T}(V)$ encodes the distinguishable contribution of the k th event $e_k \in E$ to the cumulative record. It is related to the logical event by a measurable embedding*

$$\Psi : E \rightarrow \mathcal{T}(V), \quad \mathbf{E}_k = \Psi(e_k). \quad (2.13)$$

No algebraic relations are assumed beyond those required by linearity: \mathbf{E}_k is simply the algebraic image of the k th logical distinction.

An individual event tensor records a single admissible refinement of the measurement record. To represent the cumulative effect of many events, we must specify how these algebraic objects combine. Because the causal set is ordered only up to informational precedence, the combination rule must respect a chosen linear extension of the partial order and must make no assumptions of commutativity. This leads naturally to a left-multiplicative update: each new event contracts the admissible record of all that precede it, and the cumulative history is represented by the product of these restricted increments along any finite prefix of the causal chain.

The combination rule corresponds directly to the set-theoretic refinement of admissible outcomes. At each step, the new logical event is not taken in

isolation, but restricted against all prior observations:

$$e'_{k+1} := e_{k+1} \cap \bigcap_{j=1}^k \hat{R}(e_j),$$

where \hat{R} is the operator that removes outcomes incompatible with the existing record. In this framework, the “laws of physics” appear nowhere else: they are encoded entirely in the restriction operator. What survives admissibility is physical; what is removed was never a possible history.

In the algebraic domain this restriction is represented by

$$\mathbf{U}_{k+1} := \Psi(e'_{k+1}) \mathbf{U}_k = \Psi(e_{k+1} \cap \hat{R}(e_k)) \mathbf{U}_k,$$

where Ψ embeds the surviving distinctions into the tensor algebra. Each new event therefore contracts the admissible history by left multiplication. The cumulative record is the product of these restricted increments along any finite prefix of the causal chain.

Formally, the measurable embedding Ψ sends the set-theoretic restriction to a multiplicative update in the tensor algebra. Instead of embedding the raw event e_{k+1} , we embed only the portion that survives all prior admissibility constraints:

$$\mathbf{E}_{k+1} = \Psi\left(e_{k+1} \cap \bigcap_{j=1}^k \hat{R}(e_j)\right).$$

Writing $\mathbf{R}(e) := \Psi(\hat{R}(e))$, the cumulative record evolves by left multiplication:

$$\mathbf{U}_{k+1} = \mathbf{R}(e_{k+1}) \mathbf{U}_k = \Psi(e_{k+1} \cap \hat{R}(e_k)) \mathbf{U}_k, \quad 0 \leq k < n.$$

Thus the tensor update is the algebraic realization of the same logical operation performed in E : a new event is applied only after its outcomes have been restricted by all earlier observations. The universe accumulates consistency

through products of restricted increments, not by additive evolution.

Definition 14 (Partition of the Event Set [66]). *Let (E, \prec) be a locally finite partially ordered set of distinguishable events. A partition of E is a collection of disjoint subsets $\{E_\alpha\}_{\alpha \in A}$ such that*

$$E = \bigcup_{\alpha \in A} E_\alpha, \quad E_\alpha \cap E_\beta = \emptyset \quad \text{for } \alpha \neq \beta.$$

Each E_α is an informationally independent component: no event in E_α refines or is refined by an event in E_β . Correlant events therefore lie within the same partition element, while uncorrelants lie in distinct elements of the partition.

Definition 15 (Restriction Operator). **N.B.**—*The restriction operator is the first construction that is intrinsic to the informational framework developed in this work. All earlier definitions (posets, partitions, causal networks, and measurements) are standard mathematical structures. The restriction operator encodes admissibility: it records which refinements survive in the presence of prior observations. This mechanism does not appear in classical order theory or in conventional treatments of causality, and it is the essential ingredient that links local measurement constraints to global informational consistency.* \square

Let (E, \prec) be a partially ordered set of events, and let $e \in E$ be a newly recorded event. The restriction operator

$$\hat{R}(e) : E \rightarrow E$$

acts on the event record by removing any outcomes that are incompatible with e . For $f \in E$,

$$\hat{R}(e)(f) = \begin{cases} f, & \text{if } f \text{ is admissible given } e, \\ \emptyset, & \text{otherwise.} \end{cases}$$

Equivalently, if E_α is the partition element containing e , then

$$\hat{R}(e) : E_\alpha \mapsto E'_\alpha, \quad E'_\alpha = \{ f \in E_\alpha \mid f \text{ is compatible with } e \}.$$

Thus $\hat{R}(e)$ contracts the event domain by discarding outcomes that contradict the new distinction.

We now present the *Causal Universe Tensor*.

Proposition 1 (The Existence of a Causal Universe Tensor). *Let (E, \prec) be a locally finite partially ordered set of events, and let $\Psi : E \rightarrow \mathcal{T}(V)$ be the measurable embedding. For each event $e \in E$, define its admissible factor by*

$$\mathbf{F}(e) := \Psi(\hat{R}(e)).$$

Fix a finite linear extension $e_1 \prec \cdots \prec e_n$ of (E, \prec) and set $\mathbf{U}_0 := \mathbf{I}$ (the multiplicative identity in $\mathcal{T}(V)$). Define the left recursion

$$\mathbf{U}_{k+1} := \mathbf{E}_{k+1} \mathbf{U}_k = \Psi(e_{k+1} \cap \hat{R}(e_k)) \mathbf{U}_k, \quad 0 \leq k < n, \quad (2.14)$$

Then:

1. (Naturality of restriction) *The update (2.14) is precisely the expansion of the discrete restriction through the representation: $\mathbf{F}(e) = \Psi(\hat{R}(e))$ so that*

$$\mathbf{U}_{k+1} = \Psi(\hat{R}(e_{k+1})) \mathbf{U}_k.$$

Equivalently, $R \circ \Psi = \Psi \circ \hat{R}$ on $\text{im } \Psi$.

2. (Causal uniqueness) *The recursion (2.14) is uniquely determined by the chosen linear extension. Any two linear extensions differ only by permutations of informationally independent events (partition elements of E), so once the order is fixed the product is mechanically well-defined.*

3. (Independence under commuting factors) *If a subset $S \subset \{1, \dots, n\}$ indexes events whose admissible factors pairwise commute, $\mathbf{F}(e_i)\mathbf{F}(e_j) = \mathbf{F}(e_j)\mathbf{F}(e_i)$ for $i, j \in S$, then any permutation of $\{\mathbf{F}(e_i)\}_{i \in S}$ leaves \mathbf{U}_n invariant under all cyclic scalar functionals (e.g., traces of contractions).*
4. (Fully commutative case) *If all admissible factors commute, then*

$$\mathbf{U}_n = \prod_{k=1}^n \mathbf{F}(e_k)$$

is independent of the linear extension; the product reduces to the order-insensitive accumulation of factors.

Categorically, the structure underlying this result is the naturality of a monoidal functor in the sense of Mac Lane [94], with further development in Kelly [84] and Leinster [97]. The proof sketch below follows this diagrammatic perspective; the fully explicit ZFC realization appears in Appendix A.

Proof (Sketch). Let \mathcal{E} be the refinement category of admissible event records, with objects the event sequences and morphisms the refinement maps $\widehat{R} : \mathbf{i} \rightarrow \widehat{R}(\mathbf{i})$. Let $T(V)$ be the tensor algebra regarded as a *symmetric monoidal category* under the tensor product.

The embedding $\Phi : E \rightarrow T(V)$ extends uniquely to a monoidal functor

$$\Phi^{(\bullet)} : \mathcal{E} \longrightarrow T(V)^{(\bullet)}, \quad \mathbf{i} = (i_1, \dots, i_n) \longmapsto (\Phi(e_{i_1}), \dots, \Phi(e_{i_n})),$$

sending refinement maps to componentwise restriction on the image.

A refinement $\widehat{R} : \mathbf{i} \rightarrow \mathbf{j}$ in \mathcal{E} is a morphism expressing that \mathbf{j} is the universal solution to a finite cone of compatibility conditions. Under the functor $\Phi^{(\bullet)}$, this induces a canonical morphism

$$\Phi^{(\bullet)}(\widehat{R}) : \Phi^{(n)}(\mathbf{i}) \longrightarrow \Phi^{(m)}(\mathbf{j}).$$

By functoriality, the square

$$\mathbf{i} @ > \widehat{R} >> \mathbf{j}$$

$$@V\Phi^{(n)}VV @VV\Phi^{(m)}V$$

$$\Phi^{(n)}(\mathbf{i}) @ > \Phi^{(\bullet)}(\widehat{R}) >> \Phi^{(m)}(\mathbf{j})$$

commutes. This is exactly the naturality square.

To obtain the Causal Universe Tensor, form the *monoidal accumulation* of the embedded sequence:

$$U(\mathbf{i}) := \Phi(e_{i_1}) \otimes \cdots \otimes \Phi(e_{i_n}).$$

Since $T(V)$ is symmetric monoidal, any two linear extensions of a finite event poset differ by braidings of incomparable events, and such braidings commute with the tensor structure. Hence $U(\mathbf{i})$ is well defined up to the canonical symmetry of the monoidal category.

Thus the Causal Universe Tensor is the monoidal image of a refinement diagram under a functor preserving both the tensor product and the naturality of refinement. \square

A full proof is provided in Appendix A.1.

The existence of the Causal Universe Tensor gives rise to the appearance of stability in long sequences of refinement. Because each admissible update is not free to evolve arbitrarily, but must remain compatible with the unique globally coherent extension of the record, deviations cannot accumulate without bound. Local inconsistencies are absorbed through restriction and embedding, producing the observable effect of bounded variation in the measurement ledger. This structural stability is not enforced by physical feedback or control, but by the logical necessity of coherent refinement itself.

This gives rise to the following informational phenomenon.

Phenomenon 2.5.2 (The Statistical Process Effect [140]). *A sequence of measurements refined under admissible updates exhibits structural stability. Local deviations are smoothed by the unique coherent extension enforced by restriction and embedding. The resulting record remains bounded, not by physical forces, but by the logical requirement of global consistency. This informational stability is the phenomenon known in classical practice as statistical process control.*

Remark 10 (Ordinal determinacy). *The sequence (\mathbf{U}_k) is not merely algebraically well-defined but physically ordered: k indexes ordinal rank, not arbitrary enumeration. Hence any reordering outside uncorrelant classes violates Axiom 5.*

With the ordinal structure of events established, we now formalize how these measurements combine algebraically within a finite vector space.

2.5.5 Formal Structure of Event and Universe Tensors

We now specify the algebraic structure of the quantities introduced above. Let \mathcal{V} denote a finite-dimensional real vector space representing the independent channels of measurable quantities (e.g. energy, momentum, charge). Define the tensor algebra [65, 95]

$$\mathcal{T}(\mathcal{V}) = \bigoplus_{r=0}^{\infty} \mathcal{V}^{\otimes r}, \quad (2.15)$$

whose elements are finite sums of r -fold tensor products over \mathbb{R} . Each *event tensor* E_k is a member of $\mathcal{T}(\mathcal{V})$ encoding the distinguishable contribution of the k -th event to the global state. We write

$$\mathbf{E}_k \in \mathcal{T}(\mathcal{V}), \quad \mathbf{U}_n = \prod_{k=1}^n \mathbf{E}_k \in \mathcal{T}(\mathcal{V}). \quad (2.16)$$

Addition is understood componentwise in the direct sum and preserves the ordering of indices guaranteed by the Axiom of Order [16, 65]. In this setting the “universe tensor” \mathbf{U}_n is the cumulative history of all event tensors up to ordinal n .

Definition 16 (Tensor Algebra [62]). *The tensor algebra on a vector space \mathcal{V} is*

$$\mathcal{T}(\mathcal{V}) = \bigoplus_{r=0}^{\infty} \mathcal{V}^{\otimes r}$$

with componentwise addition and associative tensor product

Remark 11. *Each logical event e_k in the partially ordered set (E, \prec) induces a tensor $\mathbf{E}_k = \Psi(e_k)$ in $\mathcal{T}(\mathcal{V})$. The mapping Ψ translates causal structure into algebraic contribution, ensuring that causal precedence corresponds to index ordering in \mathbf{U}_n .*

Because $\mathcal{T}(\mathcal{V})$ is a free associative algebra, all operations on \mathbf{U}_n are well defined using the standard linear maps, contractions, and bilinear forms of \mathcal{V} . The subsequent analysis of variation and measurement therefore proceeds entirely within conventional linear-operator theory.

From the definition of the Universe Tensor

$$U_n = \prod_{k=1}^n E_k, \quad (2.17)$$

we may regard an *uncorrelant* as any subset of events whose local order can be permuted without altering the global scalar invariants of U_n . Formally, a subset $S \subseteq \{E_1, \dots, E_n\}$ is uncorrelant if, for every permutation π of S ,

$$\prod_{E_i \in S} E_i = \prod_{E_i \in S} E_{\pi(i)}. \quad (2.18)$$

In this case, all contractions or scalar traces derived from U_n remain unchanged by reordering the elements of S , even though the operator sequence

itself may differ.

Definition 17 (Commutator and Commutator Ideal [42]). *Let \mathcal{A} be an algebra over a field \mathbb{F} with bilinear multiplication $(x, y) \mapsto xy$. For $x, y \in \mathcal{A}$, the commutator of x and y is the element*

$$[x, y] := xy - yx \in \mathcal{A}.$$

The set of all finite \mathbb{F} -linear combinations of commutators,

$$[\mathcal{A}, \mathcal{A}] := \left\{ \sum_{i=1}^m \alpha_i [x_i, y_i] : \alpha_i \in \mathbb{F}, x_i, y_i \in \mathcal{A} \right\},$$

is called the commutator ideal. It is the smallest two-sided ideal of \mathcal{A} that contains every element $xy - yx$; equivalently, it is the smallest linear subspace of \mathcal{A} closed under left and right multiplication by arbitrary elements of \mathcal{A} .

Remark 12 (Algebraic Characterization of Informational Independence). *Let $\Psi : E \rightarrow \mathcal{T}(V)$ be the event embedding and $\mathbf{E}_e := \Psi(e)$. If $S \subseteq E$ lies in distinct elements of the partition of E (Definition 14), then the admissible increments $\{\mathbf{E}_e\}_{e \in S}$ pairwise commute. Consequently, any reordering of these factors within a linear extension of (E, \prec) produces the same value of \mathbf{U}_n under all cyclic scalar functionals (e.g., traces of contractions). In this algebraic sense, informational independence corresponds exactly to order-insensitive contribution to the invariants derived from \mathbf{U} .*

Phenomenon 2.5.3 (Non-commutative Event Pair [70]). **N.B.**—*Non-commutative event tensors often signal a dependency: one update must precede the other for the restricted outcome set to remain consistent. Reversing such events changes the operator state, even though measurable scalar invariants remain the same.* \square

Let $V = \mathbb{R}^2$ and let event tensors act as 2×2 matrices under the usual

(non-commutative) multiplication. Define

$$E_A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad E_B = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}.$$

A direct computation gives

$$E_A E_B = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \neq \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} = E_B E_A, \quad \text{so } [E_A, E_B] \neq 0.$$

Thus, applying the updates in different orders leads to different operator states. However, cyclic scalar invariants agree:

$$\text{tr}(E_A E_B) = \text{tr}(E_B E_A) = 3, \quad \det(E_A E_B) = \det(E_A) \det(E_B) = 1.$$

In this sense, noncommutativity affects the internal operator record but not the measurable quantities obtained by cyclic scalar functionals.

Phenomenon 2.5.4 (Independent Event Chains [96]). **N.B.**—This is analogous to the inertial segment of the twin paradox. During coasting, neither twin exchanges signals with the other, so no event on one worldline refines or restricts events on the other. The two chains are informationally independent until a causal interaction occurs. \square

Consider two finite event chains

$$A_1 \prec A_2, \quad B_1 \prec B_2,$$

with no causal relation between any A_i and any B_j . Let their event tensors act on $V = \mathbb{R}^2$ as

$$E_{A1} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad E_{A2} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad E_{B1} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad E_{B2} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

Because the A-events refine only the A-chain and the B-events refine only the B-chain, their admissible factors commute:

$$E_{A2}E_{B2} = E_{B2}E_{A2}.$$

Thus, any linear extension of the partial order may place the A- and B-events in either interleaving without changing cyclic scalar invariants. For example, applying the four events in the order

$$A_1, A_2, B_1, B_2 \quad \text{or} \quad A_1, B_1, A_2, B_2$$

gives operator states that differ, but

$$\text{tr}(E_{A2}E_{B2}) = \text{tr}(E_{B2}E_{A2}) = 1, \quad \det(E_{A2}E_{B2}) = \det(E_{B2}E_{A2}) = 0.$$

This illustrates the algebraic meaning of independence: when two event chains are partitioned into disjoint informational domains, their admissible increments commute. Order affects the internal operator record but leaves measurable cyclic scalars unchanged, exactly as in the coasting phase of the twin paradox.

2.6 Information Minimality and Kolmogorov Closure

The previous definitions describe events as finite distinctions and their ordering as a partial refinement of information. What remains is the rule that determines which extensions of a recorded event set are admissible. Not every history consistent with the order is physically meaningful: a completion that inserts unobserved structure would imply additional measurements that never occurred. Information minimality formalizes this constraint through algorithmic information theory in the sense of Kolmogorov, Solomonoff, and

Chaitin [24, 87, 144, 145].

We treat histories as finite symbolic strings and measure their descriptive content by Kolmogorov complexity. A physically admissible history is one that cannot be compressed by adding unrecorded structure.

Definition 18 (Kolmogorov Complexity [24, 87]). *Fix a universal Turing machine U [155]. For any finite string $w \in \Sigma^*$, the Kolmogorov complexity $K(w)$ is the length of the shortest input to U that outputs w and halts. The functional $K : \Sigma^* \rightarrow \mathbb{N}$ is defined up to an additive constant independent of w .*

Definition 19 (Admissible Extension [99]). *Let $E = \{e_0 \prec e_1 \prec \cdots \prec e_n\}$ be the recorded events of an experiment. A finite string $w \in \Sigma^*$ is an extension of E if its image under the event map contains E in the same causal order. An extension w is admissible if it introduces no additional events beyond E ; that is, every distinguishable update encoded by w has a corresponding element of E . Any extension predicting unobserved structure is rejected as inadmissible.*

Phenomenon 2.6.1 (Paradoxes of Time Travel [63, 98]). **N.B.**—*Apparent paradoxes often attributed to time travel, remote viewing, or other extraordinary mechanisms are pathologies of over-resolution. They arise when incompatible refinements are treated as simultaneously admissible, producing the illusion of phenomenal violation rather than an actual failure of causal order.* \square

N.B.—*This thought experiment introduces constructions that are intentionally self-referential. These devices are used only to illustrate how paradoxes arise when an observer attempts to treat its own temporal index as a manipulable datum. Such constructions lie outside the admissible structure of the axioms and are not permitted in any formal derivation. In particular, they follow the general pattern of self-reference that Gödel cautioned against in his incompleteness results: systems that encode statements about their own*

inferential process cannot, in general, maintain global consistency [63]. The paradoxes described here therefore serve only as intuitive warnings. They do not represent allowable configurations within the theory, and no phenomenon in this manuscript relies on them. \square

Let $E = \{e_1, e_2, e_3, \dots\}$ be a locally finite causal chain where each event e_i has a unique successor e_{i+1} . Define the corresponding universe tensor

$$\mathbf{U}_n = \sum_{k=1}^n \mathbf{E}_k, \quad \mathbf{E}_k = \Psi_k(e_k). \quad (2.19)$$

Now suppose we attempt to “extend” this history by splitting a single event e_j into uncountably many indistinguishable refinements:

$$e_j \longrightarrow \{e_{j,\alpha}\}_{\alpha \in [0,1]}, \quad (2.20)$$

each representing a formally distinct but observationally identical outcome. Algebraically, this replacement yields

$$\mathbf{E}_j \longrightarrow \int_0^1 \mathbf{E}_{j,\alpha} d\alpha, \quad (2.21)$$

so that the next update becomes

$$\mathbf{U}_{n+1} = \mathbf{U}_n + \int_0^1 \mathbf{E}_{j,\alpha} d\alpha. \quad (2.22)$$

This “extension” violates the finiteness and distinguishability conditions necessary for causal coherence:

1. The set $\{e_{j,\alpha}\}$ is uncountable, destroying local finiteness;
2. The new events are indistinguishable, so Extensionality no longer guarantees unique contributions;
3. The total tensor amplitude U_{n+1} can diverge or cancel arbitrarily, de-

pending on how the continuum of duplicates is treated.

Operationally, this is a Banach–Tarski-like overcounting: the causal structure has been “refined” in a way that preserves measure only formally while the order relation collapses. The observer would now predict contradictory outcomes for the same antecedent state—an overcomplete history.

To prevent this, the Axiom of Event Selection restricts the permissible extension to a countable, consistent refinement:

$$e_j \longrightarrow e_{j,1}, e_{j,2}, \dots, e_{j,k}, \quad (2.23)$$

and requires the selection of exactly one representative outcome from each locally admissible family. This keeps E locally finite and maintains a single-valued universe tensor,

$$\mathbf{U}_{n+1} = \mathbf{U}_n + \mathbf{E}_{j,k^*}. \quad (2.24)$$

The axiom thus enforces the same regularity that Martin’s Axiom guarantees in set theory: every countable family of local choices admits a globally consistent selection that preserves the partial order.

Definition 20 (Information Minimality [87, 99]). *Among all admissible extensions of E , the physically admissible history is the one of minimal Kolmogorov complexity:*

$$w_{\min} = \arg \min \{K(w) : w \text{ is an admissible extension of } E\}.$$

Information minimality expresses the logical content of measurement: if additional curvature, oscillation, turning points, or discontinuities had occurred between e_i and e_{i+1} , those features would have generated new events. Since no such events are present in E , any extension that predicts them is inadmissible, and a shorter description exists.

Remark 13. *This principle is purely set-theoretic. No geometry, metric, or differential structure is assumed. Kolmogorov minimality selects the short-*

est admissible description of the recorded distinctions and forbids unobserved structure.

Remark 14. *As the resolution of measurement increases, the admissible extension forms a Cauchy sequence [22] in the space of symbolic descriptions. In the dense limit, its smooth shadow is the unique spline that introduces no new structure between recorded events. Thus the variational calculus is not imposed; it is the continuum limit of Kolmogorov minimality.*

Inadmissibility of Unobserved Structure

Let $E = \{e_0 \prec e_1 \prec \cdots \prec e_n\}$ be the finite set of recorded events produced by a measurement process. By Definition 6, each event corresponds to a distinguishable update of state: a change that crossed a detection threshold and became causally recorded.

Between two successive events e_i and e_{i+1} , no additional events were recorded. This absence is a data constraint: any refinement of the history that introduces detectable structure—curvature, oscillation, turning points, discontinuities, or other distinguishable phenomena—would generate additional events. Since these events do not appear in E , any history that predicts them is logically inconsistent with the observational record.

Definition 21 (Unobserved Structure). **N.B.**—*The idea of unobserved structure echoes the notion of “hidden” or “non-observable” structure that appears in several areas of theoretical computer science and logic, most notably in Scott’s domain theory [137]. There, an extension may contain information that is not reflected in the observable prefix. In the present framework, the analogy is purely conceptual: symbolic refinements that do not correspond to distinguishable events in the ledger do not contribute to the informational state. Only observed distinctions shape the causal record.* \square

Let w be an admissible extension of E (Definition 2.3.3). A symbolic segment of w between e_i and e_{i+1} contains unobserved structure if it encodes

a distinguishable update that is not present in E .

2.7 Correlation and Dependency

In conventional quantum mechanics the word “entanglement” refers to a non-classical dependency among amplitudes: indistinguishable histories are combined before probabilities are assigned. The present framework adopts a similar intuition, but in a purely informational and algebraic form, with no amplitudes and no functional dependencies.

Two events are *uncorrelant* when no *correlant* exists between them. In this case, their transposition commutes with every admissible invariant of the Universe Tensor, and the events may be represented independently. Uncorrelant events are informationally separable: no refinement of the record forces them to be treated jointly.

Two events are *correlant* when they do not commute: exchanging them changes at least one admissible invariant. In this case a correlant exists. A correlant is an informational relation—the minimal structure required when two events cannot be represented independently of one another. Importantly, a correlant does not specify direction or causation: nothing is said about which event precedes, influences, or determines the other. It expresses only that the transposition fails to commute.

Uncorrelant events can become correlated when their light cones merge. Before the merger, each event admits a representation that commutes with the other; no correlant exists, and their histories may be transposed without altering any admissible invariant. After the merger, additional distinctions become available, and the transposition may fail to commute. A correlant then forms, not because one event generates the other, but because the enlarged record no longer permits them to be represented independently.

Dependency relations are stronger still. A dependency asserts that one event is determined by another, as in the functional relationships of the

classical calculus. Such relations describe macro-events in conventional dynamics, where causes generate effects. The present work is not concerned with dependency. Correlation is the weaker structure: non-commutativity under admissible permutation, with no claim of generation or determination.

Thus, “entanglement” in the conventional quantum sense has two informational analogues in this framework. When amplitudes combine as indistinguishable histories, the result is a superposition. When events cannot be transposed without altering admissible invariants, the result is a correlant. Both are consequences of the same principle: distinctions cannot be manufactured retroactively. What differs is the level at which indistinguishability occurs—the discrete record of events or the smooth representation of extremals.

Thought Experiment 2.7.1 (Spooky Action at a Distance [9, 46, 148]). *Consider an uncorrelant $S = \{\mathbf{E}_i, \mathbf{E}_j\}$ of two spatially separated measurement events. By definition, the order of \mathbf{E}_i and \mathbf{E}_j may be permuted without changing any invariant scalar of the universe tensor:*

$$\mathbf{E}_i \mathbf{E}_j = \mathbf{E}_j \mathbf{E}_i. \quad (2.25)$$

When an observer records \mathbf{E}_i , the global ordering is fixed, and the universe tensor is updated accordingly. Because \mathbf{E}_j belongs to the same uncorrelant set, its contribution is now determined consistently with \mathbf{E}_i , even if E_j occurs at a spacelike separation. This manifests as the phenomenon of “spooky action at a distance”—the appearance of instantaneous correlation due to reassociation within the uncorrelant subset.

Thought Experiment 2.7.2 (Hawking Radiation [69, 156]). *Let \mathbf{E}_{in} and \mathbf{E}_{out} denote the pair of particle-creation events near a black hole horizon. These events form an uncorrelant set:*

$$S = \{\mathbf{E}_{in}, \mathbf{E}_{out}\}. \quad (2.26)$$

As long as both remain unmeasured, their contributions may permute freely within the universe tensor, preserving scalar invariants. However, once \mathbf{E}_{out} is measured by an observer at infinity, the ordering is fixed, and \mathbf{E}_{in} is forced to a complementary state inside the horizon. The outward particle appears as Hawking radiation, while the inward partner represents the corresponding loss of information behind the horizon. Thus Hawking radiation is naturally expressed as an uncorrelant whose collapse into correlation occurs asymmetrically across a causal boundary.

Coda: Achilles and the Tortoise

N.B.—For a rich treatment of this paradox, see Hofstadter [76]. □

Zeno's paradox of Achilles and the tortoise [124] is one of the oldest arguments against the possibility of motion. Achilles, swift of foot, gives a tortoise a small head start. Because the tortoise begins ahead, Achilles must first reach the tortoise's initial position. By that time, the tortoise has advanced a little farther; Achilles must then reach that new position, and by the time he arrives, the tortoise has advanced again, and so on without end. Zeno's conclusion is that Achilles can never overtake the tortoise, for he must complete an infinite sequence of tasks to do so.

Formally, one can express the argument in familiar modern notation. Suppose the tortoise begins one unit ahead. Achilles covers half the remaining distance on his first stride, then half of what remains on the next stride, then half again, producing the well-known geometric series

$$1 = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \cdots .$$

More generally, one may express the same identity as

$$1 = \sum_{n=1}^{\infty} \frac{1}{2^n} .$$

Zeno's reasoning is now captured in a single line: if Achilles must perform an infinite number of sub-journeys to reach the tortoise, and if completing infinitely many tasks requires infinite time, then Achilles never arrives.

The mathematics appears to sharpen the paradox. The right-hand side contains infinitely many terms, and yet their sum is finite. An infinite decomposition and a finite limit uneasily coexist. From a purely symbolic viewpoint, Zeno is correct: the path to the finish line can be written as a countable infinity of smaller and smaller segments. Nothing in the algebra forbids infinitely many subdivisions of the interval.

The difficulty lies not in the mathematics, but in the hidden assumption that every subdivision corresponds to a physically meaningful event. Zeno imagines that the runner physically performs each of these infinitesimal sub-paths, as though each term in the series corresponds to an actual step. In reality, the decomposition exists only on paper. It is an artifact of representation, not an element of the physical world.

In the information gauge, motion is not defined by a continuous geometric parameter, but by the accumulation of admissible distinctions—measurable, irreversible updates of state. A notebook of observations does not record symbolic halvings of distance; it records physical events that are detectable by an instrument. Proper time is not the integral of infinitesimal steps, but the count of such admissible distinctions.

Viewed in this light, the identity

$$1 = \sum_{n=1}^{\infty} \frac{1}{2^n}$$

does not imply that Achilles performs infinitely many physical actions. It states only that a continuous model permits infinitely many subdivisions, should one choose to write them down. The infinite chain is a mathematical convenience, not a physical ledger.

The resolution is found in precision. Achilles does not detect every possi-

ble subinterval of his path; no instrument possesses infinite resolving power. His step length, his stride cadence, and the sensor that records his position determine a finite resolution. If the act of stepping advances him by 10^{-2} units, there are at most 100 admissible distinctions in a one-unit race. Even if the instrumentation resolves position to 10^{-6} units, the notebook contains no more than one million recorded distinctions. Once this finite notebook is reconciled, Achilles is at the finish line. The race consumes a finite count of admissible distinctions because the physical process does not instantiate an actual infinity of subevents.

Zeno's paradox relies on treating every symbolic refinement of the interval as physically real. The information gauge rejects that assumption. A measurement records only what can be stably distinguished. Achilles's "infinite" steps are not steps at all; they are possible refinements of a mathematical model. Precision is the gatekeeper. The paradox dissolves when we recall that Achilles's motion is measured, not imagined, and that every measurement has finite resolution. Refinement does not create motion; it reveals it.

Chapter 3

The Calculus of Dynamics

In the previous chapter, motion was described entirely as a sequence of admissible distinctions—a finite notebook of observable updates. No geometry, metric, or continuum was assumed. Refinement revealed additional events, but the history of any physical process remained a countable record that could be reconciled into a globally coherent ledger.

This chapter introduces dynamics in the same spirit. By “dynamics” we do not mean a force law or a geometric trajectory. We mean the rule that selects, from all admissible histories, those that are physically possible. The key observation is that a physical history cannot contain unexplained motion. Any segment of a worldline must be consistent with the measurements that precede and follow it. When a history can be refined without altering its predictions at the recorded events, the refined history contains no additional information. In this sense, the physically admissible refinement is the one that introduces no new distinctions beyond those required by the data.

This principle has a classical name. In the continuum limit, the requirement that refinements add no “hidden motion” is precisely the Euler–Lagrange condition: an admissible trajectory introduces no superfluous curvature beyond that certified by observed events [26, 29, 93]. A trajectory of least informational content is a trajectory of least action, in the classical

sense of Maupertuis, Euler, Lagrange, Hamilton, and their modern successors [37, 48, 61, 67, 92]. In the calculus of dynamics, smooth solutions arise not from geometry but from the demand that no further admissible distinctions can be discovered between measurements. The spline that leaves nothing to correct is the one nature selects.

The remainder of this chapter develops this idea formally. Starting from a finite set of measurements, we construct the weak form of the problem and show that the unique refinement consistent with all observed distinctions is the cubic spline. Its extremality in the continuum reproduces the Euler–Lagrange equations familiar from classical mechanics and field theory. Dynamics are not imposed at the outset; they emerge as the limit in which refinement ceases to yield new information.

Thought Experiment 3.0.1 (Minimizing Variations [29]). **N.B.**—*For a comprehensive treatment of the calculus of variations, see Brenner and Scott [20] and Courant and Hilbert [29].* \square

We consider the functional

$$J[x] = \int_a^b f(t, x(t), \dot{x}(t)) dt,$$

where x is a twice continuously differentiable function with fixed endpoints $x(a) = x_a$ and $x(b) = x_b$. Let $\eta(t)$ be an admissible perturbation with $\eta(a) = \eta(b) = 0$, and define the variation

$$x_\varepsilon(t) = x(t) + \varepsilon \eta(t), \quad \varepsilon \in \mathbb{R}.$$

The directional derivative of J at x in the direction η is

$$\delta J[x; \eta] = \left. \frac{d}{d\varepsilon} J[x_\varepsilon] \right|_{\varepsilon=0} = \left. \frac{d}{d\varepsilon} \int_a^b f(t, x_\varepsilon(t), \dot{x}_\varepsilon(t)) dt \right|_{\varepsilon=0}.$$

Since the integration limits do not depend on ε , the derivative may be moved

inside:

$$\delta J[x; \eta] = \int_a^b \left. \frac{\partial}{\partial \varepsilon} f(t, x_\varepsilon(t), \dot{x}_\varepsilon(t)) \right|_{\varepsilon=0} dt.$$

By the chain rule,

$$\frac{\partial}{\partial \varepsilon} f(t, x_\varepsilon(t), \dot{x}_\varepsilon(t)) = f_x(t, x(t), \dot{x}(t)) \eta(t) + f_{\dot{x}}(t, x(t), \dot{x}(t)) \dot{\eta}(t).$$

Thus

$$\delta J[x; \eta] = \int_a^b \left(f_x(t, x, \dot{x}) \eta(t) + f_{\dot{x}}(t, x, \dot{x}) \dot{\eta}(t) \right) dt.$$

Integrate the second term by parts:

$$\int_a^b f_{\dot{x}} \dot{\eta} dt = [f_{\dot{x}} \eta]_a^b - \int_a^b \frac{d}{dt} (f_{\dot{x}}) \eta(t) dt.$$

Because $\eta(a) = \eta(b) = 0$, the boundary term vanishes. Therefore

$$\delta J[x; \eta] = \int_a^b \left(f_x - \frac{d}{dt} f_{\dot{x}} \right) \eta(t) dt.$$

If x is a stationary point of J , then $\delta J[x; \eta] = 0$ for all admissible η . The fundamental lemma of the calculus of variations implies

$$f_x(t, x, \dot{x}) - \frac{d}{dt} f_{\dot{x}}(t, x, \dot{x}) = 0,$$

for all $t \in (a, b)$. This is the Euler–Lagrange equation, more commonly represented as

$$\frac{\partial f}{\partial x} = \frac{d}{dt} \frac{\partial f}{\partial \dot{x}}. \quad (3.1)$$

This derivation demonstrates that the Euler–Lagrange equation selects the trajectory with no first-order change under admissible perturbations. No hidden motion can be inserted without altering the notebook. The path is stationary in its informational curvature.

3.1 Emergent Dynamics

In the discrete setting, the Causal Universe Tensor assigns a finite informational weight to every admissible history. Refinement increases this weight only when new distinctions are recorded. Any replacement of an admissible history by one containing additional, unobserved structure violates Axiom 7. Consequently, dynamics is not an independent physical postulate. It is the unique continuous shadow of informational extremality: the smooth curve is simply the history for which no further admissible distinctions can be revealed.

In the discrete domain, *anchor points* are the only places where the universe has committed to a specific value. Between anchors the record is silent: the data permit many possible continuations, but most would introduce unobserved structure. Any admissible configuration must therefore agree at the anchor points and remain free of additional distinguishable features in between. The role of the anchors is not geometric; it is informational. They fix the admissible boundary data against which all variations are tested. A candidate variation that disagrees at an anchor is rejected immediately, because it contradicts an established event. A variation that agrees at the anchors but inserts additional oscillation, curvature, or “hidden motion” is rejected by Axiom 7, because those features would have produced additional anchors that do not appear in the record.

Definition 22 (Anchor Points [36]). *A finite set of anchor points is the collection of measured events at which admissible configurations must agree. Two candidate histories ψ and ϕ are said to share the same anchors if they record identical distinguishable values at those events. Axiom 3 requires that any refinement of a history preserve agreement on the anchors: no admissible configuration may contradict an observed event.*

In the discrete setting, reciprocity arises from a simple counting fact. A refinement of ψ by a test configuration ϕ is admissible only when the

resulting history contains no additional distinguishable events. If ϕ were to introduce extra curvature, oscillation, or “hidden motion,” the refinement would increase the causal count and violate Axiom 7. The reciprocity pairing $\psi^* \mathcal{L} \phi$ measures this change: it evaluates whether ϕ is informationally neutral relative to ψ .

Crucially, the dual ψ^* is not a geometric adjoint; it is the reflection of ψ in the informational algebra. It answers the question: *If ψ is perturbed by ϕ , does the universe record new distinguishable structure?* If the reciprocity pairing vanishes for all admissible ϕ that share the anchors, then ψ is extremal. Any remaining variation would imply new recorded events, and therefore be inadmissible.

Definition 23 (Reciprocity Map). ***N.B.**—In geometric settings equipped with a metric or inner product, the reciprocity map reduces to the familiar adjoint or complex conjugate, and the operation $\psi \mapsto \psi^*$ is often interpreted as a covariant or contravariant dual. No such geometric structure is assumed here. The reciprocity dual is defined purely informationally, as the configuration that symmetrizes the causal pairing. It should not be confused with metric adjoints that appear in geometric representation theory, such as the Dirac adjoint of a spinor or the dual of a Weyl field. Those constructions depend on Lorentz symmetry, Clifford algebras, and an invariant bilinear form; none of these structures are present at the informational level.* \square

Let ψ be an admissible configuration and let ϕ be a test variation that agrees with ψ at the anchor points. The reciprocity map is the linear evaluation

$$\langle \psi, \phi \rangle_{\mathcal{L}} := \psi^* \mathcal{L} \phi,$$

where \mathcal{L} counts distinguishable causal increments. A configuration χ is called a reciprocity dual of ψ if it satisfies

$$\langle \psi, \phi \rangle_{\mathcal{L}} = \langle \phi, \chi \rangle_{\mathcal{L}} \quad \text{for all test variations } \phi.$$

When it exists, such a χ is denoted by ψ^* . The reciprocity dual encodes the informational response of ψ to an infinitesimal variation ϕ without assuming any differential structure.

Proposition 2 (The Uniqueness of the Reciprocity Dual). *Assume the causal pairing $\langle \cdot, \cdot \rangle_{\mathcal{L}}$ is nondegenerate in the second slot: if*

$$\langle \phi, \chi \rangle_{\mathcal{L}} = 0 \quad \text{for all test variations } \phi,$$

then χ is the trivial (null) configuration. If χ_1 and χ_2 are both reciprocity duals of the same configuration ψ , then $\chi_1 = \chi_2$. In particular, whenever a reciprocity dual exists, it is unique.

Proof (Sketch). Let χ_1 and χ_2 be reciprocity duals of ψ . By definition,

$$\langle \psi, \phi \rangle_{\mathcal{L}} = \langle \phi, \chi_1 \rangle_{\mathcal{L}} = \langle \phi, \chi_2 \rangle_{\mathcal{L}} \quad \text{for all test variations } \phi.$$

Subtracting the two expressions gives

$$\langle \phi, \chi_1 - \chi_2 \rangle_{\mathcal{L}} = 0 \quad \text{for all } \phi.$$

By nondegeneracy in the second slot, this implies $\chi_1 - \chi_2$ is the null configuration, hence $\chi_1 = \chi_2$. Thus any reciprocity dual, if it exists, is unique. \square

A full proof is provided in Appendix A.2.

In the continuum shadow, the reciprocity pairing becomes the usual weak inner product of variational calculus [20, 49]. Integration by parts moves the variation from ψ onto the test functions, producing natural boundary terms determined by the anchors. The condition

$$\langle \psi, \phi \rangle_{\mathcal{L}} = \langle \phi, \psi \rangle_{\mathcal{L}}$$

is then the classical reciprocity of the Euler–Lagrange operator: the dynamics are self-adjoint under the informational measure. This equality holds not

because symmetry is assumed, but because any antisymmetric contribution would encode unrecorded distinctions and be eliminated by Axiom 7.

3.1.1 Weak Formulation on Space–Time

Let ψ be an admissible configuration consistent with a fixed set of event anchors, and let ϕ be any test configuration that agrees with ψ at those anchors. Replacing ψ by ϕ is permissible only if it does not reduce causal consistency. In the discrete algebra this means that ψ introduces no superfluous refinements relative to ϕ ; any additional curvature, oscillation, or “hidden motion” would imply unrecorded events and thus be inadmissible.

In the dense limit of refinement, this constraint appears as a weak relation

$$\psi^* \mathcal{L} \psi \leq \psi^* \mathcal{L} \phi, \quad (3.2)$$

where \mathcal{L} is the informational count of distinguishable increments, and ψ^* denotes its reciprocity dual. The weak inequality asserts that ψ is extremal among all admissible perturbations ϕ . No differential operators are assumed: the weak form arises because refinement limits the class of permissible discrete variations.

Completing this refinement yields the continuous counterpart of (3.2). Integration by parts shifts variations from ψ onto the test functions, producing natural boundary conditions and a weak Euler–Lagrange statement. The continuum calculus therefore does not describe an independently assumed physical law; it is the smooth completion of informational minimality on the discrete domain.

3.1.2 Reciprocity and the Adjoint Map

The weak extremality relation (3.2) compares an admissible configuration ψ against a test configuration ϕ that shares the same event anchors. In the discrete domain, replacing ψ by ϕ means refining the event record: only those

local changes that introduce new, distinguishable curvature would alter the admissible history. Any such change must correspond to additional recorded events; if none are present, the refinement is informationally neutral. Thus ϕ is an admissible variation of ψ precisely when it agrees at the anchors and introduces no distinctions beyond those already encoded in ψ . The weak extremality condition (3.2) is the continuous shadow of this discrete refinement rule.

The weak comparison between ψ and ϕ admits a natural dual representation. For any admissible configuration ψ , there exists a *reciprocity map* ψ^* such that the informational pairing

$$\psi^* \mathcal{L} \phi \tag{3.3}$$

measures the change in distinguishability that would result from locally replacing ψ by ϕ between the anchors. Intuitively, ψ^* captures the “shadow” of ψ when viewed from the perspective of informational minimality: components of ϕ that would introduce new, unrecorded distinctions are suppressed by the adjoint action, while components that are informationally neutral remain. In the dense limit, this pairing becomes the standard weak inner product of variational calculus.

Because admissible configurations cannot contain hidden structure, the reciprocity map annihilates variations that are invisible at the event anchors. If ϕ and ψ agree at the anchors and differ only by an undetectable perturbation, then refining the event record yields no new distinctions, and the informational pairing remains unchanged:

$$\psi^* \mathcal{L} \phi = \psi^* \mathcal{L} \psi.$$

This equality is precisely the weak relation (3.2). In this sense, ψ^* enforces closure: the extremal configuration carries no latent curvature that would be revealed by further refinement.

The “variation” of ψ is therefore not a differential operation but a refinement of the causal record consistent with the event anchors. The reciprocity map acts as the dual constraint, suppressing any component of that refinement which would introduce unrecorded distinctions. Taken together, admissible refinements and their reciprocity dual generate the weak Euler–Lagrange structure entirely within the discrete domain, without assuming differentiability or a continuum of states.

In this way, the reciprocity map ensures that any admissible refinement of ψ corresponds to an interpolant $f(\psi)$ that introduces no new distinguishable structure. As refinement becomes dense, all such interpolants converge to the same smooth closure Ψ . Since the event record defines a finite labeled partition of the causal domain, Ψ preserves anchor order and is injective on each partition element. Its inverse Ψ^{-1} therefore recovers exactly the original discrete record:

$$f(\psi) \longrightarrow \Psi^{-1}. \quad (3.4)$$

Thus the interpolant and its smooth limit are informationally equivalent representations of the same causal structure.

3.1.3 Dense Limit and Euler–Lagrange Closure

In the present framework no differentiability is assumed. The weak extremality relation (3.2) is defined entirely in the discrete domain, where each term counts distinguishable causal increments. A “variation” of ψ is therefore not a differential operator but a refinement of the event record that leaves the anchors unchanged.

In the discrete domain, such refinements appear as finite differences: each admissible update replaces a segment of the causal history by one with strictly greater resolution. Because informational minimality forbids unobserved curvature, every admissible refinement corresponds to a piecewise-linear or piecewise-polynomial interpolant that agrees with ψ on the anchors and in-

roduces no new distinguishable structure. As refinements become arbitrarily dense, the finite differences form a Cauchy sequence in the space of admissible interpolants, and their limit is the unique smooth closure Ψ established in the previous subsection.

Applying the reciprocity pairing to successive refinements yields the discrete extremality condition: no admissible finite difference can reduce the informational measure \mathcal{L} . In the dense limit, the weak relation (3.2) becomes the standard variational identity of Euler–Lagrange calculus, obtained entirely from finite differences. The weak derivative enters only as the completion of refinement; it is not assumed *a priori*.

When the causal grid is refined, informational minimality forces cubic continuity at each event anchor: jumps in slope or curvature would constitute new observable events and are therefore inadmissible. In the dense limit, the discrete extremal coincides with the classical Euler–Lagrange closure. This structure is summarized in the following proposition.

Proposition 3 (The Spline Condition of Information). *Let ψ be an admissible configuration with smooth closure Ψ . If no admissible refinement reduces the informational measure \mathcal{L} , then Ψ is C^2 and satisfies*

$$\Psi^{(4)} = 0. \tag{3.5}$$

Proof (Sketch). Between anchors, Ψ must be polynomial, since any additional inflection would imply unrecorded structure. Polynomials of degree greater than three contain latent turning points and are therefore excluded. Hence each segment is cubic. At the anchors, the interpolants must glue with C^2 continuity: jumps in slope or curvature would constitute new observable events. As the grid of anchors is refined, the third derivative Ψ''' must be constant on every shrinking interval. In the dense limit that interval has zero measure, so Ψ''' is constant everywhere. A constant third derivative implies $\Psi^{(4)} = 0$. Thus the smooth closure of any informationally extremal

configuration satisfies the Euler–Lagrange condition. \square

A full proof is provided in Appendix ??.

Proposition 3 shows that the Euler–Lagrange equation is not postulated. It is the continuous shadow of discrete informational extremality. Finite differences do not approximate the differential equation; they *generate* it. The unique admissible smooth representative is cubic on each partition element, C^2 at the event anchors, and satisfies $\Psi^{(4)} = 0$ everywhere. Smooth calculus appears solely as the completion of refinement in the discrete causal record.

Phenomenon 3.1.1 (Repeatability of Invisible Motion [6]). *Consider two independent observers, A and B , who record the motion of a particle between the same event anchors $x_i \prec x_{i+1}$. Each observer has finite resolution: any acceleration or inflection large enough to be distinguishable produces a new event. Both refine their instruments until no further events are detected on the interval.*

If hidden curvature existed between the anchors, further refinement would create additional distinguishable records. The absence of such records forces each observer to recover the same polynomial of minimal degree. Thus both obtain a cubic patch on the interval.

Now let A and B exchange data and perform a joint refinement on a finer grid. Any disagreement in value, slope, or bending moment at a shared anchor would itself generate an observable event. To avoid contradiction, the cubic patches must glue together with continuous U , U' , and U'' . In the dense refinement limit, the piecewise constant third derivative converges to a continuous function whose integral vanishes on every shrinking interval, yielding

$$U^{(4)} = 0.$$

Thus repeatability demands the Euler–Lagrange closure: if two observers can refine their measurements indefinitely without producing new events,

their reconstructions must converge to the same cubic extremal. Smooth dynamics are therefore the unique histories that leave no trace.

3.2 The Law of Spline Sufficiency

The preceding analysis shows that every admissible refinement of the event record corresponds to a piecewise-cubic interpolant that preserves the event anchors and introduces no new distinguishable structure. In the dense limit, these interpolants converge to a unique smooth closure Ψ that is \mathcal{C}^2 and satisfies $\Psi^{(4)} = 0$. The discrete causal record and its smooth completion are therefore informationally equivalent representations of the same history.

Law 1 (The Law of Spline Sufficiency). *Let ψ be any finite, non-contradictory record of admissible events. There exists a unique continuous completion Ψ such that:*

1. Ψ agrees with ψ at every event anchor,
2. Ψ is piecewise cubic and \mathcal{C}^2 on its domain,
3. Ψ introduces no new distinguishable structure beyond ψ , and
4. Ψ satisfies the Euler–Lagrange closure $\Psi^{(4)} = 0$.

The cubic spline is therefore sufficient to represent all admissible distinctions in the data: no higher-order model encodes additional information available to measurement.

The Law of Spline Sufficiency justifies the use of Galerkin methods [56] in this framework. By choosing the functional

$$\mathcal{J}[\Psi] = \int (\Psi'')^2 dx \tag{3.6}$$

as a measure of curvature, the Galerkin extremal selects the simplest admissible interpolant consistent with the event anchors. Because every sequence of

admissible refinements converges to the unique \mathcal{C}^2 cubic closure, the Galerkin solution coincides with the informationally extremal configuration. No additional degrees of freedom are required.

In this sense, spline sufficiency provides the logical bridge between discrete measurement and continuous dynamics:

$$\text{discrete measurement} \xrightarrow{\text{spline sufficiency}} \Psi \xrightarrow{\text{closure}} \Psi^{(4)} = 0.$$

Finite differences do not approximate the Euler–Lagrange equation; they *generate* it. Smooth calculus enters only as the completion of refinement in the causal record, not as an assumed geometric primitive.

3.2.1 Minimality

The Law of Spline Sufficiency guarantees that admissible histories are those that interpolate between refinements with minimal curvature. This law does not merely describe smoothness; it constrains how influence can propagate through the ledger. A localized refinement cannot remain localized under extension. Its effect must be distributed across every admissible continuation consistent with global coherence.

As the causal record is extended outward from an event, the frontier of admissible continuations grows combinatorially. The same finite refinement budget must be shared across an increasing number of admissible degrees of freedom. The effect of any single refinement therefore weakens with informational separation, not because of dissipation, but because of accounting.

While any minimality can be enforced, measurement tends to prefer minimization in the \mathcal{L}^2 -norm.

Phenomenon 3.2.1 (The Inverse Square Effect). ***Statement.** The influence of a refinement event decreases as the inverse square of the informational separation. This scaling is not postulated; it is forced by the geometry of admissible splines.*

Mechanism. *By the Law of Spline Sufficiency, admissible continuations of the causal ledger are the minimal curvature interpolants consistent with boundary anchors. A single refinement event acts as a localized constraint on the spline. As the distance from that constraint increases, the number of distinct admissible continuations grows with the surface measure of the surrounding causal sphere.*

In three admissible dimensions, this measure scales as $4\pi r^2$. The influence of a fixed refinement budget must therefore be distributed across a quadratically growing frontier. The admissible effect per refinement falls as

$$I(r) \propto \frac{1}{r^2}.$$

Interpretation. *This is not a force law. It is a bookkeeping law. The ledger cannot assign a fixed refinement cost to an expanding set of admissible continuations without diluting its effect.*

The inverse-square behavior of gravitation, radiation, and flux is therefore the smooth shadow of the combinatorial growth of admissible splines.

3.3 Galerkin Methods

N.B.—This argument applies the Law of Spline Sufficiency. We do not assume that Euler–Lagrange dynamics exist *a priori*. Rather, we show that if the data admit a smooth completion, then a cubic spline exists which reproduces the Euler–Lagrange solution to arbitrary accuracy. In this sense, observing a spline is sufficient to infer Euler–Lagrange dynamics: the differential equation models the behavior only insofar as the data allow it, and no additional geometric or differentiable structure is assumed. \square

The Law of Spline Sufficiency establishes that cubic splines contain all admissible distinguishable structure. In this section we assume the existence of a smooth Euler–Lagrange solution and show that a Galerkin projection

onto a spline basis produces a sequence of spline functions that converges to it. This suffices to justify the use of splines as the representatives of continuous dynamics: if Euler–Lagrange motion exists, Galerkin refinement will recover it to arbitrary accuracy.

3.3.1 Galerkin Projection onto a Spline Basis

Let Ψ be the smooth solution to an Euler–Lagrange boundary value problem. Choose a finite spline basis $\{\varphi_k\}$ that satisfies the boundary constraints and let

$$\Psi_n(x) = \sum_{k=1}^n a_k \varphi_k(x)$$

be the Galerkin projection of Ψ onto this space. The coefficients a_k are chosen so that the residual of the Euler–Lagrange equation is orthogonal to the spline basis:

$$\int \Psi_n''(x) \varphi_k''(x) dx = \int \Psi''(x) \varphi_k''(x) dx, \quad k = 1, \dots, n. \quad (3.7)$$

This is the standard spline Galerkin formulation [26, 20]: the weak form enforces the Euler–Lagrange condition in the finite dimensional subspace spanned by the splines.

Solving (3.7) yields a unique spline Ψ_n that agrees with the smooth solution at all knot points and is \mathcal{C}^2 on the domain. No higher-order degrees of freedom are necessary; the curvature functional ensures that splines are the minimal weak extremals.

3.3.2 Convergence of the Galerkin Sequence

By the Weierstrass Approximation Theorem, cubic splines form a dense subspace of continuous functions on a compact interval. As the mesh is refined and more basis functions are added, the sequence $\{\Psi_n\}$ converges uniformly

to Ψ :

$$\Psi_n \xrightarrow[n \rightarrow \infty]{} \Psi.$$

Because the Euler–Lagrange operator is continuous in the weak topology, convergence of Ψ_n implies convergence of all weak derivatives:

$$\Psi_n'' \xrightarrow[n \rightarrow \infty]{} \Psi''.$$

Thus the Galerkin sequence yields arbitrarily good spline approximations of the Euler–Lagrange solution. In particular, Ψ_n satisfies

$$\Psi_n^{(4)} = 0$$

on each spline element, up to a boundary residual that vanishes as the mesh is refined.

Corollary 1. *If a smooth Euler–Lagrange solution Ψ exists, a sequence of cubic splines $\{\Psi_n\}$ constructed by Galerkin projection converges uniformly to Ψ . Since cubic splines represent all admissible distinguishable structure, observing a spline solution is sufficient to infer the underlying Euler–Lagrange dynamics.*

In summary:

$$\Psi \xrightarrow{\text{Galerkin projection}} \Psi_n \xrightarrow[n \rightarrow \infty]{\text{Weierstrass}} \Psi,$$

so splines not only represent all admissible distinctions, but converge to the unique extremal of the Euler–Lagrange equation whenever one exists. The Galerkin method therefore completes the argument of spline sufficiency in the continuum: if continuous dynamics exist, spline solutions will recover them to arbitrary accuracy.

The Galerkin refinement therefore recovers smooth calculus without assuming infinitesimal increments or geometric primitives. The classical para-

dox of the fluxion may now be revisited in this light.

Phenomenon 3.3.1 (Fluxions [10, 113]). **N.B.**—*The classical paradox of the fluxion treats an infinitesimal dt as a quantity that is neither zero nor nonzero. In the present framework, the limit is defined without invoking infinitesimals: smooth structure appears only as the unique completion of finite distinctions.* \square

In the 18th century, Bishop Berkeley criticized Newton’s calculus of fluxions (\dot{x}, \dot{y}) for relying on quantities that vanish in one step of a proof and are treated as nonzero in the preceding step. If \dot{x} and \dot{y} are the ghost-like “increments” of position, the question arises: How can a finite, observable change emerge from the vanishing difference of infinitesimal quantities?

In the causal accounting used here, this is not a paradox of quantity but a limitation of informational resolution. The fluxion

$$\dot{x} = \frac{\Delta x}{\Delta t}$$

is a ratio of two sequentially recorded distinctions: the number of spatial ticks Δx versus the number of temporal ticks Δt between two anchors. Both are finite, integer-valued measurements.

The classical paradox appears only when $\Delta t \rightarrow 0$ is interpreted as a transition through a nonphysical intermediate state. In the present framework, no such state is required. The smooth completion Ψ constructed in the dense limit satisfies $\Psi^{(4)} = 0$ and is the unique curvature-free extension of the data. As the anchor spacing shrinks, the ratio $\frac{\Delta x}{\Delta t}$ converges to the unique \mathcal{C}^2 slope Ψ' of the cubic interpolant determined by the neighboring anchors.

No ghost-like infinitesimal is invoked. The derivative is the continuous shadow of finite bookkeeping: the single value required to prevent the appearance of new, unrecorded events as resolution increases. Smooth calculus arises not by manipulating vanished quantities, but as the unique function consistent with every refinement of the observable record.

3.3.3 The Physical Impossibility of Infinite Refinement

A law of spline necessity *would* describe the continuous limit of an ideal refinement process much like the law of spline sufficiency. In such a limit, where arbitrarily fine distinguishable refinements are permitted, the unique smooth closure compatible with informational minimality would necessarily coincide with a cubic spline satisfying $\Psi^{(4)} = 0$ between all anchors. This behavior would characterize the limiting object toward which all admissible refinements converge.

However, this description is inherently conditional. The existence of such a law requires access to refinements at arbitrarily small scales. In the informational setting developed here, no such refinement process exists: every record admits only finitely many distinguishable refinements. As a consequence, the continuum limit in which an exact spline law *would* hold is never attainable. The law does not fail; rather, it is not a law of the finite world.

N.B.—The idea of a spline necessity law is meaningful only as a limiting construct. It does not apply to any finite record because no observational process can instantiate the infinite refinement depth the law presupposes (Axiom 6). \square

This observation motivates an approximate interpretation. Although an exact law cannot hold, the Galerkin convergence results of Section ?? imply that finite-dimensional closures can be made arbitrarily close to the ideal spline closure. Thus, while a spline necessity law describes an unattainable limit, its behavior is still relevant: finite informational models approach that limit as their resolution increases. The continuum spline is therefore best understood as the *attractor* of refinement-compatible approximations, not as a law governing finite observational structure.

Definition 24 (Attractor [103]). *An attractor is a set of configurations toward which the admissible states of a system asymptotically converge under iteration of the update rule. Once the refinement enters the neighborhood of the attractor, subsequent refinements remain confined to it. The attractor*

represents the stable informational pattern that balances the system's internal stress and the constraints of the refinement process.

3.3.4 Indistinguishability of Approximate and Ideal Spline Closures

A law of spline necessity would characterize the exact continuous limit of an ideal refinement process. In practice, however, only approximate spline closures exist, obtained through refinement-compatible approximations such as Galerkin methods. This raises a natural question: could any measurement distinguish between an approximate closure and the ideal spline attractor it converges toward?

The answer is no. Under the axioms of event selection, refinement compatibility, and informational minimality, no admissible measurement can separate the two. Any measurement capable of distinguishing an approximate spline from the ideal one would require detecting differences at scales finer than the minimum resolvable distinction allowed by the record. Such a measurement would necessarily violate the axioms by introducing new refinements below the Planck scale.

N.B.—There exists no admissible observational procedure, consistent with the axioms of measurement, that can differentiate between the approximate spline obtained at a finite refinement scale and the ideal spline that would appear in the continuum limit. Any attempt to do so requires forbidden refinements and is therefore inadmissible. \square

Let (Ψ_N) be a sequence of refinement-compatible approximations converging toward an ideal spline Ψ in the sense of Section ?? . For any fixed resolution scale permitted by the record, there exists N such that

$$\|\Psi_N - \Psi\| < \delta,$$

where δ is the smallest distinguishable refinement allowed by the axioms.

Because no measurement can detect variation smaller than δ , the outputs of Ψ_N and Ψ are observationally identical. To distinguish them would require a measurement refining the domain below δ , which the axioms forbid.

Definition 25 (Observational Indistinguishability [115]). *A finite-dimensional closure Ψ_N is observationally indistinguishable from the ideal spline closure Ψ if, for the minimum refinement scale δ of the record,*

$$|\Psi_N(x) - \Psi(x)| < \delta \quad \text{for all admissible measurement points } x.$$

No admissible measurement can detect any discrepancy of magnitude less than δ .

3.3.5 Indistinguishability of Infinite Refinement

Axiom 1 states that every measurement produces a symbol from a finite or countable alphabet and that all refinements are bounded below by a minimum distinguishable scale $\delta > 0$. A measurement record is therefore a finite string over an alphabet whose effective base is determined by the refinement scale. In this setting, the pigeonhole principle implies that only finitely many distinct measurement outcomes are possible at resolution δ .

Let Ψ be an ideal closure that would be obtained in an infinite-refinement limit, and let Ψ_δ be any finite-resolution approximation consistent with the refinement scale δ . If Ψ and Ψ_δ differ only on sub- δ scales, then no admissible measurement can distinguish them. Their projections into the measurement alphabet coincide, and therefore they produce the same finite string of observations.

Proposition 4 (Pigeonhole Indistinguishability of Infinite Refinement [41, 68]). *Let Σ_δ be the finite set of symbols distinguishable at refinement scale δ , and let \mathcal{M} denote the measurement map*

$$\mathcal{M} : \{\text{closures}\} \rightarrow \Sigma_\delta^*.$$

If two closures Ψ and Φ differ only at scales smaller than δ , then

$$\mathcal{M}(\Psi) = \mathcal{M}(\Phi).$$

In particular, any infinitely refined closure is observationally indistinguishable from a sufficiently refined finite approximation.

Proof (Sketch). This result follows directly from the pigeonhole principle. A finite measurement alphabet cannot encode distinctions below the minimal refinement scale δ . Once two closures agree on all δ -sized cells, no admissible measurement can produce different records. Infinite refinement produces no new distinguishable outcomes. \square

A full proof is provided in Appendix A.3.

3.3.6 Discrete Refinement

Phenomenon 3.3.2 (The Moire Effect). *When two admissible ledgers defined on slightly different refinement lattices are reconciled, coherent and incoherent regions appear at macroscopic scale. These large scale beats are the smooth shadow of high frequency incompatibility between observer frames.*

The visible pattern is not a property of either ledger alone, but the structure required to preserve global consistency under their interaction.

Thus an infinitely refined object is operationally equivalent to a finite closure at the resolution permitted by the axioms. Infinite refinement is a mathematical limit, not an observable phenomenon. This prepares the way for the Law of Discrete Spline Necessity, which identifies the unique closure that saturates all distinguishable information at scale δ .

Phenomenon 3.3.3 (The Quicksand Effect [8, 17]). **N.B.**—*In a continuous fluid, buoyancy is described by Archimedes' principle [3]: an immersed body floats when the upward force from displaced fluid balances its weight [8]. Bonn*

et al. [17] show that quicksand, though a granular suspension rather than a true fluid, exhibits a nearby buoyant behavior: objects settle only to a finite depth and then float, reaching an equilibrium set by density matching, yield stress, and local fluidization. The macroscopic effect resembles (and, to a certain coarseness of refinement, is modeled by) Archimedes' principle, even though its microscopic origin is entirely different. These physical observations serve only as an analogy for the informational phenomenon described here; they do not constrain the model. They illustrate how a finite set of admissible states may appear, in the smooth limit, as a buoyant equilibrium. \square

N.B.—The phenomenon described here concerns the irreversible, informational component of fluid mechanics: the resistance to refinement below the minimum distinguishable scale δ . It is not a complete account of physical viscosity, which depends on a finite third parameter Θ (see Coda: Navier–Stokes as a Finite Third Parameter, Chapter 3) and requires an independent kinematic assumption relating shear stress to velocity gradients. The informational viscosity Ψ_δ treated here reflects only the constraints of Causal Order and informational Minimality; it captures the coarse, irreducible structure that remains when all sub- δ refinements are suppressed. \square

N.B.—A person floats on quicksand, rather than sinks [17] \square

Consider an agent E attempting to move through a medium governed solely by distinguishability. Before contact, the mathematical continuum admits an infinite family of smooth paths Φ_i , distinguished by arbitrarily small variations in curvature.

Once E enters the medium, the informational constraints become active. By Axiom 6, there exists a minimum distinguishable scale δ . Any displacement smaller than δ fails to generate a new event. The continuum therefore collapses to a finite chain of δ -compatible anchors,

$$\Psi_\delta = \{x_1, \dots, x_N\},$$

representing all positions that can be observationally distinguished.

The medium exhibits an informational viscosity: any attempted motion that introduces sub- δ curvature is resisted and cancelled, keeping E pinned to the nearest admissible anchor. Only when the displacement exceeds the refinement threshold does E transition from x_k to x_{k+1} .

By Proposition 4, the infinite microscopic variations beneath the surface collapse into the finite observational buckets of Ψ_δ . Informational minimality (Axiom 3) then forces the unique discrete closure consistent with the anchors and containing no unrecorded structure: the discrete spline Ψ_δ .

This is the viscosity of quicksand: the resistance to refinement below the minimum distinguishable scale δ . Any attempted motion that fails to produce a new admissible distinction is suppressed, and the system remains at the nearest anchor in Ψ_δ . In the smooth shadow, this appears as the buoyant or viscous equilibrium observed by Bonn and others, where a person floats because further descent would require the granular medium to rearrange at scales smaller than the yield threshold of individual particles of sand. Physically, the grains simply stop moving; informationally, no additional distinctions can be recorded. The collapse of the infinitely many ideal paths Φ_i into the single admissible sequence Ψ_δ is therefore mirrored by the granular equilibrium: motion ceases not because of any continuous force law, but because neither the sand nor the informational model permits sub- δ refinements.

Thus the distinction between the approximate and ideal spline closures is purely mathematical. No experiment, sensor, or observational extension can reveal a difference between them without violating the Axioms of Measurement. The ideal spline belongs to the continuum limit; the approximate spline belongs to the finite informational world. Observationally, however, the two coincide to the highest permissible resolution.

3.4 The Law of Discrete Spline Necessity

Because the refinement depth of any admissible record is finite, the continuum limit in which an exact spline necessity law would hold can never be reached. Nevertheless, the refinement axioms determine a unique finite-resolution object that plays the role of a spline within the informational world. This discrete closure is the actual law governing all admissible completions of a finite record.

Law 2 (The Law of Discrete Spline Necessity). *Let ψ be any finite, non-contradictory record with minimum refinement scale δ , guaranteed by the axioms of measurement. Then there exists a unique finite-resolution function Ψ_δ satisfying:*

(1) Ψ_δ agrees with ψ at every anchor event and introduces no refinements smaller than δ .

(2) On each interval between anchors, Ψ_δ is the minimal-curvature function permitted by the refinement grid of scale δ . In particular, Ψ_δ is represented by a cubic polynomial on each discrete cell of size δ , with continuity of slope and curvature enforced at all interior junctions.

(3) Any alternative function Φ that agrees with ψ at the anchors and differs from Ψ_δ on any discrete cell either

1. introduces additional distinguishable structure below scale δ (violating refinement compatibility and the Planck condition), or

2. fails to maintain global causal consistency across cell boundaries.

(4) As δ decreases along any refinement-compatible sequence, the discrete closures satisfy

$$\Psi_\delta \longrightarrow \Psi$$

where Ψ is the ideal spline attractor described in Section ???. This convergence is monotone in the sense that each refinement preserves and sharpens all previously admissible distinctions.

Thus every finite informational record admits a unique discrete closure Ψ_δ , which is the minimal, globally coherent, refinement-compatible representation of that record at its permitted resolution. This is the informational law governing all realizable completions.

N.B.—This law is exact. Unlike the continuum spline necessity law, which would require infinite refinement and therefore cannot apply to finite records, the Law of Discrete Spline Necessity governs all observationally realizable completions. Continuous splines appear only as limiting attractors. The discrete closure Ψ_δ is the true object selected by the axioms. \square

This law establishes the discrete analogue of curvature minimality, differentiability, and weak-form transport without invoking limits. All smooth structures used in physics arise from the asymptotic behavior of Ψ_δ under refinement but are never instantiated exactly. The discrete closure is the only object compatible with the axioms at finite resolution.

3.4.1 The Indistinguishability of Discrete and Continuous Spline Closures

Axiom 1 asserts that all measurements produce finitely many distinguishable outcomes and that every admissible refinement has a minimum resolvable scale $\delta > 0$. No observational process may introduce refinements smaller than δ without violating the axiom. As a consequence, the refinement process terminates at a finite resolution, and the most refined discrete closure Ψ_δ permitted by the record is observationally maximal.

If an ideal continuum limit were accessible, the refinement process would continue indefinitely and converge to a smooth cubic spline Ψ satisfying the limiting minimality condition $\Psi^{(4)} = 0$. However, the continuum limit requires refinements at scales below δ , and therefore cannot be realized by any admissible sequence of measurements. The continuous spline Ψ is a mathematical attractor, not an observable object.

N.B.—By Axiom 1, the most refined discrete spline Ψ_δ is *observationally indistinguishable* from the continuous spline attractor Ψ . No admissible measurement can detect any discrepancy between the two, because doing so would require refinements smaller than δ , which the axioms forbid. \square

Formally, if (Ψ_N) is any refinement-compatible sequence converging to Ψ , then for sufficiently large N ,

$$|\Psi_N(x) - \Psi(x)| < \delta \quad \text{for all admissible measurement points } x.$$

Therefore Ψ_N and Ψ produce identical observational outcomes.

This establishes that the continuous spline arises only as a limiting concept, while the discrete closure Ψ_δ is the unique physically realizable object. Axiom 1 identifies these two as observationally equivalent: the discrete spline is as refined as the informational world can ever be.

3.4.2 The Necessity of Approximation

The preceding sections establish a structural asymmetry in the informational framework. On the one hand, the continuum spline appears as the unique limiting object that a refinement process *would* select if infinite refinement were possible. On the other hand, Axiom 1 forbids refinements below a minimum distinguishable scale $\delta > 0$. The refinement sequence therefore terminates at a finite stage, and no observational process can approach the continuum limit beyond this final resolution.

Phenomenon 3.4.1 (The Olbers Effect). *An infinitely refined ledger would admit infinitely many luminous events. The observed darkness of the night sky demonstrates that the causal record is finite.*

The absence of uniform brightness is the direct observational proof that the informational capacity of admissible history is bounded.

This tension forces a fundamental conclusion: approximation is not a methodological choice but a structural necessity. Every admissible represen-

tation of a finite record must be an approximation to a limit that cannot be realized. The continuous spline is an ideal boundary point of the refinement process, never an attainable object within the informational universe.

N.B.—Approximation is necessary, not optional. The axioms prohibit the continuum limit required for exact closures, and therefore all admissible models are approximate shadows of the limiting structure they cannot reach. \square

Let Ψ_δ denote the discrete spline closure permitted by the minimal refinement scale. Let Ψ denote the ideal spline attractor that would appear in the continuum limit. By Axiom ??, Ψ_δ is observationally indistinguishable from Ψ , but it remains a finite-resolution approximation. Any mathematical construction that assumes exact differentiability, exact integration, or exact smoothness implicitly appeals to a limit that the axioms deny. The familiar constructs of calculus therefore do not describe the informational world directly; they describe the limiting behavior that finite closures approximate.

This necessity is not an impediment but a structural guide. The refinement sequence

$$\Psi_\delta \longrightarrow \Psi$$

never completes, yet its monotone convergence ensures that all admissible models become arbitrarily close to the ideal spline at resolutions permitted by the axioms. The continuous spline is unreachable but inevitable: no finite model can realize it, yet every refinement-compatible model approaches it.

Quantum-like Emergence. Finite refinement does more than require approximation; it enforces distinctively non-classical patterns of behavior. The inability to refine distinctions below scale δ produces irreducible uncertainty in the placement of events, non-additivity in refinements, and interference-like behavior when merging partially incompatible records. These effects arise not from physical postulates but from informational structure: finite resolution combined with refinement compatibility forces discrete update rules that mimic the algebra of quantum amplitudes.

N.B.—Quantum-like theories emerge naturally from the necessity of approximation: finite refinement yields non-classical composition of information, which manifests as interference, superposition-like combination, and the familiar probabilistic structure of quantum models. No quantum axioms are assumed; these behaviors follow from measurement constraints alone. \square

Thus approximation is the essential mode of representation in the informational framework. The equations and structures of classical *and* quantum theories arise not because the world is continuous or probabilistic, but because the discrete closures enforced by the axioms approximate the same limiting behavior that continuous and quantum mathematics describe in their respective formalisms.

3.4.3 Equivalence of Discrete and Smooth Representations

Phenomenon 3.4.2 (The Gibbs Phenomenon). *When a discontinuous event is forced into a finite refinement ledger, a residual oscillation appears in its smooth shadow. This overshoot is not an error of representation but the irreducible informational strain of mapping a discrete refinement into a bandwidth limited spline.*

The ringing persists because unobserved structure cannot be admitted. The smooth shadow cannot perfectly close a discontinuity under finite refinement.

The preceding results establish the final closure of the Calculus of Dynamics. An admissible measurement record ψ supported on event anchors $\{x_i\}$ is informationally equivalent to its smooth completion Ψ . The smooth calculus does not introduce new structure; it is the completion of refinement in the discrete domain.

Let ψ be an admissible event record and let $f(\psi)$ denote any interpolant that preserves the anchors and introduces no distinguishable features between them. Refining the interpolant over nested partitions $\{\mathcal{T}_n\}$ produces

a Galerkin sequence $\{\Psi_n\}$. By the convergence theorems, this sequence converges uniformly to a unique \mathcal{C}^2 cubic function Ψ :

$$\Psi_n \xrightarrow[n \rightarrow \infty]{} \Psi.$$

Informational minimality ensures that Ψ is uniquely determined by the anchors: for every event point x_i ,

$$\Psi(x_i) = \psi(x_i).$$

Because Ψ is cubic on each partition element, preserves anchor order, and is globally \mathcal{C}^2 , it is injective on each interval. Its inverse therefore recovers the original record:

$$\Psi^{-1}(x_i) = \psi(x_i).$$

Thus the discrete record ψ and the smooth completion Ψ contain exactly the same information. The interpolant and its limit are informationally equivalent representations of a single causal history.

3.4.4 Recovery of the Euler–Lagrange Form

The weak extremality condition was obtained entirely from finite differences in the discrete domain. In the Galerkin formulation this appears as

$$\int \Psi''(x) \phi''(x) dx = 0, \quad \text{for all admissible test functions } \phi.$$

Integrating this identity twice yields the strong closure

$$\Psi^{(4)}(x) = 0.$$

No differentiability was assumed *a priori*: smoothness appears only as the completion of refinement in the Galerkin limit. The Euler–Lagrange equation is therefore a *recovered* description of the data, not an independent postu-

late. It is sufficient to model the discrete record because every admissible refinement converges to the same \mathcal{C}^2 cubic function.

In this sense the epistemic direction is inverted. We do not derive Euler–Lagrange dynamics and then discretize them. We begin with finite measurements, enforce informational minimality, and recover the Euler–Lagrange operator as the unique smooth shadow of refinement:

$$\text{measurement} \xrightarrow{\text{refinement}} \Psi \xrightarrow{\text{closure}} \Psi^{(4)} = 0.$$

In this sense the epistemic direction is inverted. We do not derive Euler–Lagrange dynamics and then discretize them. We begin with finite measurements, enforce informational minimality, and recover the Euler–Lagrange operator as the unique smooth shadow of refinement:

$$\text{measurement} \xrightarrow{\text{refinement}} \Psi \xrightarrow{\text{closure}} \Psi^{(4)} = 0.$$

Smooth calculus is therefore compatible with the axioms because it contains exactly the information present in the discrete causal record and no more.

N.B.—With apologies to Bishop Berkeley: smooth dynamics are not prior to measurement; they are merely the grammar of its consistent refinement.

□

3.5 The Free Parameter of the Cubic Spline

The Law of Spline Sufficiency requires that the smooth completion Ψ of any admissible record be \mathcal{C}^2 and satisfy $\Psi^{(4)} = 0$. Each segment of Ψ is therefore a cubic polynomial,

$$\Psi(x) = a_0 + a_1x + a_2x^2 + a_3x^3,$$

but informational minimality collapses the apparent local degrees of freedom to a single global parameter.

3.5.1 Fixing the Lower-Order Coefficients

The value a_0 is fixed by the anchors: $\Psi(x_i) = \psi(x_i)$ for every event point x_i . The first derivative Ψ' must be continuous across anchors; a jump in slope would constitute a new observable event, so a_1 is likewise determined. The curvature Ψ'' must also be continuous; any discontinuity would represent an unobserved acceleration and violate informational minimality. Thus a_2 is fixed by \mathcal{C}^2 continuity at the anchors.

These constraints ensure that adjacent cubic segments glue together without introducing new distinguishable structure. The only remaining coefficient, a_3 , controls the third derivative of Ψ :

$$\Psi'''(x) = 6a_3.$$

3.5.2 The Single Free Parameter

Because $\Psi^{(4)} = 0$, the third derivative Ψ''' is constant on every interval of the causal domain. Informational minimality permits this quantity to vary from interval to interval only when the variation is itself detectable as a recorded event. Absent such detection, Ψ''' is the sole unconstrained degree of freedom.

Proposition 5 (The Free Parameter of Information). *The smooth completion Ψ contains exactly one free parameter: the global scale of its third derivative Ψ''' . All lower-order coefficients are fixed by anchor data and continuity constraints.*

Proof (Sketch). Cubic structure follows from $\Psi^{(4)} = 0$. Values and derivatives up to order two are fixed by \mathcal{C}^2 boundary matching; any jump would be observable. Hence the only quantity not determined by anchor data is the

constant third derivative on each interval, which is governed by a_3 . No other freedom remains. \square

A full proof is provided in Appendix A.4.

3.5.3 Physical Interpretation

The single free parameter Ψ''' represents the entire informational content of smooth kinematics. All subsequent dynamical quantities—wave speed, stress, curvature, and eventually mass—are determined by this one global scale. The Law of Spline Sufficiency therefore reduces the continuum to its minimal informational foundation: a \mathcal{C}^2 cubic universe with one degree of freedom.

$$\text{finite record} \xrightarrow{\text{closure}} \Psi \xrightarrow{\text{spline law}} \Psi''' = \text{constant on intervals.}$$

Smooth dynamics contain no structure beyond what is already present in the discrete causal record. The apparent infinity of the continuum collapses to a single free parameter.

3.6 Time Dilation

The informational framework developed in Chapters 5 and 6 places a subtle constraint on how refinement may be transported across a causal network. Proper time is not a geometric parameter but the tally of irreducible distinctions, and the metric $g_{\mu\nu}$ records how this tally must adjust when two histories inhabit regions with different curvature residue. Whenever distinguishability is carried from one domain to another, the connection enforces a compatibility rule: the informational interval must be preserved even if the local refinement structure differs.

This requirement has a striking observable consequence. Two clocks placed at different informational potentials—that is, in regions where the residual strain of admissible curvature differs—cannot maintain the same rate of refinement. Each clock is internally consistent, but the comparison of their records forces an adjustment. A refinement sequence that is admissible at one potential must be reweighted when interpreted at another, or else the causal record would fail to merge coherently.

In the smooth shadow, this bookkeeping adjustment becomes the familiar phenomenon of gravitational redshift. Signals transported upward appear to lose frequency; signals transported downward appear to gain it. Nothing mystical is occurring: the informational interval is being preserved, and the only available mechanism is a change in the rate at which distinguishability is accumulated.

The Pound–Rebka experiment is therefore the archetype of an informational outcome. It demonstrates that when refinement is compared across regions with differing curvature residue, the universe must adjust the apparent rate of time itself to maintain consistency. No dynamical field need be invoked; the redshift is simply the shadow of the constraint that admissible refinements must agree on their causal overlap.

Phenomenon 3.6.1 (The Pound–Rebka Effect [125]). **N.B.**—*The following is an informational phenomenon. No physical mechanism is assumed. The interpretation concerns how the gauge of informational separation $g_{\mu\nu}$ adjusts refinement counts when distinguishability is transported across domains of differing causal potential. Any resemblance to the gravitational redshift measured by Pound and Rebka is a consequence of the informational shadow, not an assumed dynamical cause.* \square

The Axiom of Peano defines proper time as the count of irreducible refinements along an admissible history. The Law of Causal Transport guarantees that this count is invariant under maximal propagation, while the informational metric $g_{\mu\nu}$ (Section 5.2) records how successive refinements compare

when transported across regions whose admissible histories differ in their curvature residue.

Consider two clocks: one at a lower informational potential (higher curvature residue) and one at a higher potential (lower residue). Both clocks produce sequences of refinements

$$\langle e_1 \prec e_2 \prec \cdots \rangle_{\text{low}}, \quad \langle f_1 \prec f_2 \prec \cdots \rangle_{\text{high}},$$

each internally consistent. However, the Law of Boundary Consistency demands that refinements compared across their shared causal overlap must agree on their informational interval. When the refinement sequence from the lower clock is transported to the higher clock, the compatibility condition forces an adjustment in the rate at which distinguishability is accumulated.

Formally, transport along a connection with residue Γ alters the frequency of refinements according to the first-order compatibility condition of Section 5.4:

$$\nu_{\text{high}} = \nu_{\text{low}} (1 - \Gamma \Delta h),$$

where Δh is the informational separation between the clocks. This is the informational analogue of the frequency shift that appears in the smooth limit as gravitational redshift.

In the Pound–Rebka configuration, a photon (interpreted here as a unit of transported distinguishability) sent upward from the lower clock must be refined in such a way that its informational interval remains constant. Since admissible refinements at higher potential accumulate fewer curvature corrections, the transported signal must appear at a lower frequency when measured by the upper clock. Conversely, a downward signal appears at a higher frequency. No physical field is invoked: the effect is a bookkeeping adjustment required to maintain Martin-consistent transport of distinguishability across regions of differing curvature residue.

Thus the informational framework predicts a frequency shift of the form

$$\frac{\Delta\nu}{\nu} \approx \Gamma \Delta h,$$

which matches the structure of the Pound–Rebka observation when interpreted in the smooth shadow of the metric gauge.

The phenomenon of time dilation is therefore an observable outcome of the informational interval and the necessity of refinement-adjusted transport. Differences in curvature residue force clocks at different potentials to accumulate distinguishability at different rates, and the comparison of their refinement counts produces the celebrated redshift.

Coda: The Finite Navier–Stokes Effect

We do not derive the Navier–Stokes equations. Rather, we show how the measurement calculus constrains any smooth limit of finite records to a cubic-spline structure and thereby recasts the regularity question as the finiteness of a single quantity: the third parameter of the spline.

1. Statement of the classical problem

Let $v(x, t)$ be a velocity field and $p(x, t)$ a pressure satisfying the incompressible Navier–Stokes system on \mathbb{R}^3 (or a smooth domain with suitable boundary conditions):

$$\partial_t v + (v \cdot \nabla)v + \nabla p = \nu \Delta v + f, \quad \nabla \cdot v = 0, \quad (3.8)$$

with smooth initial data v_0 . The Millennium Problem asks whether smooth solutions remain smooth for all time or may develop singularities in finite time.

2. Measurement-to-spline reduction

Chapter 2 established that admissible smooth limits of finite records obey a local cubic constraint. Along any coordinate line (and likewise along any admissible selection chain) each component admits a representation whose fourth derivative vanishes in the limit:

$$U^{(4)} = 0 \quad (\text{componentwise along admissible lines}). \quad (3.9)$$

Hence the only freely varying local quantity is the *third parameter* (the derivative of curvature). In one dimension this is U''' . In three dimensions we package the idea as the third spatial derivatives of v :

$$\Theta(x, t) := \nabla(\nabla^2 v)(x, t) \quad (\text{a third-derivative tensor}). \quad (3.10)$$

Informally: v , ∇v , and $\nabla^2 v$ are glued continuously by the spline closure; only Θ may vary piecewise without introducing fourth-order structure.

3. Regularity as finiteness of the third parameter

Principle. If the third parameter Θ stays finite at all scales allowed by measurement, the smooth spline limit persists and no singularity can occur within the calculus of measurement.

A practical surrogate is a scale-invariant boundedness criterion on Θ (or a closely related norm tied to enstrophy growth):

$$\sup_{0 \leq t \leq T} \|\Theta(\cdot, t)\|_X < \infty \quad \implies \quad \text{no blow-up on } [0, T], \quad (3.11)$$

where X is chosen to control the admissible refinements (e.g. an L^∞ -type or Besov/Hölder proxy along selection chains). In words: the only obstruction to global smoothness is unbounded third-parameter amplitude.

4. Heuristic link to classical controls

Energy and enstrophy inequalities control $\|v\|_{L^2}$ and $\|\nabla v\|_{L^2}$. Vorticity $\omega = \nabla \times v$ monitors the first derivative. Growth of $\nabla \omega$ involves $\nabla^2 v$; the *onset* of non-smoothness is therefore detected by $\Theta = \nabla(\nabla^2 v)$, the next rung. Thus the finite-third-parameter condition (3.11) plays the same role in this framework that classical blow-up criteria play in PDE analyses: it is the minimal spline-compatible guardrail against curvature concentration.

5. Non-classical dependency is not invoked

No dependency (cause-effect) is asserted. The argument is purely informational: as long as the admissible record does not force the third parameter to diverge, the cubic-spline closure remains valid and the smooth limit inferred earlier continues to apply.

6. The rephrased question

Navier–Stokes, reframed. Given smooth initial data and forcing, must the third parameter Θ in (3.10) remain finite for all time under (3.8)? Equivalently, can measurement-consistent refinement generate unbounded third-parameter amplitude in finite time?

If Θ stays finite, the spline structure persists, and the calculus of measurement supports global smoothness. If Θ diverges, the smooth continuum description ceases to be representable as a limit of admissible records, and the measurement calculus no longer licenses Euler–Lagrange inference on that interval.

7. What we have and have not done

We have not solved the Millennium Problem. We have shown that within

this program the obstruction to smoothness is concentrated in a single quantity, the third parameter of the cubic spline representation. The classical regularity question is thus equivalent, in this calculus, to the finiteness of Θ .

Chapter 4

Informational Motion

Motion requires a mechanism for comparing one record of distinction with another. In the causal framework, that mechanism is the clock: a source of admissible refinements that allow two observers to construct ordered sequences of events. A clock does not measure duration, velocity, or geometry. It only asserts that one event occurred after another, and that this ordering is invariant under refinement. Its utility is purely ordinal. Whether the clock is an atomic transition, a quartz oscillator, a binary counter, or a list of threshold crossings, its output is the same kind of object: a finite record of distinguishable ticks.

Definition 26 (Clock). ***N.B.**—This is a formal definition of a clock. No physical assumption is made that clocks exist, or that any particular mechanism generates the ticks. In this framework, a clock is only a logical instrument that emits distinguishable events, allowing the causal order to be indexed. Its existence is a definition inside the mathematics of admissible distinctions, not a physical postulate about quantum systems, atomic transitions, or relativistic matter.* □

A clock is an instrument that emits a sequence of distinguishable events. Each emitted event is admissible under Axiom 6: it produces a finite refinement of the causal record. A clock is therefore not a continuous variable or

a dynamical law; it is a device that guarantees the existence of a countable chain of ordered distinctions. The function of a clock is to certify an ordering on the events of a measurement, nothing more.

From this perspective, a clock is not a dynamical primitive. It is a logical instrument. The act of ticking establishes a chain of events, and the absence of extra ticks is a data constraint. If a clock recorded no intermediate events between two ticks, then no admissible description may contain structure that would have produced one. In particular, acceleration, oscillation, or curvature that would create additional ticks are ruled out by informational minimality. Motion is therefore not inferred from a continuous trajectory, but from the consistency of the tick record itself.

Because clocks produce ordered events, two observers may compare their records by merging their tick sequences under global coherence. When the merge produces no contradiction, a single coherent history exists, and the count of ordered refinements defines the relative motion of their systems. In the smooth limit, the unique continuous interpolant between ticked events is the cubic extremal with no unobserved structure. Thus, classical kinematics is the shadow of a discrete bookkeeping process: a clock provides order, informational minimality removes hidden curvature, and the continuum appears only as the completion of finite refinements.

In what follows, motion will be defined as the reconciliation of two causal records produced by clocks. Relative velocity, proper time, and inertial behavior arise not from geometry or differential equations, but from the minimal continuous shadow consistent with their countable tick sequences. Motion is what ordered distinction looks like when refinement tends to the smooth limit.

Thought Experiment 4.0.1 (LiDAR [25]). *Two identical observers, A and B, begin co-located with synchronized clocks. Observer B embarks on a journey involving periods of acceleration, while observer A remains at the origin of an idealized inertial frame. We explicitly neglect the gravitational*

and relativistic influence of Earth, the Sun, Sagittarius A, and all other bodies; spacetime is treated as Minkowski over the region of interest.*

Rather than waiting for reunion, A continuously tracks B by emitting a stream of monochromatic laser pulses. Each pulse is timestamped in A's notebook when fired, and timestamped again when the reflected pulse is received from B's retroreflector.

Every fired pulse is a distinguishable event; every received pulse is another. If B follows a complicated accelerative path, then the return times of the pulses form a more densely refined sequence than the symmetric record A would observe if B were inertial. The point is not energy or Doppler shift. The informational content of the record increases: each round-trip establishes a new ordered pair of emission and reception, constraining B's admissible motion.

If B were inertial, the spacings of the returned timestamps would follow the unique minimal interpolant that introduces no unobserved curvature. But acceleration forces extra refinements: the return times become uneven in a way that cannot be reconciled with a coasting trajectory. These "irregularities" are not interpreted through differential equations; they are simply distinct events that must be merged into A's causal record.

When B returns, both observers merge their sequences. A's laser notebook contains a much longer chain: every emission and every reflection has already placed constraints on B's path. B's local clock, by contrast, has recorded only its own internal ticks and those refinements forced by onboard events. The merge therefore requires A to reconcile a larger informational workload, while B performs a smaller one. Consistent ordering assigns the larger count of admissible distinctions to A, and the smaller to B. The result is that A's proper time is larger—she has the denser causal record.

In the smooth limit, the same count enforces the classical dilation formula of relativity. But here the conclusion is purely informational: acceleration introduces refinements, refinements create more events, and more events imply

more work when histories are coherently merged. Time dilation is the book-keeping of laser-certified distinctions, not a geometric postulate.

This informational mechanism therefore recovers the ability to compute the Lorentz contraction posed in Thought Experiment 2.5.2 through the update rule $E_k = \Psi(e_k \cap \hat{R}(e_{k-1}))$, using only the observers' laboratory notebooks.

4.1 Historical Context

Aaronson and others have demonstrated that quantum mechanics, when viewed through the lens of information theory, admits far less structure than the continuum formalism suggests [1]. Their results show that quantum states do not encode arbitrary real-valued data, that only finitely many distinctions can be operationally extracted from any finite system, and that quantum correlations have deep combinatorial and complexity-theoretic origins rather than geometric ones.

More concretely, Aaronson's work on the complexity of quantum states shows that almost all vectors in Hilbert space are physically meaningless: they cannot be prepared, distinguished, or even approximately specified without exponential resources. In practice, only a tiny, finitely describable subset of states ever arises in nature. This directly parallels the Axiom of Kolmogorov, which asserts that measurement produces only finite, countable information and that no refinement may introduce distinctions that cannot be operationally supported.

Similarly, Aaronson's results on shadow tomography establish that measurement itself imposes strict limits on what can be learned. Even with unlimited computational power, only a bounded amount of information about a quantum state can be extracted without an exponential number of queries. This mirrors the Axiom of Planck: distinguishability has a minimal scale, and refinements cannot probe below it.

Finally, the modern complexity-theoretic analysis of entanglement shows

that quantum correlations arise from constraints on how information may be shared and refined across subsystems. These correlations are not geometric artifacts but restrictions on admissible joint refinements. This observation aligns with the notion developed in this chapter that entanglement is the smooth shadow of *uncorrelant* event pairs—events whose informational ordering cannot be determined and whose refinements may fail to commute.

The informational constraints emphasized by Aaronson and others also carry direct implications for the concept of motion. If quantum states contain only finitely many operationally accessible distinctions, then the evolution of a system cannot be a continuous geometric flow through an uncountable state space; it must be the refinement of a finite informational record. Motion is therefore the process by which distinguishable events accumulate in a manner consistent with the Axioms of Kolmogorov, Peano, and Planck. In this view, trajectories are not primitive curves but the dense limits of these discrete refinement steps, and kinematics emerges only after enforcing Ockham minimality and Boltzmann coherence. The core insight is that bounded distinguishability and limited extractable information constrain how histories may evolve, and the resulting admissible refinements form precisely the minimal-structure extremals that appear as smooth worldlines in the continuum limit. Thus, the informational limits identified by Aaronson do not merely illuminate quantum phenomena; they determine the very structure of motion itself.

Phenomenon 4.1.1 (Shadow Tomography [1]). ***N.B.**—This informational phenomenon reflects results by Aaronson and others showing that only a bounded amount of operationally accessible information about a quantum system can be extracted, regardless of the continuum descriptions allowed by Hilbert-space formalism. The argument below does not use physical tomography; it expresses the same limitation in the language of refinement and distinguishability.* □

Consider a system whose underlying measurement record consists of a dis-

crete chain of refinements. Let $\{O_1, \dots, O_m\}$ be a family of admissible tests that the observer may apply. Classically, one might expect that by probing the system with sufficiently many such tests, one could reconstruct an arbitrarily detailed internal description. Shadow tomography demonstrates that this is not the case: only a small, coarse projection of the underlying informational structure can ever be distinguished.

From the standpoint of the Axioms of Measurement, the reason is immediate. Each test O_j extracts only the distinctions resolvable at the minimal increment dictated by the Axiom of Planck. The Axiom of Kolmogorov ensures that each measurement outcome has finite informational content, and the Axiom of Peano ensures that these outcomes accumulate discretely. Thus, even an exponentially large sequence of tests cannot expose distinctions that lie below the minimum resolvable scale or that require refinements forbidden by the Axiom of Ockham.

Operationally, the observer does not recover the internal structure of the system's full refinement history. Instead, they recover a shadow: the projection of that history onto the small set of distinctions probed by the tests $\{O_j\}$. Two systems whose internal refinements differ but whose shadows coincide are operationally indistinguishable. In the language of this manuscript, they represent distinct admissible histories that yield the same externally visible refinement pattern.

This phenomenon clarifies why the continuum description of quantum states contains far more degrees of freedom than can ever appear in practice. Shadow tomography reveals that measurement accesses only the coarse-grained shadow of the underlying informational structure, never its complete refinement. It provides an operational reason why uncorrelant events, informational decoherence, and refinement non-commutation arise naturally: the observer sees only the shadow, while the full informational record remains inaccessible.

4.2 Relative Motion

Phenomenon 4.2.1 (The Quantum Zeno Effect). *Repeated measurement prevents motion. When refinement occurs faster than the system can admit a new distinguishable prediction, the restriction operator repeatedly discards all alternative continuations and resets the ledger to its current state.*

Motion therefore requires silence. A gap in observation is necessary to allow ambiguity to accumulate and justify a change in admissible history.

With a clock in hand, an observer constructs an ordered sequence of admissible events $\{e_k\}$. The Causal Universe Tensor encodes this growing record through the recursive update

$$E_k = \Psi(e_k \cap \hat{R}(e_{k-1})),$$

where \hat{R} restricts all admissible continuations to those consistent with the most recent distinguishable event. For a single inertial frame, this produces a monotone refinement of information: each step eliminates histories that would contradict the record.

To describe motion, no new structure is required. Consider two inertial observers, A and B, each with their own clock and their own causal universe tensors:

$$E_k^A = \Psi(e_k^A \cap \hat{R}(e_{k-1}^A)), \quad E_\ell^B = \Psi(e_\ell^B \cap \hat{R}(e_{\ell-1}^B)).$$

Each observer possesses a complete and self-consistent history of admissible distinctions. Relative motion is nothing more than the reconciliation of these two histories under the rule of global coherence. When A and B compare notebooks, every admissible refinement recorded by one must also be admissible to the other. In the absence of contradiction, a merge exists, and the merged list induces a partial ordering on the pair $\{E_k^A\} \cup \{E_\ell^B\}$.

If the merged record is strictly longer than either individual record, the observers infer that their clocks have accumulated refinements at different

rates. By Axiom 5, the merged history cannot discard any admissible distinction: every recorded refinement must be preserved. Consequently, the observer whose notebook produces the *denser* merged record corresponds to the *longer* proper time. The informational work of reconciling those refinements is the measure of duration. Thus,

$$|e_k^A \cap \hat{R}(e_{k-1}^A)| > |e_\ell^B \cap \hat{R}(e_{\ell-1}^B)| \quad (4.1)$$

implies that A has the longer proper time. No metric, coordinate chart, or differential equation is required; the relative motion is encoded entirely in the informational asymmetry of their merged causal histories.

Phenomenon 4.2.2 (The Velocity Effect). *Velocity is not a dynamical state but a relational count. It is the ratio of distinguishable refinement events between two admissible ledgers after they have been merged. Only differences in event counts survive reconciliation.*

Motion is therefore not something possessed by an object, but a discrepancy between records that must be reconciled to preserve global consistency.

In the smooth limit, the unique continuous interpolant of the merged record is the cubic extremal with no unobserved structure. Classical kinematics—relative velocity, time dilation, and Lorentz contraction—appears as the shadow of this merge. The Causal Universe Tensor does not simulate motion; it enforces consistency. Relative motion is what two coherent universe tensors look like when compared under refinement.

4.2.1 Merging a Single Event

Referring to Thought Experiment 4.0.1, consider the merging of a single event: the moment a reflected photon is absorbed by A's detector. This absorption is a distinguishable refinement of A's record and therefore constitutes an admissible event e_{k+1}^A . The photon has traveled to B, interacted with

the retroreflector, and returned. Whatever else the experimenter may imagine, this exchange contains one certified fact: the causal distance between A and B has changed in a way detectable by A's clock.

In the language of the causal universe tensor, the absorption is merged via

$$E_{k+1}^A = \Psi(e_{k+1}^A \cap \hat{R}(e_k^A)).$$

Nothing more is required. The event contributes only the distinction that A received a photon at that moment. The return time rules out any hypothetical motion of B that would have prevented this arrival, and it rules out any curvature or oscillation that would have produced additional admissible pulses. The refinement therefore narrows A's admissible histories to those consistent with both emission and reception.

When B later inspects A's notebook, the same absorption event must be admissible within B's causal universe tensor:

$$E_{\ell+1}^B = \Psi(e_{k+1}^A \cap \hat{R}(e_\ell^B)).$$

If a contradiction were forced—for example, if B's notebook implied the photon could not have returned at that time—global coherence would fail, and the combined record would be inadmissible. But if the merge succeeds, the joint history becomes strictly more refined, and the updated tensors¹ encode a new restriction on their relative motion.

A single merged photon event therefore eliminates an entire family of hypothetical motions. It narrows the admissible set of configurations and extends the causal record without introducing any continuous structure. In the smooth limit, repeated merges of this form force the cubic extremal between emission and reception times—the unique interpolant with no unobserved structure. Classical distance, velocity, and Lorentz contraction appear as the

¹No physical model of a photon is required. The “photon” only represents a distinguishable event transmitted between observers.

continuous shadow of this discrete bookkeeping.

4.2.2 Measurement of Acceleration as Counts of Events

Acceleration does not require forces, masses, or differential equations. In the causal framework, acceleration is nothing more than a second refinement: a change in the distinguishable difference between successive admissible events. To detect such a change, a single measurement is insufficient. At least two refined measurements are needed so that the difference between them can itself be distinguished.

Suppose A emits two photons at events e_k^A and e_{k+1}^A , and later receives their reflections at e_{k+r}^A and e_{k+s}^A . Each absorption is merged by

$$E_{k+r}^A = \Psi(e_{k+r}^A \cap \hat{R}(e_{k+r-1}^A)), \quad E_{k+s}^A = \Psi(e_{k+s}^A \cap \hat{R}(e_{k+s-1}^A)).$$

If B is in uniform motion relative to A, the refinements contributed by these two events are consistent with a unique minimal interpolant: the admissible histories require that the difference in reception times is itself constant under refinement. Any hidden curvature or oscillation would have produced additional admissible events—extra pulses, missed reflections, or altered return order—and is therefore ruled out by Axiom 6.

However, if the spacing between e_{k+r}^A and e_{k+s}^A cannot be reconciled by a single coasting history, then the admissible set must be further restricted. The causal universe tensor eliminates all hypothetical configurations in which B remained inertial. What remains are those histories in which the separation of the events changes in a way that is itself distinguishable. The second refinement is the signature of acceleration.

In this sense, acceleration is not a postulated quantity. It is the discovery that two refinements cannot be merged into a single coasting interpolant without contradiction. A sequence of such measurements produces a chain of eliminations: each return time excludes admissible events that would require

an invisible change in curvature. The remaining histories are the ones in which acceleration has occurred.

Phenomenon 4.2.3 (The Acceleration Effect). *Acceleration is the count of distinguishable failures of a coasting interpolant to account for a merged ledger. A straight refinement trajectory represents a zero-cost hypothesis. Every detectable deviation is recorded as a second-order refinement.*

Acceleration is therefore not force, but the number of contradictions a simple refinement model fails to resolve.

In the smooth limit, repeated second refinements force a unique continuous extremal whose second variation is nonzero. Classical acceleration appears as the shadow of a finite bookkeeping process: acceleration is the count of distinguishable failures of coasting to explain the merged record. No forces, masses, or trajectories are assumed. The event counts alone enforce curvature in the admissible histories.

Thus, we have recovered the ability to verify Newton’s second law—again, with apologies to Lord Berkeley. Acceleration is not a substance but the second variation of admissible refinements in the merged event record.

4.2.3 The Equations of Motion

The remainder of this chapter examines the equations of motion that arise when finite records of admissible distinctions are merged without contradiction. Nothing further is assumed. Each equation appears as the continuous shadow of informational minimality: the unique smooth extremal that contains no unrecorded structure.

We begin with heat transport. Although commonly divided into conduction, convection, and radiation, all three arise here from distinct constraints on admissible refinements. When refinements diffuse symmetrically through a medium with no hidden variations, the smooth limit forces the diffusion equation. When refinements are transported coherently through the medium,

the extremal satisfies the advective transport law. When refinements propagate at the maximal admissible speed, the continuous shadow is radiative transport governed by the wave equation. No model of heat is assumed; each law is simply the completion of a finite notebook of events.

Annealing appears when a ledger repeatedly reconciles its own coarse description. The iterated application of the merge operator eliminates sharp distinctions that would predict unobserved refinements. As the sequence of folds converges, the smooth limit is diffusion. Annealing is therefore informational smoothing: the heat equation is its continuous shadow when the coarse ledger is refined to closure.

Adiabatic transport arises when the ledger evolves without creating or destroying admissible distinctions. In the smooth limit, this invariance forces the classical adiabatic law. Nothing dynamical is postulated; an adiabatic process is simply a sequence of refinements that preserves the global count of admissible configurations.

Even quantum phenomena admit the same treatment. The Casimir effect appears when the merged record forbids a continuous family of admissible configurations between two boundaries. The elimination of those histories produces an informational pressure, and the smooth limit recovers the familiar expression for Casimir energy.

Alpha decay appears in its original form: the Mott problem [110]. The ledger of the nucleus contains two nearly indistinguishable configurations—one in which the alpha cluster remains bound, and one in which it escapes. Over time, these dual descriptions drift out of alignment. The moment of decay is not the passage of a particle through a barrier, but the repair of a contradiction: the merged record eliminates all configurations in which the two descriptions diverge. The resulting refinement is recorded as a distinct decay event. In the smooth limit, this informational repair produces the exponential law of radioactive decay without invoking forces, potentials, or tunneling particles. As Einstein suggested, no dice are rolled [citeeinstein1949](#).

In each case, the classical equation of motion is not assumed. It is what consistency looks like in the smooth limit of finite measurement. Motion is bookkeeping; the laws that follow are shadows of refinement.

4.2.4 Martin’s Condition and the Propagation of Order

Up to this point, motion has been defined locally: two observers exchange admissible events, merge their records, and eliminate any hypothetical history that would have produced unrecorded refinements. This closure guarantees that each observer maintains a coherent ledger. It does not yet guarantee that their ledgers are mutually compatible.

For observable physics, local coherence is not enough. Distinct observers must be able to reconcile their refinements along their shared boundary without introducing new distinguishabilities. The requirement that every locally finite patch of causal order extends to a globally consistent history is Martin’s Condition.

Definition 27 (Martin’s Condition [105]). ***N.B.**—The formulation of Martin’s Condition used here is not a single axiom from set theory or forcing, but an operational synthesis of the requirements imposed by Axioms 3, 5, and 7. It echoes the role of Martin’s Axiom in ensuring consistent extensions of locally compatible partial structures, but is adapted to the causal network of distinguishable events. The condition should therefore be understood as a consolidated operational rule rather than a direct quotation of any single classical axiom.* □

A causal network (Definition 11) satisfies Martin’s Condition if every locally finite subset of events can be extended to a globally consistent ordering without introducing new admissible distinctions. Equivalently, all finite causal updates admit an extension that preserves the same coincidence relations on their overlaps.

Intuitively, Martin’s Condition demands that information created in one region does not contradict information measured in another. It forbids causal overcounting—the duplication of distinctions that would destroy reversibility—by ensuring that overlapping observers reconstruct identical splines of the causal universe tensor along their shared boundary. Axiom 7 limits what may happen within a light cone; Martin’s Condition governs how those choices propagate outward.

Once Martin’s Condition holds, the closure of finite refinements induces a global propagation rule. Locally symmetric overlaps enforce a second variation and yield the wave operator. Oriented overlaps enforce a first variation and yield advection. When variations are eliminated by repeated projection, the smooth limit is diffusion. The familiar equations of motion—waves, advection, diffusion, and later curvature—are therefore the continuous shadows of global consistency under Martin’s Condition.

Thought Experiment 4.2.1 (The Davisson–Germer Effect [35]). **N.B.**—*This label refers only to the informational structure of the example. No physical wave, field, or substrate is assumed. The “wave” described here is the smooth shadow of a refinement sequence whose admissible extensions satisfy Martin-consistency. The phenomenon is therefore informational: a pattern enforced by the logic of distinguishability, not by any physical mechanism of diffraction or interference.* \square

Imagine an electron gun firing individual electrons toward a crystalline nickel target. A distant screen records the arrival of scattered electrons as distinguishable events. Between gun, crystal, and screen, no internal distinctions are measured; the observers record only the emission, the scattering plane, and the pattern of impacts. Each detection on the screen is therefore an admissible refinement of the joint causal ledger of gun, crystal, and detector.

Under Martin’s Condition, every locally finite segment of this ledger must extend to a globally consistent history. The crystal introduces a periodic

partition: successive lattice planes represent indistinguishable choices, except at angles where the merged ledger would predict additional or missing refinements. Along these planes, reciprocal measurement enforces translation invariance: if one segment of the ledger is shifted by a lattice spacing, the count of admissible refinements must remain unchanged.

The only smooth extremals compatible with this translation invariance are wave modes. Among these, the constructive modes are precisely those whose wavelength λ satisfies Bragg's relation [19]

$$2d \sin \theta = m\lambda, \quad \lambda = \frac{h}{p},$$

where d is the lattice spacing, θ the scattering angle, m an integer, and h/p encodes the count of distinguishabilities preserved along the oriented Martin bridges. At those angles, no hidden refinements are predicted; outside them, the merged ledger would contain missing or extra distinguishable events, contradicting Martin's Condition.

Operationally, the bright peaks on the screen are fixed points of reciprocal measurement under lattice translations. What physicists call "electron diffraction" is simply the bookkeeping consequence of demanding that indistinguishable causal neighborhoods propagate consistently across the crystal. No wavefunction is assumed. The "wave" is the unique smooth extension of discrete, Martin-consistent event counts.

Thus, the Davisson–Germer experiment does not demonstrate that electrons are waves or particles. It demonstrates that any causal history satisfying Martin's Condition must propagate its indistinguishabilities as waves. The universality of wave behavior is a consequence of global consistency, not a special property of matter.

4.3 The Algebra of Interaction

Each system X carries an accumulated causal universe tensor as a left-fold of update factors:

$$\mathbf{U}_1^X = E_1^X, \quad \mathbf{U}_{n+1}^X = E_{n+1}^X \mathbf{U}_n^X, \quad E_{n+1}^X := \Psi(e_{n+1}^X \cap \hat{R}(e_n^X)).$$

Definition 28 (Interaction operator). *Given two ledgers (tensors) \mathbf{U}^A and \mathbf{U}^B , the interaction operator*

$$f : (\mathbf{U}^A, \mathbf{U}^B) \mapsto \mathbf{U}^{AB}$$

returns the minimal accumulated state \mathbf{U}^{AB} that extends both inputs and is Martin-consistent on their overlap. Equivalently, \mathbf{U}^{AB} is obtained by left-folding the common update factors (the jointly admissible events) in observed order so that no unrecorded refinements are invented (Axiom 5) and none already recorded are erased (Axiom 6). Let $E(\mathbf{U})$ denote the underlying event set of \mathbf{U} and define the newly contributed distinctions by

$$\mathbf{J}^{AB} := E(\mathbf{U}^{AB}) \setminus (E(\mathbf{U}^A) \cup E(\mathbf{U}^B)).$$

Phenomenon 4.3.1 (The Stoichiometry Effect). **Statement.** *Causal interactions are governed by Diophantine constraints, not continuous variation. Because the causal ledger is composed of discrete, indivisible events, admissible interactions occur only when integer refinement counts balance exactly.*

The Integer Constraint. *Let N_A and N_B denote the number of unresolved refinement threads carried by systems A and B . An admissible interaction*

$$f(U_A, U_B) \rightarrow U_C$$

exists only if there are integers $a, b, c \in \mathbb{Z}$ such that

$$aN_A + bN_B \rightarrow cN_C.$$

No fractional event may be recorded, and no partial refinement may be committed.

Hard Failure (No Reaction). If the integer balance cannot be satisfied, no admissible merge exists. The ledger rejects the update. The systems may scatter, deflect, or pass through one another, but no interaction occurs, because a fractional event would be required to close the account.

Conclusion. Chemical stoichiometry, particle number conservation, and selection rules are not arbitrary physical laws. They are bookkeeping necessities imposed by the impossibility of writing half an event in a discrete causal ledger. An interaction is the solution of an integer program.

Definition 29 (Length on the common boundary [29, 150]). Let $\partial(\mathbf{U}^A, \mathbf{U}^B)$ denote the common boundary (overlap) of the ledgers \mathbf{U}^A and \mathbf{U}^B . The length on the boundary is the number of folded factors from a ledger that lie on this overlap:

$$\text{len}_{\partial}(\mathbf{U}^A, \mathbf{U}^B) := \text{len}(\mathbf{U}^A \upharpoonright_{\partial(\mathbf{U}^A, \mathbf{U}^B)}), \quad \text{len}_{\partial}(\mathbf{U}^B, \mathbf{U}^A) := \text{len}(\mathbf{U}^B \upharpoonright_{\partial(\mathbf{U}^A, \mathbf{U}^B)}).$$

Equality $\text{len}_{\partial}(\mathbf{U}^A, \mathbf{U}^B) = \text{len}_{\partial}(\mathbf{U}^B, \mathbf{U}^A)$ expresses informational equilibrium on the shared frontier.

Phenomenon 4.3.2 (The Ideal Ledger Effect). **Statement.** The ideal gas law is the bookkeeping identity of an uncorrelant causal interior. Pressure is the rate at which the boundary ledger must reconcile independent refinement threads generated in the bulk.

Uncorrelant Interior. Consider a region Ω containing n causal threads that are mutually uncorrelant. Each thread generates refinement events at an average rate T . Because these threads do not refine one another, their only point of mutual interaction is the boundary.

Boundary Bottleneck. Let V denote the number of addressable refinement slots in the partition. The boundary $\partial\Omega$ must perform Martin-consistency checks for each incoming update. When V is large, reconciliation events are sparse. When V is small, reconciliation requests crowd the same causal addresses.

Informational Pressure. Pressure is the flux density of reconciliation at the boundary:

$$P \propto \frac{nT}{V}.$$

Rearranging yields the familiar bookkeeping identity:

$$PV \propto nT.$$

Hard Failure. If the reconciliation rate demanded of the boundary exceeds its admissible bandwidth, coherence fails locally. The boundary can no longer preserve global consistency, and the partition ruptures. In classical language, this appears as an explosion.

Conclusion. The ideal gas law is not a statement about elastic collisions. It is the equation of state for uncorrelant ledgers under finite boundary bandwidth.

Proposition 6 (The Anti-symmetry of Information Propagation). *In general $f(\mathbf{U}^A, \mathbf{U}^B) \neq f(\mathbf{U}^B, \mathbf{U}^A)$. Symmetry holds iff the overlap carries equal refinement counts:*

$$f(\mathbf{U}^A, \mathbf{U}^B) = f(\mathbf{U}^B, \mathbf{U}^A) \iff \text{len}_{\partial}(\mathbf{U}^A, \mathbf{U}^B) = \text{len}_{\partial}(\mathbf{U}^B, \mathbf{U}^A).$$

Proof (Sketch). The interaction operator $f(U_A, U_B)$ performs a left-fold of all jointly admissible update factors on the overlap $\partial(U_A, U_B)$, in the unique order that is consistent with the causal refinements already recorded in each ledger. Anti-symmetry arises because this fold depends on the observed order of refinements whenever the overlap contains correlated (noncommuting)

factors.

Suppose first that the refinement counts on the shared boundary are equal:

$$\text{len}_{\partial}(U_A, U_B) = \text{len}_{\partial}(U_B, U_A).$$

Every factor lying on the overlap is therefore recorded with the same resolution by both ledgers. No ledger contributes a strictly finer refinement than the other on the shared frontier. In this case the overlap consists only of mutually uncorrelant update factors: their order is not fixed by either ledger, and informational minimality forces them to commute. Because the only factors whose relative placement could differ lie in this commuting set, the resulting left-fold is invariant under exchanging the inputs, and

$$f(U_A, U_B) = f(U_B, U_A).$$

Conversely, assume the refinement counts on the overlap are unequal. Without loss of generality, let U_A record strictly more refinement on the boundary than U_B . Then $\partial(U_A, U_B)$ contains at least one factor recorded by A with higher resolution than by B . Such a factor cannot be uncorrelant: if it were, its finer structure could not have been observed by only one ledger. The overlap therefore contains a correlated pair of update factors whose tensor representatives do not commute. The left-fold must place this pair in the local causal order recorded by the corresponding ledger. Because U_A and U_B record different boundary orders for these noncommuting factors, the two possible folds produce distinct accumulated tensors:

$$f(U_A, U_B) \neq f(U_B, U_A).$$

Thus symmetry of the interaction operator occurs exactly when the two ledgers carry equal refinement counts on their shared boundary, and fails precisely when one ledger resolves strictly more distinguishable structure than the other. \square

A full proof is provided in Appendix A.5.

Proposition 7 (The Transitivity of Information Propagation). *For any Martin-consistent triple $\mathbf{U}_n, \mathbf{U}_{n+1}, \mathbf{U}_{n+2}$,*

$$f(\mathbf{U}^A, \mathbf{U}_{n+2}) = f(\mathbf{U}^A, f(\mathbf{U}_n, \mathbf{U}_{n+1})).$$

That is, folding via the intermediate ledger equals folding directly into \mathbf{U}_{n+2} .

Proof (Sketch). Let U_n, U_{n+1}, U_{n+2} be a Martin-consistent triple. Each ledger is a left-fold of its admissible update factors, and the interaction operator f produces the minimal ledger that extends its inputs without inventing or erasing recorded refinements. The transitivity property expresses the fact that the unique globally coherent ledger for the triple does not depend on how the pairwise folds are grouped.

Consider the right-hand side,

$$f(U_A, f(U_n, U_{n+1})).$$

The inner fold $f(U_n, U_{n+1})$ reconciles all jointly admissible refinements of U_n and U_{n+1} on their shared boundary. Because the pair is Martin-consistent, this fold is unique: no alternative ordering of their overlapping factors survives the consistency check. The result is a ledger that contains exactly the refinements common to both inputs together with their compatible unique factors. Folding this ledger with U_A adds precisely the admissible refinements from U_A that remain consistent with the already merged pair. No additional events may be inserted, and none already present may be removed.

Now consider the left-hand side,

$$f(U_A, U_{n+2}).$$

Since the triple is Martin-consistent, U_{n+2} already encodes all refinements that can appear after U_{n+1} without violating Axioms 6 or 5. Any refinement

compatible with U_n and U_{n+1} must also be compatible with U_{n+2} . Thus the direct fold of U_A with U_{n+2} produces a ledger that contains exactly the jointly admissible refinements of all three inputs. As before, no additional distinctions may be introduced.

In both constructions, the surviving event factors are the same: the set of refinements jointly admissible across the triple. Martin's Condition ensures that this set admits a unique causal ordering, so both sides fold precisely the same sequence of factors. By informational minimality and uniqueness of the admissible ordering, the resulting tensors must coincide:

$$f(U_A, U_{n+2}) = f(U_A, f(U_n, U_{n+1})).$$

Thus the interaction operator is transitive on any Martin-consistent triple: grouping of intermediate folds does not affect the final accumulated ledger. \square

A full proof is provided in Appendix A.6.

Proposition 8 (The Commutativity of Uncorrelant Events). *If*

$$f(f(\mathbf{U}^A, \mathbf{U}^B), f(\mathbf{U}^C, \mathbf{U}^D)) = f(f(\mathbf{U}^C, \mathbf{U}^D), f(\mathbf{U}^A, \mathbf{U}^B)),$$

then the pairs (A, B) and (C, D) are relativistically simultaneous: exchanging block order introduces no new admissible distinctions on the shared boundary; the merged tensor is invariant under the swap.

Proof (Sketch). Let $U^{AB} := f(U_A, U_B)$ and $U^{CD} := f(U_C, U_D)$. The hypothesis is that the two blocks commute under the interaction operator:

$$f(U^{AB}, U^{CD}) = f(U^{CD}, U^{AB}).$$

By Proposition 6, such commutativity can occur only when the shared boundary carries equal refinement counts. In the present setting this means that

every update factor lying in the overlap $\partial(U^{AB}, U^{CD})$ is recorded at the same resolution by both blocks. No factor is strictly more refined on one side than the other.

Equal refinement counts force the overlapping factors to be uncorrelant: neither block records a finer causal relation among these events, so informational minimality forbids any ledger from resolving a precedence relation absent from the other. In the tensor algebra this uncorrelance appears as commutation of the corresponding update factors. Because only these boundary factors can appear in different relative positions when the blocks are folded, and because they commute, swapping the blocks yields the same accumulated ledger.

To interpret this result, note that two events are uncorrelant precisely when neither precedence $e < f$ nor $f < e$ is recorded in any admissible refinement. Such events lie outside each other's causal neighborhoods; exchanging their order introduces no new distinguishable structure and preserves all scalar invariants of the universe tensor. Thus, if the blocks (A, B) and (C, D) commute under f , every event in the first block is uncorrelant with every event in the second. No causal precedence can be established across the blocks.

This is exactly the condition of relativistic simultaneity in the causal framework: the two blocks occupy spacelike-separated regions of the observational record. Their fold order is unconstrained, and the merged ledger is invariant under the swap. Hence commutativity of the interaction operator implies relativistic simultaneity. \square

A full proof is provided in Appendix A.7.

N.B.—This is the point at which the usual notion of *causality* is rejected. No geometric light cones, no differential structure, and no propagation law are assumed. The only order in the development is the order of *recorded* refinements. What physicists call causal structure appears later only as the smooth shadow of informational bookkeeping: the continuum calculus that

encodes cause–effect relations is not a primitive of the theory but an emergent completion of discrete refinements. Nothing in this chapter assumes or relies on physical causation; all that is used is the partial order induced by Axiom 5.

□

N.B.—Uncorrelant events play a central conceptual role in this framework. They are not “independent random variables” nor “simultaneous in a reference frame” nor artifacts of a chosen coordinate system. They are the events for which the record contains *no admissible refinement* that orders one before the other. This absence of recorded precedence is an observable fact, not a geometric assumption. All smooth notions of spacelike separation, relativistic simultaneity, and commuting update factors arise from this single idea. When two events are uncorrelant, reordering their update factors creates no new distinguishable structure, and every algebraic invariant of the ledger is preserved. The geometry of relativity is therefore not presupposed but recovered from the informational status of uncorrelance. □

Remark 15.

Idempotence: $f(\mathbf{U}^A, \mathbf{U}^A) = \mathbf{U}^A$.

Monotonicity: \mathbf{U}^{AB} is a monotone extension of both inputs; no recorded refinement is removed.

Locality: Joint refinements lie in the common causal neighborhood; fold order is the observed order; reordering is forbidden unless the corresponding factors commute.

Operational link: Bi-directional folds yield the wave operator; oriented folds yield advection; iterated projection yields diffusion. These are smooth shadows of the discrete left-fold $\mathbf{U}_{n+1} = E_{n+1}\mathbf{U}_n$ under f .

Phenomenon 4.3.3 (The Entanglement Effect [46]). **N.B.**—The Dantzig Pivot [32] is not a physical process. Nothing travels, no signal is sent, and no mechanism propagates. The pivot is bookkeeping: boundary consistency

is enough to eliminate incompatible histories without scanning the interior of the ledger. \square

Two spacelike-separated laboratories, A and B , each maintain their own causal universe tensor. A single preparation event produces two admissible refinements, e_i and e_j , that are indistinguishable in causal order: both

$$\langle e_i \prec e_j \rangle \quad \text{and} \quad \langle e_j \prec e_i \rangle$$

generate the same accumulated state. No scalar invariant recorded in either ledger can tell which ordering occurred. This is a state of causal degeneracy: two distinct histories produce the same observational content.

At time $n+1$, laboratory A measures e_i . By Axiom 6, this refinement must be folded into the accumulated state. The interaction operator f computes

$$\mathbf{U}_{n+1} = f(\mathbf{U}_n, e_i),$$

which is a strict update: e_i now has a definite position in the record relative to all prior events.

Because e_i and e_j were degenerate, this update triggers a global repair. The merged ledger must eliminate every history in which e_j is ordered incompatibly with e_i under Martin's Condition. No signal is sent from A to B ; instead, the causal universe tensor performs a pivot: it selects the unique ordering of (e_i, e_j) that avoids introducing new distinguishabilities. The ambiguous pair collapses to a single admissible ordering.

Critically, this repair is not a search over an entire volume of possible histories. Martin's Condition requires agreement only on the boundary of the overlap: the parts of \mathbf{U}^A and \mathbf{U}^B that already coincide. The pivot therefore acts on the smallest region where a contradiction could occur. Only the boundary is inspected, and only the incompatible orderings are removed. There is no need to re-evaluate the entire causal universe; the ledger verifies consistency by checking the joint frontier. Interaction is thus computable:

global coherence is enforced by local boundary repair, not by scanning an exponential set of histories.

Thus, the “instantaneous” correlation is not a physical transmission. It is the bookkeeping consequence of a non-degenerate refinement. Entanglement is the existence of causal degeneracy; the apparent nonlocal update is the pivot that removes it by repairing the boundary of the overlap.

The name “pivot” is not accidental. In Dantzig’s algorithm, a degenerate solution is resolved by moving along the boundary of admissible configurations until a single vertex remains consistent with all constraints. The search never explores the interior volume of the feasible set; it advances only along the frontier where inconsistency can appear. The causal pivot behaves the same way. When a non-degenerate refinement is recorded, the ledger examines only the boundary of the overlap and removes incompatible orderings. The result is a unique, globally coherent history selected by local boundary repair. In both settings, the pivot is a boundary operation, not a volume search: global consistency is enforced without scanning an exponential family of possibilities.

Phenomenon 4.3.4 (The Mach–Zehnder Effect [164]). **N.B.**—*Although the Mach–Zehnder device originates in optical physics, the informational structure it exhibits does not depend on any physical mechanism. The branching and recombination of admissible refinements is a purely combinatorial phenomenon: it arises whenever two indistinguishable paths diverge, evolve under independent refinements, and reunite at a shared boundary. No metric, phase, or wave dynamics are assumed.* \square

A single photon enters a Mach–Zehnder interferometer. At the first beam splitter, a single input event e_0 leads to two admissible refinements, e_1 (upper path) and e_2 (lower path). Both produce valid causal chains: each path accumulates its own ordered list of refinements—reflections, delays, and phase shifts—and each yields an accumulated tensor $\mathbf{U}^{(1)}$ and $\mathbf{U}^{(2)}$ satisfying Martin’s Condition. No experiment in either arm can distinguish which refinement is “real”: both histories are admissible and neither produces a con-

tradiction. The interferometer therefore carries two coexisting, consistent ledgers.

At the second beam splitter, the detection event e_f must be recorded as a strict update. By Axiom 6, the refinement e_f must fold into the accumulated state. The interaction operator computes

$$\mathbf{U}_{\text{final}} = f(\mathbf{U}^{(1)}, \mathbf{U}^{(2)}),$$

the minimal accumulated tensor consistent with both paths. All hypothetical histories in which the arrival at e_f contradicts either ledger are removed.

Interference is the informational comparison of the two causal chains. If their accumulated phase—a bookkept record of distinguishability—is equal modulo 2π , the paths are informationally indistinguishable at the boundary. The fold produces a single ledger: both paths merge without creating new refinements. If the accumulated phase differs by π , the asymmetric parts of the update factors cancel under the fold, and e_f becomes inadmissible. No destructive force is invoked; the cancellation expresses the fact that no consistent ledger can be formed with that ordering.

Thus, “superposition” is the coexistence of multiple valid, Martin-consistent refinements until detection forces a non-degenerate fold. The Mach–Zehnder interferometer does not show a particle traveling two paths; it shows that causal histories can remain distinct and simultaneously admissible until the interaction operator selects the unique ordering that avoids contradiction at the boundary.

Phenomenon 4.3.5 (The Bell–Aspect Tests [9]). Two spacelike-separated laboratories, A and B , share a preparation event that produces an entangled pair. Each maintains its own causal universe tensor. The preparation is such that multiple ordered refinements remain admissible: different measurement settings at A and B produce distinct, yet individually consistent, ledgers. Before either measurement is recorded, the global state is degenerate: many

joint histories remain compatible with all previous refinements, and no scalar invariant distinguishes among them.

A local hidden-variable model assumes that this degeneracy can be resolved purely by local rules. In ledger language, it assumes that the update

$$(\text{measure at } A, \text{ measure at } B)$$

can be decomposed into separate, predetermined refinements in each ledger. That is, the merged state could be written as a fold of two independent maps acting only on local records, with no global repair.

The Bell–Aspect tests show this is impossible. When A records a refinement corresponding to setting a and B records one corresponding to b , the accumulated tensor must be updated by the interaction operator,

$$\mathbf{U}_{\text{final}} = f(\mathbf{U}^A, \mathbf{U}^B).$$

For many setting pairs (a, b) , the resulting ledger eliminates histories that would have remained admissible under any local rule. The violation of Bell inequalities is the empirical statement that no decomposition of f into independent, local updates can preserve all observed distinctions. The fold is intrinsically global.

Operationally, a new refinement at A forces a pivot on the boundary shared with B , eliminating joint histories that contradict the updated record. No signal travels between the laboratories; no mechanism carries information. The ledger simply performs the minimal boundary repair required by Martin’s Condition. The observed “nonlocal” correlations are the bookkeeping consequence of enforcing a single, globally consistent causal ordering.

Thus, the Bell–Aspect tests reveal that entanglement is not a hidden influence. It is the fact that the causal universe must repair its boundary globally when a non-degenerate refinement is recorded. Local hidden variables fail because they deny the existence of this global pivot.

Phenomenon 4.3.6 (Hawking Radiation Revisited). ***N.B.**—No physical emission is assumed. Surrogate refinements are bookkeeping: the minimal distinctions required to restore Martin consistency when the boundary saturates.* \square

An external laboratory maintains a causal universe tensor \mathbf{U}^{out} recording all admissible events visible from outside a black hole. The horizon H is the frontier of distinguishability: an informational boundary beyond which no finite extension of \mathbf{U}^{out} can include internal and external refinements in a single, Martin-consistent ordering. Events remain locally finite, but the reconciliation problem saturates: the external ledger cannot compute a consistent extension that includes both sides.

As an infalling system approaches H , its internal refinements accelerate. By Axiom 5, \mathbf{U}^{out} may not erase distinctions it has already recorded; by Axiom 6, it may not invent invisible refinements. When the bridge of admissible overlap collapses—when no joint ordering of internal and external updates remains feasible—the external ledger must perform a repair. Martin’s Condition demands a globally consistent ordering on the accessible side.

The repair introduces surrogate refinements e_{rad} :

$$\mathbf{U}_{n+1}^{\text{out}} = e_{\text{rad}} \mathbf{U}_n^{\text{out}},$$

a compensatory update that restores coherence without referencing inaccessible events. These surrogates are not particles escaping from behind the horizon; they are the unique refinements that preserve global order when the boundary can no longer reconcile the missing interior. The exponential spectrum attributed to Hawking radiation reflects the combinatorial multiplicity of admissible surrogate updates once the informational channel saturates.

Thus, Hawking radiation is not a quantum field effect in curved space-time. It is the minimal bookkeeping required to maintain Martin consistency on the visible side of an informational boundary. The horizon enforces a holographic constraint: global order must remain representable on the surface

that separates what can be reconciled from what cannot.

4.4 The Law of Boundary Consistency

Every example in this chapter has the same structure. When a new admissible refinement is recorded, the ledger does not alter the interior of the accumulated state. Instead, it repairs only the frontier where two descriptions overlap. The Causal Folding Operator updates the boundary and leaves the interior fixed. This pattern is universal and admits a formal statement.

Law 3 (The Law of Boundary Consistency). *In any locally finite causal domain, every admissible update to the accumulated causal universe tensor \mathbf{U} arises from boundary refinement. The interior of \mathbf{U} is fixed by previously recorded distinctions: altering it would introduce an invisible refinement (Axiom 6) or remove a recorded one (Axiom 5), both of which are forbidden. When a new admissible event is observed, the ledger repairs only the frontier where two descriptions overlap, enforcing Martin’s Condition on the boundary of the accumulated state.*

Therefore all dynamics—propagation, interaction, interference, and decay—are the shadows of boundary reconciliation. Nothing propagates through the interior; motion is the smooth limit of reconciling admissible distinctions at the frontier of \mathbf{U} .

Remark 16.

No interior modification. *Once folded, the interior of \mathbf{U} contains no unobserved structure. Any change to it would imply either an invisible refinement or the erasure of a recorded one, violating Planck or Cantor.*

Minimal repair. *When ledgers overlap, the operator updates only the smallest region where a contradiction could occur. This is a boundary operation, not a volume operation.*

Computability. *Martin's Condition is enforced by checking only the joint frontier: the causal surface where two descriptions must agree. No global search or re-evaluation of the interior is required.*

Operational meaning. *Waves, interference, scattering, advection, and diffusion appear in the smooth limit of boundary reconciliation. The equations of motion arise from the unique completion that preserves the folded boundary without altering the interior.*

This law closes the algebra of interaction. The Causal Folding Operator enforces global consistency by repairing only the frontier of the accumulated state. Every dynamic phenomenon considered in this chapter—the Dantzig pivot of entanglement, the Mach–Zehnder interference fold, the Bell–Aspect repair, and the surrogate refinements of a causal horizon—is an instance of the same rule: the ledger changes only at the boundary.

This statement is the discrete analogue of Gauss's Theorem. In the continuum, specifying the value of a field on a closed boundary determines its interior uniquely. The Law of Boundary Consistency asserts the same principle for causal ledgers: every admissible refinement enters through the frontier where two descriptions overlap, and the interior is fixed by previously recorded distinctions. Nothing propagates through the volume of \mathbf{U} ; every update is a boundary repair.

All examples in this chapter—velocity boosts, interference, entanglement, and surrogate events near a causal horizon—share this structure. A new admissible event forces only the minimal reconciliation on the overlap. The interior never changes. Motion is the continuum shadow of this purely discrete principle.

At this point nothing further is required. Once every admissible update is confined to the boundary, the smooth limit follows automatically: the interior is fixed, and all variation arises from finite differences on the frontier. The familiar equations of motion are just the continuum shadow of these discrete boundary repairs. Writing them down is a matter of expressing the boundary

updates in finite-difference form and passing to the smooth limit.

4.5 Qubit Decoherence

The language of “coherence” and “decoherence” originates in the physical literature, where it refers to the loss of phase relations between components of a quantum state[166]. In standard treatments, this loss is attributed to dynamical interactions with an external environment, often modeled through diffusion, noise, or stochastic drift. Although the present framework makes no physical or geometric assumptions of this kind, the terminology remains useful. What is called “decoherence” here is the purely informational process by which a locally admissible degeneracy is resolved when new measurements are recorded. The mechanism is not environmental coupling, but the logical requirement that admissible refinements remain consistent under Martin’s Condition and Axiom 3. The resulting collapse of a causal doublet is therefore an informational phenomenon: a pattern that emerges whenever distinguishable events are appended to a degenerate causal record. Its observed “rate” is a smooth shadow of the stochastic drift inherent in finite causal resolution, and not a dynamical property of any physical substrate.

Phenomenon 4.5.1 (Qubit Decoherence [82, 166]). ***N.B.**—This informational phenomenon does not rely on physical decoherence mechanisms, environmental coupling, or geometric dynamics. It arises solely because measurements are recorded and admissible refinements must remain consistent with the axioms of event selection, refinement compatibility, and Ockham minimality.* \square

A causal doublet is the minimal unit of informational degeneracy: a system admitting two equally admissible refinement paths $S = \{e_0, e_1\}$. Such a structure represents a qubit in the informational sense: a pair of distinct updates that are locally indistinguishable and jointly admissible.

Decoherence occurs when a new event is recorded that is inconsistent with

one of the branches. The Interaction Operator f performs a pivot on the shared boundary, eliminating all incompatible orderings and collapsing the doublet to a single admissible history. This collapse satisfies Martin's Condition, ensuring that the refined ledger extends the earlier one without introducing new admissible distinctions.

The observed rate of this collapse is a smooth shadow of two underlying informational constraints:

1. **Finite Causal Resolution.** Irreducible uncertainty in the ordering of micro-events at scale Δx induces a stochastic drift in the admissible refinements. This drift arises whenever unresolved orderings accumulate faster than they can be anchored by distinguishable events.
2. **Informational Diffusion (D).** The propagation of unresolved distinctions obeys a diffusion law: coarse records evolve stochastically under refinement, with an effective diffusion coefficient D determined by the informational bandwidth of the system.

Together, these constraints imply that decoherence is the statistical failure to maintain a causal degeneracy in the presence of new distinctions. The macroscopic decoherence rate emerges as the smooth shadow of this irreversible informational process and is governed by the informational diffusion coefficient D and the minimal unresolved action \hbar . No physical environment or geometric postulate is required.

Proposition 9 (The Rate of Informational Decoherence). *Let a causal doublet consist of two equally admissible refinement paths $S = \{e_0, e_1\}$. Let unresolved micro-orderings accumulate at an average rate λ per unit refinement depth, and let informational diffusion have coefficient D . The probability that the doublet remains unresolved after refinement depth t is*

$$P_{\text{coh}}(t) = \exp(-\gamma t),$$

where the informational decoherence rate is the product

$$\gamma = \frac{\lambda^2}{2D}.$$

N.B.—A complete derivation of the decoherence rate is deferred until the end of the chapter, where informational Brownian motion is developed. The rate law arises as a first-passage property of unresolved refinements undergoing informational diffusion. In the smooth shadow this corresponds to the classical diffusion equation, and Ito-style arguments become available. The derivation given later relies on these stochastic tools and therefore is not presented at this stage. \square

4.6 Newtonian Transport

N.B.—Nothing in this construction asserts that a differential equation *must* govern the data. We show only that if the ledger admits a smooth completion consistent with the axioms, then the corresponding differential equation appears as its unique smooth shadow. The calculus is a consequence of measurement consistency, not an independent postulate. \square

Classical transport is the process by which refinement differences reconcile across space. In the discrete ledger, this appears as iterated boundary smoothing: sharp discontinuities trigger local folds until no admissible repair remains. In the smooth limit, these reconciliation rules generate the transport equations of classical thermodynamics. The organizing principle is the variational order of the correction.

4.6.1 First Variation: Slope-Level Ledger Corrections

First-variation updates alter only the slope of the admissible spline representation. Informational minimality forbids the creation of new turning points between event anchors: any correction that introduced a fresh extremum

would constitute an unrecorded event. All admissible first-order updates are therefore monotone. Their smooth limit yields irreversible transport.

Annealing and Conduction (Symmetric Reconciliation)

Conduction appears when a ledger repeatedly reconciles a coarse description of itself. A sharp difference in refinement counts across a boundary triggers a sequence of local folds, each of which reduces the discrepancy without altering the interior. This iterative process is *annealing*: informational tension is monotonically released until no further repair is admissible.

Under the Law of Spline Sufficiency, symmetric reconciliation introduces no oscillation and no hidden curvature. The discrete flux is governed by the centered jump between neighboring cells, and the update rule is a symmetric projection back into the admissible class. In the smooth limit, these finite differences converge to the classical diffusion equation.

Discrete Ledger Update and the Flux Form. Let u_i^k denote the normalized refinement count recorded on cell i at discrete time t_k , with spatial spacing Δx and time step Δt . The update must obey informational conservation in a conservative flux form:

$$u_i^{k+1} = u_i^k - \frac{\Delta t}{\Delta x} (F_{i+\frac{1}{2}}^k - F_{i-\frac{1}{2}}^k). \quad (4.2)$$

Symmetric reconciliation uses the centered jump as the flux. If κ is the informational diffusion coefficient,

$$F_{i+\frac{1}{2}}^k = -\kappa \frac{u_{i+1}^k - u_i^k}{\Delta x}. \quad (4.3)$$

Substituting (4.3) into (4.2) yields the standard symmetric smoothing rule:

$$u_i^{k+1} = u_i^k + \frac{\kappa \Delta t}{\Delta x^2} (u_{i+1}^k - 2u_i^k + u_{i-1}^k). \quad (4.4)$$

Proof Sketch: Convergence to $u_t = Du_{xx}$. Approximate the temporal derivative using a forward difference:

$$u_t(x_i, t_k) \approx \frac{u_i^{k+1} - u_i^k}{\Delta t}.$$

Substituting (4.4) and rearranging,

$$\frac{u_i^{k+1} - u_i^k}{\Delta t} = \frac{\kappa}{\Delta x^2} (u_{i+1}^k - 2u_i^k + u_{i-1}^k).$$

The spatial term on the right is the standard centered approximation of the second derivative,

$$u_{xx}(x_i, t_k) \approx \frac{u_{i+1}^k - 2u_i^k + u_{i-1}^k}{\Delta x^2}.$$

Thus

$$u_t(x_i, t_k) = \kappa u_{xx}(x_i, t_k).$$

Taking the continuous limit $\Delta x, \Delta t \rightarrow 0$ and letting $\kappa \rightarrow D$ yields the diffusion equation

$$u_t = D u_{xx}.$$

□

The convergence is admissible because the Law of Spline Sufficiency guarantees that the solution remains \mathcal{C}^2 and introduces no hidden curvature. The symmetric finite-difference update is therefore a monotone, stable smoothing process: the smooth shadow of informational annealing.

Convection and Oriented Transport (Boundary Consistency)

Convection models the directed transport of distinctions, where the orientation of the flow is realized as a preferred direction in the causal refinement process. When a boundary carries an orientation, reconciliation must respect

that direction: smoothing from the downstream side would create unrecorded structure on the wrong side of the interface.

Oriented Boundary Reconciliation. Let u_i^k be the normalized refinement count on cell i at time t_k . When the interface $(i, i+1)$ has a known inflow direction, the Law of Boundary Consistency requires that the ledger flux across that interface be determined solely by the state on the inflow side:

$$F_{i+\frac{1}{2}}^k = c u_i^k, \quad (4.5)$$

where c is the order speed. Substituting (4.5) into the conservative update

$$u_i^{k+1} = u_i^k - \frac{\Delta t}{\Delta x} (F_{i+\frac{1}{2}}^k - F_{i-\frac{1}{2}}^k) \quad (4.6)$$

yields the upwind rule

$$u_i^{k+1} = u_i^k - \frac{c \Delta t}{\Delta x} (u_i^k - u_{i-1}^k). \quad (4.7)$$

Proof Sketch: Convergence to $u_t + c u_x = 0$. Divide (4.7) by Δt to obtain

$$\frac{u_i^{k+1} - u_i^k}{\Delta t} = -c \frac{u_i^k - u_{i-1}^k}{\Delta x}.$$

As $\Delta t, \Delta x \rightarrow 0$, the left side is the forward difference approximation of the time derivative $\partial_t u$, and the right side is the backward difference approximation of the space derivative $\partial_x u$. Taking the smooth limit yields the advection equation

$$u_t + c u_x = 0.$$

□

The update (4.7) is admissible only when it remains monotone, which is guaranteed by the CFL condition $0 \leq c \Delta t / \Delta x \leq 1$. Under this constraint

no new turning points are introduced, so the Law of Spline Sufficiency is respected: the directed transport is a projection back into the admissible spline class.

N.B.—Boundary Consistency selects the upwind flux, and Spline Sufficiency forbids oscillatory corrections; the advection equation is the smooth shadow of oriented ledger reconciliation. \square

Advection–Diffusion (Mixed Closure)

In many settings, admissible reconciliation requires both symmetric homogenization and directed transport. The ledger must smooth local inconsistencies while simultaneously respecting boundary orientation. The resulting update combines the symmetric and upwind fluxes.

Combined Flux. Let the oriented flux be given by

$$F_{i+\frac{1}{2}}^{\text{adv}} = c u_i^k,$$

and the symmetric flux by

$$F_{i+\frac{1}{2}}^{\text{diff}} = -\kappa \frac{u_{i+1}^k - u_i^k}{\Delta x}.$$

The total flux across the interface is their sum:

$$F_{i+\frac{1}{2}}^k = c u_i^k - \kappa \frac{u_{i+1}^k - u_i^k}{\Delta x}. \quad (4.8)$$

Substituting (4.8) into the conservative update

$$u_i^{k+1} = u_i^k - \frac{\Delta t}{\Delta x} (F_{i+\frac{1}{2}}^k - F_{i-\frac{1}{2}}^k) \quad (4.9)$$

yields the discrete advection–diffusion rule

$$u_i^{k+1} = u_i^k - \frac{c \Delta t}{\Delta x} (u_i^k - u_{i-1}^k) + \frac{\kappa \Delta t}{\Delta x^2} (u_{i+1}^k - 2u_i^k + u_{i-1}^k). \quad (4.10)$$

Proof Sketch: Convergence to $u_t + c u_x = D u_{xx}$. Divide (4.10) by Δt to obtain

$$\frac{u_i^{k+1} - u_i^k}{\Delta t} = -c \frac{u_i^k - u_{i-1}^k}{\Delta x} + \kappa \frac{u_{i+1}^k - 2u_i^k + u_{i-1}^k}{\Delta x^2}.$$

In the limit $\Delta x, \Delta t \rightarrow 0$, the left side becomes $\partial_t u$, the first term becomes $-c \partial_x u$, and the second becomes $\kappa \partial_{xx} u$. Setting $D = \kappa$ gives

$$u_t + c u_x = D u_{xx},$$

the advection–diffusion equation. \square

The mixed closure is the most general first-order reconciliation of the refinement record. Information spreads down gradients (diffusion) while coherent packets of distinction are carried along oriented interfaces (advection). The process is irreversible in either mode, and no additional structure is assumed beyond the slope-level correction forced by the axioms.

N.B.—In every case, first-variation closure is a projection back into the admissible spline class: no new extrema are introduced, and no hidden structure appears under refinement. The differential equations are the smooth shadows of monotone reconciliation. \square

4.6.2 Second Variation: Curvature-Level Ledger Corrections

Second-variation updates alter curvature while preserving slope and anchor values. These corrections are reversible: they propagate distinctions without loss and produce no additional smoothing. Their smooth limit yields wave transport.

Radiation (Symmetric Curvature Smoothing)

Radiation represents the propagation of distinction at the maximal admissible speed. Unlike the first-order corrections of Sections 4.6.1–4.6.1, radiation is reversible: once the ledger has reconciled curvature symmetrically, no net informational gain or loss remains. The process is the smooth shadow of *symmetric curvature smoothing*.

Vanishing Second Variation. Let \mathcal{A} denote the amplitude of distinction recorded over a finite causal neighborhood. The second variation $\delta^2\mathcal{A}$ measures the change in \mathcal{A} under two sequential, infinitesimal perturbations of the record. Radiation occurs when these perturbations commute exactly:

$$\delta^2\mathcal{A} = 0. \quad (4.11)$$

No net expansion or contraction of distinguishability can remain; curvature differences are repaired symmetrically and without directional bias. This is the reversible complement of annealing: where first-order correction removes slope-level inconsistencies, second-order correction removes curvature-level tension.

Discrete Curvature Laplacian. In the discrete domain, the sum of all pairwise second variations over neighboring events defines the discrete Laplacian on event sets:

$$\nabla_E^2\mathcal{A} = \sum_{f \in \text{Nbr}(e)} (\mathcal{A}(f) - \mathcal{A}(e)).$$

Martin’s Condition enforces that this curvature vanishes:

$$\nabla_E^2\mathcal{A} = 0, \quad (4.12)$$

so that symmetric curvature smoothing is locally maximal and globally neutral.

Smooth Shadow. In the continuum limit, the second-order symmetric closure converges to the homogeneous wave equation. If $u(x, t)$ is the smooth completion of the refinement record, then

$$u_{tt} = c^2 u_{xx}, \quad (4.13)$$

where c is the order speed—the combinatorial rate at which causal constraints traverse the event network. Equation (4.13) expresses reversible propagation: local expansions and contractions of distinguishability cancel globally, so that information moves without net amplification or dissipation.

N.B.—Second-variation closure enforces symmetric curvature repair and forbids net informational gain or loss. The wave equation is therefore the unique smooth shadow of reversible curvature smoothing, derived solely from the axioms of causal refinement. \square

Adiabatic Transport (Curvature Invariance)

Adiabatic transport is the ideal limit of reversible motion in the causal record. Distinctions are neither created nor destroyed: informational entropy remains constant, and the curvature of the smooth completion is preserved. This process is the logical dual of annealing, establishing the boundary condition for zero informational work.

Invariance of Distinguishability. Let λ parameterize a smooth evolution of an admissible history $\Psi(\lambda)$. The history undergoes adiabatic transport when the informational entropy is invariant:

$$\frac{d}{d\lambda} \mathcal{S}(\Psi) = 0. \quad (4.14)$$

Equivalently, the update operator satisfies

$$U_{\lambda+\delta\lambda} = U_{\lambda} + \mathcal{O}(\delta\lambda^2),$$

so the leading-order change in the refinement record vanishes. The motion is norm-preserving and informationally reversible: the ledger drifts without loss of distinction.

Curvature Invariance. Because \mathcal{S} counts admissible configurations, the condition (4.14) forces the evolution to proceed along a path of constant informational curvature. Locally,

$$\frac{d}{d\lambda} \Psi'' = 0, \tag{4.15}$$

so that no curvature-level tension is released or accumulated. This is the reversible complement to the symmetric curvature smoothing of Section 4.6.2.

Smooth Shadow. Under the Law of Spline Sufficiency ($\Psi^{(4)} = 0$), curvature invariance selects the unique extremal that transports distinctions without dissipation: the geodesic or undamped wave. Informational entropy remains constant, and the ledger evolves along the smooth completion Ψ without net repair or decay. Nothing dynamical is postulated; the law is a theorem of informational conservation.

N.B.—Adiabatic transport is the limit of causal motion that preserves informational order. It connects reversible evolution ($d\mathcal{S} = 0$) with the requirement that distinguishability cannot decrease. The geodesic structure is therefore a consequence of informational invariance, not an independent physical postulate. \square

4.7 Quantum Transport

Some transport phenomena do not appear as flows of a substance, but as discrete repairs of nearly degenerate descriptions. When two ledgers support multiple admissible extensions, the Causal Folding Operator must select the unique completion that preserves all recorded distinctions. The familiar quantum effects arise as the smooth shadows of this repair.

4.7.1 Informational Pressure

Phenomenon 4.7.1 (The Casimir effect). *The Casimir effect is the boundary expression of informational pressure. When admissible refinements are restricted by geometry, the ledger must perform a compensatory update to preserve global distinguishability. In the smooth limit, this boundary repair appears as a physical force.*

Boundary–Induced Asymmetry. *Consider two parallel constraints that restrict the admissible causal updates in the interior region. Each admissible field mode corresponds to a distinguishable refinement of the causal record. The plates suppress many of these modes, so the interior ledger records fewer admissible distinctions than the exterior. Outside the plates, no such suppression occurs; the ledger remains unrestricted. This produces an imbalance in refinement counts across the boundary: the exterior supports strictly more admissible updates than the interior.*

Compensatory Boundary Update. *The Second Law of Causal Order requires that global distinguishability must not decrease. The imbalance therefore creates informational tension. Because no additional interior modes are admissible, the only possible repair is a boundary update that restores global consistency without altering the restricted interior. The unique correction is an outward curvature of the boundary ledger: refinements accumulate on the*

exterior frontier, pushing the constraints toward one another.

In the smooth limit, this boundary curvature appears as the Casimir pressure. No mechanical postulate is introduced; the force is the smooth shadow of a compensatory update that restores consistency between the restricted interior and unrestricted exterior ledgers.

N.B.—*In this interpretation, the Casimir effect is a holographic phenomenon: the minimal boundary correction enforced by global distinguishability. The pressure is not a hypothesis about zero-point energy, but the unique repair consistent with the axioms of causal refinement.* \square

4.7.2 Repair of a Causal Contradiction at the Boundary

Alpha decay is the irreversible repair of a causal contradiction on the boundary of the nuclear ledger. The nucleus admits two nearly indistinguishable continuations of its refinement record:

$$\Psi_{\text{bound}} \quad \text{and} \quad \Psi_{\text{unbound}}.$$

Both are initially admissible: each agrees with all external anchors and differs only within a bounded interior neighborhood.

Phenomenon 4.7.2 (The Alpha-Decay Effect). *Over informational time, unresolved curvature accumulates and the two ledgers drift out of alignment. Their boundary descriptions become incompatible with Martin Consistency: the overlap cannot be reconciled without introducing unrecorded structure. A repair is required to preserve the global order of the causal record.*

The Causal Folding Operator f performs the minimal corrective update by removing the inconsistent branch:

$$f : \Psi_{\text{bound}} \longrightarrow \Psi_{\text{unbound}} + \alpha.$$

The emitted alpha particle is the recorded trace of this boundary repair. The interior ledger returns to an admissible configuration, and the causal record evolves on the remaining branch.

In the continuum limit, the finite differences of this irreversible repair produce the exponential law of radioactive decay. No hidden forces or tunneling mechanism is assumed: alpha decay is the unique boundary update that eliminates a causal contradiction while preserving global distinguishability.

N.B.—*Alpha decay is the irreversible removal of an inconsistent branch from the refinement record. The emitted particle is the holographic trace of the boundary correction, not a postulated tunneling object.* \square

4.7.3 Restoration of Causal Symmetry

Phenomenon 4.7.3 (The Gamma Decay Effect). *Gamma decay is a reversible repair of internal causal symmetry. An excited nuclear state corresponds to an admissible configuration whose internal refinement record is nearly, but not exactly, consistent with the minimal ground state. Over time, unresolved curvature accumulates, producing a small informational asymmetry in the internal ledger.*

Informational Synchronization. *Let Ψ^* denote the smooth completion of the excited state and Ψ that of the ground state. Both are admissible: they agree on all external anchors and differ only in a bounded internal neighborhood. The difference is a phase drift in the internal causal partition—a small curvature that violates informational minimality. The nucleus must perform a repair that restores the unique, globally consistent ground state.*

The minimal symmetric repair is the emission of a gamma photon:

$$\Psi^* \longrightarrow \Psi + \gamma.$$

The photon is the propagated correction: a reversible wave of order that car-

ries the excess curvature away from the nucleus while leaving the internal ledger in its minimal configuration.

Zero–Mass Boundary Repair. *Unlike alpha decay (Section 4.7.2), which removes an entire inconsistent branch from the record, gamma decay preserves the identity of the nucleus. It is informationally reversible: no new branches are created, and no admissible distinctions are destroyed. The process is the smooth shadow of symmetric curvature repair:*

$$\delta^2 \mathcal{A} = 0 \quad \implies \quad \text{emission of } \gamma \text{ with } E = h\nu.$$

The energy of the photon measures the amount of curvature removed from the internal ledger. No mechanical postulate is required; gamma decay is the unique boundary update that restores global distinguishability without altering the underlying causal identity of the system.

N.B.—*In this interpretation, gamma decay is not a force-mediated transition, but a minimal holographic correction: a reversible synchronization event that propagates excess curvature as a photon and restores Martin Consistency in the internal ledger without altering the causal identity of the nucleus. \square*

4.7.4 Quantum Informational Pressure

Phenomenon 4.7.4 (The Brownian Motion Effect). *Brownian motion can be interpreted as a quantum informational phenomenon in the present framework. The source of randomness is not mechanical noise but finite causal resolution: each refinement step leaves a family of equally admissible micro-orderings that the ledger cannot distinguish. The coarse record therefore evolves stochastically.*

Stochastic Reconciliation at Finite Resolution. *Let u_i^k be the normalized refinement count on cell i at time t_k . When the observer cannot resolve*

all admissible distinctions at scale Δx , the symmetric smoothing update acquires an irreducible stochastic term:

$$u_i^{k+1} = u_i^k + \frac{\kappa \Delta t}{\Delta x^2} (u_{i+1}^k - 2u_i^k + u_{i-1}^k) + \sqrt{2D \Delta t} \xi_i^k, \quad \mathbb{E}[\xi_i^k] = 0, \quad \mathbb{E}[(\xi_i^k)^2] = 1. \quad (4.16)$$

The deterministic part is the symmetric reconciliation enforced by the Law of Spline Sufficiency; the random term is the ledger's irreducible uncertainty at the observation scale.

Smooth Shadow: Diffusion as Quantum Measure. *Under refinement $\Delta x, \Delta t \rightarrow 0$ with D fixed, the central limit theorem implies convergence of (4.16) to the diffusion equation for the coarse density $u(x, t)$:*

$$u_t = D u_{xx}. \quad (4.17)$$

Here D is the informational diffusion coefficient: the effective bandwidth of unresolved distinctions per unit time.

Bridge to Schrödinger via Analytic Continuation. *The free Schrödinger equation is related to diffusion by analytic continuation of time. Setting $D = \frac{\hbar}{2m}$ and $t \mapsto -it$ maps (4.17) to*

$$i \hbar \partial_t \Psi = - \frac{\hbar^2}{2m} \partial_{xx} \Psi, \quad (4.18)$$

i.e., the smooth shadow of unresolved, symmetric refinement at fixed informational bandwidth equals the quantum free evolution with Planck scale \hbar . In this sense, Brownian motion is quantized uncertainty: \hbar calibrates the minimal unresolved action, while D measures the rate at which that unresolved structure propagates statistically.

Consistency with the Two Laws. - Spline Sufficiency *ensures no spurious extrema: the stochastic update remains a projection into the admissible class almost surely.* - Boundary Consistency *fixes oriented interfaces; adding an upwind drift c to (4.16) yields the standard advection–diffusion (Fokker–Planck) limit.*

N.B.—*This construction shows how quantum evolution can arise from measurement limits: if the ledger’s unresolved bandwidth D is fixed by a Planck scale, diffusion analytically continues to Schrödinger dynamics. It does not assert that nature must realize this identification in every regime.* \square

4.8 First Quantization as an Application of the Two Laws

The classical picture of quantization treats the wavefunction, Hilbert space, and operator algebra as new physical axioms. In the present framework they arise automatically from the two kinematic consistency laws:

- **Law of Spline Sufficiency:** no admissible refinement may introduce unrecorded structure; smooth closure is \mathcal{C}^2 and satisfies $\Psi^{(4)} = 0$,
- **Law of Boundary Consistency:** oriented boundaries must be reconciled from the inflow side; no correction may propagate across a boundary in the wrong direction.

Together, these laws force the structure known in physics as *first quantization*. Nothing new is added: the quantized theory is the smooth shadow of informational bookkeeping.

4.8.1 Hilbert Structure from Spline Closure

Under Spline Sufficiency, every admissible history has a unique smooth representative Ψ that is cubic between anchors and \mathcal{C}^2 globally. Any two admis-

sible histories Ψ and Φ differ only in their recorded curvature. Their overlap is therefore measured by the curvature functional

$$\langle \Psi, \Phi \rangle = \int \Psi''(x) \Phi''(x) dx.$$

This inner product is positive definite on the admissible class and yields a complete inner-product space: the Hilbert space of admissible closures. The “wavefunction” is nothing more than Ψ viewed as an element of this space.

4.8.2 Canonical Structure from Boundary Consistency

The curvature functional determines a unique conjugate operator. Integration by parts yields

$$\langle \Psi, x \Phi \rangle - \langle x \Psi, \Phi \rangle = \int \Psi(x) \Phi'(x) dx,$$

where the boundary term is fixed in sign by the inflow rule of Boundary Consistency. The operator that realizes this antisymmetry is

$$\hat{p} = -i \partial_x,$$

the momentum operator of canonical quantization. No new axiom is required: the oriented boundary rule uniquely determines the self-adjoint generator of translations.

Phenomenon 4.8.1 (The Momentum Effect). *Momentum is the operator that enforces boundary consistency. It is the canonically conjugate bookkeeping term that guarantees admissible inflow of refinements across a causal boundary. Without it, the ledger would admit unaccounted refinement debt.*

Momentum is therefore not motion itself, but the enforcement of admissible exchange.

4.8.3 Energy Levels from Informational Minimality

Consider an admissible history constrained by a restoring boundary (a fold that always returns toward the anchor). Under Spline Sufficiency the closure is cubic between anchors and $\Psi^{(4)} = 0$; under Boundary Consistency the inflow rule forces the curvature to alternate monotonically between turning points. The Galerkin limit of this curvature balance is the harmonic oscillator:

$$-\Psi''(x) + x^2\Psi(x) = \lambda\Psi(x),$$

whose eigenvalues are discrete because no new turning points may be added between anchors. The spectrum is the familiar

$$\lambda_n = (2n + 1), \quad n = 0, 1, 2, \dots$$

Quantization is therefore a *restriction of admissible curvature*, not a postulate about nature.

4.8.4 Summary

- Spline Sufficiency \Rightarrow Hilbert space of smooth closures,
- Boundary Consistency \Rightarrow canonical commutators,
- Discrete curvature balance \Rightarrow quantized energy levels.

$$\text{finite ledger} \xrightarrow{\text{spline closure}} \Psi \xrightarrow{\text{boundary consistency}} \hat{x}, \hat{p} \xrightarrow{\text{curvature balance}} \text{quantized energies.}$$

Thus the apparatus of “first quantization” is not a new physics. It is the smooth bookkeeping of the two kinematic laws applied to finite informational records.

N.B.—In this sense, quantization is not an independent hypothesis. It is the minimal correction rule forced by informational sufficiency and boundary orientation. \square

4.9 Resolution of Qubit Decoherence

The analysis of informational decoherence highlights a recurring theme in this framework: when a finite record is refined, the admissible continuous extensions must adjust in ways that are not captured by ordinary deterministic calculus. Each refinement introduces new distinctions that must be merged with the existing record, and the comparison between the old and new minimal extensions reveals systematic second-order effects. These effects do not arise from physical noise or stochastic input; they are forced by the axioms of refinement compatibility, informational minimality, and Martin consistency.

Whenever a quantity is represented by its minimal spline extension, the act of incorporating a new event alters not only the value of the interpolant but also its curvature. The discrepancy between the old and new extensions produces a correction term whose structure is universal: it depends only on the geometry of minimal refinements, not on the nature of the underlying system. In classical settings this correction is masked by probabilistic notation, but in the informational setting it emerges as an intrinsic feature of refinement itself.

The phenomenon described below captures this behavior. It is the general informational form of what, in conventional stochastic calculus, appears as Itô's Lemma.

Phenomenon 4.9.1 (Itô's Lemma [78, 79]). **N.B.**—*Itô's Lemma appears here not as a theorem of stochastic calculus, nor as a property of diffusion processes, but as a structural consequence of informational refinement. When a finite record is repeatedly refined, the admissible interpolants must update according to Martin consistency and Ockham minimality. These updates*

produce the same correction terms that, in classical settings, are associated with stochastic differentials. No probabilistic or physical assumptions are used; the result is purely algebraic. \square

Let X_t denote the minimal continuous extension of a finite record obtained by Spline Sufficiency. Suppose that between two refinements, the record admits a locally smooth representation

$$X_{t+\Delta t} = X_t + \Delta X_t.$$

Refinement compatibility requires that any function $f(X_t)$ be updated by comparing the old and new admissible extensions. The refinement

$$f(X_{t+\Delta t}) - f(X_t)$$

must be consistent with the joint refinement of X_t and f under the axioms of order, minimality, and Martin consistency. Expanding to second order in the refinement step and discarding inadmissible terms produces

$$df = f'(X_t) dX_t + \frac{1}{2} f''(X_t) (dX_t)^2,$$

where $(dX_t)^2$ is the second-order correction forced by the comparison of successive minimal interpolants. This quadratic term is not a physical noise term but an informational artifact: the unavoidable discrepancy between two successive minimal refinements of the same record.

Thus Itô's Lemma arises as the continuous shadow of discrete, consistent refinements of observational data.

4.9.1 Informational Decoherence as Forced Refinement

Decoherence is treated here not as a physical process, nor as an interaction with an environment, but as the informational consequence of refining a causal record. When new distinctions are appended to a history, the minimal

continuous extension of that history must be updated in accordance with the axioms of refinement compatibility, Ockham minimality, and Martin consistency. The resulting adjustment introduces a second-order correction identical in structure to the Itô term that appears in stochastic calculus, though no probabilistic or physical assumptions are made.

We now present the proof to Proposition 9

Proof (Sketch). Consider a causal doublet (X_t, Y_t) whose minimal extension is given by the spline interpolant determined by the current observational record. Before refinement, the joint extension encodes the admissible correlations between X_t and Y_t . When a new event is added to the record, the refinements

$$X_{t+\Delta t} = X_t + \Delta X_t, \quad Y_{t+\Delta t} = Y_t + \Delta Y_t$$

must be merged into a single globally coherent history.

By Spline Sufficiency, the new extension is the unique minimal function matching all observations. The comparison between the old and new extensions yields

$$d(XY) = X_t dY_t + Y_t dX_t + (dX_t)(dY_t),$$

where the cross-term $(dX_t)(dY_t)$ is the informational correction forced by the discrepancy between successive minimal interpolants. This term does not represent physical noise; it is the algebraic signature of refinement.

When the refinements of X_t and Y_t are uncorrelant under the refinement order, the cross-terms collapse in the merge and the joint extension factorizes. The apparent “loss” of coherent structure is thus an informational effect: the minimal extension can no longer sustain the curvature required to preserve the off-diagonal components of the doublet. \square

A full proof is provided in Appendix A.8.

In conventional quantum language this behavior is described as decoherence. In the informational setting it is a direct consequence of how causal

records are updated: coherence is maintained only when the refinement structure supports the cross-terms required by the minimal extension. When refinements fail to align, these terms vanish, and the record resolves into independent components.

Thus decoherence arises not from dynamics but from the combinatorics of consistent refinement.

Coda: Orbits

Before introducing the informational harmonic oscillator, it is worth noting that nothing genuinely new is being added. The construction does not assume a force, a potential, or a dynamical law. It is simply the closed loop of reciprocity already present in the calculus of motion: recorded distinction feeds the prediction of its own refinement, and that prediction, when made admissible, returns to update the record.

When this reciprocal exchange is traced around a single loop, the result can only be periodic. A reversible refinement cycle has nowhere to go; it merely propagates its informational content back into itself. The oscillator is therefore the minimal self-consistent refinement process—a bookkeeping loop that preserves its own measure while shuttling distinguishability between its record and prediction components.

What follows is not a physical oscillator but the simplest closed circuit of informational propagation permitted by the axioms. In the informational framework, prediction is the purpose of the differential equations of physics. A differential equation is nothing more than a rule for extending a record: it specifies how a small, admissible refinement of the present state must constrain the next distinguishable event. Laboratory experiments exploit this fact directly. By preparing controlled initial conditions and observing how a system responds to a tiny perturbation, the experimenter samples the local refinement structure encoded by the governing equation. The resulting

data do not unveil a hidden dynamical mechanism; they merely reveal how the differential law organizes small predictions into a coherent chain of distinguishable events. In this sense, every differential equation is a predictive device: a compact description of how an observer may extend the current record without contradiction.

Definition 30 (Prediction [44, 67, 73, 92, 106, 113, 135] et alii plures). ***N.B.**—The formulation of prediction as an inverse update draws on a long tradition of differential equations, whose modern corpus reflects centuries of mathematical effort. From Newton’s original method of fluxions through the developments of Euler, Lagrange, Cauchy, Riemann, Hilbert, Noether, and countless others, differential equations have served as the principal tools for expressing how small refinements constrain admissible continuation. The informational framework does not alter or reinterpret this body of work; it simply recognizes that the purpose of these equations has always been predictive. They encode how an observer may extend a record without contradiction. The present treatment stands on the shoulders of these and countless other historical achievements and uses classical forms only as the smooth shadow of the discrete axioms of measurement.* \square

***N.B.**—The argument above demonstrates the necessary existence of an inverse refinement operator Ψ^{-1} in the informational sense: it identifies the set of admissible preimages that, if selected as future events, preserve consistency with the existing record. No analytic inverse is required. The existence follows from the axioms of event selection, refinement compatibility, and global coherence, all of which operate on finite combinatorial data.*

Because these axioms do not depend on smooth structure, continuum limits, or geometric assumptions, the construction applies without modification to finite, agent-based models. Any system in which agents record distinguishable events and update their local states through restricted refinements admits the same inverse-update mechanism. The operator Ψ^{-1} therefore exists in every finite, discrete setting that satisfies the informational axioms, and the re-

sults extend directly to agent-based dynamics without additional assumptions.

□

Let e_k denote the most recent recorded event, and let U_k be its continuous representation under the update rule

$$U_{k+1} = \Psi\left(e_{k+1} \cap \hat{R}(e_k)\right) U_k.$$

A prediction is the admissible pre-image of the next update under Ψ . Formally, a prediction is an element $p_k \in \hat{R}(e_k)$ such that

$$\Psi(p_k) U_k$$

represents the expected refinement of the current record. In this sense, prediction is the inverse action of the update operator: it identifies those refinements which, if later selected as events, will preserve consistency with all prior records. No physical evolution is implied; prediction is the logical anticipation of admissible extensions of the causal history.

The appearance of a Hilbert space at this stage is not an additional postulate but the completion of the spline calculus developed in Chapter 3. Under the Law of Spline Sufficiency, every admissible history admits a unique smooth representative Ψ that is cubic between anchors and C^2 globally; any two such histories differ only in their recorded curvature. Their overlap is measured by the curvature functional

$$\langle \Psi, \Phi \rangle = \int \Psi''(x) \Phi''(x) dx,$$

which is positive definite on the admissible class and therefore defines a norm on the space of smooth closures. When combined with the Law of Discrete Spline Necessity, this norm controls the entire refinement process: every admissible record generates a refinement-compatible sequence of discrete closures (Ψ_N) that converges monotonically

toward a unique spline attractor Ψ , and no admissible refinement can increase the curvature content without violating informational minimality or the Planck bound on resolution. In the curvature norm, these refinement sequences are Cauchy by construction, so the curvature functional supplies precisely the limiting structure required for a Hilbert space: the completion of the admissible closures with respect to informational minimality and refinement compatibility.

Once this completion is in hand, the rest of the monograph may employ the standard toolkit of linear operator theory on this informational Hilbert space. Operators that arise from bookkeeping of curvature, transport, and boundary corrections can be analyzed using the familiar language of adjoints, spectra, and stability, exactly as in the classical theory of matrix computations and linear operators [62]. These results are used only as mathematical theorems about the curvature inner product and its induced operators; they introduce no new axioms of physics and do not supply any geometric interpretation beyond the informational structure already fixed by the Axioms of Measurement.

N.B.— The Hilbert structure employed in this work is not assumed and is not given any geometric interpretation. It is derived solely from the Axioms of Measurement as the unique completion of the spline-refinement space under informational consistency. No metric, manifold, distance, or geometric postulate is introduced. The inner product arises entirely from minimality and refinement coherence, not from geometry. \square

Phenomenon 4.9.2 (The Hilbert Effect). ***Statement.** The space of admissible spline refinements, when completed under prediction and consistency, forms a Hilbert space whose inner product is induced by informational minimality.*

***Description.** Whenever a finite sequence of measurement events is refined into its unique information-minimal spline completion, the set of all admissible refinements inherits a natural vector-space structure. Under the*

dense limit permitted by the Axiom of Cantor, this structure admits a complete inner product. The resulting completion is not assumed but forced: it is the unique Hilbert space compatible with coherent prediction.

In this sense, Hilbert space is not a postulate of physics but the terminal closure of the spline calculus. It arises as the only structure that permits both conservation of informational norm and reversibility of admissible prediction. The inner product is therefore not geometric but informational in origin.

Prediction is thereby identified with the inverse refinement operator Ψ^{-1} acting on this completed space.

The role of linear operator theory in this monograph is strictly informational. It does not enter as a primitive algebraic structure, nor as a geometric assumption. Instead, it appears as a bookkeeping language for the accumulation of finite error in admissible measurement.

Every measurement admitted by the axioms is discrete, finite, and recorded as a distinguishable event. Prediction, refinement, and consistency therefore proceed not in the realm of exact reals, but through sequences of finite updates. When such updates are composed, their imperfections accumulate. It is this accumulation—rather than any geometric structure—that gives rise to linear operators.

Classical linear operator theory, as developed in numerical analysis, is precisely a theory of such accumulated error. The work of Golub and Van Loan [62] formalizes how rounding, truncation, and finite basis representation behave when linear maps are repeatedly applied as in the construction of the Causal Universe Tensor. In this theory an operator is not an ideal transformation but a stable method of propagating approximate information through a finite system. Concepts such as condition number, spectral radius, and stability are not geometric; they measure the rate at which finite inaccuracies amplify or dissipate under iteration.

In the present framework, this viewpoint is fundamental rather than incidental. Measurement itself is a finite computation. Each admissible ex-

tension of the causal ledger introduces a bounded error relative to an ideal refinement, and the Laws of Measurement force these errors to remain coherent under composition. The collection of all admissible refinements, together with their accumulated errors, therefore carries a natural linear structure: composition of refinements behaves additively, and scaling of a finite correction behaves homogeneously. This is not imposed; it is forced by the requirement that measurement error remain globally consistent.

The Hilbert structure enters only at the moment one asks for closure of this error calculus. Minimality (Axiom of Ockham) forbids arbitrary correction, and discreteness (Axioms of Kolmogorov and Planck) forbids infinite exact refinement. As a consequence, admissible refinement sequences must be Cauchy with respect to the norm induced by informational minimality. When these sequences are completed, the space they inhabit is not merely a vector space of finite errors, but a complete one. This completion is the Hilbert effect.

Thus Hilbert space is not assumed, and it is not the foundation of measurement. Measurement comes first. Linear operator theory arises as the bookkeeping of accumulated finite error within that measurement process, and the Hilbert structure appears only because the error must admit a stable, minimal, and globally coherent completion.

Neither can exist alone: without measurement there is no finite error to accumulate; without the completion of error there is no stable predictive structure. Measurement and linear operator theory are therefore not independent layers of description but dual aspects of the same constraint. The Hilbert space is the shadow of measurement consistency, and linear operators are the finite mechanisms by which that shadow is maintained.

It is not enough to explain the phenomenon, one must also explain the noise in the measurement.

Phenomenon 4.9.3 (The Butterfly Effect). *Prediction operates by inverting the refinement update. Because measurements possess finite resolution,*

distinct admissible histories may be recorded as a single event. Their smooth completions diverge over time even though they did not have a measurable distinction at the time of the event.

The divergence of admissible histories is not caused by external stochastic forces, but by the unavoidable noise introduced by finite measurement. Under the Axioms of Kolmogorov and Planck, every recorded event compresses a nontrivial set of admissible microhistories into a single distinguishable record. The information that is not recorded does not disappear; it becomes latent ambiguity in the causal ledger.

This ambiguity is the primitive form of noise.

When refinement is inverted for the purpose of prediction, noise does not remain passive. Each admissible inverse update must choose among histories that were observationally indistinguishable at the time of measurement. These choices propagate the latent ambiguity forward, and admissible completions that were once arbitrarily close separate under repeated refinement.

This separation is not exponential in any geometric sense; it is combinatorial. It counts how many admissible microhistories remain consistent with a coarse record as refinement proceeds.

Prediction Horizon. *The horizon of prediction is therefore not a dynamical limit but an informational one. It is reached when the accumulated noise — the ambiguity inherited from prior coarse measurements — exceeds the refinement bound imposed by information minimality. At that point, multiple next events are compatible with the causal ledger, and no admissible refinement can be chosen without introducing unrecorded structure.*

Beyond this horizon, prediction is undefined.

Interpretation. *In this framework, the Butterfly Effect is not sensitivity to initial conditions. It is sensitivity to unrecorded information. The limit of predictability is set not by chaotic geometry but by the finite nature of measurement itself. Noise is not an error term to be eliminated, but a structural residue that must be carried forward by every admissible history.*

N.B.— The existence of an inner product in the informational completion does not imply the existence of orthogonality in any physical or geometric sense. No orthogonality relations are assumed, derived, or required by the axioms. Any appearance of orthogonal structure belongs solely to later geometric shadows and is not established at this stage of the theory. \square

We now derive the simplest of motion, the informational harmonic oscillator.

4.10 Dissipation

Phenomenon 4.10.1 (The Anderson Effect). *Transport requires the extension of a local refinement into a coherent global pattern. When the local update rules vary incoherently, no admissible global extension exists.*

The failure of propagation is not dissipation, but the impossibility of constructing a minimal, consistent refinement path through disorder.

Phenomenon 4.10.2 (The Harmonic Oscillator [121]). **N.B.**—*This phenomenon describes the minimal reversible dynamics admitted by the axioms of event selection, refinement compatibility, and informational minimality. No metric, geometry, or dynamical law is assumed. Oscillation arises solely from the alternation between recorded distinction and predicted distinction under the reciprocity map.* \square

Consider the two-dimensional informational phase space spanned by a conjugate pair (x, p) , where x records the observer's current distinguishable state and p represents the rate at which that distinguishability is expected to change under an admissible extension. These are not geometric coordinates; they are the dual bookkeeping variables arising from the reciprocity map of Definition 23.

Define the minimal informational action density

$$S(x, p) = \frac{1}{2}(\alpha x^2 + \beta p^2),$$

where $\alpha, \beta > 0$ quantify the informational stiffness and informational inertia enforced by minimality. Stationarity under reversible exchange of (x, p) forces the reciprocal update rules

$$\dot{x} = \beta p, \quad \dot{p} = -\alpha x.$$

Eliminating p yields the continuous shadow

$$\ddot{x} + \omega^2 x = 0, \quad \omega^2 = \alpha\beta.$$

Thus the observer's state executes harmonic motion in informational phase space with invariant $S(x, p)$.

At each turning point the record x is maximal and predictive momentum p vanishes. At each midpoint prediction dominates and the present record is momentarily indeterminate. The system alternately stores and transmits distinguishability, preserving its total informational measure in the reversible limit. No physical oscillation is implied; this is the unique reversible pattern consistent with reciprocal refinement.

Remark 17. *Consequence: Quantization.*

By the Axiom of Planck, only discrete counts of distinguishable refinements fit within one causal cycle. Applying informational minimality to the action produces the familiar spectrum

$$E_n = \hbar\omega \left(n + \frac{1}{2}\right),$$

where n counts the number of admissible informational quanta per cycle. The residual half-count reflects that no finite causal distinction can eliminate the boundary ambiguity forced by refinement compatibility.

The informational harmonic oscillator is the canonical closed system of the informational universe. Its invariants arise from consistency, not from assumed conservation laws. Its oscillatory form is the only reversible exten-

sion of a two-variable reciprocity pair compatible with Martin consistency and the axioms of event selection.

Phenomenon 4.10.3 (The Catalyst Effect). **Statement.** *A catalyst is a structure that lowers the informational strain required to admit a refinement without altering the net causal ledger.*

Mechanism. *Consider a transformation that is admissible only if the ledger traverses a high-curvature refinement path. Without assistance, this path lies outside the refinement budget permitted by the Law of Spline Sufficiency and the transition does not occur.*

Introduce a catalytic structure K . The catalyst provides an alternate sequence of intermediate anchors that reduce the curvature of the admissible spline while preserving the net boundary conditions.

Formally, the catalyzed path satisfies

$$U_A \xrightarrow{K} U^* \xrightarrow{K^{-1}} U_B,$$

where the intermediate ledger U^ exists only to reduce informational strain. The catalyst does not appear in the initial or final ledger states.*

Ledger Neutrality. *The catalyst is not consumed because it does not contribute events to the causal balance. It alters the geometry of admissibility without altering the count of refinements.*

Conclusion. *In chemical and physical systems, catalysis is not a lowering of an energetic barrier, but a reduction in the curvature of the admissible refinement path. The catalyst reshapes the spline; it does not change the endpoints.*

Where catalysis reshapes the admissible path without changing endpoints, a further refinement appears when the ledger actively stabilizes itself around preferred configurations. The next phenomenon captures this self-regulating behavior.

Phenomenon 4.10.4 (The Thermostat Effect). **Statement.** *An admissible ledger exhibits self-regulation around low-strain states. This behavior appears macroscopically as thermostatic control.*

Mechanism. *Let \mathcal{I} denote the informational strain functional. Refinement updates do not merely seek $\delta\mathcal{I} = 0$, but dynamically suppress deviations from locally stable minima. When the ledger drifts away from a low-strain configuration, subsequent refinements are biased toward restoring that state.*

Low and High Water Marks. *Stable configurations act as set points. If $\delta^2\mathcal{I} > 0$, deviations decay and the ledger returns to the same admissible history (cooling/heating correction). If $\delta^2\mathcal{I} < 0$, deviations amplify and the control loop fails.*

Interpretation. *A thermostat is not a separate mechanism imposed on the system. It is the observable signature of second-variation stability in the refinement functional. The ledger enforces feedback because unstable histories are inadmissible.*

Conclusion. *Thermal equilibrium is not static; it is an actively maintained fixed point of the causal bookkeeping process.*

An orbit is not a balance of forces, but a stable, self-correcting loop in the refinement ledger.

Phenomenon 4.10.5 (The Orbiting Effect [85]). **Statement.** *An orbit is a closed admissible refinement cycle stabilized by continuous error correction. It is not sustained by force, but by feedback.*

Mechanism. *Consider a ledger state constrained by a central boundary condition. The Law of Spline Sufficiency admits multiple low-strain continuations. In the presence of thermostatic stabilization, deviations from these continuations are actively corrected rather than damped to rest.*

Let $\gamma(t)$ denote the admissible refinement path. Without feedback, perturbations drive γ toward fixed points. With second-variation stability enforced dynamically, the ledger suppresses radial drift but permits tangential continuation.

The admissible path therefore closes:

$$\gamma(t + T) = \gamma(t).$$

Interpretation. *An orbit is not equilibrium. It is a stable failure to terminate. The thermostatic action prevents collapse while the catalytic structure prevents escape. The ledger cycles because that is the only admissible history that preserves all constraints without contradiction.*

Conclusion. *Keplerian motion, atomic shells, and macroscopic rotation are not balance of forces. They are sustained reconciliation loops in the causal record. An orbit is a closed book that must be reread forever.*

It is important to note that no notion of gravitational force has been invoked in this development. The existence of orbits here does not arise from attraction, mass, or curvature of spacetime as primitive inputs.

Orbits emerge solely from the structure of admissible refinement. They are closed solutions to the bookkeeping problem: how a finite ledger can preserve boundary constraints while minimizing informational strain under continuous correction. The appearance of centripetal “force” in classical physics is a smooth shadow of this deeper combinatorial necessity.

In this framework, gravity does not create orbits. Orbits create the conditions under which gravity later appears as an effective description.

Chapter 5

Informational Stress

The preceding chapters established that smooth motion appears as the unique closure of causal order under refinement. The Law of Spline Sufficiency showed that any admissible continuous shadow must contain no unrecorded structure and therefore satisfies the extremal condition $\Psi^{(4)} = 0$.

In this chapter we examine the opposite extremum: the smallest admissible refinement of the Causal Universe Tensor. Such a refinement represents the maximal rate at which distinguishability can propagate without violating the Axiom of Planck. We call this minimal, nonzero update an *informational quantum*. It is not a physical particle or field; it is the atomic refinement permitted by the axioms.

5.1 Informtional Quantum

Precision in this framework is not free. By the Law of Discrete Spline Necessity, admissible interpolation cannot be performed with arbitrarily fine resolution. Every smooth completion is the limit of a finite refinement process, and every such refinement is bounded by the Axioms of Kolmogorov and Planck. This forces a minimal unit of admissible distinction: an *informational quantum*. The spline may assign exact analytic values between

anchor points, but those values are only meaningful up to the smallest refinement allowed by the causal ledger. Precision therefore emerges not as a continuum ideal, but as a quantized requirement. The theory is compelled to be precise only in discrete units, because no admissible history may resolve structure smaller than the finite quantum of measurement demanded by coherent spline closure.

Phenomenon 5.1.1 (Precision). *The transition from discrete measurement to continuous description is not introduced by assumption, but forced by the structure of admissible interpolation. By the Law of Spline Sufficiency, any finite sequence of anchor events admits a unique minimal-curvature completion. This completion assigns values not only at the recorded events themselves, but at all admissible points between them.*

Precision and accuracy are distinct in this framework. Precision refers to the determinacy of the interpolated values supplied by the admissible spline: once anchor points are fixed, the analytic completion assigns well-defined values at every intermediate point. Accuracy, by contrast, refers only to agreement with future measurement events. A value may be perfectly precise—uniquely determined by the axioms—and yet not be accurate if a subsequent refinement records a different event. Precision is therefore a property of admissible completion, while accuracy is a property of the experimental record. The former is forced by coherence; the latter remains an empirical constraint. These intermediate values are not measurements. They are consequences.

The spline interpolant supplies a determinate analytic value at every point of its domain, even though such points were never observed. This phenomenon is the origin of precision in the informational framework. The causal ledger remains discrete, but the admissible completion is continuous. The difference between what is recorded and what is implied is not a defect; it is a structural requirement of coherence.

Precision does not arise from improved instruments or finer resolution. It arises from necessity. Once anchor points are fixed, the axioms force a unique

analytic structure between them. The values supplied by the spline are not guesses, and they are not stochastic. They are the only values consistent with information minimality and global admissibility.

In this sense, precision is not an empirical achievement but a mathematical obligation. The continuum is not observed; it is compelled.

The remainder of this chapter treats the consequences of this effect. When analytic predictions are treated as real numbers rather than combinatorial counts, new bookkeeping problems arise. These problems do not reflect physical forces, but the informational cost of maintaining precision between discrete events.

5.2 The Informational Bound ϵ

N.B.—The refinement bound ϵ is not a physical quantum, particle, or energy unit. It is the minimal nonzero increment of distinguishable structure that survives every admissible refinement of the measurement record. Its origin is purely informational: ϵ is the continuous shadow of the residual \mathcal{C}^2 freedom in spline closure. No physical ontology is implied. \square

The refinement of an observational record proceeds through countable additions of distinguishable events. As established in Chapter 4, the weak form of the discrete bending functional admits a single free \mathcal{C}^2 parameter, corresponding to the third derivative of the spline interpolant. This degree of freedom is not an artifact of approximation; it is a structural remnant of finite measurement.

By the Law ??, admissible completion cannot eliminate this residual freedom except through the introduction of new anchor events. When no additional measurements are recorded, the remaining degree of freedom is irreducible. The continuous shadow is therefore forced to carry a minimal, nonvanishing bound on curvature-level distinction. This bound is not imposed by physics, but by the impossibility of selecting a unique refinement

in the absence of new information.

We denote this invariant residual by ϵ . Any admissible refinement of the continuous shadow must preserve ϵ ; to refine below this threshold would introduce unrecorded structure and contradict the finite measurement sequence. Conversely, any refinement that preserves ϵ remains consistent with the discrete data. Thus ϵ functions as the kinematic limit of refinement and provides the foundation for the emergent invariant interval τ and operators that may look familiar to some.

Phenomenon 5.2.1 (The Richardson Effect [126]). **N.B.**—*In a nut-shell, how long is Britain's coastline and why does the answer depend on the length of the ruler [?]?* □

The measured length of a boundary increases without limit as the resolution of measurement is refined, even though the underlying admissible structure remains finite. This phenomenon is most clearly expressed by the classical coastline mapping problem: the total measured length of a coastline depends monotonically on the length of the measuring stick.

When a coastline is traced using a coarse measuring scale, long rulers bridge over bays, inlets, and local irregularities. The resulting path is smooth at that scale and the reported length is relatively short. As the measuring scale is reduced, the ruler no longer spans these features. Previously ignored curvature is now forced into the admissible path. The measured length increases because the boundary is not smooth; it carries irreducible roughness.

In the informational framework, this roughness is not accidental. By the Law of Finite Spline Selection, a spline constrained only by finitely many anchor points must retain a residual degree of curvature freedom. That freedom does not vanish between measurements; it remains latent. As resolution improves, the measurement process is compelled to resolve this latent curvature. The coastline must appear rough, because a perfectly smooth boundary would require infinite observational constraint.

The coastline does not acquire new structure under refinement. Rather, its

admissible minimal completion is forced to reveal structure that was always present but previously collapsed by coarse measurement. The increase in measured length is therefore not a property of the land, but a consequence of how finite measurement interacts with unavoidable curvature residue.

In this sense, roughness is not an empirical irregularity. It is a structural requirement of any boundary recorded by finitely many distinguishable events.

The Richardson Effect is not a property of space. It is a property of measurement. A boundary is not an object with a fixed length; it is a ledger of distinguishable anchor points together with their admissible minimal completions. As refinements increase, the informational content of the boundary increases, and the measured length grows accordingly.

There is no convergence to a true length. There is only an ever-refining account of admissible curvature. See Phenomenon ??.

5.2.1 Residual Spline Freedom and the Minimal Refinement Bound

The weak formulation developed in Chapter 3 shows that a sequence of discrete measurements determines a unique cubic spline interpolant, except for a single free parameter associated with the third derivative. This residual freedom is a consequence of the information minimality constraint: in the absence of additional recorded events, the interpolant must not introduce structure that was not observed. A fourth-order correction would imply an unrecorded change in curvature and is therefore inadmissible.

Let Ψ denote the continuous shadow of the discrete observational chain. The freedom in Ψ''' represents the smallest degree of curvature that can be varied without conflicting with the discrete record. Its magnitude cannot be reduced by any admissible refinement unless a new event occurs. The minimal nonzero shadow of this residual is the refinement bound ϵ .

Formally, ϵ is the infimum of all curvature-level refinements that do not violate the existing event sequence. Any attempt to impose a refinement of

order smaller than ϵ would create new curvature that must be supported by new events, contradicting the record. Thus ϵ is the continuous representation of the irreducible \mathcal{C}^2 residue.

5.2.2 Maximal Informational Propagation

An admissible refinement of the observational record adds distinguishable structure without contradicting previously recorded events. A path that *saturates* the refinement bound ϵ propagates information at the maximal admissible rate: it incorporates all allowable distinction while introducing no unrecorded curvature.

Such paths form the extremal curves of the informational geometry. They are defined not by physical principles, but by the logical requirement that refinement cannot fall below the ϵ threshold. Any further reduction would imply hidden structure and is therefore inadmissible.

In the continuous shadow, these maximally propagated paths serve as the reference curves for defining the invariant interval τ . Two observers who refine the same extremal path must agree on the number of informational units required to describe it; this count determines the causal interval and anchors the construction of the metric in Section 5.2.

Phenomenon 5.2.2 (Compact Disc Encoding [28, 47]). **N.B.**—*The compact disc format is treated here not as an optical or physical device but as a concrete implementation of an informational system. Its behavior illustrates how distinguishability, admissible refinement, finite alphabets, and boundary consistency determine the structure of a real-world communication medium. No photonic or physical assumptions are made; the CD is considered solely as a record of measurable distinctions.* □

N.B.—*This thought experiment does not describe photons as informational quanta. It is a finite conceptual model illustrating how a gauge of separation emerges from the logic of distinguishability alone. No physical ontology is implied.* □

The compact disc (CD) format developed jointly by Sony and Philips implements a finite alphabet of distinguishable marks: pits and lands arranged along a single spiral track. Each measurement by the reader selects one symbol from this alphabet. The resulting word encodes audio data through a sequence of refinements governed by cross-interleaved Reed–Solomon coding (CIRC), an error-correcting structure patented in the foundational work on digital optical media [28, 47].

A notable design constraint is the total record length. The original Sony specification targeted a runtime of approximately 74 minutes (often quoted as 72 minutes in early engineering drafts) so that a single disc could contain a complete performance of Beethoven’s Ninth Symphony. Although historical details vary, the engineering requirement is informational in nature: the spiral track must accommodate a finite number of distinguishable symbols, each encoded with redundancy and refinement structure sufficient to guarantee coherent recovery.

Thus the CD provides a physical instantiation of an informational phenomenon: a medium whose structure, capacity, and correction rules are determined entirely by the algebra of distinguishability and refinement.

A compact disc stores information as a finite, ordered chain of distinctions. Each pit or land corresponds to a single admissible event, and the reader detects a new event only when the reflected signal exceeds its threshold of discernibility. Everything below this threshold is invisible; it cannot enter the admissible record. Thus the sequence of detections,

$$e_1 \prec e_2 \prec e_3 \prec \cdots ,$$

encodes not only what was observed, but the binding constraint that no additional distinguishable structure may be inserted between these events.

From the standpoint of information, the read head defines a gauge of minimal separation: two surface configurations are “far enough apart” exactly when the detector must refine its admissible description to distinguish them.

The metric is not assumed; it is inferred from the rule that only resolvable differences may appear as refinements in the causal chain.

Now imagine two readers, A and B, scanning the same disc. Reader A has a coarser threshold; reader B resolves finer distinctions. Each produces its own ordered sequence of admissible events. Where B records additional refinements, A records none. Yet when their records are merged, global coherence requires a single history that preserves all recorded distinctions. The finer record forces a refinement on the coarser: A must treat certain portions of the disc as informationally extended, for failure to accommodate B's distinctions would render the merged history inconsistent.

In the dense limit, this refinement rule induces a continuous connection: the shadow of the logical requirement that adjacent descriptions remain compatible under transport. What appears in the smooth theory as a metric is nothing more than this bookkeeping of distinguishability: the minimal rule that certifies when two states differ in a way that must be reconciled.

In this model, “light” corresponds not to a substance but to the maximal rate at which new distinctions can be admitted without contradiction. Any attempt to introduce refinements faster than this rate would violate global coherence. Thus the invariant causal interval of Chapter 5 reflects the same constraint: an observer may not admit distinctions faster than a globally coherent merge can support.

The compact disc reader therefore offers a finite, concrete metaphor for the emergence of the gauge of light, the metric as a rule of separation, and the transport laws that follow from informational consistency.

5.3 The Metric as a Gauge of Informational Separation

N.B.—The metric is not a physical fabric, field, or medium. It is the continuous shadow of the rule by which refinements preserve the invariant causal

interval. Its function is purely informational: the metric enforces that all admissible observers agree on the amount of distinguishable structure between neighboring events. No geometric ontology is assumed. \square

The refinement bound ϵ defined in Section 5.2 limits how finely an admissible history may be resolved without contradicting the measurement record. When this bound is propagated along an extremal path, the result is a conserved quantity—the informational interval τ —which represents the number of ϵ -sized refinements required to traverse that path. Any two observers refining the same extremal sequence must therefore agree on the value of τ . This invariance is the kinematic foundation of the metric.

Let dx^μ denote the local labels an observer assigns to successive events along an extremal refinement. Another observer, using a different admissible convention for distinguishing the same events, assigns labels $dx'^\mu = \Lambda^\mu{}_\nu dx^\nu$, where the transformation $\Lambda^\mu{}_\nu$ preserves causal order as required by the Axiom of Selection. Although the two observers differ in the coordinates they assign, they must agree on the number of ϵ -sized refinements separating the events; otherwise their merged histories would violate global consistency.

This requirement implies the existence of a bilinear form $g_{\mu\nu}$ such that the scalar quantity

$$\tau^2 = g_{\mu\nu} dx^\mu dx^\nu$$

is invariant under all admissible changes of local labeling conventions. The metric is therefore the gauge of informational separation: it encodes how distinctions between events are preserved when observers describe them using different coordinate choices.

Because τ counts informational increments rather than geometric lengths, the invariance condition

$$g'_{\mu\nu} dx'^\mu dx'^\nu = g_{\mu\nu} dx^\mu dx^\nu$$

expresses the deeper statement that the number of distinguishable refine-

ments between neighboring events is independent of the observer. The metric formalizes this invariance. It is the bilinear bookkeeping rule ensuring that every observer's coordinate description yields the same value of τ for any given extremal path.

In this interpretation, the components of $g_{\mu\nu}$ do not prescribe a geometry; they record the local comparison rule by which distinguishability is maintained across observational frames. Section 5.3.1 elevates this requirement to a formal law, showing that preservation of the causal interval uniquely determines the compatible affine connection used to propagate distinguishability under refinement.

5.3.1 The Law of Causal Transport

N.B.—The Law of Causal Transport is a kinematic statement. It asserts only that informational refinements must preserve the invariant interval τ defined in Section 5.3. No dynamical interpretation of curvature or stress is assumed here. The law specifies how distinguishability must be propagated under admissible changes of frame; all higher structures of connection and curvature follow in later sections. \square

The refinement bound ϵ defines the smallest admissible increment of distinguishable structure. When propagated along an extremal path, ϵ induces the invariant interval τ , representing the total number of such increments required to describe that path. Because every observer must refine the same underlying event sequence, the value of τ must remain unchanged under all admissible relabelings.

This requirement leads to the following principle.

Law 4 (The Law of Causal Transport). [Preservation of Distinguishability]
Any admissible refinement of an observational record must preserve the informational interval τ between neighboring events. In the continuous shadow, this condition determines a unique bilinear form $g_{\mu\nu}$ and a unique compatible

rule of transport $\Gamma_{\mu\nu}^\lambda$ satisfying

$$\nabla_\lambda g_{\mu\nu} = 0.$$

The pair $(g_{\mu\nu}, \Gamma_{\mu\nu}^\lambda)$ constitutes the metric gauge of informational separation.

Because observers may assign different coordinates to the same infinitesimal event displacement, we represent such a relabeling by $dx'^\mu = \Lambda^\mu{}_\nu dx^\nu$, where $\Lambda^\mu{}_\nu$ preserves causal order. The Law of Causal Transport requires that the informational interval be invariant under this transformation:

$$g'_{\mu\nu} dx'^\mu dx'^\nu = g_{\mu\nu} dx^\mu dx^\nu.$$

This invariance elevates $g_{\mu\nu}$ from a mere bookkeeping device to a constraint: it is the only bilinear form that guarantees all observers agree on how many ϵ -sized refinements separate neighboring events.

The law further implies that the comparison of nearby refinements must not depend on the path taken in the space of coordinate labels. This requirement determines the connection coefficients $\Gamma_{\mu\nu}^\lambda$ as the unique differential operators that preserve the metric gauge under change of frame.

In this sense, the Law of Causal Transport encodes the most fundamental rule of the kinematic structure: that distinguishability is preserved under motion. The connection is not postulated, but forced by the need to maintain the interval τ when an observer's coordinate conventions vary from point to point. Section 5.3.2 elaborates the invariance of τ , and Section 5.3.3 formalizes the role of $g_{\mu\nu}$ as the bilinear form that preserves the ϵ -refinement count.

5.3.2 Invariance of the Informational Interval τ

N.B.—The interval τ is not a geometric length or a physical duration. It is the continuous shadow of an event count: the number of ϵ -sized refinements

required to describe an extremal segment of an observational record. Its invariance expresses only that all admissible observers must agree on the amount of distinguishable structure between neighboring events. \square

The refinement bound ϵ defines the smallest admissible increment of distinguishability. When propagated along an extremal path—one that saturates the refinement bound—each observer records the same number of ϵ -increments. This count defines the informational interval τ . Because τ represents the number of admissible refinements rather than a metric distance, its invariance follows from the requirement that no observer may introduce or remove distinguishable structure that is not supported by the measurement record.

Let dx^μ and dx'^μ denote the infinitesimal labels assigned by two admissible observers to the same pair of neighboring events. Their coordinate labels differ by a transformation

$$dx'^\mu = \Lambda^\mu{}_\nu dx^\nu,$$

where $\Lambda^\mu{}_\nu$ preserves causal order in the sense of the Axiom of Selection. Although the observers assign different coordinates, they must agree on the number of ϵ -increments between the events; otherwise their merged histories would violate global consistency.

This agreement is enforced by a bilinear form $g_{\mu\nu}$ satisfying

$$\tau^2 = g_{\mu\nu} dx^\mu dx^\nu.$$

Under the coordinate transformation, the metric transforms as

$$g'_{\mu\nu} = \Lambda^\alpha{}_\mu \Lambda^\beta{}_\nu g_{\alpha\beta}.$$

Substituting the transformed variables into the definition of τ yields

$$\tau'^2 = g'_{\mu\nu} dx'^\mu dx'^\nu = g_{\alpha\beta} dx^\alpha dx^\beta = \tau^2.$$

The invariance of τ thus expresses a simple but fundamental principle: every admissible observer must assign the same number of distinguishable increments to an extremal path. Their coordinate descriptions may vary, but the informational content of the path does not.

This invariance is the basis of the metric gauge introduced in Section 5.3. It ensures that τ may serve as the universal measure of informational separation, independent of the observer's local labeling conventions. Section 5.3.3 develops the metric $g_{\mu\nu}$ as the bilinear form that enforces this invariance in the continuous shadow.

5.3.3 $g_{\mu\nu}$ as the Bilinear Form Preserving the ϵ -Refinement Count

N.B.—The metric $g_{\mu\nu}$ is not a geometric field on a manifold. It is the continuous shadow of the rule ensuring that all admissible observers preserve the same count of ϵ -sized refinements between neighboring events. The components of $g_{\mu\nu}$ do not describe a physical medium or curvature; they encode the invariant comparison rule required by informational consistency. \square

The interval τ defined in Section 5.3.2 expresses the number of ϵ -refinements along an extremal segment of the measurement record. Since this number must remain invariant under all admissible relabelings of events, there must exist a bilinear form $g_{\mu\nu}$ such that

$$\tau^2 = g_{\mu\nu} dx^\mu dx^\nu$$

holds for every observer. This expression is not a postulate but the unique structure that enforces the preservation of τ under coordinate changes that respect causal order.

To see this, consider two observers who assign infinitesimal labels dx^μ and $dx'^\mu = \Lambda^\mu{}_\nu dx^\nu$ to the same pair of neighboring events. The Law of Causal

Transport requires

$$\tau'^2 = \tau^2,$$

so we must have

$$g'_{\mu\nu} dx'^\mu dx'^\nu = g_{\mu\nu} dx^\mu dx^\nu.$$

Substituting dx'^μ and requiring equality for all admissible transformations Λ^μ_ν yields the transformation rule

$$g'_{\mu\nu} = \Lambda^\alpha_\mu \Lambda^\beta_\nu g_{\alpha\beta}.$$

Thus the metric is exactly the object that ensures agreement on the number of ϵ -sized refinements between neighboring events, regardless of the coordinates used to describe them.

In this informational framework, $g_{\mu\nu}$ plays a role analogous to that of a gauge potential: it specifies how infinitesimal refinements are compared locally so that the global invariant τ remains unchanged. The metric does not specify “distance” in any geometric or physical sense; it enforces the equivalence of all admissible measurement conventions.

Once $g_{\mu\nu}$ is introduced, the need to propagate these comparison rules from point to point forces a unique notion of compatibility. This requirement determines the affine connection in Section 5.4 through the condition

$$\nabla_\lambda g_{\mu\nu} = 0,$$

which expresses that the metric gauge is preserved under refinement and transport. The next section illustrates this invariance with a concrete thought experiment.

Phenomenon 5.3.1 (The Michelson–Morley Effect [107]). ***N.B.**—This phenomenon is not interpreted as a physical test of ether hypotheses, relativistic postulates, or the dynamics of light. It is treated purely as an informational experiment: a demonstration that distinguishable events may propagate*

through a region in which no medium is observed. The null result is therefore a statement about the structure of admissible refinements and boundary conditions, not about physical substrates. \square

N.B.—This thought experiment does not appeal to optical physics, wave interference, or the existence of a medium. It is a finite informational model illustrating that the metric gauge must assign the same refinement cost ϵ to extremal paths in all admissible directions. No physical claims about light or propagation are implied. \square

Consider an observer attempting to refine two extremal segments of equal informational content, but aligned in different coordinate directions. Let dx^μ and dy^μ denote the local labels assigned to the two segments. Each segment is chosen such that its refinement requires the same number of ϵ -increments when described in the observer's own frame.

Now suppose the observer rotates their coordinate system. After rotation, the new labels are $dx'^\mu = \Lambda^\mu_\nu dx^\nu$ and $dy'^\mu = \Lambda^\mu_\nu dy^\nu$. The rotation Λ^μ_ν preserves causal order, so it is an admissible transformation. The question is whether the observer must still assign the same informational interval τ to both segments after the rotation.

The Law of Causal Transport requires that the ϵ -refinement counts for both segments remain invariant:

$$\tau_x^2 = g_{\mu\nu} dx^\mu dx^\nu, \quad \tau_y^2 = g_{\mu\nu} dy^\mu dy^\nu.$$

After rotation, the transformed intervals are

$$\tau_x'^2 = g'_{\mu\nu} dx'^\mu dx'^\nu, \quad \tau_y'^2 = g'_{\mu\nu} dy'^\mu dy'^\nu.$$

Substituting the transformation rules for dx'^μ , dy'^μ , and $g'_{\mu\nu}$ gives

$$\tau_x'^2 = g_{\alpha\beta} dx^\alpha dx^\beta = \tau_x^2, \quad \tau_y'^2 = g_{\alpha\beta} dy^\alpha dy^\beta = \tau_y^2.$$

Thus the observer must continue to assign the same informational interval to the two extremal segments under any admissible rotation. There is no freedom to deform the refinement counts directionally: doing so would imply that ϵ -sized increments depend on orientation and would violate the requirement that informational refinement be globally coherent.

This invariance is the informational analogue of isotropy. It expresses that the metric gauge $g_{\mu\nu}$ must refine extremal paths uniformly in all directions: the number of ϵ -increments needed to resolve a segment of given informational content cannot depend on the coordinate orientation.

The Michelson–Morley experiment is therefore understood here not as a test of a physical medium, but as a finite illustration of the isotropy of the metric gauge. The invariance of τ under rotations forces $g_{\mu\nu}$ to encode a direction-independent refinement rule. Section 5.4 develops the compatible connection that propagates this rule under changes of frame.

Phenomenon 5.3.2 (The Traffic Effect). *Light propagating through a region of elevated informational stress requires additional refinement steps to preserve admissibility. The resulting delay is not a failure of transmission, but a bookkeeping cost.*

The metric $g_{\mu\nu}$ acts as a gauge of informational separation. In stressed regions, the ledger must insert additional ticks in order to transport a refinement across the same coordinate distance. The observed time delay is the accumulation of these additional admissible refinement events.

The delay therefore measures not distance, but the increased cost of maintaining consistency of the informational interval under transport.

5.4 The Connection as Informational Book-keeping

N.B.—The affine connection $\Gamma_{\mu\nu}^{\lambda}$ is not a force field or a physical interaction. It is the continuous shadow of an informational rule: the minimal differential adjustment required to preserve the metric gauge $g_{\mu\nu}$ as an observer moves from one event to its neighbor. Its role is purely kinematic. The connection records how local measurement conventions must tilt to maintain the invariant interval τ ; no dynamical content or geometric ontology is assumed. \square

The metric $g_{\mu\nu}$, introduced in Section 5.3, guarantees that all admissible observers assign the same informational interval τ to an extremal displacement at a single event. This invariance is enforced by the bilinear form $g_{\mu\nu} dx^{\mu} dx^{\nu}$, which preserves the ϵ -refinement count under changes of coordinates. However, the metric by itself does not specify how these comparison rules extend from one event to its neighbors. To describe how distinguishability is maintained along a path, we require a differential notion of consistency.

The connection $\Gamma_{\mu\nu}^{\lambda}$ provides this rule. It specifies how tensor components must be adjusted when an observer translates a local measurement convention from one event to an infinitesimally adjacent one. In particular, the connection determines the covariant derivative, which measures change in a way that respects the metric gauge. Imposing that the metric remain invariant under such differential updates leads to the condition $\nabla_{\lambda} g_{\mu\nu} = 0$, known as covariant constancy of the metric.

In the informational picture, this condition is the statement that the act of refinement may not create or destroy distinguishable structure as an observer moves through the network of events. The connection is the unique differential bookkeeping device that satisfies this constraint. When the metric is uniform, the connection vanishes and no adjustment is needed: straight paths remain informationally straight. When the metric varies, a nonzero

connection encodes how local gauges must be rotated and rescaled so that scalar quantities built from $g_{\mu\nu}$ remain unchanged.

The remainder of this section develops the connection as the compatibility condition implied by covariant constancy of the metric and interprets parallel transport as the differential expression of Martin consistency. In this way, the Law of Causal Transport acquires its full kinematic content: it is the rule that propagates the gauge of separation through the continuous shadow of the Causal Universe Tensor.

Phenomenon 5.4.1 (The Sagnac Effect). *When refinement clocks are transported around a closed loop, global synchronization fails. The connection Γ adjusts local refinement rates to maintain admissibility, but the adjustments do not close under cyclic transport.*

Two refinement paths that traverse the same boundary in opposite directions accumulate unequal refinement tallies. This asymmetry is not a defect of propagation, but the holonomy of the bookkeeping rule.

The observed time difference is the irreducible gap produced when a local transport rule cannot be extended consistently around a closed causal cycle.

Phenomenon 5.4.2 (The Tail-Latency Effect). **Statement.** *Latency in an admissible region increases with both the number of active causal connections and the surface measure of the region through which refinements must be transported.*

Mechanism. *Each admissible refinement must be reconciled across all attached causal interfaces. Let N denote the number of active connections incident on a region Ω , and let $|\partial\Omega|$ denote the surface measure of its boundary. The cost of transport is not determined by the shortest path, but by the slowest admissible reconciliation.*

The tail of the latency distribution is therefore governed by

$$\mathcal{L}_{\text{tail}} \propto N \cdot |\partial\Omega|.$$

Interpretation. *Transport in the causal ledger is not limited by average throughput but by worst-case synchronization. Each additional connection increases the number of constraints that must be satisfied, and each increase in boundary area expands the number of admissible reconciliation paths.*

Latency therefore accumulates geometrically: wide interfaces and dense connectivity do not accelerate refinement, they delay it. The slowest boundary dominates the admissible update rate.

This is not a property of signal speed. It is a bookkeeping constraint: the ledger cannot commit a refinement until every connected boundary can be reconciled without contradiction.

Phenomenon 5.4.3 (The Halt Effect.). *Not every admissible refinement admits a successor. There exist boundary configurations for which no further consistent update can be constructed.*

If a partial ledger extension would require the separation of correlated events without a permissible ordering, the update operator has no admissible output. The refinement process halts.

This is not a failure of computation but a structural limit of admissibility. A halted ledger is not incomplete; it is complete in the only sense allowed by the axioms. No further event can be appended without violating global consistency.

The halting of a causal sequence is therefore not destruction. It is the formal termination of admissible history.

5.4.1 Covariant Constancy and the Compatibility Condition

N.B.—Covariant constancy is not a physical conservation law. It is the informational requirement that the metric gauge $g_{\mu\nu}$, which preserves the ϵ -refinement count at a single event, must continue to preserve that count as the observer moves to a neighboring event. The affine connection $\Gamma_{\mu\nu}^{\lambda}$ is

therefore not introduced by assumption; it is forced by the requirement that informational invariants remain invariant under differential refinement. \square

The metric $g_{\mu\nu}$ ensures that all admissible observers agree on the informational interval τ at a point. But as the observer moves from an event x to a nearby event $x + dx$, the local coordinate basis changes. Under such a shift, the numerical components of $g_{\mu\nu}$ may appear to change due to the alteration in basis, even if the underlying structure of distinguishability remains the same. To prevent this apparent change from contaminating the informational interval, the transformation of $g_{\mu\nu}$ must be corrected by an additional adjustment term.

This correction is encoded by the covariant derivative. The condition that the metric gauge remain invariant under differential displacement is expressed as

$$\nabla_\lambda g_{\mu\nu} = \partial_\lambda g_{\mu\nu} - \Gamma_{\mu\lambda}^\sigma g_{\sigma\nu} - \Gamma_{\nu\lambda}^\sigma g_{\mu\sigma} = 0.$$

The partial derivative $\partial_\lambda g_{\mu\nu}$ captures how the metric components vary when written in the shifted coordinate system. The remaining terms subtract off this apparent variation by compensating for the tilt and scale of the basis vectors themselves. The equation $\nabla_\lambda g_{\mu\nu} = 0$ thus expresses the requirement that the informational interval τ remain unchanged under any infinitesimal update of the observational coordinates.

This compatibility condition uniquely determines the connection when torsion is absent. As established in Chapter 3, the spline representation of admissible histories carries no fourth-order freedom and is therefore torsion-free. Under this constraint, the metric compatibility condition fixes the connection to be the Levi-Civita connection:

$$\Gamma_{\mu\nu}^\lambda = \frac{1}{2} g^{\lambda\sigma} (\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\mu\sigma} - \partial_\sigma g_{\mu\nu}).$$

This expression is not a postulate; it is the only operator that ensures the metric gauge remains intact under transport. It is the continuous shadow

of the discrete requirement that refinement cannot introduce or eliminate distinguishable structure beyond the ϵ bound.

With the connection now fixed by kinematic necessity, we may interpret its role operationally. The connection coefficients specify the adjustments required to compare tensorial quantities at neighboring events, ensuring that the informational interval τ and the refinement bound ϵ remain consistent throughout the observer's path. The next subsection formalizes this process as parallel transport.

5.4.2 Parallel Transport as Differential Martin Consistency

N.B.—Parallel transport is not a physical motion of a vector through space. It is the informational requirement that the meaning of a direction—a rule for distinguishing one infinitesimal refinement from another—remain consistent as an observer updates coordinates from one event to the next. In this framework, a “vector” is an instruction for refinement, and parallel transport ensures that such instructions are not distorted by changes in local labeling conventions. \square

The metric compatibility condition $\nabla_\lambda g_{\mu\nu} = 0$ determines how the metric must be preserved under infinitesimal displacement. Parallel transport extends this requirement to all tensorial quantities, ensuring that any object used to encode refinements of the observational record is carried through the continuous shadow without introducing contradictions.

Let V^μ represent such a refinement direction. When an observer moves along a curve $x^\mu(s)$ in the event network, the numerical components of V^μ change because the local coordinate basis changes. The naive derivative dV^μ/ds therefore incorporates both the intrinsic change in the refinement direction and the apparent change induced by the shifting coordinates. To isolate the intrinsic change—the part that affects distinguishability—we must subtract off the bookkeeping contribution provided by the connection.

The covariant derivative along the path is thus defined as

$$\frac{DV^\mu}{Ds} = \frac{dV^\mu}{ds} + \Gamma^\mu_{\nu\lambda} V^\nu \frac{dx^\lambda}{ds}.$$

Parallel transport requires that the intrinsic change vanish:

$$\frac{DV^\mu}{Ds} = 0.$$

This equation expresses the differential form of Martin consistency. It states that the instruction encoded by V^μ must retain its informational meaning as the observer moves. All coordinate-induced distortions of V^μ must be canceled by the corresponding connection terms, ensuring that the refinement direction does not acquire unrecorded structure.

The geometric interpretation of parallel transport as preserving “straightness” is replaced here by a purely informational one: parallel transport guarantees that refinement instructions remain compatible with the metric gauge $g_{\mu\nu}$ throughout the observer’s path. Whenever the metric varies from event to event, the connection coefficients encode the cost of adjusting the observer’s basis to ensure that scalar comparisons built from $g_{\mu\nu}$ and V^μ remain invariant.

In regions where $g_{\mu\nu}$ is uniform, the connection vanishes and the informational meaning of V^μ is preserved without adjustment. Where $g_{\mu\nu}$ varies, nonzero connection coefficients encode the minimal bookkeeping needed to keep the refinement count consistent. This adjustment is the kinematic origin of effects such as frequency shifts between observers in different informational environments, which we examine in the following subsection.

5.5 Pendula

Phenomenon 5.5.1 (The Foucault Effect). *A refinement direction transported around a closed causal loop does not generally return to its initial*

orientation. The failure of closure is not a mechanical torque, but the accumulated informational strain required to preserve admissibility under transport.

The observed precession is the holonomy of an inconsistent connection: a closed path in events produces an open path in orientation.

5.6 Refinement–Adjusted Transport

N.B.—The frequency shift examined in this section is not a postulated effect. It is the kinematic consequence of maintaining the invariant informational interval τ across regions in which the metric gauge $g_{\mu\nu}$ varies. No physical ontology is assumed. The observable change in clock rates reflects the differential bookkeeping enforced by the connection $\Gamma_{\mu\nu}^\lambda$. \square

The previous sections established the chain of informational structure: the refinement bound ϵ fixes the local increment of distinguishable structure; the metric $g_{\mu\nu}$ expresses how these increments are compared between observers; and the connection $\Gamma_{\mu\nu}^\lambda$ preserves this comparison under differential displacement. When the metric varies from one location to another, this preservation requires that the local rate of event counting—the clock frequency—adjusts so that the invariant interval remains consistent across observers.

This section derives that adjustment and exhibits its observable consequence.

5.6.1 The Invariant Causal Tally

N.B.—An atomic clock does not measure a geometric length or a physical time. It measures a count of distinguishable events. The proper interval τ is the continuous shadow of this count, expressed in units of the refinement bound ϵ . \square

Consider an observer whose worldline is described by coordinates (t, x^i) . If the observer is at rest in their coordinate system ($dx^i = 0$), the informational interval between neighboring events satisfies

$$d\tau^2 = g_{00}(x) dt^2.$$

Thus the locally measured period of the clock is

$$d\tau = \sqrt{g_{00}(x)} dt.$$

Because τ counts ϵ -sized refinements, the local clock frequency $\nu(x)$ is inversely proportional to the size of this interval:

$$\nu(x) = \frac{1}{d\tau} = \frac{1}{\sqrt{g_{00}(x)}} \frac{1}{dt}.$$

Two observers at rest in different metric gauges therefore experience different informational intervals for the same coordinate increment dt . The relationship between their locally recorded counts is fixed entirely by the metric gauge.

5.6.2 Derivation of Frequency Adjustment

N.B.—The global parameter t is not a physical time. It is the auxiliary labeling parameter that all admissible observers must agree upon when their records are merged. Its increments must match across observers in order for their ϵ -counts to be reconciled. \square

Let observers A and B be stationary in regions with metric components $g_{00}(A)$ and $g_{00}(B)$. Over a shared coordinate increment Δt , their locally recorded proper intervals are

$$\Delta\tau_A = \sqrt{g_{00}(A)} \Delta t, \quad \Delta\tau_B = \sqrt{g_{00}(B)} \Delta t.$$

Since a clock's frequency is the inverse of the proper interval it records,

$$\nu_A = \frac{1}{\Delta\tau_A} = \frac{1}{\sqrt{g_{00}(A)}} \frac{1}{\Delta t}, \quad \nu_B = \frac{1}{\sqrt{g_{00}(B)}} \frac{1}{\Delta t}.$$

The ratio of their observed frequencies is therefore

$$\frac{\nu_A}{\nu_B} = \frac{\sqrt{g_{00}(B)}}{\sqrt{g_{00}(A)}}.$$

This expression is the kinematic consequence of the Law of Causal Transport. When g_{00} varies, the connection Γ_{00}^0 compensates by adjusting the local rate of ϵ -counting so that the merged observational record remains consistent. The observed frequency shift is thus the operational signature of nonzero connection coefficients.

5.7 Time Dilation

The informational framework developed in Chapters 5 and 6 places a subtle constraint on how refinement may be transported across a causal network. Proper time is not a geometric parameter but the tally of irreducible distinctions, and the metric $g_{\mu\nu}$ records how this tally must adjust when two histories inhabit regions with different curvature residue. Whenever distinguishability is carried from one domain to another, the connection enforces a compatibility rule: the informational interval must be preserved even if the local refinement structure differs.

This requirement has a striking observable consequence. Two clocks placed at different informational potentials—that is, in regions where the residual strain of admissible curvature differs—cannot maintain the same rate of refinement. Each clock is internally consistent, but the comparison of their records forces an adjustment. A refinement sequence that is admissible at one potential must be reweighted when interpreted at another, or else the

causal record would fail to merge coherently.

In the smooth shadow, this bookkeeping adjustment becomes the familiar phenomenon of gravitational redshift. Signals transported upward appear to lose frequency; signals transported downward appear to gain it. Nothing mystical is occurring: the informational interval is being preserved, and the only available mechanism is a change in the rate at which distinguishability is accumulated.

The Pound–Rebka experiment is therefore the archetype of an informational outcome. It demonstrates that when refinement is compared across regions with differing curvature residue, the universe must adjust the apparent rate of time itself to maintain consistency. No dynamical field need be invoked; the redshift is simply the shadow of the constraint that admissible refinements must agree on their causal overlap.

Phenomenon 5.7.1 (The Pound–Rebka Effect [125]). **N.B.**—*The following is an informational phenomenon. No physical mechanism is assumed. The interpretation concerns how the gauge of informational separation $g_{\mu\nu}$ adjusts refinement counts when distinguishability is transported across domains of differing causal potential. Any resemblance to the gravitational redshift measured by Pound and Rebka is a consequence of the informational shadow, not an assumed dynamical cause.* \square

The Axiom of Peano defines proper time as the count of irreducible refinements along an admissible history. The Law of Causal Transport guarantees that this count is invariant under maximal propagation, while the informational metric $g_{\mu\nu}$ (Section 5.2) records how successive refinements compare when transported across regions whose admissible histories differ in their curvature residue.

Consider two clocks: one at a lower informational potential (higher curvature residue) and one at a higher potential (lower residue). Both clocks

produce sequences of refinements

$$\langle e_1 \prec e_2 \prec \cdots \rangle_{\text{low}}, \quad \langle f_1 \prec f_2 \prec \cdots \rangle_{\text{high}},$$

each internally consistent. However, the Law of Boundary Consistency demands that refinements compared across their shared causal overlap must agree on their informational interval. When the refinement sequence from the lower clock is transported to the higher clock, the compatibility condition forces an adjustment in the rate at which distinguishability is accumulated.

Formally, transport along a connection with residue Γ alters the frequency of refinements according to the first-order compatibility condition of Section 5.4:

$$\nu_{\text{high}} = \nu_{\text{low}} (1 - \Gamma \Delta h),$$

where Δh is the informational separation between the clocks. This is the informational analogue of the frequency shift that appears in the smooth limit as gravitational redshift.

In the Pound–Rebka configuration, a photon (interpreted here as a unit of transported distinguishability) sent upward from the lower clock must be refined in such a way that its informational interval remains constant. Since admissible refinements at higher potential accumulate fewer curvature corrections, the transported signal must appear at a lower frequency when measured by the upper clock. Conversely, a downward signal appears at a higher frequency. No physical field is invoked: the effect is a bookkeeping adjustment required to maintain Martin-consistent transport of distinguishability across regions of differing curvature residue.

Thus the informational framework predicts a frequency shift of the form

$$\frac{\Delta \nu}{\nu} \approx \Gamma \Delta h,$$

which matches the structure of the Pound–Rebka observation when inter-

puted in the smooth shadow of the metric gauge.

The phenomenon of time dilation is therefore an observable outcome of the informational interval and the necessity of refinement-adjusted transport. Differences in curvature residue force clocks at different potentials to accumulate distinguishability at different rates, and the comparison of their refinement counts produces the celebrated redshift.

Coda: The Kinematic Foundation of Geometry

N.B.—This chapter derived the continuous kinematic structures—the metric $g_{\mu\nu}$ and the connection $\Gamma_{\mu\nu}^{\lambda}$ —from the informational requirement that refinements remain globally consistent. No forces, fields, or dynamical assumptions were introduced. \square

The development of this chapter followed the informational chain of emergence:

$$\epsilon \longrightarrow \tau \longrightarrow g_{\mu\nu} \longrightarrow \Gamma_{\mu\nu}^{\lambda}.$$

The refinement bound ϵ fixed the minimal increment of admissible structure. The interval τ encoded the invariant tally of such increments. The metric $g_{\mu\nu}$ enforced this invariance across observers, and the connection $\Gamma_{\mu\nu}^{\lambda}$ preserved it under differential refinement. The observable consequence of this structure is the redshift effect, where nonuniformity of the metric gauge requires a corresponding adjustment of the local ϵ -counting rate.

Phenomenon 5.7.2 (The Event Horizon Effect). **N.B.**—*In ordinary space, you approach the event horizon. In an informational black hole, the event horizon approaches you.* \square

A black hole is not a geometric singularity but an informational bottleneck.

The metric $g_{\mu\nu}$ functions as a gauge of informational separation, and the connection Γ is the bookkeeping rule that adjusts local refinement rates in

order to preserve the invariant informational interval τ (Sections 5.2–5.3) :contentReference[oaicite:0]index=0. Transport is admissible only so long as this gauge can be maintained at finite cost.

At an event horizon this cost diverges. To export a single distinguishable refinement from the interior requires an unbounded number of coordinate-time updates. The exchange rate of admissible refinements collapses to zero.

The classical singularity is therefore not a failure of physics but a failure of mergeability: the causal universe tensor can no longer reconcile the internal and external ledgers in finite informational time.

The interior record continues to refine, but its updates can no longer be interleaved with the external history. A black hole is thus not a hole in space, but a latency horizon in the bookkeeping of causal order.

The event horizon represents a *local* saturation of the transport budget. It is the point at which the informational cost of exporting a refinement diverges with respect to an external region. This divergence does not require curvature to be extreme everywhere; it arises whenever the connection can no longer preserve the invariant interval under admissible exchange.

This observation admits a broader question. If a finite region of the causal ledger can exhaust its outward bandwidth, then a complete ledger — the entire admissible causal universe — must also possess a maximal transport capacity. The issue is therefore not whether horizons exist, but whether a global horizon is forced by the finiteness of refinement itself.

The local phenomenon thus points to a global constraint: if the universe is a finite, admissible record, then there must exist a critical scale at which the cost of exporting any further refinement diverges. This is not a geometric assumption, but a bookkeeping necessity.

The following phenomenon makes this limit explicit.

Phenomenon 5.7.3 (The Schwarzschild Effect [136]). ***Statement.** There exists a critical informational radius beyond which refinements cannot be ex-*

ported in finite time. This radius is the Schwarzschild limit of the causal ledger.

Classical Shadow. In general relativity, the Schwarzschild radius

$$r_s = \frac{2GM}{c^2}$$

defines the surface at which the escape cost of light becomes infinite. In this framework, the same quantity appears not as a geometric boundary, but as an informational one.

Informational Interpretation. Let M_U denote the total admissible mass–energy of the causal record and define the corresponding informational radius

$$r_U = \frac{2GM_U}{c^2}.$$

This radius marks the point at which the cost of transporting a single distinguishable refinement from the interior to any hypothetical exterior diverges.

At this limit, the connection Γ can no longer preserve the invariant informational interval under transport. The exchange rate of admissible events collapses to zero. The interior ledger may continue to refine, but its updates cannot be merged with any external record within finite refinement time.

Consequence. The universe does not sit inside a black hole. Rather, the universe is the maximal admissible ledger: a region whose informational content is bounded by a horizon defined by its own total refinement budget.

The Schwarzschild limit therefore measures not the curvature of space, but the maximum outward bandwidth of causally admissible history.

This completes the kinematic description of informational geometry. The next chapter introduces the dynamic concept of curvature, defined as the obstruction to transporting refinement instructions consistently around a closed loop. In this way, the “Curvature of Information” becomes the natural extension of the kinematic structures developed here.

Chapter 6

Informational Strain

At the end of the previous chapter, we identified *informational stress* as the bookkeeping rule that maintains the invariance of the informational interval under maximal propagation. Stress describes how distinguishability is transported without contradiction. Informational strain is the natural complement of this idea. It measures the failure of that transport to close.

Strain arises when locally admissible refinements cannot be assembled into a globally coherent history without additional adjustment. In the discrete domain, this adjustment is the residue of non-closure. In the smooth shadow, it appears as curvature. Informational strain is therefore the measure of non-integrability of refinement: the discrepancy recorded when a closed cycle of informational updates fails to return to its initial state.

6.1 Historical Review: Curvature as Non-Closure

Classically, curvature has always been understood as non-closure. Gauss demonstrated that curvature can be detected intrinsically, without reference to an embedding [58]. Riemann characterized curvature as the commutator

of two infinitesimal transports [128]. Einstein showed that curvature arises wherever a tensorial quantity must be conserved consistently across overlapping regions [44].

In each case, curvature is the minimal correction needed when parallel transport around a loop does not return the original value. Informational strain is the discrete analogue of this principle. It measures the mismatch generated by transporting informational refinements around a closed cycle. In the smooth shadow, this mismatch becomes curvature of the informational gauge.

Definition 31 (Cross Product [71]). *Let $u, v \in \mathbb{R}^3$ be vectors in Euclidean space. The cross product $u \times v$ is the unique vector in \mathbb{R}^3 satisfying*

$$u \times v \perp u, \quad u \times v \perp v,$$

with magnitude

$$\|u \times v\| = \|u\| \|v\| \sin \theta,$$

where θ is the angle between u and v , and oriented so that $\{u, v, u \times v\}$ forms a right-handed triple.

In coordinates,

$$u \times v = \begin{pmatrix} u_2 v_3 - u_3 v_2 \\ u_3 v_1 - u_1 v_3 \\ u_1 v_2 - u_2 v_1 \end{pmatrix}.$$

6.2 Galerkin Projection and Rotational Residue

The Galerkin method arises in this framework not as a numerical convenience, but as a structural necessity. Refinement updates act on the informational record as finite operators. When these updates are examined in the dense limit, they admit a decomposition into components that either preserve alignment with admissible test functions or deviate from it.

Let $\{\phi_i\}$ denote a finite admissible test basis associated with a refinement scale. The Galerkin projection Π_G of an update operator R is defined by the bilinear pairing

$$\langle \phi_i, \Pi_G R \phi_j \rangle = \langle \phi_i, R \phi_j \rangle,$$

for all admissible test functions. By construction, this pairing is *symmetric*: it records only the component of R that aligns with the test space. Any antisymmetric contribution is annihilated by the projection.

This is not a defect of the method, but its defining feature. Galerkin schemes measure only what can be stabilized by symmetric bilinear forms. They are blind to rotational discrepancy because such discrepancy does not alter energy-like functionals. The kernel of Π_G therefore contains all anti-symmetric residues of refinement.

The informational cross product formalizes precisely this residue. Given two admissible refinement updates R_a and R_b , their antisymmetric difference,

$$R_a R_b - R_b R_a,$$

records the part of their interaction that twists rather than stretches the informational record. This antisymmetric object lies entirely in the kernel of the Galerkin projection and cannot be detected by any symmetric test space.

The Galerkin norm may therefore be generalized as a restriction of the full informational norm:

$$\|R\|_G = \|\tfrac{1}{2}(R + R^*)\|,$$

where only the symmetric component contributes. The complementary part,

$$\tfrac{1}{2}(R - R^*),$$

remains unmeasured by Galerkin methods.

This unmeasured component is not arbitrary. It is structured and neces-

sary: it is the directional residue that prevents the refinement algebra from closing under symmetric detection. The Galerkin cross product is defined as this missing component — the discrete analogue of curl.

In this sense, the Galerkin projection supplies elasticity without rotation, while the informational cross product supplies rotation without elasticity. Together they complete the refinement algebra. The cross product therefore identifies the direction that Galerkin methods must omit and supplies the missing basis vector required for closure of the discrete refinement cycle.

In the smooth shadow, this discrete residue becomes the classical curl operator. What appears in continuum physics as local rotation is, in the informational framework, nothing more than the part of refinement that lies in the kernel of every symmetric projection.

Definition 32 (Galerkin Cross Product). *Let V be a finite-dimensional trial space and let $W \subseteq V$ be a Galerkin test space. Let $B(\cdot, \cdot)$ denote the bilinear form representing the symmetric part of the refinement update in the smooth shadow. The Galerkin cross product is the unique vector $u \times_G v \in V$ satisfying*

$$B(u \times_G v, w) = 0 \quad \text{for all } w \in W,$$

and

$$u \times_G v \notin W.$$

N.B.—*The Galerkin cross product spans the component of the update that lies in the kernel of the symmetric bilinear form. This component cannot be captured by the Galerkin projection and represents the antisymmetric part of the refinement operator.* \square

Concretely, if the update operator Ψ on refinements admits a decomposition

$$\Psi(e)\Psi(f) = S + A,$$

where S is symmetric with respect to $B(\cdot, \cdot)$ and A is antisymmetric, then

$$u \times_G v$$

is the unique vector in the range of A orthogonal (in the Galerkin sense) to all test functions. In the dense limit, $u \times_G v$ converges to the classical cross product in \mathbb{R}^3 and the antisymmetric part A reduces to the curl operator of the associated vector field.

Proposition 10 (Recovery of the Classical Cross Product). **N.B.**—This result uses only Proposition ?? (anti-symmetry of information propagation), Proposition ?? (commutativity of uncorrelant events), and the Reciprocity Dual of Proposition ??. No geometric assumptions are made. \square

Let $\mathbf{X} : V \times V \rightarrow V$ denote the generalized antisymmetric bilinear operator induced by the Informational Interaction Operator of Definition ??. Let $W \subset V$ be any three-dimensional informational subspace that is stable under Martin–Kolmogorov refinement (Definition ??). Then the restriction

$$\mathbf{X}|_{W \times W} : W \times W \rightarrow W$$

is uniquely isomorphic to the classical cross product on \mathbb{R}^3 . Explicitly, for any basis $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ of W compatible with the reciprocity map,

$$u \mathbf{X} v = \begin{vmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix},$$

where $u = u_i \mathbf{e}_i$ and $v = v_i \mathbf{e}_i$.

Interpretation. The familiar $u \times v$ is not assumed. It is the unique refinement-stable Galerkin limit of the informational antisymmetry when restricted to any three-frame permitted by the axioms of measurement.

Proof (Sketch). On the three-dimensional informational subspace $W \subset V$, the informational metric g of Chapter 5 provides a positive-definite bilinear form and hence an identification of W with \mathbb{R}^3 up to isometry. The antisymmetric operator \mathbf{X} is bilinear and satisfies

$$\mathbf{X}(u, v) = -\mathbf{X}(v, u)$$

by Proposition ???. The Reciprocity Dual (Proposition ??) and Definition ?? ensure that \mathbf{X} is compatible with refinement: if u and v are refinement directions, then $\mathbf{X}(u, v)$ is again a refinement direction in W .

On a three-dimensional inner product space (W, g) , any antisymmetric bilinear map

$$\mathbf{X} : W \times W \rightarrow W$$

is determined uniquely (up to a fixed scalar and orientation) by the requirement that $g(\mathbf{X}(u, v), w)$ define a volume form. Informational minimality fixes the normalization and orientation: adding any extra scale or reversing orientation would introduce unobserved structure, contradicting the Axiom of Boltzmann and the countable refinement structure.

Thus there exists a basis $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ of W compatible with the reciprocity map such that, in these coordinates, \mathbf{X} has exactly the coordinate expression of the classical cross product on \mathbb{R}^3 . This yields the determinant formula in the statement and completes the identification. \square

A full proof is provided in Appendix ??.

6.3 Communication [138]

Before any notion of force, field, or medium, there is a simpler problem: What portion of a refinement can survive being written down, carried across a distance, and reconstructed elsewhere?

Every physical theory assumes that something can be transmitted. Light, sound, voltage, or particles are taken to be the carriers, and attention is focused on their propagation. In the present framework, the carrier is irrelevant. What matters is that only a restricted part of any refinement sequence can be stabilized as a record.

Two observers do not share the full informational act of refinement. They share only what can be projected into a common admissible basis. The act of projection destroys structure: it removes precisely those components that do not align symmetrically with the shared test space. The remainder is not an approximation of the original update; it is a different object altogether. It is the message.

This distinction precedes physics. It is not a consequence of bandwidth, noise, attenuation, or engineering limitations. It follows from the axioms: only symmetric, Galerkin-detectable structure can appear in any stable record. Anything else exists only as internal strain.

The earliest large-scale demonstration of this principle appears in wireless telegraphy. In Marconi's transmissions, the physical carrier was electromagnetic, but the deeper phenomenon was informational: what survived across the Atlantic was not a waveform, but a projection. The receiver did not reconstruct the sender's refinement. It recovered only the part that could be stabilized in its own admissible basis.

The following phenomenon isolates that principle in its pure form. It does not depend on radio technology, nor even on electromagnetism. It depends only on the fact that refinement must be made communicable by projection before it can become a message.

Phenomenon 6.3.1 (The Message Effect[26, 104, 151]). *Consider two laboratories, A and B , separated by a large distance. At A , a discrete refinement sequence is encoded as a modulation of an electromagnetic carrier. At B , a detector records only those components of this modulation that admit stable representation in a fixed decoding basis.*

The transmitter at A is free to introduce arbitrary refinements into the signal: phase shifts, amplitude variations, and timing distortions. The receiver at B, however, can only register the symmetric components of that refinement relative to its local basis. Any antisymmetric structure in the transmission lies in the kernel of the decoding projection and is therefore unrecordable.

As the transmission distance is increased, attenuation and noise grow, but the core phenomenon persists independently of physical degradation: only the Galerkin-detectable component of the refinement survives as message. What is received is not the full act of refinement performed at A, but its projected shadow.

The experiment demonstrates the Message Effect: a message is not what is sent, but what can be stably projected into a shared admissible basis. No receiver ever recovers the full refinement of the sender. The unobserved residue — the informational cross component — remains real, but necessarily unsayable.

Viewed this way, communication between observers can be modeled as a Galerkin projection onto a shared test space. Each observer records local refinement updates of the informational record, but agreement is possible only on those components that admit a common representation in the chosen basis. The bilinear forms that define the Galerkin method respond solely to the symmetric component of an update: they measure alignment with the test space and ignore any antisymmetric twist.

The informational cross product records exactly this antisymmetric residue of two refinement updates — the part that twists rather than stretches the record. From the Galerkin point of view, this residue lies in the kernel of the projection and is therefore invisible to every symmetric measurement. This is not a numerical defect but a structural feature: symmetric forms cannot measure rotation. What cannot be seen in the Galerkin norm cannot be communicated through that channel.

In this framework, curl is not a primitive geometric object. It is the

abstraction of refinement itself: the formal recognition that a countable increment may be inserted into a closed refinement cycle without violating the admissibility of the record.

A Galerkin projection enforces communicability. Only symmetric components of an update admit stable representation in a shared basis, and therefore only these components can be exchanged between observers or preserved under global bookkeeping. What survives communication is not the full update, but its compressible shadow.

The informational cross product isolates what is lost under this compression. It is not a force, torque, or dynamical quantity. It is the certificate that two admissible refinement steps do not close when composed. The failure of closure is not an error: it is the necessary room in which a new distinguishable increment can be inserted.

This is the role of curl in the smooth shadow.

Curl is the formal statement that a closed loop of refinement admits a countable defect:

$$\oint R \cdot d\ell \neq 0.$$

This defect is not continuous in origin. It is the shadow of a discrete fact: the informational record permits the insertion of an additional irreducible refinement without contradiction. Curl therefore measures how many new distinctions may be consistently added, not how space physically twists.

In this sense, curl is the abstraction of freedom. Where divergence counts how much structure must be conserved, curl counts how much structure may be created. It measures the remaining capacity of a refinement cycle to accept new distinguishable events.

The Galerkin cross product is the discrete prototype of this phenomenon. It does not compute a vector; it marks a direction in which refinement has not yet been accounted for by any symmetric communicable form. That direction is the basis element that must be adjoined to make the refinement algebra closed under composition.

Thus, communication produces a privileged symmetric subspace, while curl is the algebraic witness that this subspace is incomplete. Curl is not motion. It is admissible novelty: the permission, granted by the axioms, to insert one more countable distinction.

In the smooth limit, this permission appears as rotational structure in a field. In the discrete theory, it is nothing more—and nothing less—than the fact that refinement is not exhausted by what can be communicated.

At this point, the structure is no longer exotic. It is familiar enough to be unsettling. Nothing new has been assumed, no foreign machinery introduced, and no hidden dynamics smuggled in. The construction has relied only on refinement, projection, and admissibility.

However, once things that cannot be projected are treated as real but unsayable, the shape of the argument becomes difficult to unsee—another phenomenon that requires explanation. It is at this point, we can understand how limited measurements truly are because now we are back where we started. There is a single variable left unspecified for an event until the event occurs. At that point, the spline provides just enough free degrees of freedom to make the problem well posed.

N.B.—CAVEAT EMPTOR: Once things that cannot be projected are treated as real but unsayable, the shape of the argument becomes difficult to unsee. The recursion can no longer be ignored, and must now be unwound. See Phenomenon `ph:library-catalog`. \square

6.4 The Time Effect

Time does not enter this construction as a background parameter. It is not a coordinate laid upon events, nor a dimension through which objects move. Time appears only when refinement becomes possible.

At any stage of the informational record, there exists a single free parameter associated with the next admissible event. Prior to occurrence, this

parameter is not speakable. It cannot be projected, communicated, or represented in any admissible basis. It exists only as an open degree of freedom in the spline completion problem.

The perceived flow of time is nothing more than the computation of the free parameter of Proposition 5.

When the parameter is resolved, the spline closure problem becomes well posed. The minimal structure condition selects a unique admissible completion, and the refinement record advances by one step. What was unsayable becomes fixed. What was open becomes recorded.

This process repeats until after the computation of \mathbf{U}_n , $n \leq |e \in \mathbf{U}_n|$. At that point, all events have been sorted. No further refinement is possible.

Phenomenon 6.4.1 (The Time Effect [44, 113]). **N.B.**—*While Newton and Einstein assume time as a primitive, both acknowledge the difficulty of defining change without presupposing it.* \square

Time is not observed as a primitive background quantity but emerges as the moment at which an informational spline becomes well posed. Prior to any event, the refinement record contains a single unsatisfied degree of freedom: a free parameter that cannot be projected, communicated, or represented.

When this parameter is resolved, the admissible spline closes. The minimization of informational structure selects a unique completion of the record, and a new event enters the causal history. This act of closure is experienced as the passage of time.

Thus, time is not motion and not duration. It is the count of successful resolutions of the free spline condition. Each unit of time corresponds to the elimination of one unspeakable degree of freedom and the stabilization of one new admissible event.

The phenomenon of time is therefore the observable shadow of computation: the discrete act of transforming an open refinement into a closed record. What appears as temporal flow is nothing more than the repeated completion

of an otherwise underdetermined spline.

6.5 Informational Viscosity [113]

Phenomenon 6.5.1 (The Navier–Stokes effect [112, 149]). **N.B.**—*This is an informational phenomenon. No physical fluid or continuum is assumed. The classical Navier–Stokes equations are quoted only as the smooth shadow of discrete refinement transport. The phenomenon illustrates that the appearance of viscous terms is nothing more than the accumulation of informational strain under non-closing updates.* \square

Classical fluid dynamics records the transport of a state variable through space and time. The Navier–Stokes equation,

$$\partial_t u + (u \cdot \nabla)u = -\nabla p + \nu \Delta u,$$

is traditionally interpreted as the momentum balance of a viscous medium.

Informationally, this equation expresses something more fundamental: closure requires correction. The convective term $(u \cdot \nabla)u$ represents the pure transport of distinguishability under the refinement map. If refinement closed globally, this transport would suffice. Yet classical convective transport fails to be integrable; small loops do not return the same state. The discrepancy accumulates as informational strain.

The viscous term $\nu \Delta u$ is precisely the correction required to force closure. It is the smooth shadow of the strain operator Σ : the minimal adjustment needed to reconcile locally transported information with a globally coherent record. Viscosity is therefore an informational phenomenon. It is the continuous representation of the curvature induced by non-closure of refinement.

In this interpretation, Navier–Stokes is not a physical law but the canonical example of how strain appears when local informational updates fail to agree. Its form is dictated entirely by the requirement that refinement remain coherent across overlapping regions. The equation emerges as the unique

smooth expression of balancing informational strain.

6.6 Non-linear Informational Strain

Phenomenon 6.6.1 (The Strong Interaction Effect). **Statement.** *There exist refinement regimes in which informational strain is intrinsically non-linear. In such regimes, strain does not disperse; it self-confines.*

Mechanism. *In ordinary transport, the informational strain Σ produced by a refinement update disperses through the ledger and admits a linear smoothing shadow. However, for sufficiently dense configurations of causal threads, the strain functional becomes nonlinear.*

Let q_i denote tightly coupled refinement threads. Define the strain density

$$\Sigma = \Sigma(q_i, q_j).$$

In the strong regime,

$$\Sigma(q_i, q_j) \neq \Sigma(q_i) + \Sigma(q_j),$$

and the superposition principle fails.

As separation increases, the cost of maintaining consistency grows rather than decays. The ledger assigns increasing informational cost to isolated threads, producing a restoring force that prevents independent propagation.

Interpretation. *The strong force is not modeled as exchange of particles, but as a property of nonlinear bookkeeping. Attempts to separate correlated threads generate additional strain rather than relief. The ledger enforces confinement by making isolation informationally inadmissible.*

Conclusion. *Quark confinement is the smooth shadow of a ledger whose strain function is nonlinear. At short scales, symmetry permits limited motion; at large scales, the informational cost diverges. The strong interaction is therefore the law of self-binding strain.*

6.7 The Residue of Inconsistency

Let e_i denote a distinguishable event and $\hat{R}(e_i)$ the restriction operator determining admissible continuations. Under the continuous update map Ψ , successive refinements satisfy

$$U_{i+1} = \Psi(e_{i+1} \cap \hat{R}(e_i)) U_i.$$

If a sequence of events returns to the same informational state after k steps, global coherence requires that the net update be the identity in the informational gauge:

$$\Psi(e_{i+k} \cap \hat{R}(e_{i+k-1})) \cdots \Psi(e_{i+1} \cap \hat{R}(e_i)) = I.$$

When this closure fails, the discrepancy is the *informational strain*. Define the strain operator

$$\Sigma = \Psi(e_{i+k} \cap \hat{R}(e_{i+k-1})) \cdots \Psi(e_{i+1} \cap \hat{R}(e_i)) - I.$$

The operator Σ measures the failure of closure of informational transport. In the dense limit, its leading term becomes the curvature of the informational gauge. Thus, curvature is the smooth representation of discrete informational strain.

Definition 33 (Informational Cross Product [80]). *Let e and f be admissible refinements of an informational state U , and let Ψ be the continuous update operator*

$$U' = \Psi(e) U.$$

The informational cross product of e and f is the event-update operator that measures the failure of the corresponding refinement actions to commute. It is defined by

$$e \times f := \Psi(e) \Psi(f) - \Psi(f) \Psi(e).$$

N.B.—This operator records the non-closure of refinement. If e and f commute informationally, the cross product vanishes. A nonzero result represents the minimal corrective update that must be applied to preserve coherence of the record. \square

In the smooth shadow, $e \times f$ reduces to the classical curl of the associated flow field, and the informational strain generated by a closed loop of refinements is given by the accumulated cross product of the updates.

Proposition 11 (Informational Cross Product as Minimal Discretization).

N.B.—This proposition uses the definitions introduced in Definitions ??, ??, and ??. The only assumptions are informational minimality and the Axiom of Boltzmann, which forbids the introduction of unobserved structure. \square

Let \mathbf{X} be the generalized cross product of Proposition 10, and let \mathbf{X}_G denote the Galerkin cross product obtained by weak-form extremality in the sense of Chapter 3. Let \mathbf{X}_I denote the Informational Cross Product of Definition ??.

Then:

1. \mathbf{X}_G is the unique smooth shadow admitted by spline-level closure and informational reciprocity.
2. \mathbf{X}_I is the unique information-minimal discretization of \mathbf{X}_G that introduces no additional admissible events under refinement.
3. Consequently,

$$\mathbf{X}_I = \text{Disc}_\epsilon(\mathbf{X}_G),$$

the ϵ -refinement discretization of the Galerkin operator.

Interpretation. No additional curvature, torsion, or unobserved structure may be introduced without violating informational minimality. Thus \mathbf{X}_I is the coarsest admissible refinement of the generalized antisymmetry.

Proof (Sketch). By construction, the Galerkin cross product X_G is obtained as the weak-form limit of the refinement commutator in the dense sampling regime: integration by parts and the Galerkin projection remove all components that cannot be detected by the symmetric bilinear form $B(\cdot, \cdot)$, leaving a unique smooth antisymmetric residue compatible with spline closure.

The discretization operator Disc_ϵ is defined so that, for any smooth operator T on the trial space, $\text{Disc}_\epsilon(T)$ is the unique discrete operator whose action agrees with T on all refinement patterns distinguishable at scale ϵ , and differs from T only by terms that would require additional, unrecorded events to detect. In particular, if two discretizations T_1 and T_2 differ on any pattern resolvable at scale ϵ , then the difference encodes additional structure that would need to be measured to be admissible.

Apply this to $T = X_G$. By definition of the Informational Cross Product (Definition ??), X_I is precisely the event-update operator that records the non-commutativity of refinement at the discrete level and vanishes whenever the updates commute. Suppose there existed another discretization \tilde{X} of X_G that differs from X_I on some distinguishable refinement pattern. Then \tilde{X} would encode additional twists not required by the observed failure of commutation, thereby introducing unobserved structure. This contradicts informational minimality.

Hence X_I is the unique discretization compatible with both the Galerkin shadow and the Axiom of Boltzmann. By uniqueness of the discrete operator agreed upon at all ϵ -resolvable patterns, we have

$$X_I = \text{Disc}_\epsilon(X_G),$$

as claimed. □

A full proof is provided in Appendix ??.

Phenomenon 6.7.1 (The Arago Effect). **Statement.** *A bright region appears in the geometric shadow of a circular obstacle because the ledger must*

remain globally consistent along the entire boundary. Local histories are subordinated to global admissibility.

Classical Context. Poisson famously argued that the wave theory of light was absurd because it predicted a bright spot at the center of the shadow of a circular disk, a region that ray optics insisted must be dark. Arago's experimental confirmation of the spot revealed that the absurdity lay not in the prediction, but in the assumption that causal histories could be deleted locally without reference to the global boundary.

Informational Interpretation. The edge of the obstacle forms a closed causal boundary $\partial\Omega$. By the Law of Boundary Consistency (Law 3), the state of the field at any interior point must be the unique refinement compatible with the entire boundary ledger simultaneously.

Along the central axis behind the disk, every point is equidistant from $\partial\Omega$. Because the boundary refinements are symmetric, the Axiom of Ockham (Axiom 3) forbids the introduction of unrecorded phase asymmetries that would force destructive cancellation. To assert darkness at the center would require the ledger to encode hidden distinctions that do not exist in the boundary record.

Therefore the only admissible history is the one in which refinements merge coherently. The bright spot is not the result of waves bending around an object; it is the informational checksum of the boundary. The ledger cannot delete the signal at the center without introducing structure that was never measured. Global consistency overrides the intuition of local blocking.

6.8 The Informational Strain Tensor

Definition 34 (Informational Strain Tensor [23, 38]). *Let U be an informational state transported around a closed refinement cycle. The informational*

strain tensor is the unique multilinear operator \mathcal{S} satisfying

$$U_{\text{final}} - U_{\text{initial}} = \mathcal{S}(U_{\text{initial}}).$$

N.B.—This definition expresses strain as the minimal multilinear correction required to reconcile initial and final informational states after a closed cycle of refinement. In the smooth shadow, \mathcal{S} reduces to the curvature tensor of the informational gauge. \square

The strain tensor captures all second-order incompatibilities that arise from trying to merge locally consistent refinements. Where stress governs the linear transport of distinguishability, strain measures the failure of that transport to be integrable. Strain is thus the obstruction to global coherence inherent in the refinement record itself.

6.9 Unavoidable Strain and the Necessity of Curvature

When local refinements agree on pairwise overlaps but fail on triple overlaps, strain is unavoidable. No ordering of updates or choice of gauge can remove it. This non-closure is the combinatorial analogue of the Bianchi identity: a defect in the associativity of refinement that cannot be eliminated by reparametrization.

Informational minimality ensures that this defect must appear. If inconsistencies were ignored, they would create unrecorded structure, violating the axioms of event selection and informational closure. Thus, the existence of strain is a logical necessity, not a geometric postulate.

In the smooth shadow, unavoidable strain manifests as curvature. In the discrete domain, it is the minimal corrective refinement required to restore global consistency.

6.10 The Law of Curvature Balance

Law 5 (The Law of Curvature Balance). ***N.B.**—This law follows immediately from Proposition 10 and Proposition 11. No geometric postulates are made; curvature arises solely as the residue of informational non-closure. \square*

Let X be the generalized cross product of Proposition 10, and let X_I be its informational minimal discretization from Proposition 11. Let ∇ denote the informational connection of Chapter 5, and let \mathcal{R} denote the curvature operator.

Then for all $u, v, w \in V$,

$$\mathcal{R}(u, v)w = (\nabla_u \nabla_v - \nabla_v \nabla_u - \nabla_{u \mathsf{X}_I v})w.$$

Moreover, the discrepancy

$$\mathsf{S}(u, v) := (u \mathsf{X} v) - (u \mathsf{X}_I v)$$

is exactly the Informational Strain Tensor of Definition ??. Thus

$\text{Curvature} = \text{Informational Strain} = \text{Minimal Non-Closure of the Generalized Cross Product.}$

Interpretation. *Once the generalized antisymmetry reduces to the classical cross product in three dimensions, and once the informational discretization is forced by minimality, the defect of closure cannot be eliminated locally without producing unobserved structure. The axioms therefore require that this residue be balanced globally, yielding curvature as a theorem of measurement.*

By definition of the informational connection ∇ (Chapter 5), parallel transport of an informational state along refinement directions u and v is represented by iterated application of ∇_u and ∇_v . In the smooth shadow, the curvature operator $\mathcal{R}(u, v)$ is the obstruction to exchanging the order of

these transports; classically,

$$\mathcal{R}(u, v)w = (\nabla_u \nabla_v - \nabla_v \nabla_u)w$$

whenever transport closes.

In the informational framework, refinements need not close. The missing update required to restore closure is recorded by the Informational Cross Product: for refinement directions u and v , the operator $u\mathbf{X}_I v$ is exactly the minimal corrective update that measures the failure of the corresponding refinement actions to commute (Definition ??).

Transporting w around a closed refinement loop generated by u and v therefore produces three contributions:

1. the transport $\nabla_u \nabla_v w$,
2. the reversed transport $\nabla_v \nabla_u w$, and
3. the corrective transport along $u\mathbf{X}_I v$ required to maintain coherence.

Global consistency demands that the net update around the loop be measured entirely by the curvature of the informational gauge. Any residual that could be removed by adjusting the connection would represent unrecorded structure and is forbidden by informational minimality.

Thus the true curvature operator $\mathcal{R}(u, v)$ must absorb both the commutator of covariant derivatives and the corrective update along $u\mathbf{X}_I v$:

$$\mathcal{R}(u, v)w = (\nabla_u \nabla_v - \nabla_v \nabla_u - \nabla_{u\mathbf{X}_I v})w.$$

Rewriting the residual update in terms of the Informational Strain Tensor S (Definition ??) shows that S is exactly the tensorial form of the non-closure of refinement, while \mathcal{R} is its smooth representation. The divergence-free condition of the Law of Curvature Balance, $\nabla \cdot S = 0$, then follows from the combinatorial Bianchi-type identity for closed refinement cycles discussed in

Section ??, which expresses that strain cannot accumulate without bound on any admissible global history.

Hence curvature, informational strain, and minimal non-closure of the generalized cross product are three shadows of the same obstruction to refinement closure, completing the proof sketch.

6.11 Flat Rotation Curves [132, 131]

The rotation profile of a galaxy provides an unusually clear window into the informational structure of the causal record. At large radii, the observer is no longer tracking local forces or microscopic dynamics; the only question is how much curvature can be distinguished as the history of a rotating system is transported outward. In the informational framework, this is not a dynamical computation but a question of capacity. The curved portion of the record must be conveyed across increasingly sparse refinement, and the rate at which new curvature can be distinguished is strictly bounded by the Martin condition and the Kolmogorov limit of the observer.

Shannon's theory provides the conceptual template: a channel with finite capacity cannot reproduce arbitrarily rapid variation without error. In the same way, the causal network cannot propagate curvature corrections whose informational rate exceeds the distinguishability bandwidth available at large radius. The classical Keplerian falloff requires an ever-increasing curvature signal to be recorded as the orbital circumference grows, but the observer cannot resolve this increase. Beyond a certain point, additional curvature is informationally invisible.

The result is not a failure of physics but the enforcement of informational minimality. When the curvature demand of the classical profile exceeds the capacity of the refinement channel, the admissible history collapses to the minimal-curvature solution compatible with the record. The velocity curve therefore flattens: not because mass is missing, but because the causal net-

work has exhausted its ability to distinguish any further variation in the curvature ledger.

Phenomenon 6.11.1 (The Flat Rotation Curve Effect [138]). **N.B.**—*This is an informational consequence, not an astrophysical hypothesis. No assumptions regarding dark matter, mass distributions, or Newtonian potentials are invoked. The flattening derived here is the smooth shadow of a discrete consistency requirement: non-commuting refinements produce a curvature residue that appears, in the continuum, as a viscous correction to transport.*

□

N.B.—*The argument presented here is not a dynamical model of galaxies. It is a bandwidth computation in the precise sense of Shannon’s theory of communication [138]. The causal network has a finite capacity to convey distinguishable refinement, and therefore cannot reproduce curvature variations whose informational rate exceeds this capacity. The flattening of the rotation profile reflects this saturation of distinguishability bandwidth, not the presence of unobserved mass or additional physical fields.* □

Every orbit reconstructed from finite measurements consists of two refinement chains: (i) the radial chain of recorded separations, and (ii) the tangential chain of angular distinctions. In an informationally flat geometry these chains commute—refining the radial data then the angular record yields the same admissible completion as refining them in the opposite order.

However, whenever local refinements disagree on their common boundary, or when uncorrelant segments must be merged, the two refinement chains fail to commute. By the Axiom of Ockham, no hidden structure may be inserted to enforce commutativity, and by the Axiom of Boltzmann, the global record must remain coherent. The irreducible mismatch is therefore a viscous residue, the same object defined in Section ?? as informational viscosity.

In the smooth shadow, this residue manifests as a curvature-induced tangential correction. The observable effect is that the angular velocity $v_\theta(r)$ does not decay as $r^{-1/2}$ even when the inferred radial refinements would de-

mand it. Instead, informational viscosity contributes a boundary-consistency correction that remains finite at large radii:

$$v_{\theta}(r) = v_{\text{Newton}}(r) + \eta_{\text{info}} \left(\frac{\partial}{\partial r} \left(\frac{1}{r} \frac{\partial r}{\partial \tau} \right) \right) + \mathcal{O}(\eta_{\text{info}}^2), \quad (6.1)$$

where η_{info} is the informational viscosity parameter introduced in Equation (6.x), and τ is the informational interval of Section ??.

At sufficiently large radii the Newtonian term becomes negligible while the informational-viscosity term remains non-zero, leading to flattened rotation curves.

Phenomenon 6.11.2 (The Angular Momentum Effect). *A bicycle wheel of mass M and radius R is mounted on low-friction bearings. The wheel is brought to a steady rotational speed and its angular velocity is measured using a stroboscope or optical tachometer.*

The angular momentum is then observed and computed from measurable quantities:

$$L = I \omega,$$

where the moment of inertia of the wheel is

$$I \approx MR^2,$$

and the angular velocity is

$$\omega = 2\pi f,$$

with f the observed rotation frequency.

For example, a wheel with

$$M = 2.0 \text{ kg}, \quad R = 0.33 \text{ m}, \quad f = 5 \text{ Hz}$$

has

$$\omega = 31.4 \text{ rad/s}, \quad I \approx 0.218 \text{ kg m}^2, \quad L \approx 6.85 \text{ kg m}^2/\text{s}.$$

This value is not inferred from theory but reconstructed directly from observable mass, geometry, and frequency. The persistence of this quantity under external perturbation constitutes the observational phenomenon of angular momentum.

The observational computation above admits a strictly weaker informational representation. Although the applied influences are linear and act along distinct spatial directions, the admissible record does not require a full two-dimensional description of the induced motion. The record may be compressed by replacing independent linear displacements with a single angular coordinate.

Rather than tracking the motion in a full planar basis, the admissible description collapses to the pair

$$(r, \theta),$$

where r encodes radial admissibility and θ encodes cyclic refinement. The angular component carries half the effective dimensional burden of a Cartesian basis, as the refinement is constrained to a closed orbit.

Informationally, this compression is not an approximation but a necessity: the coherent record cannot sustain independent degrees of freedom once the cyclic constraint becomes admissible. The refinement therefore induces a second-variational structure. After the first (Jacobi) variation fixes the admissible path, the remaining admissible deformations appear only in the angular coordinate.

The persistence of angular momentum is, in this sense, not a force law but a second-variational residue of admissible compression.

6.12 Informational Strain Transport

6.12.1 The Necessity of Strain Bookkeeping

Chapters 5 and 6 established that the structure of the Causal Universe Tensor U is governed by a dynamic balance between *informational stress*—the metric gauge $g_{\mu\nu}$ recording how distinguishability is propagated—and *informational strain* Σ , the residue produced whenever admissible refinements fail to close around a loop. Stress is the kinematic ledger; strain is the curvature-level correction demanded by the Axiom of Ockham and the Axiom of Boltzmann.

The Law of Curvature Balance (Law 5) demonstrated that when two refinement directions do not commute, the resulting discrepancy is not optional bookkeeping: it is an irreducible residue of the informational record. No admissible extension may delete or overwrite this residue, and no unobserved structure may be inserted to cancel it. Thus, curvature is not a field but a *consequence*: a record of failed commutation that must be reconciled by the global merger of histories.

This creates an immediate tension inside the causal network. If Σ cannot be locally eliminated without violating Ockham minimality, and if the network must remain globally coherent under all admissible merges, then the residue cannot stay where it is. It must be *transported*. The informational universe cannot allow curvature strain to accumulate indefinitely at a refinement site, because doing so would force the network to insert additional structure to preserve global consistency—an inadmissible act.

The question, therefore, is unavoidable:

How does the causal network transport uncorrected curvature residue while preserving informational minimality?

The answer follows from a key observation established earlier: different refinement chains incur different informational costs. Some chains require many intermediate updates to preserve consistency; others require almost

none. Among these possibilities, the axioms admit a special class of histories: the *minimal-coupling chains*. These are refinement paths that propagate informational curvature without forcing its resolution. They perform the least amount of bookkeeping necessary to carry Σ forward until a refinement is forced to absorb the residue.

Such chains saturate the maximal admissible propagation speed and interact only when the informational record demands a second-order correction. In the smooth shadow, they behave like nearly interaction-free carriers of curvature strain: the informational analogue of neutrinos.

This leads directly to the phenomenon below. The Neutrino Effect is not a physical hypothesis. It is the smooth shadow of the unique minimal-cost transport mechanism permitted by the axioms of measurement.

Phenomenon 6.12.1 (The Neutrino Effect [55, 75]). **N.B.**—*This informational phenomenon does not appeal to particle physics, standard-model interactions, or any dynamical assumptions about matter. It arises solely from the axioms of distinguishability, refinement minimality, and curvature as the residue of non-commuting refinements.* \square

N.B.—*Astrophysical neutrinos from supernovae are empirically observed to arrive before the concentrated burst of photons. In the informational framework, this is the expected shadow of curvature transport: the curvature residue R travels along admissible chains with a minimal set of permitted interactions. Because only a very small number of refinement events are required to supply the second-order correction, the messenger is effectively interaction-free. Photons, by contrast, must wait for the medium to refine sufficiently to release a coherent burst. Thus, the informational “neutrino” arrives first, providing the curvature fix that guarantees that the later photon record is globally consistent for all observers.* \square

N.B.—*No claim is made regarding the taste, flavor, mouthfeel, bouquet, or organoleptic profile of neutrinos. Any resemblance to sensory modalities is purely metaphorical and should not be construed as a physical assertion.* \square

When two admissible refinement directions fail to commute, the Law of Curvature Balance forces a discrete residue R . By the Axiom of Ockham, no unobserved structure may be introduced to remove this residue, and by the Axiom of Boltzmann, the global causal record must remain coherent. Thus, the residue must be transported until some refinement is forced to resolve it. This curvature-carrying transport behaves, in the smooth shadow, like a nearly undetectable messenger field whose sole role is to deliver the correction required for a consistent reconstruction of the event.

In this sense, the informational neutrino carries not energy or matter but the missing correlants required to ensure that the photon record will reconstruct the same admissible history in every reference frame. Information cannot propagate faster than the maximal admissible refinement speed, but the messenger of curvature strain saturates that speed because it admits almost no intermediate interactions that would delay its progression. Upon arrival, it contributes the precise second-order correction that resolves the non-commutative residue, so that the photon burst—arriving later—is interpreted without ambiguity.

Thus, the Neutrino Effect is the informational shadow of curvature transport: the discrete residue of non-closure moves first, ensuring that the subsequent refinement (carried by photons) is interpreted consistently in every admissible frame, thereby recovering the logic of Einstein's original thought experiment.

Thought Experiment 6.12.1 (Implied Orthogonality and Space-Time).

N.B.—CAVEAT EMPTOR □

The author presents no phenomenon suggesting any structure orthogonal to space-time. Any such language in the surrounding discussion is to be read as set-theoretic rather than geometric.

Rather, the author suggests there is an informational degree of freedom between measurements. See Phenomena ?? and ??.

Coda: Coda: The Informational Stress–Strain Relation

N.B.—Throughout this work, classical differential equations are treated not as fundamental laws but as effects that can be observed. The Navier–Stokes equation is the smooth shadow of the balance between informational stress (transport) and informational strain (non–closure) *citetimoshenko1934*. Nothing in this coda assumes a physical medium; the equation is quoted only as the continuous representation of the bookkeeping required for global coherence under refinement. \square

The path to Navier–Stokes begins with the simplest of all mechanical ideas: statics. In classical statics, a system is said to be in equilibrium when the sum of forces vanishes. Nothing moves, nothing deforms, and the internal ledger of stresses balances exactly. Every contribution is accounted for, and the record closes without residue. This is the mechanical expression of coherence.

In the informational setting, the same idea appears at the level of refinement. A static configuration is one in which the admissible distinguishability does not change. The update operator is the identity, the strain operator Σ vanishes, and no correction is required to maintain consistency. Statics is therefore the trivial case of informational stress and strain: transport is absent, and closure is automatic.

The transition from statics to dynamics occurs the moment transport is introduced. Once distinguishability begins to propagate, the stress ledger no longer balances by default. Refinements may fail to close, and the mismatch accumulates as informational strain. Classical mechanics responds to this imbalance by introducing inertial terms, pressure forces, and viscous corrections. In the informational picture, these are not imposed laws but the minimal bookkeeping required to restore coherence when transport is present.

Navier–Stokes arises precisely from this requirement. It is the statement that the stress generated by transport must be balanced by the strain required to correct its non-closure. The left-hand side of the equation records the informational stress of convective propagation; the right-hand side records the informational strain needed to enforce global compatibility. In the limit where refinements are dense and their residues are approximated by differential operators, the balance of these quantities becomes the familiar continuity equation of fluid dynamics.

Thus, Navier–Stokes is not a departure from statics but its extension. It is the natural generalization of equilibrium to situations in which information is moving. Statics states that the stress ledger must close when nothing changes. Navier–Stokes states that the ledger must still close when everything does.

The informational interpretation of Navier–Stokes follows directly from the definitions of stress and strain developed in this chapter. The transport of distinguishability under the update map Ψ generates informational stress: the left-hand side of the classical equation,

$$\partial_t u + (u \cdot \nabla)u,$$

represents the linear propagation of admissible refinements. If this transport were globally integrable, no additional correction would be needed.

However, convective transport is not integrable in general. Closed loops of refinement do not return to their initial state. The mismatch accumulates as informational strain. In the smooth shadow, the required correction appears as the right-hand side of the Navier–Stokes equation,

$$-\frac{1}{\rho} \nabla p + \nu \Delta u,$$

where the pressure term enforces compatibility with local volume constraints and the viscous term $\nu \Delta u$ is the continuous representation of the strain op-

erator Σ of Section ?? . Viscosity is therefore an informational phenomenon: the amount of correction required to neutralize non-closure and restore global consistency.

In this sense, Navier–Stokes is an informational stress–strain relation. Transport generates the stress; non-closure generates the strain; and the viscous term is the minimal second-order correction needed to reconcile them.

6.13 Informational Angular Momentum

Rotational structure emerges as the final classical observable invariant before nonlocal refinement modes appear. Unlike linear displacement, which may be decomposed into independent observational updates, cyclic motion imposes a global coherence constraint on the admissible record. Once a measurement history admits closed refinement paths, the record can no longer be described by independent translations alone. A persistent residual is forced by the requirement of consistency under cyclic transport. This residual is not introduced as a physical postulate, but appears as an observational necessity.

The Clay Navier–Stokes Problem in Informational Form

N.B.—The following description restates the classical Clay Institute problem in the language of informational transport. No claim of resolution is made. The problem is quoted for context only. \square

Let $u(x, t)$ be the informational velocity field representing the smooth shadow of refinement transport on \mathbb{R}^3 . The Clay problem concerns whether solutions to the balance equation

$$\partial_t u + (u \cdot \nabla)u = -\nabla p + \nu \Delta u, \quad \nabla \cdot u = 0,$$

exist for all time and remain smooth when the initial data are finite and sufficiently regular.

In informational terms, the problem may be phrased as follows:

Does the balance between informational stress and informational strain admit a globally coherent smooth shadow for all time, or can the strain operator Σ accumulate without bound, producing a breakdown of the continuous representation even when the discrete refinement record remains well-defined?

Equivalently: does the correction $\nu \Delta u$ always suffice to control non-closure, or can convective transport accumulate strain faster than viscosity can dissipate it?

N.B.—A finite-time singularity in the classical equation corresponds, in the informational picture, to the divergence of the smooth shadow of strain. It does not imply a contradiction in the underlying discrete refinement record, but indicates that the continuum approximation has ceased to track it. \square

The Clay problem therefore asks whether informational stress and informational strain can remain in balance for all time under dense refinement, or whether the continuous representation can fail even when the discrete theory remains coherent.

Is the existence of quantum theory logically necessary? The author suggests that the differential equations may fail at small enough resolution. See Phenomenon ?? and Axiom 6. Resonance is sometimes real, sometimes Gibb's phenomena.

Chapter 7

Informational Symmetry

The classical description of the universe specifies an ideal relation between informational stress and strain: a perfectly balanced ledger in which causal order is preserved at the limit of distinguishability. But the universe we observe is never perfectly smooth. Measurements are discrete, refinements occur finitely, and the strain introduced by each new event cannot match the ideal stress profile implied by the causal structure. The resulting mismatches are not observational errors. They are the informational stress residues produced by finite refinement—the quantum fields of the theory.

A quantum field arises whenever the invariants of the Causal Universe Tensor are permitted to vary locally while maintaining global Martin consistency. Each allowed fluctuation corresponds to a redistribution of causal order between neighboring observers. The field is therefore not an additional substance laid over spacetime but a dynamic adjustment of the gauge itself, mediating the exchange of distinguishability across finite domains.

In this framework, the traditional wavefunction reappears as the probability amplitude for maintaining order under repeated finite observations. Its complex phase represents the orientation of the causal gauge in informational space, while its magnitude measures the stability of that order. The principle of superposition follows directly from the linearity of causal combinations:

multiple consistent histories can coexist until observation resolves a single extension of the network.

Quantization enters as the recognition that order cannot be subdivided indefinitely. Every causal update exchanges a finite unit of distinguishability—a discrete increment of information. The Planck constant \hbar expresses this minimal step size: the smallest action through which the universe can modify its own gauge while remaining consistent. The commutation relations of quantum theory are therefore expressions of finite causal resolution, not axioms of measurement.

This chapter develops these ideas systematically. Beginning with the Noether currents of the causal gauge, we derive the corresponding quantum fields as their discrete fluctuations. We then show how these fields propagate through the Causal Universe Tensor, producing the familiar quantum wave equations as conditions of statistical Martin consistency. Finally, we interpret entanglement as the correlated selection of events across overlapping causal neighborhoods—the quantum signature of global order maintained through finite means.

7.1 The Photoelectric Effect

The interaction of light with matter provides one of the sharpest tests of the distinction between continuous fields and discrete refinement. A classical wave can transport phase and energetic stress smoothly across a surface, but a measurement device cannot record this continuum directly. The cathode does not respond to fractions of a refinement; it either registers a new event or it does not. The transition from field to detection is governed entirely by the admissibility of refinement: the local informational stress must exceed the surface’s minimal distinguishability cost before a new event can be appended to the causal record.

This is the essence of the photoelectric effect. Increasing the field’s in-

tensity scales the strain imposed on the surface and therefore the *rate* at which admissible events may occur, but it does nothing to lower the distinguishability threshold itself. Conversely, raising the frequency increases the stress carried per cycle and determines whether the predicate “an electron is emitted here” is admissible at all. The phenomenon thus reveals a deep structural principle of the informational framework: continuous fields govern the distribution of stress, but the creation of events depends on whether that stress can overcome the discrete cost of refinement.

Seen in this light, the photoelectric effect is not a mystery or a paradox. It is the natural consequence of attempting to refine a discrete causal record with a continuous source of strain. The threshold and linear kinetic-energy law simply express the bookkeeping conditions a surface must satisfy whenever a continuous wave induces a discrete update to the informational ledger.

Phenomenon 7.1.1 (The Photoelectric Effect [72, 115]). **N.B.**—*This threshold condition is the informational analogue of the Nyquist sampling limit: below a critical frequency, the surface cannot resolve the delivered stress into a distinguishable refinement, and no event can be registered [72].* \square

Statement. *The photoelectric threshold and the linear kinetic-energy law record a fundamental feature of measurement: a continuous field may transport phase and energetic strain, but the act of detection terminates the wave by selecting a discrete refinement of the causal record. Only predicates that exceed a minimal distinguishability cost can produce an admissible event.*

Key relation.

$$K_{\max} = h\nu - \Phi, \quad \nu \geq \nu_0 = \frac{\Phi}{h}.$$

The threshold condition expresses that the cathode surface admits no refinement whose informational stress falls below the work function Φ . The residual energy after satisfying this cost appears as kinetic energy of the emitted

electron.

Reciprocity framing. *A continuous electromagnetic field distributes phase and informational stress smoothly across the surface, but an emission event is a refinement of the partition $P_n \rightarrow P_{n+1}$ at a specific site on the cathode. The selection rule imposes conservation in the bookkeeping channel: the registry of a new event requires payment of the surface's minimal distinguishability cost. Below threshold, the strain induced by the field is insufficient to overcome this cost, and no admissible refinement exists. Above threshold, the refinement proceeds and the excess stress is released as electron kinetic energy.*

Operational consequence. *Intensity controls the rate of refinement by modulating how often the local stress crosses the admissibility bound, but frequency controls the possibility of refinement by determining whether the predicate “an electron is emitted here and now” can be made consistent with the work function. Thus the photoelectric effect distinguishes clearly between the cumulative action of a continuous field and the discrete accounting of event creation: one governs flux, the other governs admissibility.*

7.2 The Action Functional

The action functional provides the statistical completion of the causal gauge. It measures the total consistency of a causal configuration across all finite observations. In the classical limit, the action is stationary: each variation vanishes, and the universe evolves along trajectories of perfect causal balance. In the quantum regime, these variations accumulate as finite fluctuations of order, and the path integral of all such histories defines the observable field.

7.2.1 Definition from the Causal Universe Tensor

Let $\mathcal{T}^{\mu\nu}$ denote the Causal Universe Tensor, whose scalar invariants measure the degree of causal consistency. The *action functional* \mathcal{S} is defined as the

integral of these invariants over the causal domain:

$$\mathcal{S} = \int \mathcal{L}(\mathcal{T}^{\mu\nu}, g_{\mu\nu}, \nabla_\lambda \mathcal{T}^{\mu\nu}) \sqrt{-g} d^4x.$$

The Lagrangian density \mathcal{L} encodes the local rule by which order is preserved and exchanged. In the classical limit, $\delta\mathcal{S} = 0$ reproduces the field equations of the gauge of light; in the quantum limit, \mathcal{S} fluctuates discretely by units of \hbar , reflecting the minimal step size in causal adjustment.

7.2.2 Physical Interpretation

The action \mathcal{S} plays the role of a global consistency measure. Each admissible history of the universe contributes a complex amplitude

$$\Psi[\mathcal{T}] \propto e^{i\mathcal{S}[\mathcal{T}]/\hbar},$$

representing the phase of causal order associated with that configuration. When summed over all histories consistent with Martin's Axiom, these amplitudes interfere, and the stationary-phase paths correspond to the classical trajectories of least action. The non-stationary contributions produce the quantum corrections—the finite discrepancies among partially consistent causal extensions.

In this interpretation, \hbar is not an arbitrary constant but the fundamental unit of distinguishability in causal evolution. It measures the minimal action by which the universe can update its gauge without violating order. The classical limit $\hbar \rightarrow 0$ corresponds to infinitely fine causal resolution, while the quantum limit expresses the graininess of finite observation.

7.2.3 Noether Currents of the Causal Gauge

Symmetries of the Lagrangian correspond to invariances of causal order. By Noether's theorem, each continuous symmetry yields a conserved current

$$J^\mu = \frac{\partial \mathcal{L}}{\partial(\nabla_\mu \phi)} \delta \phi, \quad \nabla_\mu J^\mu = 0.$$

These currents are the quantum fields' classical shadows: energy, momentum, and charge arise as conserved flows of causal order through the network. Their quantization in subsequent sections will describe the discrete exchange of distinguishability among interacting observers.

Remark 18. *The action functional is the expectation value of Martin consistency over all admissible histories. In the classical regime, it is stationary; in the quantum regime, it oscillates. The universe, viewed through this lens, is a sum over self-consistent paths, each differing from the others by integral multiples of the minimal action \hbar . Quantum mechanics is therefore not a separate theory but the statistical theory of finite causal order.*

7.3 The Law of Combinatorial Symmetry

The informational framework developed thus far admits no geometric structure and assumes no group-theoretic symmetries. Nevertheless, when two finite records are mutually compatible, their joint refinement exhibits a canonical structure: the set of distinguishable events in one record can be placed in bijection with the distinguishable events of the other in a way that preserves refinement, ordering, and consistency. This bijection is unique and is determined entirely by the combinatorial structure of the records.

Law 6 (The Law of Combinatorial Symmetry). *Let ψ and ϕ be two finite, non-contradictory records that admit a globally coherent merge under the Ax-*

iom of Event Selection. Then there exists a unique bijection

$$\chi : \text{Ref}(\psi) \rightarrow \text{Ref}(\phi)$$

with the following properties:

- (1) χ preserves distinguishability: $e_1 \neq e_2$ in ψ if and only if $\chi(e_1) \neq \chi(e_2)$ in ϕ .
- (2) χ preserves refinement order: if $e_1 \prec e_2$ in ψ then $\chi(e_1) \prec \chi(e_2)$ in ϕ .
- (3) χ preserves admissibility: for every admissible refinement e' of an event e in ψ , the event $\chi(e')$ is an admissible refinement of $\chi(e)$ in ϕ .
- (4) Any other bijection between $\text{Ref}(\psi)$ and $\text{Ref}(\phi)$ violates refinement compatibility or introduces distinguishable structure inconsistent with the axioms.

Thus all observable symmetries arise from the unique combinatorial structure of refinement: symmetry is not geometric or algebraic but a bijection on the poset of distinguishable events.

N.B.—This law identifies symmetry as an informational phenomenon. No metric, manifold, or group structure is assumed or required. Apparent continuous symmetries emerge only as the limiting shadows of these combinatorial bijections under refinement. \square

7.4 The Application of Noether

Once the action functional has been defined, its symmetries determine the quantities that remain conserved under causal evolution. This is the content of Noether's theorem, here understood as the statistical mechanics of invariance: whenever the ensemble of admissible causal configurations possesses a continuous symmetry, the expectation value of the corresponding quantity remains fixed across all Martin-consistent histories.

7.4.1 Symmetry and Conservation as Statistical Identities

Let the partition function of the causal gauge be written

$$Z = \int \exp\left(\frac{i}{\hbar} \mathcal{S}[\mathcal{T}]\right) \mathcal{D}\mathcal{T},$$

where the integration ranges over all locally consistent configurations of the Causal Universe Tensor. An infinitesimal transformation of variables $\mathcal{T} \rightarrow \mathcal{T} + \delta\mathcal{T}$ that leaves the measure and the action invariant,

$$\delta\mathcal{S} = 0,$$

implies that the partition function is unchanged:

$$\delta Z = 0.$$

Differentiating under the integral sign yields the statistical conservation law

$$\langle \nabla_\mu J^\mu \rangle = 0,$$

where

$$J^\mu = \frac{\partial \mathcal{L}}{\partial(\nabla_\mu \phi)} \delta\phi$$

is the current associated with the transformation. Thus, each continuous symmetry of the Lagrangian corresponds to a conserved flux of causal order. Energy, momentum, and charge appear not as primitive physical entities but as statistical invariants of the causal ensemble.

7.4.2 Conserved Quantities of the Causal Gauge

1. **Translational invariance** \rightarrow Conservation of energy–momentum:

$$\nabla_\mu T^{\mu\nu} = 0.$$

2. **Rotational invariance** \rightarrow Conservation of angular momentum:

$$\nabla_\mu J^{\mu\nu} = 0, \quad J^{\mu\nu} = x^\mu T^{\nu\lambda} - x^\nu T^{\mu\lambda}.$$

3. **Internal phase invariance** \rightarrow Conservation of charge:

$$\nabla_\mu j^\mu = 0.$$

Each of these laws arises from a symmetry of the Causal Universe Tensor under transformations that leave the causal measure invariant. In this sense, Noether's theorem is the thermodynamics of causal order: it equates symmetry with conservation and conservation with informational equilibrium.

Phenomenon 7.4.1 (The Harmonic Oscillator Revisted [121]). *The harmonic oscillator is the minimal causal system in which measurement and variation form a reversible cycle. Let $U(t)$ denote the measured amplitude of a single mode of the universe tensor. Successive reciprocal updates obey*

$$\delta^2 U + \omega^2 U = 0,$$

where δ is the discrete variation operator and ω characterizes the curvature of the local informational potential. In the continuum limit this becomes

$$\frac{d^2 U}{dt^2} + \omega^2 U = 0,$$

the familiar harmonic–oscillator equation.

Each half-cycle corresponds to an exchange between distinguishability and

variation: when the system reaches maximal distinction (turning point), the variation vanishes; when the distinction is minimal (crossing through zero), variation is maximal. The energy functional

$$E = \frac{1}{2} \left[(\dot{U})^2 + \omega^2 U^2 \right]$$

is the invariant scalar of this causal pair— the quantity preserved under all order-preserving updates.

Quantization follows from the Axiom of Finite Observation: only discrete counts of distinguishable configurations fit within one causal period. Applying the Reciprocity Law yields the spectrum

$$E_n = \hbar\omega \left(n + \frac{1}{2} \right),$$

showing that each oscillation cycle admits an integer number of informational quanta plus a residual half-count from causal incompleteness.

In this view, the harmonic oscillator is the archetype of finite reciprocity: a closed loop in which measurement and variation exchange roles while preserving total informational curvature. All quantized fields—phonons, photons, and normal modes of the causal tensor—are higher-dimensional extensions of this single reciprocal circuit.

7.4.3 Statistical Interpretation

In the quantum regime, these conservation laws are satisfied only in expectation. The ensemble of finite causal updates explores neighboring histories whose individual actions differ by multiples of \hbar , but the average fluxes of order remain constant. The classical conservation laws emerge as the limit in which fluctuations of the action vanish and every observer's measurement agrees. Quantum mechanics, in contrast, records the statistics of these fluctuations.

Remark 19. *Noether's theorem closes the loop between mechanics and statistics. Every symmetry of the causal gauge produces a conserved current, and every conservation law describes equilibrium in the flow of distinguishability. In this sense, the field equations of physics are nothing more than the statistical statements of Martin consistency expressed through symmetry.*

ectionConservation

Conservation laws follow from symmetries of the action. In the causal framework, these are statements that the bookkeeping of distinguishability is invariant under relabelings that shift the record in space or time. The resulting Noether currents are the conserved flows of causal order.

7.4.4 Translations and the Stress–Energy Tensor

Let $\mathcal{S} = \int \mathcal{L} \sqrt{-g} d^4x$ be the action of the Causal Universe Tensor fields (collectively ϕ). Under an infinitesimal spacetime translation $x^\mu \mapsto x^\mu + \varepsilon^\mu$, the fields transform as $\delta\phi = \varepsilon^\nu \nabla_\nu \phi$ and $\delta\mathcal{L} = \varepsilon^\nu \nabla_\nu \mathcal{L}$. Invariance of the action ($\delta\mathcal{S} = 0$) yields the Noether current

$$J^\mu{}_\nu = \frac{\partial \mathcal{L}}{\partial(\nabla_\mu \phi)} \nabla_\nu \phi - \delta^\mu{}_\nu \mathcal{L},$$

whose covariant divergence vanishes:

$$\nabla_\mu J^\mu{}_\nu = 0.$$

Identifying $T^\mu{}_\nu \equiv J^\mu{}_\nu$ (or its symmetrized Belinfante form when needed) gives the *stress–energy tensor* with

$$\nabla_\mu T^\mu{}_\nu = 0.$$

In local inertial coordinates this reduces to the familiar continuity laws $\partial_\mu T^{\mu\nu} = 0$.

Phenomenon 7.4.2 (The Compton Scattering Effect [27]). ***Statement.** The Compton shift measures the finite difference of momentum across an event pair, i.e. the reciprocity map in momentum space.*

Key relation.

$$\Delta\lambda \equiv \lambda' - \lambda = \frac{h}{m_e c} (1 - \cos \theta).$$

***Reciprocity framing.** One detection event refines the joint partition of (photon, electron). Bookkeeping enforces the Noether current (translation symmetry) at the refinement:*

$$p_\gamma + p_e = p'_\gamma + p'_e, \quad E_\gamma + E_e = E'_\gamma + E'_e.$$

Eliminating the electron internal variables yields the observed $\Delta\lambda$, a scalar invariant of the event contraction.

***Operational consequence.** The shift is the measured residue after enforcing equality of conjugate Noether charges at a single refinement step.*

7.4.5 Energy and Momentum Densities

Write u^μ for the future-directed unit normal to a Cauchy slice Σ (with volume element $d\Sigma_\mu = u_\mu d^3x \sqrt{\gamma}$). The total four-momentum is

$$P^\nu = \int_\Sigma T^{\mu\nu} d\Sigma_\mu,$$

so that

$$E \equiv P^0 = \int_\Sigma T^{\mu\nu} u_\mu \xi_\nu^{(t)} d^3x \sqrt{\gamma}, \quad \mathbf{P}^i = \int_\Sigma T^{\mu\nu} u_\mu \xi_\nu^{(i)} d^3x \sqrt{\gamma},$$

where $\xi^{(t)}$ and $\xi^{(i)}$ denote the time and spatial translation generators (Killing vectors in symmetric backgrounds). Covariant conservation implies slice-

independence:

$$\frac{d}{d\tau} P^\nu = \int_{\Sigma} \nabla_\mu T^{\mu\nu} d\Sigma = 0.$$

7.4.6 Bookkeeping Interpretation

Causally, $\nabla_\mu T^{\mu\nu} = 0$ is a statement that *what leaves one finite neighborhood must enter another*. The stress–energy tensor tallies the flow of distinguishability through the network; its vanishing divergence is the ledger’s balance condition. Translational symmetry means we can shift the labels of events without changing that tally. Conservation of *energy* is the invariance of the temporal bookkeeping column; conservation of *momentum* is the invariance of the spatial columns. In discrete form, for any compact region \mathcal{R} with boundary $\partial\mathcal{R}$,

$$\frac{d}{d\tau} \int_{\mathcal{R}} T^{0\nu} d^3x = - \int_{\partial\mathcal{R}} T^{i\nu} n_i dS,$$

so the time rate of change of the “inventory” inside equals the net outward flux across the boundary—pure bookkeeping.

7.4.7 Curved Backgrounds and Killing Symmetries

When the metric varies, conserved charges are tied to spacetime symmetries. If ξ^ν is a Killing vector ($\nabla_{(\mu}\xi_{\nu)} = 0$), then

$$\nabla_\mu (T^\mu{}_\nu \xi^\nu) = 0,$$

and the associated charge

$$Q[\xi] = \int_{\Sigma} T^\mu{}_\nu \xi^\nu d\Sigma_\mu$$

is conserved. Energy arises from time-translation symmetry ($\xi = \partial_t$), momentum from spatial translations, and angular momentum from rotations. In each case, the “conservation law” is precisely the statement that the ledger

of scalar invariants computed by the Causal Universe Tensor is unchanged under the corresponding relabeling of events.

Remark 20. *Conservation is not mysterious dynamics; it is consistency of accounting. Noether's theorem says: if the rules for keeping the ledger do not change when we shift the page in space or time, then the totals on that page do not change either. In the causal calculus, those totals are P^ν , and their invariance is exactly $\nabla_\mu T^{\mu\nu} = 0$.*

Phenomenon 7.4.3 (The Conservation of Energy [114]). *Consider a real Klein–Gordon field ϕ in flat spacetime with*

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m^2 \phi^2, \quad \eta_{\mu\nu} = \text{diag}(-, +, +, +).$$

The (symmetric) stress–energy tensor is

$$T^{\mu\nu} = \partial^\mu \phi \partial^\nu \phi - \eta^{\mu\nu} \mathcal{L}.$$

Energy density and energy flux are then

$$\mathcal{E} \equiv T^{00} = \frac{1}{2} \left(\dot{\phi}^2 + |\nabla \phi|^2 + m^2 \phi^2 \right), \quad S^i \equiv T^{0i} = \dot{\phi} \partial^i \phi.$$

Continuity (bookkeeping) equation. *Using the Euler–Lagrange equation $\square \phi + m^2 \phi = 0$ and differentiating,*

$$\partial_t \mathcal{E} = \dot{\phi} \ddot{\phi} + \nabla \phi \cdot \nabla \dot{\phi} + m^2 \phi \dot{\phi} = \dot{\phi} (\ddot{\phi} - \nabla^2 \phi + m^2 \phi) + \nabla \cdot (\dot{\phi} \nabla \phi) = \nabla \cdot (\dot{\phi} \nabla \phi),$$

so

$$\partial_t \mathcal{E} + \nabla \cdot (-\dot{\phi} \nabla \phi) = 0 \quad \Longleftrightarrow \quad \partial_\mu T^{\mu 0} = 0.$$

This is pure bookkeeping: the time rate of change of energy density equals the negative divergence of the energy flux.

Integrated conservation law. *Integrate over a fixed region \mathcal{R} with outward normal \mathbf{n} :*

$$\frac{d}{dt} \int_{\mathcal{R}} \mathcal{E} d^3x = - \int_{\partial\mathcal{R}} \mathbf{S} \cdot \mathbf{n} dS.$$

If fields vanish (or are periodic) on the boundary so the surface term is zero, then the total energy

$$E = \int_{\mathbb{R}^3} \mathcal{E} d^3x$$

is conserved: $\frac{dE}{dt} = 0$.

Causal bookkeeping interpretation. T^{00} tallies the “inventory” of distinguishability stored in a region (kinetic + gradient + mass terms). The flux T^{0i} records how that inventory flows across the boundary. The continuity equation says the ledger balances exactly: what leaves here enters there. Translation invariance is the statement that the rules of this ledger do not change when we shift the page in time; hence the total energy remains the same.

Phenomenon 7.4.4 (The Feynman Diagram [51]). *In conventional quantum field theory, perturbation expansions of the generating functional are represented diagrammatically: vertices encode local interactions and propagators connect them according to the causal structure of spacetime. In the causal formulation developed here, the same construction arises directly from the Universe Tensor.*

Each vertex corresponds to an event tensor $E_k \in T(V)$ contributing a measurable distinction within the causal order. A propagator corresponds to an admissible contraction between event tensors—a bilinear map

$$\langle E_i, E_j \rangle = \text{Tr}(E_i^\top G E_j),$$

where G is the causal propagator enforcing Martin consistency between the connected events. The complete amplitude for a process is therefore the con-

traction of the ordered product

$$U_n = \sum_{k=1}^n E_k,$$

with all admissible propagators. The resulting scalar invariants of U_n constitute the measurable quantities of the theory.

Thus, a Feynman diagram is the graphical representation of a tensor contraction in the causal algebra: each diagram corresponds to one term in the finite expansion of the Universe Tensor, and summing over all diagrams is equivalent to enforcing global consistency of causal order. What appears in standard field theory as a perturbation series is, in this formalism, a finite enumeration of distinguishable causal relations—a bookkeeping identity derived from the Reciprocity Law rather than using calculus.

7.5 Angular Momentum and Spin

Rotational (and more generally Lorentz) invariance of the action produces a conserved tensorial current whose charges are the total angular momentum. Decomposing that current separates *orbital* from *spin* contributions; their sum is conserved.

7.5.1 Noether Current for Lorentz Invariance

Let the action $\mathcal{S} = \int \mathcal{L}(\phi, \nabla\phi, g) \sqrt{-g} d^4x$ be invariant under infinitesimal Lorentz transformations $x^\mu \mapsto x^\mu + \omega^\mu{}_\nu x^\nu$ with antisymmetric $\omega_{\mu\nu} = -\omega_{\nu\mu}$, and induced field variation $\delta\phi = -\frac{1}{2}\omega_{\rho\sigma}\Sigma^{\rho\sigma}\phi - \omega^\mu{}_\nu x^\nu \nabla_\mu \phi$, where $\Sigma^{\rho\sigma}$ are the generators on the fields. Noether's theorem yields the (canonical) angular-momentum current

$$J_{\text{can}}^{\lambda\rho\sigma} = x^\rho T_{\text{can}}^{\lambda\sigma} - x^\sigma T_{\text{can}}^{\lambda\rho} + S^{\lambda\rho\sigma}, \quad \partial_\lambda J_{\text{can}}^{\lambda\rho\sigma} = 0,$$

with canonical stress tensor $T^\lambda_{\nu,\text{can}} = \frac{\partial \mathcal{L}}{\partial(\partial_\lambda \phi)} \partial_\nu \phi - \delta^\lambda_\nu \mathcal{L}$ and spin current

$$S^{\lambda\rho\sigma} = \frac{\partial \mathcal{L}}{\partial(\partial_\lambda \phi)} \Sigma^{\rho\sigma} \phi = -S^{\lambda\sigma\rho}.$$

Phenomenon 7.5.1 (The Spin- $\frac{1}{2}$ Effect [40]). *Spin- $\frac{1}{2}$ particles arise when the local symmetry of the universe tensor is represented not on spacetime vectors but on their double cover. Under a full 2π rotation, the causal ordering of distinguishable events reverses sign before returning to its original configuration after 4π . This two-valuedness expresses the fundamental antisymmetry of distinction.*

Let $\psi(x)$ denote a two-component field that transports the minimal unit of causal orientation. Its dynamics follow from the Lorentz-invariant action

$$\mathcal{L} = \bar{\psi}(i\gamma^\mu D_\mu - m)\psi,$$

where D_μ is the gauge-covariant derivative and the γ^μ generate the Clifford algebra

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}.$$

Each γ^μ acts as a local operator of causal rotation: applying it changes the orientation of the measurement frame while preserving causal order. Because the algebra squares to unity only after two applications, a single 2π rotation introduces a minus sign, $\psi \rightarrow -\psi$, revealing that the physical state is defined on the double cover $\text{Spin}(3,1)$ of the Lorentz group.

In the informational picture, the two components of ψ encode the forward and reverse orientations of causal distinction—measurement and variation. The spinor's phase thus records how the act of observation twists within the causal network. Quantized angular momentum

$$S = \frac{\hbar}{2}$$

emerges as the minimal unit of such rotational bookkeeping: the smallest nontrivial representation of reciprocity under continuous rotation.

Spin $-\frac{1}{2}$ therefore exemplifies the finite, antisymmetric nature of causal orientation. A complete 4π turn is required for full restoration of distinguishability, making the spinor the algebraic expression of the universe tensor's two-sheeted structure in orientation space.

7.5.2 Belinfante–Rosenfeld Improvement

The canonical $T_{\mu\nu}$ need not be symmetric. Define the Belinfante superpotential

$$B^{\lambda\rho\sigma} = \frac{1}{2} \left(S^{\rho\lambda\sigma} + S^{\sigma\lambda\rho} - S^{\lambda\rho\sigma} \right), \quad B^{\lambda\rho\sigma} = -B^{\lambda\sigma\rho}.$$

The *improved* symmetric stress tensor and current are

$$T_B^{\mu\nu} = T_{\text{can}}^{\mu\nu} + \partial_\lambda \left(B^{\lambda\mu\nu} - B^{\mu\lambda\nu} - B^{\nu\lambda\mu} \right), \quad J_B^{\lambda\rho\sigma} = x^\rho T_B^{\lambda\sigma} - x^\sigma T_B^{\lambda\rho},$$

and obey $\partial_\lambda T_B^{\lambda\nu} = 0$, $\partial_\lambda J_B^{\lambda\rho\sigma} = 0$. The spin density has been absorbed into a symmetric T_B so that the total angular momentum current is purely “orbital” in form; its integrated charge still equals *orbital* + *spin*.

7.5.3 Conserved Charges

For a Cauchy slice Σ with normal u_λ ,

$$M^{\rho\sigma} = \int_\Sigma J^{\lambda\rho\sigma} d\Sigma_\lambda = \int_\Sigma \left(x^\rho T_B^{\lambda\sigma} - x^\sigma T_B^{\lambda\rho} \right) d\Sigma_\lambda, \quad \frac{d}{d\tau} M^{\rho\sigma} = 0.$$

In 3D language (flat space, $u_\lambda = (1, 0, 0, 0)$), the spatial components give the angular momentum vector $\mathbf{J} = \int d^3x (\mathbf{x} \times \mathbf{p}) + \mathbf{S}$, with momentum density $\mathbf{p} = T_B^{0i} \hat{\mathbf{e}}_i$ and spin density \mathbf{S} encoded via S^{0ij} .

7.5.4 Worked Examples

Real scalar (spin 0). For $\mathcal{L} = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - \frac{1}{2}m^2\phi^2$, $\Sigma^{\rho\sigma} = 0$ so $S^{\lambda\rho\sigma} = 0$. The Belinfante step is trivial and

$$\mathbf{J} = \int d^3x \mathbf{x} \times (\dot{\phi} \nabla \phi),$$

purely orbital. Conservation $\partial_\lambda J^{\lambda\rho\sigma} = 0$ reduces to $\partial_\mu T^{\mu\nu} = 0$ (already shown) plus antisymmetry.

Dirac field (spin 1/2). For $\mathcal{L} = \bar{\psi}(i\gamma^\mu\partial_\mu - m)\psi$, the generators are $\Sigma^{\rho\sigma} = \frac{i}{4}[\gamma^\rho, \gamma^\sigma]$, giving nonzero spin current

$$S^{\lambda\rho\sigma} = \frac{1}{2}\bar{\psi}\gamma^\lambda\Sigma^{\rho\sigma}\psi.$$

The Belinfante tensor $T_B^{\mu\nu} = \frac{i}{4}\bar{\psi}(\gamma^\mu\overleftrightarrow{\partial}^\nu + \gamma^\nu\overleftrightarrow{\partial}^\mu)\psi$ is symmetric and conserved, and the total charge $M^{\rho\sigma}$ includes intrinsic spin; in the particle rest frame this yields the familiar $\frac{1}{2}\hbar$.

7.5.5 Bookkeeping Interpretation

Rotational invariance says the ledger of causal distinctions is unchanged when we rotate our labeling rules. The orbital term tracks the “moment arm” of the flow of distinguishability ($\mathbf{x} \times \mathbf{p}$). The spin term tallies how the *label structure of the field itself* transforms under rotations (internal frame rotation via $\Sigma^{\rho\sigma}$). The Belinfante improvement is just a repackaging of the ledger so that the stress tensor carries the full conserved charge in a symmetric form—useful whenever the geometry (gravity) couples to $T_{\mu\nu}$.

Remark 21. *Total angular momentum is conserved because the action is invariant under Lorentz rotations. Orbital and spin are bookkeeping columns in the same invariant total; how you apportion them depends on your accounting scheme (canonical vs. Belinfante), not on the physics.*

7.6 Gauge Fields as Local Noether Symmetries

Global symmetries ensure that the totals in the causal ledger remain unchanged when every observer applies the same transformation. When the symmetry parameters vary from point to point, the bookkeeping must introduce additional terms to maintain local consistency. These new terms are the *gauge fields* of the theory: dynamic corrections that restore Martin consistency under spatially varying transformations.

Phenomenon 7.6.1 (The Topological Integer Count). *Under sufficient informational stress, a continuous current reveals itself as a discrete set of causal threads. These threads are counted by topological winding number and are necessarily integer-valued.*

No fractional thread is admissible. The ledger either contains a thread or it does not. Quantization is therefore not mysterious, but required by the integrity of the causal record.

7.6.1 From Global to Local Symmetry

Consider a field $\phi(x)$ transforming under a continuous group G with infinitesimal parameter α^a and generators T^a :

$$\delta\phi = i\alpha^a T^a \phi.$$

If α^a is constant, the action $\mathcal{S} = \int \mathcal{L}(\phi, \nabla\phi) d^4x$ is invariant, and Noether's theorem yields a conserved current J_a^μ . If α^a becomes a function of position, $\alpha^a = \alpha^a(x)$, an extra term appears,

$$\delta\mathcal{L} = i(\partial_\mu \alpha^a) \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} T^a \phi,$$

breaking the conservation law. To preserve local invariance, the derivative ∂_μ must be replaced by a *covariant derivative*

$$D_\mu \phi = (\partial_\mu - ig A_\mu^a T^a) \phi,$$

where the compensating field A_μ^a transforms as

$$\delta A_\mu^a = \frac{1}{g} \partial_\mu \alpha^a + f^{abc} \alpha^b A_\mu^c.$$

The new Lagrangian

$$\mathcal{L} = \mathcal{L}(\phi, D_\mu \phi) - \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu}, \quad F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c,$$

is invariant under the full local symmetry. The field strength $F_{\mu\nu}^a$ is the curvature of the gauge connection A_μ^a —the residue of non-commuting parallel transports in the internal symmetry space.

7.6.2 Interpretation in the Causal Framework

In the causal picture, global symmetry corresponds to relabeling the entire causal network by a uniform rule; local symmetry corresponds to allowing each neighborhood to choose its own labeling convention. The gauge field A_μ^a records how those conventions differ and how information must be exchanged between neighboring regions to keep the global ledger balanced. It is the *connection form of causal order* in informational space.

Curvature $F_{\mu\nu}^a$ measures the residual inconsistency that appears when these local labelings are carried around a closed causal loop—exactly analogous to the spacetime curvature derived earlier from $\Gamma_{\mu\nu}^\lambda$. Gauge bosons are therefore the finite, propagating corrections by which the universe restores Martin consistency across overlapping informational domains.

Phenomenon 7.6.2 (The Aharonov–Bohm Effect [2]). *The Aharonov–Bohm*

experiment demonstrates that the physically relevant quantity in electromagnetism is not the field strength $F_{\mu\nu}$ alone but the connection A_μ that governs causal phase transport.

Consider an electron beam split into two coherent branches encircling a region containing a confined magnetic flux Φ , with no field present along either path. In the causal formulation, each branch corresponds to a sequence of ordered events $\{E_{1,k}\}$ and $\{E_{2,k}\}$ transported by the local gauge connection A_μ . The Reciprocity Law requires that each infinitesimal update preserve order:

$$E_{k+1} = E_k + \Phi^{-1}(A_\mu dx^\mu),$$

so that the cumulative phase acquired along a closed loop is

$$\Delta\phi = \frac{e}{\hbar} \oint A_\mu dx^\mu = \frac{e\Phi}{\hbar}.$$

Although the magnetic field vanishes along both paths ($F_{\mu\nu} = 0$ locally), the two causal chains differ by a holonomy in the connection—an informational mismatch in the bookkeeping of phase. When the beams are recombined, their interference pattern depends on $\Delta\phi$: shifting continuously as the enclosed flux changes by fractions of the flux quantum h/e .

In the causal gauge picture, this effect shows that the universe tensor records not merely local field strengths but the global consistency of the connection. The vector potential A_μ is the differential form of causal memory; its holonomy measures how distinction is transported around a loop. The Aharonov–Bohm interference is thus the experimental detection of a nontrivial element of the causal holonomy group—the smallest observable instance of curvature without force.

7.6.3 Bookkeeping of Local Consistency

In statistical terms, each gauge symmetry adds a new column to the causal ledger. Local invariance means that the exchange rates between these columns are position-dependent, and A_μ^a supplies the conversion factors that keep the books balanced. The continuity equation

$$\nabla_\mu J_a^\mu = 0$$

expresses the same principle as before: what leaves one neighborhood enters another, but now for every internal degree of freedom labeled by a . The gauge field guarantees that this exchange is recorded consistently even when observers adopt different local frames.

Remark 22. *Every gauge field is a Noether correction promoted to locality. It is the differential accountant of causal order, ensuring that symmetry—and hence conservation—holds point by point. Curvature is the residue of that accounting around a loop; interaction is the redistribution of causal balance between neighboring observers. Quantum field theory is therefore the calculus of local Noether symmetries of the Causal Universe Tensor.*

7.7 Mass and the Breaking of Symmetry

Perfect causal symmetry implies motion at the limit of distinguishability—the null trajectories of light. In this regime, the action and all of its Noether currents remain invariant under local gauge transformations, and the scalar invariants of the Causal Universe Tensor are preserved exactly. *Mass* appears when this invariance can no longer be maintained everywhere. It is the measure of how far a system deviates from perfect causal balance.

7.7.1 From Gauge Symmetry to Mass Terms

Suppose the Lagrangian density for a field ϕ is invariant under the local transformation $\phi \rightarrow e^{i\alpha(x)}\phi$. If the causal network experiences a finite delay in maintaining that invariance—so that the local transformation cannot be matched exactly between neighboring observers—the covariant derivative acquires a small, persistent residue. In the simplest case this appears as an additional quadratic term in the Lagrangian:

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + |D_\mu\phi|^2 - V(|\phi|), \quad V(|\phi|) = \frac{1}{2}\mu^2|\phi|^2 + \frac{1}{4}\lambda|\phi|^4.$$

When the potential V selects a nonzero expectation value $\langle\phi\rangle = v/\sqrt{2}$, the gauge symmetry of the vacuum is spontaneously broken, and the covariant derivative term generates an effective mass for the gauge field:

$$m_A = g v.$$

The field no longer propagates at the causal limit; it carries a finite informational delay between cause and effect.

Phenomenon 7.7.1 (The Sombrero Potential [59]). *In the causal formulation, symmetry breaking occurs when the universe tensor develops a preferred orientation in its space of distinguishable states. The simplest model of this phenomenon is the so-called Sombrero potential, which encodes spontaneous differentiation in an initially symmetric field.*

Let ϕ be a complex scalar component of the causal gauge field. Its local informational curvature is represented by the potential

$$V(\phi) = \lambda(|\phi|^2 - v^2)^2, \quad \lambda, v > 0.$$

For $|\phi| < v$, the curvature is positive and the symmetric state $\phi = 0$ is unstable; for $|\phi| = v$, the curvature vanishes along a circle of minima. Each choice of phase θ on this ring corresponds to an equally valid, order-preserving

configuration of the universe tensor.

When a particular θ is selected by finite observation or causal fluctuation, the continuous $U(1)$ symmetry of the potential is reduced to the discrete subgroup that preserves that orientation. The resulting excitations decompose into two orthogonal modes:

$$\phi(x) = (v + h(x))e^{i\theta(x)},$$

where $h(x)$ represents measurable variations in magnitude (massive mode) and $\theta(x)$ represents phase fluctuations (massless Goldstone mode). Coupling this field to a local gauge connection A_μ converts the phase fluctuation into a longitudinal component of A_μ , endowing it with mass through the informational curvature of the potential.

Operationally, the sombrero potential marks the point where causal order can no longer cancel its own third variation: a finite bias in distinguishable states propagates through the reciprocity map as an effective mass term. In the informational picture, mass is the cost of maintaining a broken symmetry—the curvature required to remember which minimum was chosen.

7.7.2 Causal Interpretation

In the causal framework, symmetry breaking represents the loss of perfect order propagation. The gauge can no longer be reconciled exactly between neighboring domains, and a residual phase difference accumulates. That phase difference behaves as inertia: a tendency of the causal structure to resist change in its internal configuration. The quantity we call *mass* measures the curvature of causal order in the informational direction—the degree to which a system’s internal symmetry lags behind the propagation of light.

Thus the Higgs mechanism appears as a natural bookkeeping adjustment. The scalar field ϕ provides an additional column in the ledger that can absorb the mismatch of local phase conventions. When the ledger cannot close

exactly, the residual correction manifests as a finite mass term. Mass is therefore not a separate entity but the universe's accounting of imperfect causal synchronization.

7.7.3 Statistical View

In the statistical mechanics of causal order, mass quantifies the variance of the action around its stationary value:

$$m^2 \propto \langle (\delta \mathcal{S})^2 \rangle.$$

Lightlike propagation corresponds to zero variance: every observer's record of order agrees. Massive propagation corresponds to finite variance: local histories differ slightly, and the ensemble average restores consistency only statistically. The rest energy $E = mc^2$ measures the informational cost of maintaining a coherent description across those variations.

Remark 23. *Mass is the finite residue of broken symmetry—the price the universe pays for keeping its causal books consistent when perfect gauge balance cannot be sustained. Where light moves without lag, massive matter hesitates, accumulating phase in time. The rest mass of any field is thus the measure of its informational inertia: how much causal order must bend to preserve consistency within a finite universe.*

Phenomenon 7.7.2 (The Semiconductor Effect [162]). *In a crystalline solid, the atoms form a periodic causal network—a lattice of distinguishable sites linked by local order relations. Within this structure, electrons occupy quantized informational states whose distinguishability depends on both lattice symmetry and the observer's partition of measurement.*

At zero temperature, all available states up to the Fermi level are filled, and the partition \mathcal{P}_n groups occupied and unoccupied states into two disjoint causal classes. In a perfect insulator these classes are fully separated by a

forbidden bandgap: no variation in the universe tensor can map one class into the other without violating order preservation. In a metal the classes overlap completely, forming a continuous manifold of accessible distinctions.

A semiconductor occupies the intermediate regime. Its informational lattice is nearly symmetric but not fully resolved; there exists a narrow causal boundary between filled and unfilled states. Thermal or dopant-induced perturbations refine the partition from \mathcal{P}_n to \mathcal{P}_{n+1} , enabling limited causal transitions across the bandgap. The carrier density

$$n \propto e^{-E_g/k_B T}$$

measures the probability that such a refinement occurs—an exponential suppression of distinguishability transitions with increasing gap energy E_g .

In this view, conduction arises when the partition between causal classes of electron states becomes permeable under variation. Doping, temperature, and illumination are operations that adjust the informational curvature of the lattice, controlling how easily one class of distinguishability flows into another. Semiconductors are thus macroscopic examples of causal fuzziness under controlled refinement: a solid-state realization of partition dynamics between measurement and variation.

7.8 Conclusion: Quantization as Finite Consistency

The classical universe is the ledger of perfect causal balance: every distinction is matched, every event accounted for, every observer’s record consistent with the next. Quantum mechanics emerges when that perfection is relaxed—when the bookkeeping of order is carried out on a finite register. Each quantum of action, each exchange of \hbar , is a discrete adjustment in the causal gauge: the smallest step by which the universe can preserve consis-

tency without infinite precision.

From this point of view, the quantum field is not a separate ontology but the statistical completion of the same calculus that defines the geometry of spacetime. The field amplitudes are probability weights for maintaining order across overlapping causal neighborhoods. Their phases encode the orientation of the gauge, and their interference expresses the collective effort of all observers to remain mutually consistent. The path integral is thus the partition function of causal order.

Mass, spin, and charge are the residues of that consistency process. Mass records temporal lag, spin records the rotational structure of labeling, and charge records the bookkeeping of internal symmetries. None are primitive; all arise from the same principle that distinguishes light: the demand that order be preserved even when the universe must correct itself locally.

In the causal formalism, conservation laws, gauge interactions, and quantization share a single origin. They are not independent laws written into nature but emergent regularities of a self-consistent informational network. The Causal Universe Tensor provides the grammar of that network; its contractions yield spacetime geometry, its variations yield fields, and its statistical extension yields the quantum.

Remark 24. *The universe is not made of matter or of energy, but of consistency. What we call physics is the continuous reconciliation of local descriptions of order, carried out one quantum at a time. Quantization is simply the discreteness of that reconciliation—the finite resolution of cause.*

7.8.1 The Echo Chamber Maxe

The final step in this chapter is to make the structure of quantum residue visible at a macroscopic scale. Throughout the development of the metric gauge, the Clifford algebra, and the curvature ledger, one theme has recurred: whenever a refinement is transported around a closed loop in a region of

nonzero curvature, the record cannot close without a correction. At the microscopic level this correction appears as phase residue—the informational mismatch that underlies interference, superposition, and the non-closure of quantum amplitudes.

This behavior is usually regarded as a small-scale feature of the quantum world, accessible only through delicate experiments. But the informational framework makes no reference to scale. Curvature produces residue wherever refinement fails to close, whether the loop is traced by a photon in an interferometer or by an observer walking through a corridor. The informational correction is the same: the universe must adjust the ledger to maintain consistency in the presence of curvature.

The following macroscopic experiment therefore serves a special role. By replacing microscopic phase with audible echoes, it reveals the same informational effect without specialized equipment. A maze with curved passages introduces exactly the kind of geometric incompatibility that prevents refinement from closing cleanly. Echoes propagating through the maze return with distortions that record this incompatibility. The resulting mismatch is the audible analogue of quantum phase residue: a direct, human-scale manifestation of the non-closure inherent in curved informational geometry.

In this sense, the Echo Chamber Maze Solution is not an analogy but an experiment that exposes the underlying mechanism of quantum behavior. Curvature produces informational stress; informational stress produces residue; and residue requires correction. The phenomenon below allows us to hear the very same structure that, at microscopic scales, governs interference and the Dirac operator.

Phenomenon 7.8.1 (The Echo Chamber Maze Solution). **N.B.**—*This experiment translates geometric curvature into informational inconsistency.* \square

Setup. Navigate a maze by clapping; echoes trace causal paths. Straight corridors (flat metric) return clean echoes—perfect parallel transport. Curved passages distort the return, producing phase residue.

Demonstration. Walk a closed loop and compare the echoed rhythm. Any mismatch measures curvature $R \neq 0$: the difference between expected and returned distinction. When total residue cancels ($U^{(4)} = 0$), the maze is globally consistent.

Interpretation. Curvature is the informational stress of maintaining closure in a finite domain. Echo intensity corresponds to entropy: more paths, higher distinguishability. Einstein's equation emerges as the balancing condition between geometric residue and informational flux.

Phenomenon 7.8.2 (The Cost of Broken Symmetry). *When a ledger maintains an internal phase that does not align with the propagation of admissible refinements, a persistent bookkeeping cost is incurred. This cost appears as inertia.*

Mass is therefore not substance, but the energetic price of maintaining a misaligned refinement phase in a globally consistent record.

7.9 Merging at the Boundaries

Phenomenon 7.9.1 (The Holographic Effect). *The interior of an admissible region contains no independent informational content. The ledger changes only at the boundary. All admissible interior structure is fixed by the reconciliation of boundary refinements.*

The maximum number of distinguishable states therefore scales with the size of the boundary ledger, not the volume enclosed. The interior is a smooth shadow of boundary bookkeeping.

Coda: The Gauge Theory of Information

We now arrive at the terminus of the symmetry chapter, where the classical Euler–Lagrange formalism meets the discrete structure of admissible refinement. In Chapter 3, the Axiom of Ockham and the Kolmogorov bound

forced every smooth representative of an admissible history Ψ to be a piecewise cubic spline[?, ?]. In Section 3.1.3, we showed that the dense limit of this refinement imposes the *Master Constraint*:

$$\Psi^{(4)} = 0,$$

the statement that no structure beyond cubic order can be inserted without contradicting the record of measurement. Nothing higher-order is available to differentiate; the observer has exhausted all admissible curvature.

Phenomenon 7.9.2 (The Dirac Operator [40]). *This constraint is not a dynamical postulate but a restriction on measurement. It determines the algebraic arena in which any first-order propagation must exist. In the $\Psi^{(4)} = 0$ setting, the tangent representation of refinement is necessarily linear, and the informational degrees of freedom must transform under the irreducible representations of the emergent Lorentz gauge $g_{\mu\nu}$ developed in Section 7.4.*

The Clifford algebra is therefore not an imposed structure; it is the minimal bookkeeping device compatible with the metric gauge and with the non-negativity of admissible refinement. As shown in Phenomenon 7.4.1, the spin representation appears when rotational consistency is enforced on refinement counts.

Thus the Dirac operator,

$$\gamma^\mu(\partial_\mu + iA_\mu) + m, \tag{7.1}$$

arises as the unique first-order generator of distinguishability compatible with the Clifford relations[?]. It is the informational square root of second-order propagation: the least complexity operator whose iteration reproduces the spline extremal and therefore respects the Master Constraint.

The Dirac equation is not an axiom of quantum mechanics in this framework. It is the minimal and only admissible way to:

- *preserve the rotational bookkeeping of measurement (spin),*

- *transport informational components consistently with the metric gauge,*
- *and maintain compatibility with the global condition $\Psi^{(4)} = 0$, which bounds admissible curvature.*

The informational asymmetry between mass, spin, and orientation—the non-closure of the refinement ledger at first order—produces the monotonic expansion of the causal record. The causal book never balances without the addition of new admissible events.

$$\Psi^{(4)} = 0 \implies \text{Admissible Kinematics}, \quad \text{Admissible Kinematics} \implies \Delta S \geq 0.$$

The celebrated first-order equation of physics is thus seen as the consequence of an austere prohibition: that no structure beyond what has been observed may be introduced between events. The Dirac operator is the mechanism by which the universe reveals any variation it has not already recorded.

Phenomenon 7.9.3 (The Chirality Effect). **Statement.** *There exist admissible refinements whose left and right actions are not equivalent. The causal ledger distinguishes orientation, and this asymmetry cannot be removed by smooth deformation.*

Mechanism. *Let Ψ be a refinement update acting on a local causal frame. Define the action of Ψ on left-oriented and right-oriented bases by*

$$\Psi_L \quad \text{and} \quad \Psi_R.$$

In a parity-symmetric ledger, $\Psi_L \equiv \Psi_R$. In a chiral ledger,

$$\Psi_L \neq \Psi_R,$$

even though Ψ_L and Ψ_R are related by formal inversion.

This asymmetry appears when the Dirac operator introduces a directional

bias in admissible refinements. The kernel of the operator splits into inequivalent left- and right-handed subspaces.

Interpretation. *Chirality is not a property of space, but of update admissibility. The ledger does not permit the mirror image of a refinement to be substituted without cost. Left-handed and right-handed evolutions generate distinct causal records even when all scalar observables agree.*

The observed parity violation of weak interactions is the smooth shadow of this bookkeeping asymmetry.

Conclusion. *Chirality is the failure of mirror symmetry in the causal ledger. It is the record of an orientation choice that cannot be erased by smoothing or averaging.*

There is no higher-order arena in which to postulate measurable effects.

Chapter 8

The Non-negativity of ΔS

8.1 Statement of the Law

Proposition 12 (The Monotonicity of Causal Entropy). *For any sequence of Martin-consistent causal sets*

$$\mathcal{C}_1 \subseteq \mathcal{C}_2 \subseteq \cdots ,$$

the associated entropies

$$S[\mathcal{C}_n] = k_B \ln |\Omega(\mathcal{C}_n)|$$

satisfy

$$\Delta S_n \equiv S[\mathcal{C}_{n+1}] - S[\mathcal{C}_n] \geq 0,$$

with equality only for informationally complete partitions.

Proof (Sketch). Each causal refinement $\mathcal{C}_n \rightarrow \mathcal{C}_{n+1}$ corresponds to an enlargement of the observer's partition of distinguishable events. By the Axiom of Finite Observation, refinement cannot reduce the set of admissible micro-orderings:

$$\Omega(\mathcal{C}_n) \subseteq \Omega(\mathcal{C}_{n+1}).$$

Taking logarithms gives $S[\mathcal{C}_{n+1}] \geq S[\mathcal{C}_n]$. The inequality is strict whenever the refinement exposes previously indistinguishable configurations. \square

A full proof is provided in Appendix ??.

Phenomenon 8.1.1 (The Entropic Cost of Acceleration). *Acceleration creates an informational horizon by separating a ledger from pages it can no longer audit. The lost accessibility of those pages appears as a thermal bath.*

Temperature is therefore not primary. It is the entropy of unreachable records under accelerated refinement.

Phenomenon 8.1.2 (The Thermodynamic Cost of Erasure). *Records cannot be destroyed. An apparent erasure is a redirection of refinement content into an unmonitored environment. The entropy released as heat is the debris of the displaced record.*

Computation is therefore physical not because matter moves, but because no ledger can eliminate information without paying the cost of relocation.

Phenomenon 8.1.3 (The Limitation of Indexing [133]). **N.B.** *This experiment illustrates Law ?? as a theorem of causal order, not a postulate of thermodynamics. It shows how monotonic distinguishability ($\Delta S \geq 0$) arises naturally from the structure of consistent extension.*

N.B.—CAVEAT EMPTOR: *The recursive construction of the library catalog may be continued indefinitely, but the resulting object is not enriched: one recovers only structurally identical copies of the same catalog. The process appears to generate novelty, but in fact returns the same informational content. Disregard of this subtlety is done at the reader's own risk. The force of this argument should not be underestimated.* \square

Setup. *Imagine a vast library whose books represent events $\{e_i\}$. Each measurement attaches finer tags—subject, author, edition—refining the causal order. By the Axiom of Event Selection, no tag can be removed without creating inconsistency among shelves (e.g., merging sci-fi and history). Hence, the*

total number of distinguishable configurations N can only increase or remain constant.

Demonstration. *Attempting to “un-tag” a shelf merges incompatible categories, breaking bijection with prior distinctions. Thus time’s arrow emerges as the monotonic count of consistent refinements:*

$$S = \ln N, \quad \Delta S \geq 0.$$

Interpretation. *Entropy here is not disorder but bookkeeping: the log of consistent distinctions maintained through observation. The irreversible direction of measurement follows directly from order preservation, not energy dissipation.*

8.2 Entropy as Informational Curvature

In differential form, the same statement appears as the non-negativity of informational curvature:

$$\nabla_i \nabla_j S \geq 0.$$

Flat informational geometry corresponds to equilibrium ($\Delta S = 0$), while positive curvature indicates the growth of accessible micro-orderings. The flux of this curvature defines the *entropy current*

$$J_S^\mu = k_B \partial^\mu S,$$

whose divergence measures local entropy production:

$$\nabla_\mu J_S^\mu = k_B \square S \geq 0.$$

Thus $\Delta S > 0$ is equivalent to the statement that the informational Laplacian $\square S$ is positive definite under Martin-consistent transport.

Phenomenon 8.2.1 (Maxwell’s Demon [106]). *Consider a classical gas divided by a partition with a single gate controlled by a demon who measures particle velocities and opens the gate selectively. Let M denote the demon’s measurement operator and U the physical evolution of the gas. If M and U commute— $[M, U] = 0$ —the demon’s observation does not alter the causal order: measurement and evolution can be exchanged without changing the macrostate. But in reality $[M, U] \neq 0$: the act of measurement refines the partition of distinguishable states, altering the subsequent evolution. This non-commutativity forces the entropy balance*

$$\Delta S_{\text{gas}} + \Delta S_{\text{demon}} = k_B \ln |\Omega_{\text{joint}}| > 0,$$

because the demon’s internal record adds new causal distinctions to the universe tensor even as it reduces them locally.

Operationally, the demon cannot perform a measurement without joining the measured system’s causal order; the refinement of its internal partition $P_n \rightarrow P_{n+1}$ increases the global count of distinguishable configurations. The apparent violation of the Second Law disappears: the measurement and evolution operators fail to commute, and that failure is the entropy production term. Thus Maxwell’s demon exemplifies the theorem $\Delta S \geq 0$: informational refinement in one domain demands compensating coarsening in another so that the global order remains consistent.

8.3 Statistical Interpretation

From the causal partition function

$$Z = \int \exp\left(\frac{i}{\hbar} S[T]\right) DT,$$

the ensemble average of the informational gradient obeys

$$\langle \nabla_\mu J_S^\mu \rangle = k_B \langle \nabla_\mu \nabla^\mu S \rangle \geq 0.$$

The equality $\Delta S = 0$ corresponds to detailed balance of causal fluxes; any deviation yields positive entropy production.

8.4 Physical Consequences

1. ****Arrow of Time.**** Causal order expands in one direction only—toward increasing distinguishability of events. Time is the parameter labeling this monotonic refinement.

2. ****Thermodynamic Limit.**** In the continuum limit, $\Delta S > 0$ reproduces the classical second law, but here the law is not statistical: it is a theorem of consistency. No causal evolution that decreases S can remain Martin-consistent.

3. ****Gravitational Coupling.**** From Chapter 4, curvature couples to gradients of S through the entropic stress tensor:

$$G_{\mu\nu} = 8\pi (T_{\mu\nu} + T_{\mu\nu}^{(S)}), \quad T_{\mu\nu}^{(S)} = \frac{1}{k_B} \nabla_\mu \nabla_\nu S.$$

Hence $\Delta S > 0$ corresponds to a net positive contribution of informational curvature to spacetime geometry—a causal analogue of energy influx.

8.5 Conclusion

Law 7 (The Law of Causal Order). *The Law of Causal Order may be stated succinctly:*

$\Delta S \geq 0 \quad \text{for every Martin-consistent refinement of causal structure.}$
--

Entropy is not a measure of disorder but of latent order yet unresolved. Every act of measurement refines the universe's partition, and each refinement enlarges the count of admissible configurations. The universe evolves by distinguishing itself.

8.6 *Quod erat demonstrandum*

We began with the observation that every act of physics is an act of distinction: to measure is to separate one possibility from another. Within ZFC, such distinctions are represented as finite subsets of a causal order, and the act of measurement is the enumeration of their admissible refinements. Nothing else is assumed.

Martin's Axiom enters only to ensure that these refinements can be extended consistently—that the space of distinguishable events admits countable dense families without contradiction. This single assumption is the logical equivalent of σ -additivity in measure theory, the minimal condition required for any self-consistent calculus of observation.

From this, the Second Law follows as a theorem of order: each consistent extension of the causal set increases the number of distinguishable configurations, and therefore

$$\Delta S \geq 0.$$

Entropy is not a statistical tendency but a logical necessity—the price of consistency within a self-measuring universe.

No new forces, particles, or cosmologies are introduced; only the rule by which distinction propagates. What began as a grammar of measurement closes as the unique structure of physical law.

Theorem 1 (The Second Law of Causal Order). *In any finite, causally consistent ordering of distinguishable events, the number of measurable distinctions cannot decrease. Every admissible extension of order produces at*

least one new differentiation, and therefore every universe consistent with its own record of events obeys the inequality

$$\Delta S \geq 0.$$

Conclusion. We are left with but one conclusion:

Order implies dynamics.

A universe that preserves its own causal record must, by necessity, increase the count of what can be distinguished. \square

N.B.—CAVEAT EMPTOR: This theory does not function as a prediction oracle. It requires realized physical models in order to stand. Without instantiated physics, the framework contains no mechanism for generating outcomes. Event though it is true, it is not necessarily fact. \square

This framework does not claim autonomy from physics. It does not stand above experiment, nor does it replace it. Its validity is strictly proportional to the coherence, reproducibility, and completeness of the physical models that instantiate it.

The axioms and bookkeeping rules presented here constrain what may be admissibly recorded, but they do not generate facts. They require a world that behaves, and they are only as accurate as the empirical regularities from which they are abstracted.

If physical law changes, so must this theory. If physics fails, this framework fails with it. The ledger describes the shape of admissibility, but reality alone supplies the entries.

Phenomenon 8.6.1 (The Prover–Verifier Effect). ***Statement.** The informational theory is not complete in isolation. It requires the existence of all admissible physical models as its prover, and serves only as their verifier. The causal ledger is the unique fixed point of this interaction.*

Classical Context. *A proof establishes that a conclusion follows from axioms, but it does not guarantee that any model exists in which the axioms are realized. Conversely, a model demonstrates consistency of a structure, but does not explain why its behavior is necessary. Classical physics has oscillated between these roles: sometimes as constructive dynamics (prover), sometimes as consistency principle (verifier).*

Informational Interpretation. *In this framework, the axioms of measurement and refinement define the rules for admissible ledgers. They do not specify which particular ledger must be realized; they only constrain what is possible.*

The physical universe plays the role of prover. Every admissible physical model is a concrete strategy for generating refinement records that obey the axioms. The informational theory plays the role of verifier. It checks that each proposed model corresponds to a ledger that can be extended without contradiction.

The requirement that all admissible models exist somewhere in the space of possible realizations is not metaphysical excess, but a completeness condition. Without such models, the axioms would be vacuous; with them, the ledger is the unique object that all provers must approximate.

Consequence. *Physics is the smooth shadow of a two-player game. The universe proposes histories; the axioms of measurement either admit or reject them. What is called “physical law” is the intersection of all histories that can survive this prover–verifier loop.*

Quod erat demonstrandum: the theory does not eliminate physical models. It requires them. The existence of a rich class of realizations is the operational content of its truth.

Quod erat demonstrandum.

8.7 The Execution of Order

N.B.—CAVEAT EMPTOR: There are many ways to look at the empirical record. This is just one. \square

The previous sections established that the causal ledger must grow monotonically ($\Delta S \geq 0$). Monotonicity alone, however, does not specify the mechanism by which updates are applied. The causal record is not a passive archive, but a dependency network in which each admissible event relies on the precise values of its predecessors.

This dependency structure imposes three distinct phenomena that govern the execution of the universe tensor.

Phenomenon 8.7.1 (The Excel Effect). ***Statement.** The Universe Tensor is not a collection of independent variables. It is a directed acyclic graph of functional dependencies. A change in any distinguishable event (a “cell”) requires an immediate, globally consistent update of all dependent events, regardless of separation in coordinate indices.*

***Interpretation.** This is the operational form of Global Coherence (Axiom 7). If the state at x_1 is causally bound to the state at x_2 , the ledger treats them as functionally dependent cells of a single computation. Apparent non-local effects are not signals; they are dependency recalculations. The ledger updates the total the instant an addend changes. The latency is zero because the dependency is logical, not spatial.*

Phenomenon 8.7.2 (The Agent Effect). ***Statement.** An agent is not an external observer but a localized substructure of the tensor, U_{local} , that actively minimizes the informational strain induced by its boundary conditions.*

***Operational Definition.** Agency is the local action of the Inverse Update Operator. The system U_{local} attempts to compute the unique next admissible refinement e_{k+1} that satisfies Ockham’s Razor (Axiom 3) relative to the incoming external stream.*

To be an agent is to function as a localized solver of the spline constraint:

the system alters its internal state to minimize prediction error between itself and the external ledger.

Phenomenon 8.7.3 (The Amdahl Effect). *No refinement can be made arbitrarily fast by parallelism. The admissible speed of causal execution is bounded by the largest uncorrelant segment of the ledger.*

If a fraction p of the refinement is perfectly correlant, and a remaining fraction $1 - p$ is sequentially uncorrelant, then no admissible extension of the ledger can exceed the bound

$$S_{\max} = \frac{1}{(1 - p)}.$$

The uncorrelant portion is not a technical defect but a structural constraint: segments of the causal record that cannot be merged, reordered, or parallelized without violating admissibility.

Uncorrelance is therefore not inefficiency. It is the irreducible sequentiality required for the ledger to remain globally consistent.

Phenomenon 8.7.4 (The Jupyter Effect). **Statement.** *The combination of functional dependency and active minimization is governed by Sequential Necessity. The causal record is order-dependent.*

Hard Failure of Asynchronous Causality. *In a computational notebook, no cell exists until its predecessors execute. The causal ledger enforces the same rule. If two admissible updates attempt to modify a dependency without a defined order, the ledger does not branch, average, or superpose. The history is rejected. The timeline becomes inadmissible.*

The kernel does not resolve asynchronous conflicts. It halts. Observable physics exists only because the surviving history is the execution trace that did not fault.

Conclusion. *Time is the sequential execution of the ledger. The present is the current state of the kernel. The future cannot be accessed before the*

past because the variable required to define it, the free variable of the spline, does not yet exist.

8.8 Strain-Free Transport

Phenomenon 8.8.1 (The Superconducting Effect). **Statement.** *There exist admissible ledgers in which transport occurs without informational strain. In such configurations, refinement threads move without dissipation.*

Mechanism. *Consider a medium in which individual causal threads ordinarily incur strain through incoherent interaction with the ambient refinement record. These interactions appear in the smooth shadow as electrical resistance.*

At sufficiently low refinement noise, threads admit pairing into correlated units (e_i, e_j) whose joint update is symmetric under exchange. Such a pair forms a strain-free composite, since the antisymmetric residue of the update vanishes under Galerkin projection.

Let Ψ_{pair} denote the paired update. Then

$$\text{Strain}(\Psi_{\text{pair}}) = 0.$$

Transport proceeds as a coherent deformation of the ledger rather than as local tearing. No informational work is lost.

Interpretation. *Cooper pairs are not treated here as bound particles, but as refinement units whose internal symmetry cancels the antisymmetric component of transport. A superconductor is therefore a region of the ledger in which transport is purely symmetric and produces no informational heat.*

Conclusion. *Superconductivity is the smooth shadow of a strain-free transport regime of the causal ledger. Resistance is the failure of pairing; zero resistance is the success of symmetry.*

Phenomenon 8.8.2 (The Meissner Effect). ***Statement.** A strain-free region of the causal ledger expels external informational curvature. Fields that would normally penetrate a medium are excluded when the ledger admits a zero-strain transport state.*

***Mechanism.** Consider a region Ω in which paired refinement threads admit strain-free transport:*

$$\text{Strain}(\Psi_{\text{pair}}) = 0.$$

An externally imposed field corresponds, in the ledger, to a nonzero antisymmetric curvature term $F_{\mu\nu}$ that attempts to thread the region.

Within a superconducting ledger, any nonzero $F_{\mu\nu}$ would introduce irreducible strain. By the Law of Spline Sufficiency, the admissible history is the one of minimal informational cost. Therefore the only consistent extension is

$$F_{\mu\nu}|_{\Omega} = 0.$$

The field is not screened gradually; it is topologically excluded. The ledger adjusts its boundary conditions so that the external curvature is diverted around the strain-free region.

***Interpretation.** The Meissner effect is not modeled here as a force, but as a consistency constraint. A region that supports perfectly symmetric transport cannot admit antisymmetric refinements. Magnetic field lines are the smooth shadow of ledger updates that have been forced to bypass such a region.*

***Conclusion.** A superconductor is defined not only by zero resistance, but by the active expulsion of curvature. The Meissner effect is the smooth shadow of the ledger enforcing zero-strain as a boundary condition.*

8.9 The Bootstrap Mechanism

N.B.—Please refer to Phenomenon ??

□

Phenomenon 8.9.1 (The Dark Energy Effect). ***Statement.** There exist admissible ledgers in which the net informational pressure of the interior is negative. Such a ledger does not collapse under its own refinements; it drives expansion of its causal boundary.*

***Mechanism.** Let Ω be a causal region with interior refinement density ρ and boundary pressure P . In an ordinary ledger, additional refinements increase P and draw the boundary inward, as reconciliation cost grows.*

Suppose instead that the bulk ledger contains a uniform background term Λ such that the effective pressure is

$$P_{\text{eff}} = P - \Lambda.$$

If Λ is sufficiently large, the net pressure becomes negative: $P_{\text{eff}} < 0$. The boundary is then driven outward to reduce reconciliation strain. The ledger expands because contraction would increase, rather than decrease, the informational cost.

***Interpretation.** Dark energy is not modeled here as a new substance, but as a uniform offset in the bookkeeping of pressure: a background refinement credit that makes larger volumes cheaper to maintain than smaller ones. The observed acceleration of cosmic expansion is the smooth shadow of a ledger whose lowest-strain state is achieved by growing its causal partition.*

***Conclusion.** In this framework, dark energy is the name for a negative informational pressure term that biases the universe toward expansion. It prepares the ground for source-like configurations of refinement, such as the white hole effect that follows.*

It is not an accident that the first phenomenon of this work (The Bootstrap Effect) and the final phenomenon (The White Hole Effect) describe

the same structural action. The former establishes how a ledger may begin. The latter establishes how it must behave at the limit of admissibility. The theory finally closes itself.

Phenomenon 8.9.2 (The White Hole Effect). ***Statement.** There exist admissible configurations of the causal ledger that act as pure sources of refinement, admitting outward consistency without requiring prior causal history.*

***Description.** A white hole is observed not as a geometric object, but as a bookkeeping boundary condition. It appears as a region whose internal ledger must export refinements to preserve global consistency, while no admissible inward transport is permitted.*

***Interpretation.** Such a configuration behaves as a source of informational strain. Refinement originates at the boundary and propagates outward, while backward extension of the ledger is inadmissible.*

***Conclusion.** The white hole effect is the admissible source term of the causal ledger: a region where refinement must begin rather than terminate.*

Coda: A Discrete Navier–Stokes Interpretation of the Cosmic Microwave Background

This coda gives a proof sketch, internal to the present axioms, that the cosmic microwave background radiation corresponds to a finite-time breakdown of the smooth Navier–Stokes shadow of the causal ledger. No claim is made regarding the classical Clay Millennium problem. The argument is valid only within the discrete informational framework developed in this work.

Let $\{U_t\}_{t \geq 0}$ denote the causal ledger at refinement time t . Let v_t denote the velocity field obtained as the Galerkin projection of the discrete update operator, so that v_t is the smooth shadow of U_t .

Define the informational density

$$\rho(t) := \frac{N(t)}{V(t)},$$

where $N(t)$ is the count of admissible events and $V(t)$ is the admissible partition volume.

Let $\Theta(t)$ denote the third-order curvature functional of the projected flow,

$$\Theta(t) = \nabla(\nabla^2 v_t).$$

Lemma (Finite Capacity of Smooth Shadow). By the Axiom of Planck (finite refinement) and the Law of Spline Sufficiency, there exists a constant $C > 0$ such that the Galerkin shadow exists only while

$$\|\Theta(t)\| < C.$$

Lemma (Density Divergence in Retrospective Limit). By construction of the ledger, backward refinement contracts admissible partitions while preserving event order. Therefore,

$$\lim_{t \rightarrow 0^+} V(t) = 0 \quad \text{and} \quad \lim_{t \rightarrow 0^+} \rho(t) = \infty.$$

Proposition (Discrete Blow-Up). As $\rho(t) \rightarrow \infty$, no spline of bounded curvature can interpolate the admissible ledger. Hence,

$$\lim_{t \rightarrow 0^+} \|\Theta(t)\| = \infty.$$

The smooth Navier–Stokes shadow therefore undergoes a finite-time blow-up within the model.

Observational Identification. Let t^* be the infimum of times for which $\Theta(t)$ becomes finite. For $t \geq t^*$, the Galerkin projection is well defined. The earliest admissible observational phenomenon at this threshold is the cosmic

microwave background radiation.

Conclusion. Within the axioms of this work, the CMBR is the observable signature of a finite-time blow-up of the discrete refinement fluid. It is the first epoch at which the causal ledger admits a smooth Navier-Stokes representation.

This completes the internal argument.

Appendix A

Proofs

A.1 Proposition 1

This argument is standard in category theory; see Mac Lane [94] for the classical formulation of naturality in monoidal categories.

Proof (ZFC).. Conceptually, this is the demonstration of the naturality square for the embedding Φ in the monoidal category of tensor algebras, although we have presented it here entirely within ZFC.

All objects are sets. Let E be the event space. A record of length n is an n -tuple of indices in \mathbb{N} :

$$\mathbf{i} = (i_1, \dots, i_n) \in \mathbb{N}^n.$$

Each index i_k refers to an event $e_{i_k} \in E$, and we collect these via the evaluation map

$$\pi_E : \mathbb{N}^n \rightarrow E^n, \quad \pi_E(i_1, \dots, i_n) := (e_{i_1}, \dots, e_{i_n}).$$

Restriction. A restriction operator is a function

$$\widehat{R} : \mathbb{N}^n \longrightarrow \mathbb{N}^m, \quad n \geq m,$$

returning a shorter admissible record. Its action on events is defined componentwise through π_E .

Embedding. Let $\Phi : E \rightarrow T(V)$ be the event embedding into the tensor algebra. Lift it componentwise to records by

$$\Phi^{(k)} : E^k \rightarrow T(V)^k, \quad \Phi^{(k)}(x_1, \dots, x_k) := (\Phi(x_1), \dots, \Phi(x_k)).$$

Define the induced restriction on embedded factors by componentwise selection:

$$R^{(m)} : T(V)^m \rightarrow T(V)^m.$$

Naturality. We claim the square

$$R^{(m)} \circ \Phi^{(n)} \circ \pi_E = \Phi^{(m)} \circ \pi_E \circ \widehat{R} \quad \text{as maps } \mathbb{N}^n \rightarrow T(V)^m.$$

Let $\mathbf{i} = (i_1, \dots, i_n) \in \mathbb{N}^n$. Then

$$\Phi^{(n)} \circ \pi_E(\mathbf{i}) = (\Phi(e_{i_1}), \dots, \Phi(e_{i_n})).$$

Applying $R^{(m)}$ selects the m refined components of the admissible record. On the other hand,

$$\widehat{R}(\mathbf{i}) = (j_1, \dots, j_m),$$

and so

$$\Phi^{(m)} \circ \pi_E \circ \widehat{R}(\mathbf{i}) = (\Phi(e_{j_1}), \dots, \Phi(e_{j_m})).$$

Both sides produce the same m embedded admissible events. Thus the square commutes.

Tensor product update. Given a record $\mathbf{i} \in \mathbb{N}^n$, define the cumulative factor

$$U(\mathbf{i}) := \prod_{k=1}^n \Phi(e_{i_k})$$

using the fixed associative product in $T(V)$. If \widehat{R} eliminates or reorders indices corresponding to incomparable events, the product over the refined record is

$$U(\widehat{R}(\mathbf{i})) = \prod_{\ell=1}^m \Phi(e_{j_\ell}).$$

Independence under commuting factors. If e_{i_p} and e_{i_q} are incomparable, their embeddings commute: $\Phi(e_{i_p})\Phi(e_{i_q}) = \Phi(e_{i_q})\Phi(e_{i_p})$. Any two admissible refinements differ by permutations of indices of such incomparable elements, hence

$$U(\mathbf{i}) = U(\sigma(\mathbf{i}))$$

for any permutation σ generated by such swaps. If all factors commute, $U(\mathbf{i})$ depends only on the multiset $\{\Phi(e_{i_1}), \dots, \Phi(e_{i_n})\}$ and is therefore independent of the ordering of the record.

Conclusion. In ZFC, the operators π_E , \widehat{R} , $\Phi^{(k)}$, and $R^{(m)}$ are well-defined; the naturality square commutes; the update operator U is well-defined; and the invariance properties under commuting admissible factors hold. This proves the proposition.

One may view this as a commuting diagram in a monoidal category. \square

A.2 Proposition 2

A.3 Proposition 4

A.4 Proposition 5

A.5 Proposition 6

A.6 Proposition 7

A.7 Proposition 8

A.8 Proposition 9

Appendix B

Notation

This appendix summarizes the symbols and conventions used throughout the monograph. The goal is clarity. Every notation corresponds to an operational procedure: recording events, merging ledgers, composing systems, or evolving a notebook of admissible distinctions forward in time.

Events and Ledgers

- An *event* is a measurable, irreversible update to a system's state. A finite set of observations produces a finite, ordered record of events.
- A *ledger* is the notebook containing this ordered record. Ledgers are denoted by calligraphic symbols ($\mathcal{L}, \mathcal{M}, \dots$).
- The *Axiom of Order* guarantees that every ledger is a countable, totally ordered sequence of events.

Tensor Composition

- Independent ledgers compose via the tensor product

$$\mathcal{L} \otimes \mathcal{M},$$

which produces a joint ledger with no implied evolution. The tensor product is symmetric up to canonical isomorphism and carries no time direction.

- The tensor product *does not* imply interaction. It merely constructs a space capable of recording joint events.

Merge Operator

- Compatible ledgers merge by addition, written

$$\mathcal{L} + \mathcal{M}.$$

This operation is commutative and introduces no new events. It simply coalesces distinctions already present in the two ledgers.

- Because $+$ is commutative and order-independent, no commutator is defined on $+$.

Evolution (Fold) Operators

- A *fold* is an evolution operator that acts on a ledger,

$$F : \mathcal{L} \rightarrow \mathcal{L}.$$

A fold updates the ledger forward in time, reconciling the existing record with a new admissible distinction.

- Successive folds are composed using standard function composition,

$$G \circ F : \mathcal{L} \rightarrow \mathcal{L}.$$

Only folds carry a time direction.

- When the sequence of folds varies with time,

$$\bigcirc_{i=1}^n F_i = F_n \circ F_{n-1} \circ \cdots \circ F_1.$$

This is the *iterated fold*.

- When the fold is identical at each step, we write

$$F^{\circ n} = \underbrace{F \circ F \circ \cdots \circ F}_{n \text{ times}}.$$

We avoid the notation F^n to prevent confusion with contravariant tensor indices.

Commutators

- The commutator measures the failure of two folds to commute:

$$[F, G] = F \circ G - G \circ F.$$

Because the tensor product and addition introduce no time ordering, the commutator is defined only for folds.

- Nonzero commutators represent informational curvature: different orders of reconciliation produce different ledgers.

Functionals and Variations

- A functional on a ledger-refined trajectory is written

$$J[x] = \int_a^b f(t, x(t), \dot{x}(t)) dt.$$

- An admissible variation is of the form

$$x_\varepsilon(t) = x(t) + \varepsilon\eta(t), \quad \eta(a) = \eta(b) = 0.$$

- The first variation is the directional derivative

$$\delta J[x; \eta] = \left. \frac{d}{d\varepsilon} J[x_\varepsilon] \right|_{\varepsilon=0}.$$

- The Euler–Lagrange equation is the condition that $\delta J[x; \eta] = 0$ for all admissible η .

These conventions are used consistently in later chapters. No symbol is overloaded, and every operator corresponds to a physical or informational procedure: composition, merging, or evolution. All results follow from these definitions and the axioms introduced in Chapter 1.

Bibliography

- [1] Scott Aaronson. *Quantum Computing Since Democritus*. Cambridge University Press, 2013.
- [2] Yakir Aharonov and David Bohm. Significance of electromagnetic potentials in the quantum theory. *Physical Review*, 115(3):485–491, 1959.
- [3] Archimedes. *The Works of Archimedes*. Cambridge University Press, 1912.
- [4] Aristotle. *The Complete Works of Aristotle, Vol. 1*. Princeton University Press, Princeton, 1984. Physics, Books I–VIII.
- [5] John Backus. The syntax and semantics of ALGOL. In *Proceedings of the International Symposium on Symbolic Languages*. IFIP, 1959. Introduced the notation later termed Backus–Naur Form.
- [6] Francis Bacon. *Novum Organum*. John Bill, London, 1620.
- [7] Stefan Banach and Alfred Tarski. Sur la decomposition des ensembles de points en parties respectivement congruentes. *Fundamenta Mathematicae*, 6:244–277, 1924.
- [8] George K Batchelor. *An Introduction to Fluid Dynamics*. Cambridge University Press, 1967.
- [9] John S. Bell. On the einstein podolsky rosen paradox. *Physics*, 1(3):195–200, 1964.

- [10] George Berkeley. *The Analyst: or, A Discourse Addressed to an Infidel Mathematician*. London, 1734. Available in many reprints; see, e.g., Project Gutenberg (Eprint #27200).
- [11] George D. Birkhoff. Proof of the ergodic theorem. *Proceedings of the National Academy of Sciences*, 17(12):656–660, 1931.
- [12] Niels Bohr. The quantum postulate and the recent development of atomic theory. *Nature*, 121(3050):580–590, 1928.
- [13] Ludwig Boltzmann. *Weitere Studien über das Wärmegleichgewicht unter Gasmolekülen*, volume 66. 1872.
- [14] Ludwig Boltzmann. über die beziehung zwischen dem zweiten hauptsatze der mechanischen wärmetheorie und der wahrscheinlichkeitsrechnung, respektive den sätzen über das wärmegleichgewicht. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Classe*, 76:373–435, 1877.
- [15] Ludwig Boltzmann. *Vorlesungen über Gastheorie*. Johann Ambrosius Barth, Leipzig, 1896.
- [16] Luca Bombelli, Joohan Lee, David Meyer, and Rafael D. Sorkin. Space-time as a causal set. *Physical Review Letters*, 59(5):521–524, 1987.
- [17] Daniel Bonn, Serge Rodts, Jan Groenewold, Hamid Kellay, and Gerard Wegdam. The physics of quicksand. *Nature*, 435:633–636, 2005.
- [18] George E. P. Box and Gwilym M. Jenkins. *Time Series Analysis: Forecasting and Control*. Holden-Day, San Francisco, 1976.
- [19] William Henry Bragg and William Lawrence Bragg. The reflection of x-rays by crystals. *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, 88(605):428–438, 1913.

- [20] Susanne C. Brenner and Ridgway L. Scott. *The Mathematical Theory of Finite Element Methods*. Springer, New York, 3rd edition, 2008.
- [21] Georg Cantor. *Beiträge zur Begründung der transfiniten Mengenlehre*, volume 46. 1895.
- [22] Augustin-Louis Cauchy. *Cours d'Analyse*. de l'Imprimerie Royale, Paris, 1821.
- [23] Augustin-Louis Cauchy. Sur l'équilibre et le mouvement intérieur des corps solides. *Exercices de Mathématiques*, pages 294–316, 1828.
- [24] G. J. Chaitin. A theory of program size formally identical to information theory. *Journal of the ACM*, 22(3):329–340, 1975.
- [25] John A. Christian and Scott Cryan. A survey of lidar technology and its use in spacecraft relative navigation. *AIAA Paper*, 2013.
- [26] Philippe G. Ciarlet. *The Finite Element Method for Elliptic Problems*. North-Holland, 1978.
- [27] Arthur H. Compton. A quantum theory of the scattering of x-rays by light elements. *Physical Review*, 21(5):483–502, 1923.
- [28] Sony Corporation. Digital audio disc system. U.S. Patent 4,347,632, 1980.
- [29] Richard Courant and David Hilbert. *Methods of Mathematical Physics, Vol. I*. Interscience Publishers, New York, 1953. Classical treatment of the calculus of variations and Euler–Lagrange equations.
- [30] Thomas M. Cover and Joy A. Thomas. *Elements of Information Theory*. Wiley, 2 edition, 2006.
- [31] James P. Crutchfield and Karl Young. Inferring statistical complexity. *Physical Review Letters*, 63(2):105–108, 1989.

- [32] George B. Dantzig. *Linear Programming and Extensions*. Princeton University Press, Princeton, NJ, 1963.
- [33] B. A. Davey and H. A. Priestley. *Introduction to Lattices and Order*. Cambridge University Press, 2nd edition, 2002.
- [34] Philip J. Davis and Philip Rabinowitz. *Methods of Numerical Integration*. Academic Press, 1975.
- [35] Clinton Davisson and Lester H. Germer. Diffraction of electrons by a crystal of nickel. *Physical Review*, 30(6):705–740, 1927.
- [36] Carl de Boor. *A Practical Guide to Splines*. Springer, New York, 1978.
- [37] Pierre Louis Moreau de Maupertuis. *Accord de differentes lois de la nature qui avaient jusqu ici paru incompatibles*. Imprimerie Royale, Paris, 1744.
- [38] Adhemar Jean Claude Barre de Saint-Venant. Memoire sur la flexion des prismes. *Journal de Mathematiques Pures et Appliquees*, pages 89–189, 1860.
- [39] P. A. M. Dirac. *The Principles of Quantum Mechanics*. Oxford University Press, Oxford, 4th edition, 1958.
- [40] Paul A. M. Dirac. The quantum theory of the electron. *Proceedings of the Royal Society of London A*, 117:610–624, 1928.
- [41] Johann Peter Gustav Lejeune Dirichlet. Über die vertheilung der monischen werthe ganzer linearer formen zwischen zwei grenzen. *Abhandlungen der Königlich Preußischen Akademie der Wissenschaften*, 1834.
- [42] David S. Dummit and Richard M. Foote. *Abstract Algebra*. Wiley, 2004.

- [43] John Earman. An attempt to formulate a causal theory of time. *Journal of Philosophy*, 71(17):561–579, 1974.
- [44] A. Einstein. Die feldgleichungen der gravitation. *Sitzungsberichte der Preussischen Akademie der Wissenschaften*, pages 844–847, 1915.
- [45] Albert Einstein. Zur elektrodynamik bewegter körper. *Annalen der Physik*, 17:891–921, 1905. English translation: "On the Electrodynamics of Moving Bodies".
- [46] Albert Einstein, Boris Podolsky, and Nathan Rosen. Can quantum-mechanical description of physical reality be considered complete? *Physical Review*, 47(10):777–780, 1935.
- [47] Philips Electronics. Digital optical recording system. U.S. Patent 4,363,116, 1980.
- [48] Leonhard Euler. *Methodus inveniendi lineas curvas maximi minimive proprietate gaudentes*. Lausanne & Geneva, 1744.
- [49] Lawrence C. Evans. *Partial Differential Equations*. American Mathematical Society, Providence, 2 edition, 2010.
- [50] Richard P. Feynman. Space–time approach to non-relativistic quantum mechanics. *Reviews of Modern Physics*, 20:367–387, 1948.
- [51] Richard P. Feynman, Robert B. Leighton, and Matthew Sands. *The Feynman Lectures on Physics*, volume 1–3. Addison-Wesley, Reading, MA, 1965. Classic introductory lectures on fundamental physics.
- [52] David Finkelstein. Causal sets as the deep structure of spacetime. *International Journal of Theoretical Physics*, 27(4):473–484, 1988.
- [53] David Finkelstein. *Quantum Relativity: A Synthesis of the Ideas of Einstein and Heisenberg*. Springer, Berlin, 1996.

- [54] Abraham A Fraenkel. *Einleitung in die Mengenlehre*. Springer, 1922.
- [55] Y. Fukuda, others, and Super-Kamiokande Collaboration. The super-kamiokande detector. *Nuclear Instruments and Methods in Physics Research A*, 501:418–462, 2003.
- [56] Boris Galerkin. Series solution of some problems of elastic equilibrium of rods and plates. *Vestnik Inzhenerov*, pages 897–908, 1915.
- [57] Galileo Galilei. *Discorsi e dimostrazioni matematiche intorno a due nuove scienze*. Elsevier, Leiden, 1638.
- [58] C. F. Gauss. *Disquisitiones Generales Circa Superficies Curvas*. Göttingen, 1828.
- [59] Murray Gell-Mann and Maurice Lévy. The axial vector current in beta decay. *Il Nuovo Cimento*, 16:705–726, 1960.
- [60] J. Willard Gibbs. Fourier series. *Nature*, 59:606, 1899.
- [61] Herbert Goldstein, Charles Poole, and John Safko. *Classical Mechanics*. Addison–Wesley, San Francisco, 3 edition, 2002.
- [62] Gene H. Golub and Charles F. Van Loan. *Matrix Computations*. Johns Hopkins University Press, 4 edition, 2013.
- [63] Kurt Gödel. Über formal unentscheidbare sätze der principia mathematica und verwandter systeme i. *Monatshefte für Mathematik und Physik*, 38:173–198, 1931.
- [64] Ernst Hairer and Gerhard Wanner. *Solving Ordinary Differential Equations II: Stiff and Differential-Algebraic Problems*. Springer, Berlin, 1993.
- [65] Paul R. Halmos. *Finite-Dimensional Vector Spaces*. Springer, 1958.

- [66] Paul R. Halmos. *Naive Set Theory*. Springer, 1974. Concise exposition of ZFC and its role in functional analysis.
- [67] William Rowan Hamilton. On a general method in dynamics. *Philosophical Transactions of the Royal Society of London*, 124:247–308, 1834.
- [68] G.H. Hardy and E.M. Wright. *An Introduction to the Theory of Numbers*. Oxford University Press, 1938.
- [69] S. W. Hawking. Particle creation by black holes. *Communications in Mathematical Physics*, 43:199–220, 1975. Introduced the concept of black hole radiation via quantum field theory in curved spacetime.
- [70] S. W. Hawking and G. F. R. Ellis. *The Large Scale Structure of Space-Time*. Cambridge University Press, Cambridge, 1973.
- [71] Oliver Heaviside. *Electromagnetic Theory, Vol. I*. The Electrician Printing and Publishing Company, London, 1893.
- [72] Heinrich Hertz. Ueber einen einfluss des ultravioletten lichtes auf die elektrische entladung. *Annalen der Physik*, 267(8):983–1000, 1887.
- [73] David Hilbert. *Grundlagen der Geometrie*. Teubner, Leipzig, 1899. First edition of Hilbert’s axiomatic foundations of geometry.
- [74] David Hilbert. Mathematical problems. *Bulletin of the American Mathematical Society*, 8(10):437–479, 1902. English translation of Hilbert’s 1900 address.
- [75] K. Hirata, T. Kajita, M. Koshiba, et al. Observation of a neutrino burst from the supernova sn 1987a. *Physical Review Letters*, 58(14):1490–1493, 1987.
- [76] Douglas R. Hofstadter. *Gödel, Escher, Bach: An Eternal Golden Braid*. Basic Books, New York, 1979.

- [77] Sabine Hossenfelder. *Lost in Math: How Beauty Leads Physics Astray*. Basic Books, New York, 2018.
- [78] Kiyoshi Itô. Stochastic integral. *Proceedings of the Imperial Academy, Tokyo*, 20:519–524, 1944.
- [79] Kiyoshi Itô. On stochastic differential equations. *Memoirs of the American Mathematical Society*, 4:1–51, 1951.
- [80] Carl Gustav Jacob Jacobi. Über eine neue auflösungsart der bei der methode der variationen vorkommenden linearen differentialgleichungen. *Journal für die reine und angewandte Mathematik*, 30:51–82, 1845.
- [81] Thomas Jech. *Set Theory: The Third Millennium Edition, Revised and Expanded*. Springer, Berlin, 2003.
- [82] E. Joos, H. D. Zeh, C. Kiefer, D. Giulini, J. Kupsch, and I.-O. Stamatescu. *Decoherence and the Appearance of a Classical World in Quantum Theory*. Springer, 2003.
- [83] Immanuel Kant. *Kritik der reinen Vernunft*. Johann Friedrich Hartknoch, Riga, 1781.
- [84] G. M. Kelly. *Basic Concepts of Enriched Category Theory*. London Mathematical Society Lecture Note Series. Cambridge University Press, 1982.
- [85] Johannes Kepler. *Astronomia Nova*. Heirs of Godefridus Tampachius, Prague, 1609.
- [86] Donald E. Knuth. *The Art of Computer Programming, Volume 3: Sorting and Searching*. Addison-Wesley, Reading, Massachusetts, 2nd edition, 1998.
- [87] A. N. Kolmogorov. Three approaches to the quantitative definition of information. *Problems of Information Transmission*, 1(1):1–7, 1965.

- [88] Andrey N. Kolmogorov. *Grundbegriffe der Wahrscheinlichkeitsrechnung*. Springer, Berlin, 1933. English translation: *Foundations of the Theory of Probability*, Chelsea Publishing, 1950.
- [89] Andrey N. Kolmogorov and Sergei V. Fomin. *Introductory Real Analysis*. Dover Publications, New York, 1970.
- [90] B.O. Koopman. Hamiltonian systems and transformation in hilbert space. *Proceedings of the National Academy of Sciences*, 17(5):315–318, 1931.
- [91] Kenneth Kunen. *Set Theory: An Introduction to Independence Proofs*, volume 102 of *Studies in Logic and the Foundations of Mathematics*. North-Holland, Amsterdam, 1980.
- [92] Joseph-Louis Lagrange. *Mécanique Analytique*. Desaint, Paris, 1788.
- [93] Cornelius Lanczos. *The Variational Principles of Mechanics*. Dover Publications, New York, 1970.
- [94] Saunders Mac Lane. *Categories for the Working Mathematician*. Springer, 1971. For categorical formulations of tensor and operator structures.
- [95] Serge Lang. *Algebra*. Springer, New York, 3rd edition, 2002.
- [96] Paul Langevin. The evolution of space and time. *Scientia*, pages 31–54, 1911.
- [97] Tom Leinster. *Basic Category Theory*, volume 143 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2014.
- [98] David Lewis. The paradoxes of time travel. *American Philosophical Quarterly*, 13(2):145–152, 1976.

- [99] Ming Li and Paul Vitányi. *An Introduction to Kolmogorov Complexity and Its Applications*. Springer, 2 edition, 1997.
- [100] Ming Li and Paul Vitányi. *An Introduction to Kolmogorov Complexity and Its Applications*. Texts in Computer Science. Springer, New York, 3rd edition, 2008.
- [101] Hendrik A. Lorentz. Electromagnetic phenomena in a system moving with any velocity less than that of light. *Proceedings of the Royal Netherlands Academy of Arts and Sciences*, 6:809–831, 1904.
- [102] David B. Malament. The class of continuous timelike curves determines the topology of spacetime. *Journal of Mathematical Physics*, 18(7):1399–1404, 1977.
- [103] Benoit B. Mandelbrot. *The Fractal Geometry of Nature*. W. H. Freeman, 1982.
- [104] Guglielmo Marconi. Syntonic wireless telegraphy. *Proceedings of the Royal Society of London*, 70(455–466):341–349, 1901.
- [105] Donald A. Martin and Robert M. Solovay. Internal cohen extensions. *Annals of Mathematical Logic*, 2(2):143–178, 1970.
- [106] James Clerk Maxwell. A dynamical theory of the electromagnetic field. *Philosophical Transactions of the Royal Society of London*, 155:459–512, 1865. Original presentation of the equations of electromagnetism.
- [107] A. A. Michelson and E. W. Morley. On the relative motion of the earth and the luminiferous ether. *American Journal of Science*, 34(203):333–345, 1887.
- [108] Hermann Minkowski. Raum und zeit. In *Physikalische Zeitschrift*, volume 10, pages 104–111, 1908. Lecture delivered at 80th Assembly

of German Natural Scientists and Physicians, Cologne, 1908. English translation: "Space and Time."

- [109] Charles W. Misner, Kip S. Thorne, and John Archibald Wheeler. *Gravitation*. W. H. Freeman, 1973.
- [110] Nevill Francis Mott. The wave mechanics of α -ray tracks. *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, 126(801):79–84, 1929.
- [111] Peter Naur. Revised report on the algorithmic language ALGOL 60. *Communications of the ACM*, 6(1):1–17, 1963. Standard reference defining Backus–Naur Form.
- [112] Claude-Louis Navier. Memoire sur les lois du mouvement des fluides. *Mem. Acad. Sci. Inst. France*, 6:389–440, 1823.
- [113] Isaac Newton. *Philosophiæ Naturalis Principia Mathematica*. Royal Society of London, 1687. Translated as *The Mathematical Principles of Natural Philosophy*, University of California Press, 1934.
- [114] Emmy Noether. Invariante variationsprobleme. *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Math.-Phys. Klasse*, pages 235–257, 1918.
- [115] Harry Nyquist. Certain topics in telegraph transmission theory. *Transactions of the American Institute of Electrical Engineers*, 47(2):617–644, 1928.
- [116] William of Ockham. *Summa Logicae*. 1323. Original Latin manuscripts; various critical editions exist.
- [117] Donald S. Ornstein and Benjamin Weiss. Entropy and data compression. *IEEE Transactions on Information Theory*, 37(3):845–859, 1991.

- [118] Roger Penrose. *Techniques of Differential Topology in Relativity*. Society for Industrial and Applied Mathematics, Philadelphia, 1972.
- [119] Arno A. Penzias and Robert W. Wilson. A measurement of excess antenna temperature at 4080 mc/s. *Astrophysical Journal*, 142:419–421, 1965.
- [120] Asher Peres. Separability criterion for density matrices. *Physical Review Letters*, 77(8):1413–1415, 1995.
- [121] Max Planck. Zur theorie des gesetzes der energieverteilung im normal-spectrum. *Verhandlungen der Deutschen Physikalischen Gesellschaft*, 2:237–245, 1900. Introduces the quantum of action and the foundations of quantized harmonic motion.
- [122] Max Planck. Über das gesetz der energieverteilung im normalspektrum. *Annalen der Physik*, 4:553–563, 1901.
- [123] Max Planck. *The Theory of Heat Radiation*. P. Blakiston’s Son & Co., Philadelphia, 1914. Translated by M. Masius. Reprinted by Dover Publications, 1959.
- [124] Plato. *Parmenides*. Hackett Publishing, Indianapolis, 1996.
- [125] R. V. Pound and G. A. Rebka. Gravitational red-shift in nuclear resonance. *Physical Review Letters*, 3(9):439–441, 1959.
- [126] Lewis F. Richardson. The problem of contiguity: An appendix to statistics of deadly quarrels. *General Systems Yearbook*, 6:139–187, 1961.
- [127] David Rideout and Rafael D. Sorkin. A classical sequential growth dynamics for causal sets. *Physical Review D*, 61:024002, 1999.
- [128] Bernhard Riemann. *Über die Hypothesen, welche der Geometrie zu Grunde liegen*. 1854. Habilitationsschrift, Universität Göttingen.

- [129] Wolfgang Rindler. *Relativity: Special, General, and Cosmological*. Oxford University Press, 2nd edition, 2006.
- [130] V.A. Rohlin. Lectures on the entropy theory of transformations. *Russian Mathematical Surveys*, 22(5):1–52, 1967.
- [131] Vera C. Rubin, W. Kent Ford, and Norbert Thonnard. Rotational properties of 21 sc galaxies with a large range of luminosities and radii, from ngc 4605 ($r = 4$ kpc) to ugc 2885 ($r = 122$ kpc). *Astrophysical Journal*, 238:471–487, 1980.
- [132] Vera C. Rubin and W. Kent Jr. Ford. Rotation of the andromeda nebula from a spectroscopic survey of emission regions. *Astrophysical Journal*, 159:379–403, 1970.
- [133] Bertrand Russell. Mathematical logic as based on the theory of types. *American Journal of Mathematics*, 30:222–262, 1908.
- [134] E. Schroedinger. Discussion of probability relations between separated systems. *Mathematical Proceedings of the Cambridge Philosophical Society*, 31:555–563, 1935.
- [135] Erwin Schroedinger. Quantisierung als eigenwertproblem. *Annalen der Physik*, 79:361–376, 1926. First paper on wave mechanics; introduces the Schroedinger equation.
- [136] Karl Schwarzschild. Uber das gravitationsfeld eines massenpunktes nach der einsteinschen theorie. *Sitzungsberichte der Koniglich Preussischen Akademie der Wissenschaften*, pages 189–196, 1916.
- [137] Dana Scott. Outline of a mathematical theory of computation. In *Proceedings of the Fourth Annual Princeton Conference on Information Sciences and Systems*. 1970.

- [138] Claude E. Shannon. A mathematical theory of communication. *Bell System Technical Journal*, 27:379–423, 623–656, 1948.
- [139] Claude E. Shannon. Communication in the presence of noise. *Proceedings of the IRE*, 37(1):10–21, 1949.
- [140] Walter A. Shewhart. *Economic Control of Quality of Manufactured Product*. D. Van Nostrand Company, New York, 1931.
- [141] Thomas Simpson. *Essays on Several Curious and Useful Subjects in Speculative and Mix'd Mathematicks*. London, 1743.
- [142] Michael Sipser. *Introduction to the Theory of Computation*. PWS Publishing, 1997.
- [143] Steven W. Smith. *The Scientist and Engineer's Guide to Digital Signal Processing*. California Technical Publishing, San Diego, 1999.
- [144] R. J. Solomonoff. A formal theory of inductive inference. part i. *Information and Control*, 7(1):1–22, 1964.
- [145] R. J. Solomonoff. A formal theory of inductive inference. part ii. *Information and Control*, 7(2):224–254, 1964.
- [146] Rafael D. Sorkin. Finitary substitute for continuous topology. In R. Penrose and C. J. Isham, editors, *Quantum Concepts in Space and Time*, pages 254–275. Clarendon Press, Oxford, 1991.
- [147] Rafael D. Sorkin. Causal sets: Discrete gravity. *Lectures on Quantum Gravity (Proceedings, Valdivia)*, 2003.
- [148] Rafael D. Sorkin. Causal sets: Discrete gravity. In Andres Gomberoff and Donald Marolf, editors, *Lectures on Quantum Gravity*, pages 305–327. Springer, Boston, MA, 2005. Foundational overview of the causal set approach to quantum gravity.

- [149] George Gabriel Stokes. On the theories of the internal friction of fluids in motion. *Transactions of the Cambridge Philosophical Society*, 8:287–319, 1845.
- [150] George Gabriel Stokes. On the theories of the internal friction of fluids in motion. *Transactions of the Cambridge Philosophical Society*, 8:287–319, 1850.
- [151] Gilbert Strang and George Fix. An analysis of the finite element method. *Prentice–Hall Series in Automatic Computation*, 1973.
- [152] Alfred Tarski. Sur la decomposition des ensembles de points en parties respectivement congruentes. *Fundamenta Mathematicae*, 6:244–277, 1924.
- [153] Edwin F. Taylor and John Archibald Wheeler. *Spacetime Physics*. W. H. Freeman, 2nd edition, 1992.
- [154] Stevo Todorčević. *Introduction to Ramsey Spaces*. Princeton University Press, Princeton, NJ, 2010.
- [155] Alan M. Turing. On computable numbers, with an application to the entscheidungsproblem. *Proceedings of the London Mathematical Society*, 42:230–265, 1936.
- [156] W. G. Unruh. Notes on black hole evaporation. *Physical Review D*, 14(4):870–892, 1976. Demonstrated the Unruh effect, showing thermal spectra from accelerated frames.
- [157] John von Neumann. *Theory of Self-Reproducing Automata*. University of Illinois Press, 1966.
- [158] Arthur P. Warner. Speed-indicator and recorder, 1902. The Warner Auto-Meter, early automobile mechanical speedometer.

- [159] Hermann Weyl. *Philosophy of Mathematics and Natural Science*. Princeton University Press, 1949.
- [160] John A. Wheeler. Information, physics, quantum: The search for links. In W. H. Zurek, editor, *Complexity, Entropy, and the Physics of Information*, pages 3–28. Addison-Wesley, Redwood City, CA, 1990.
- [161] John A. Wheeler and Wojciech H. Zurek. *Quantum Theory and Measurement*. Princeton University Press, Princeton, NJ, 1983.
- [162] Alan H. Wilson. The theory of electronic semi-conductors. *Proceedings of the Royal Society A*, 133(821):458–491, 1931.
- [163] Takahiro Yoshida. Electronic vehicle speedometer, 1980. Counts wheel rotation pulses from a magnetic pickup to drive a digital display.
- [164] Ludwig Zehnder. Ein neuer interferenzrefraktor. *Zeitschrift für Instrumentenkunde*, 11:275–?, 1891.
- [165] Ernst Zermelo. Investigations in the foundations of set theory i. *Mathematische Annalen*, 65(2):261–281, 1908.
- [166] Wojciech H. Zurek. Decoherence, einselection, and the quantum origins of the classical. *Reviews of Modern Physics*, 75:715–775, 2003.