Hinting for fully applicative STLC

Terms

```
Terms: e ::= x \mid (e : H) \mid \lambda x. e \mid e_1 e_2 \mid \text{pair} \mid \text{outl} \mid \text{outr} \mid \text{inl} \mid \text{inr} \mid \text{case} \mid \text{unit} \mid \mathbf{0}\text{-elim}
```

Judgements:

 $\Gamma \vdash e \Leftarrow H \Rightarrow A$ – in context Γ , term e checks with hint H and infers type A

Declarative typing - basics

$$\frac{(x:A) \in \Gamma}{\Gamma \vdash x:A} VAR$$

$$\frac{\Gamma \vdash e : A \quad H \sqsubseteq A}{\Gamma \vdash (e : H) : A} Annor$$

Declarative typing – type-directed rules

$$\frac{\Gamma, x : A \vdash e : B}{\Gamma \vdash \lambda x. e : A \to B} \qquad \frac{\Gamma \vdash a : A \quad \Gamma \vdash f : A \to B}{\Gamma \vdash f : a : B}$$

$$\overline{\Gamma \vdash \text{pair} : A \to B \to A \times B}$$

$$\overline{\Gamma \vdash \text{outl} : A \times B \to A} \qquad \overline{\Gamma \vdash \text{outr} : A \times B \to B}$$

$$\overline{\Gamma \vdash \text{inl} : A \to A + B} \qquad \overline{\Gamma \vdash \text{inr} : B \to A + B}$$

$$\Gamma \vdash \mathtt{case} : (A \rightarrow C) \rightarrow (B \rightarrow C) \rightarrow A + B \rightarrow C$$

$$\overline{\Gamma \vdash \text{unit} : \mathbf{1}}$$
 $\overline{\Gamma \vdash \mathbf{0}\text{-elim} : \mathbf{0} \rightarrow A}$

Hinting – basic rules

$$\frac{(x:A) \in \Gamma \quad H \sqsubseteq A}{\Gamma \vdash x \Leftarrow H \Rightarrow A} VAR$$

$$\frac{\Gamma \vdash e \Leftarrow H_1 \sqcup H_2 \Rightarrow A}{\Gamma \vdash (e:H_1) \Leftarrow H_2 \Rightarrow A} \text{Annor}$$

Hinting – type-directed rules

$$\frac{\Gamma, x : A \vdash e \Leftarrow H \Rightarrow B}{\Gamma \vdash \lambda x. \ e \Leftarrow A \rightarrow H \Rightarrow A \rightarrow B}$$

$$\frac{\Gamma \vdash a \Leftarrow ? \Rightarrow A \quad \Gamma \vdash f \Leftarrow A \to H \Rightarrow A \to B}{\Gamma \vdash f \ a \Leftarrow H \Rightarrow B}$$

$$H \sqcup ? \rightarrow ? \rightarrow ? \times ? = H_1 \rightarrow H_2 \rightarrow H_3 \times H_4$$
 $H_1 \sqcup H_3 = A \quad H_2 \sqcup H_4 = B$

$$\Gamma \vdash \text{pair} \Leftarrow H \Rightarrow A \rightarrow B \rightarrow A \times B$$

$$\frac{H_1 \sqcup H_2 = A}{\Gamma \vdash \mathtt{outl} \Leftarrow H_1 \times B \to H_2 \Rightarrow A \times B \to A}$$

$$H_1 \sqcup H_2 = B$$

Hinting – type-directed rules

$$\frac{H_1 \sqcup H_2 = A}{\Gamma \vdash \mathtt{inl} \Leftarrow H_1 \to H_2 + B \Rightarrow A \to A + B}$$

$$\frac{H_1 \sqcup H_2 = B}{\Gamma \vdash \text{inr} \Leftarrow H_1 \to A + H_2 \Rightarrow B \to A + B}$$

$$H \sqcup (? \to ?) \to (? \to ?) \to ? + ? \to ? = (H_1 \to H_2) \to (H_3 \to H_4) \to H_5 + H_6 \to H_7 \underline{H_1 \sqcup H_5 = A} \quad H_3 \sqcup H_6 = B \quad H_2 \sqcup (H_4 \sqcup H_7) = C \overline{\Gamma \vdash \mathsf{case} \Leftarrow H \Rightarrow (A \to C) \to (B \to C) \to A + B \to C}$$

$$\frac{H \sqsubseteq \mathbf{1}}{\Gamma \vdash \mathtt{unit} \Leftarrow H \Rightarrow \mathbf{1}} \qquad \frac{H \sqsubseteq \mathbf{0}}{\Gamma \vdash \mathbf{0} - \mathtt{elim} \Leftarrow H \to A \Rightarrow \mathbf{0} \to A}$$