

1 Introduction

references for packing [1] [2] references for anisotropic packing problems The packing problem for centrally symmetric bodies in the plane is mostly understood. [3] [6] [4] [7] references for mathematica [8] [5] references for linear programming

We prove the Kuperberg and Kuperberg double lattice construction for polygons is an isolated local maximal for density in the full space of packings. That is, the density of nearby packings are of strictly lower density.

The structure of the paper is as follows.

We review the required results from [] lattices, packing...

We review Kuperberg and Kuperberg

We give a local parametrization of the neighborhood of the double lattice in the space of packings and give a characterization of the a correction function and show that the optimality of the local configuration problem implies the global density result.

We prove a stability result for non-linear programming problems of the type described.

the density function can be replaced with a function of the correct type and which implies the density result.

2 Local Stability

2.1 Background/Motivation/Main theorems

Definition 2.1. Let $\Xi = (\xi_i)_{i \in \mathbb{N}}$ be a sequence of isometries. Its mean volume is the limit

$$d(\Xi) = \lim_{r \rightarrow \infty} \frac{\text{vol}B(0, r)}{|\{i : \xi_i(0) \in B(0, r)\}|}. \quad (1)$$

The upper and lower mean volumes are the corresponding limits superior and inferior.

Definition 2.2. Let K be a compact set with interior. We say that Ξ is admissible for K if the interiors of $\xi_i(K)$ and $\xi_j(K)$ are disjoint whenever $i \neq j$.

Definition 2.3. Given two sequences Ξ and Ξ' of isometries, we define the premetric

$$\delta_R(\Xi, \Xi') = \sup\{ \|\xi_i^{-1}\xi_j - \xi_i'^{-1}\xi_j'\| : \quad (2)$$

$$i, j \text{ such that } \|\xi_i(0) - \xi_j(0)\| < R \text{ or } \|\xi_i'(0) - \xi_j'(0)\| < R \}.$$

If R is large enough so that the edges $\{i, j\}$ satisfying the condition in the definition of δ_R connect all the indices, then $\delta_R(\Xi, \Xi') = 0$ if and only if $\xi_i = \phi \xi_i'$ for some $\phi \in E(n)$.

Consider the space Ω of sequences of isometries whose first term is the identity. We consider the topology on Ω generate by the base of balls $\mathcal{B}_R(\Xi, \rho) = \{\Xi' : \delta_R(\Xi, \Xi') < \rho\}$.

Definition 2.4. We say Ξ is strongly extreme for K if it minimizes the mean volume among admissible elements in a neighborhood of Ξ .

Remark 2.1. If a lattice Λ is strongly extreme for K , then Λ is extreme for K [?]. That is, Λ is a local maximum for density in the topological space Ω .

Remark 2.2. If a periodic set $\Xi = \{T_1 \xi_i : 1 \in \Lambda, i = 1, \dots, N\}$ is strongly extreme, then it is periodic extreme for K [?].

Definition 2.5. Let Ξ be a sequence of isometries and let \mathcal{P} be a polyhedral complex whose underlying space is \mathbb{R}^n . For every face F of \mathcal{P} , let $I_F = \{i : \xi_i(0) \in F\}$. We say \mathcal{P} is a honeycomb of Ξ if each n -face (cell) P is the convex hull of $\{\xi_i(0) : i \in I_P\}$.

Theorem 2.1. Let Ξ be admissible for K and let \mathcal{P} be a honeycomb of Ξ . For every cell P , consider the optimization problem of minimizing $f_P(\Xi_P) = \text{vol conv}_{i \in I_P} \xi'_i(0)$ over admissible finite sequences of isometries $\Xi_P = (\xi'_i)_{i \in I_P}$. If $(\xi_i)_{i \in I_P}$ is a local minimum for each cell P , then Ξ is strongly extreme.

Theorem 2.2. Let $g_F(\Xi_F)$ be a real-valued function over $\Xi_F = (\xi'_i)_{i \in I_F}$ for each oriented $(n-1)$ -faces (ridge) of \mathcal{P} , such that $g_F(\Xi_F) = -g_{-F}(\Xi_F)$, where $-F$ is the orientation-reversed version of F . If we replace $f_P(\Xi_P)$ in the previous theorem with $f'_P(\Xi_P) = f_P(\Xi_P) + \sum_{F \in \partial P} g_F(\Xi_F)$, then again, if $(\xi_i)_{i \in I_P}$ is a local minimum for each cell P , then Ξ is strongly extreme.

comment: restrict to saturated packings, fix R large,
Full summary of below

2.1.1 Summary of [6, §2] on double lattices.

Definition 2.6. affine diameter of K in a particular direction is the length of the longest chord in K parallel to that direction.

Definition 2.7. Given a convex body K , extensive parallelogram in K is an inscribed parallelogram satisfying the conditions that the edges lengths of the parallelogram are at least $1/2$ the affine diameter of K in the direction of the edges.

Definition 2.8. a double lattice packing is generated by an extensive parallelogram if it a packing ...

Theorem 2.3 (Kuperberg and Kuperberg). If K is strictly convex and P is a maximum density double lattice packing of the plane with copies of K , then P is generated by a minimum area extensive parallelogram in K .

By a sequence of approximations, this gives that for convex bodies K that are not strictly convex, there exists a double lattice packing of maximal density that generated a minimum area extensive parallelogram in K .

This also gives an algorithm for finding the (uniqueness, does it matter?) densest double lattice packing for a convex body K , in particular, for a regular $(2n+1)$ -gons in the plane. The goal is to show that this configuration is in fact a local maximum for density in a broader sense, namely among configurations with nearby Delaunay triangulations.

2.1.2 parametrization

We are considering the configuration space of four polygons in the plane, in a neighborhood of a Kuperberg and Kuperberg configuration. Note that there are multiple choices of fundamental domains for the initial configuration, we choose the

2.1.3 constraints

Since we are working in the space of packings, the configuration space has a non-overlap constraint. Locally, this can be described by a non-negativity condition on the signed volume of triangles defined by oriented triples of vertices.

This requires a lemma to show that we are still exploring a full neighborhood.... (point point contacts in the polygons)

Lemma 2.1. *In a neighborhood of the Kuperberg Kuperberg rhomb, the non-negativity of the signed area condition is equivalent to the non-overlap condition*

picture of a possible violation of the lemma...

Signed area gives the non-intersection constraints for the conjectured optimal configuration, note that this is true only when contact is in the interior of the edges. Lemma, there are no corner to corner contacts in K+K construction?

The lemma should be a consequence of the existence of the extensive parallelogram. or is it true?

2.1.4 other theorems

Theorem 2.4. *Programs satisfying conditions (*) have a local maximum at 0.*

2.2 Pentagons

rewrite in terms of section 1

In the case of packings by regular pentagons, one can consider the configuration space of four pentagons with respect to the density function taken with respect to the Delaunay triangulation and parametrized by This can be shown to satisfy the conditions of Section ??....

2.2.1 exhibit the construction given by 2.3

2.2.2 parametrize neighborhood of four pentagons in correct configuration

the parametrization is important, should be consistent with the later sections. Can this be hidden in the code, and to just state that there exists a parametrization that satisfies the linear program theorem? Still need a sketch here.

2.2.3 constrained optimization problem

2.2.4 density function

2.2.5 verification

2.3 Heptagons

Outline of the method for heptagons.

For heptagons, the cost function is non-trivial. This is because there is a motion in the configuration space of four heptagons that increases the double Delaunay density.
(include figure)

2.3.1 find the construction given by Theorem 2.3.

2.3.2 parametrize neighborhood of four heptagons in correct configuration

include figures

2.3.3 constrained optimization problem

2.3.4 the cost functions

Construct a cost function that satisfies the condition of correction theorems and makes the program satisfy the conditions of the LP theorems.

YOU GET THE COST FUNCTION FROM THE PROGRAM, AND IT SATISFIES THE PROPERTIES OF SECTION 1....

This area could be confusing as the objective functions become fairly complicated. Need to justify replacing objective functions with modifications. i.e.,

Sketch: Minimize the area of the Delaunay triangles. If there is no nearby configuration that decreases area done. (can we increase area and still increase density? possibly locally, so need to trade between nearby double triangles. But by symmetry, introduce a cost function between the "outer" heptagons, i.e. penalize rotation in opposite directions.) In general, average area of double triangles is minimized implies average density is maximized.

The function optimized in this case is the (modified) area of the Delaunay triangles.

2.4 general (2n+1)-gons

failure in the case of the enneagon, the construction of K+K is problematic. This requires additional analysis as

four polygons with corrected area function and an $SL(2, \mathbb{R})$ motion hessian? The $SL(2, \mathbb{R})$ motion is required for the solution part, i.e. finding the initial configuration for regular polygons. Check this makes sense.

3 Formal methods

3.1 symbolic computation in an extension field

The Wolfram System supports symbolic computation over extension fields via it's pattern matching system.

4 Slicing nonlinear programs

A non-linear programming problem satisfying certain assumptions can be certified as locally optimal by a linear programming problem. For the geometric problems considered, there are *a priori* configurations given by the maximal density configurations on a subspace of configuration space, namely subsets of the double lattice packings. To produce a certificate of local optimality for this type of problem, parametrize a neighborhood of the conjectured optimal configuration and analyze the associated non-linear programming problem

$$\max_{x \in \mathbb{R}^n} f(x) \text{ subject to } g_r(x) \geq 0, r \in I$$

in a neighborhood of 0.

An appropriate choice of parametrization allows the full non-linear program to be sliced into a one-parameter family of non-linear programs that are subordinate to the linearization of the main program at 0. The following assumptions are required.

Assumptions 4.1.¹

1. Let I be a finite index set.
2. Let e_1 be the standard unit vector $\{1, 0, \dots, 0\}$ in \mathbb{R}^n .
3. For r in I , let f and g_r be analytic functions on a neighborhood of 0.
4. Assume $f(0) = g_r(0) = 0$ for all r in I .

¹These are the assumptions that are required for the pentagon packing problem. There are a number of ways they might be weakened, e.g. the condition that E be 1-dimensional is not essential.

5. Let $F(t) = \nabla f(te_1)$.

6. Let $G_r(t) = \nabla g_r(te_1)$.

7. Assume the linear program

$$\max_{x \in \mathbb{R}^n} F(0) \cdot x \text{ subject to } G_r(0) \cdot x \geq 0, r \in I$$

has a bounded solution and that the maximum is attained at 0.

8. Assume that the set of solutions in \mathbb{R}^n to

$$F(0) \cdot x = 0 \text{ subject to } G_r(0) \cdot x \geq 0, r \in I$$

is

$$E := \{te_1 : t \in \mathbb{R}\}.$$

9. Let H be the orthogonal complement of E so that $\mathbb{R}^n = E \oplus H$.

10. Assume there is an $\epsilon > 0$ so the functions $g_r(te_1) = 0$ for all $t \in (-\epsilon, \epsilon)$, for all r in I .

11. Assume $\frac{\partial}{\partial t} f(0) = 0$, $\frac{\partial^2}{\partial t^2} f(0) < 0$.

Lemma 4.1. *Given 4.1, the linear program*

$$\max_{x \in H} F(0) \cdot x \text{ subject to } G_r(0) \cdot x \geq 0, r \in I$$

has a unique maximum at $x = 0$

Proof. By assumptions 7 and 8, the linear program

$$\max_{x \in \mathbb{R}^n} F(0) \cdot x \text{ subject to } G_r(0) \cdot x \geq 0, r \in I$$

is maximized exactly on E . The feasible set $\{x : G_r(0) \cdot x \geq 0, r \in I \text{ and } x \in H\}$ is a subset of the feasible set $\{x : G_r(0) \cdot x \geq 0, r \in I\}$. Thus, the program

$$\max_{x \in H} F(0) \cdot x \text{ subject to } G_r(0) \cdot x \geq 0, r \in I$$

is maximized exactly on the non-empty intersection

$$E \cap \{x : G_r(0) \cdot x \geq 0, r \in I\} \cap H = \{0\}.$$

□

Definition 4.1. A finitely generated cone is a subset of \mathbb{R}^n which is the non-negative span of a finite set of non-zero vectors $\{v_1, \dots, v_m\}$ in \mathbb{R}^n , which are called the generators of the cone.

Definition 4.2. A conical linear program is a linear program with a constraint set that is a finitely generated cone.

The linear programs described throughout this section are always constrained to be on the intersection of half-spaces with 0 on the boundary. These are conical programs.

Definition 4.3. For a cone C , the set $C^p := \{x \in \mathbb{R}^n : v \cdot x \leq 0 \text{ for all } v \in C\}$ is the polar cone of C .

Lemma 4.2. A conical linear program with $F \neq 0$ given by

$$\max_{x \in \mathbb{R}^n} F \cdot x \text{ subject to } G_r \cdot x \geq 0, r \in I$$

(a) has a unique² maximum at $x = 0$ iff F is in the interior of the polar cone C^p of $C = \{x : G_r \cdot x \geq 0, r \in I\}$ (b) has a bounded solution iff F is in the polar cone C^p of $C = \{x : G_r \cdot x \geq 0, r \in I\}$ and attains its maximum exactly on the span of the generators v_i such that $F \cdot v_i = 0$.

Proof. If F is in the interior of the polar cone C^p , then $F \cdot v_i < 0$ for all generators v_i . Therefore $F \cdot x$ is uniquely maximized in C at the vertex. If F is on the boundary of the polar cone, then $F \cdot x$ is maximized in C exactly on the span of the generators v_i for which $F \cdot v_i = 0$ as $F \cdot v_j < 0$ otherwise. If F is outside the polar cone, then $F \cdot v_i > 0$ for some generator v_i . Then $F \cdot x$ is unbounded in C . \square

Lemma 4.3. Given 4.1, there exists $\epsilon > 0$ such that for all t in $(-\epsilon, \epsilon)$, the linear program

$$\max_{y_t \in H} F(t) \cdot y_t$$

subject to

$$G_r(t) \cdot y_t \geq 0, r \in I$$

has a unique maximum at $y_t = 0$.³

Proof. The program for $t \in (-\epsilon, \epsilon)$, for y_t in H , for each fixed t in $(-\epsilon, \epsilon)$, for some $\epsilon > 0$, can be written as a conical program on all of \mathbb{R}^n with a cone C_t in \mathbb{R}^n of co-dimension ≥ 1 by introducing further constraints $e_1 \cdot y_t \geq 0$ and $-e_1 \cdot y_t \geq 0$. By 4.1 and 4.2, $F(0)$ is in the polar cone of $C_0 = \{y_0 : G_r(0) \cdot y_0 \geq 0, e_1 \cdot y_0 \geq 0, -e_1 \cdot y_0 \geq 0\}$. As $f, g_r \in C^\omega$, the condition of $F(t)$ being in the interior of the polar cone C_t^p is open and the condition of

²The maximum satisfies a stronger uniqueness condition. It is stable under perturbations of F and G_k .

³Here y_t is a dummy variable and does not depend on t . It is labeled y_t to ease later exposition.

the feasible set $C_t = \{y_t : G_r(t) \cdot y_t \geq 0, e_1 \cdot y_t \geq 0, -e_1 \cdot y_t \geq 0\}$ being conical is open.⁴ Therefore, by 4.2 the program has a unique maximum at $y_t = 0$ for each fixed t in $(-\epsilon, \epsilon)$ for some $\epsilon > 0$. \square

Lemma 4.4. *Given 4.1 and ϵ as in 4.3, for all $t \in (-\epsilon, \epsilon)$ there exists $\delta(t) > 0$ and a cube $Q(t) \subset \mathbb{R}^n$ of side length $2\delta(t)$ such that*

$$\{(F(t) + Q(t)) \cap (\partial(C_t^p) + Q(t))\} = \emptyset.$$

Proof. This follows from 4.3, which shows $F(t)$ is in the interior of the polar cone C_t^p . Then $F(t)$ and the boundary of C_t^p can be separated and the existence of Q is trivial. \square

Corollary 4.1. *Given 4.1 and ϵ as in 4.3, for all $t \in (-\epsilon, \epsilon)$,*

$$(F(t) + \Delta) \cdot y_t \leq 0$$

whenever y_t satisfies

$$(G_r(t) + \Delta_r) \cdot y_t \geq 0, r \in I \text{ and } e_1 \cdot y_t \geq 0, -e_1 \cdot y_t \geq 0$$

where Δ and Δ_r are any points in the $2\delta(t)$ -cube $Q(t)$ and y_t is in H .

Proof. By 4.4, $F(t) + \Delta$ is in the interior of the polar cone $C_{t,\Delta}^p$, where $C_{t,\Delta} = \{y_t : (G_r(t) + \Delta_r) \cdot y_t \geq 0, e_1 \cdot y_t \geq 0, -e_1 \cdot y_t \geq 0, r \in I\}$. \square

Lemma 4.5. *Given 4.1 and ϵ as in 4.3, for all $t \in (-\epsilon, \epsilon)$, let $y_t = x - te_1 \in H$. Choose $\Delta = \Delta(y_t)$ and $\Delta_r = \Delta_r(y_t)$ in the $2\delta(t)$ -cube $Q(t)$ to be the corner given by the sign of $x - te_1 = y_t$. Then there is an ϵ_t for which*

$$(F(t) + \Delta(y_t)) \cdot y_t \leq 0 \implies f(x) - f(te_1) \leq 0$$

and

$$(G_r(t) + \Delta_r(y_t)) \cdot y_t \leq 0 \implies g_r(x) - g_r(te_1) = g_r(x) \leq 0$$

for all $\|y_t\| \leq \epsilon_t$.

Proof. This follows from the local expansions of the nonlinear program. By this choice of $\Delta(y_t)$ and $\Delta_r(y_t)$,

$$\begin{aligned} f(x) - f(te_1) &= F(t) \cdot (x - te_1) + O(t^2) = F(t) \cdot y_t + O(t^2) \\ &\leq F(t) \cdot y_t + \delta(t)\|y_t\|_1 = (F(t) + \Delta(y_t)) \cdot y_t \end{aligned}$$

⁴The relationships between the constraint cone, the generators v_i and the constraint gradients G_k is subtle, but the condition being open essentially follows from the continuity of the distance function.

and using assumption 10,

$$\begin{aligned} g_r(x) &= g_r(x) - g_r(te_1) = G_r(t) \cdot (x - te_1) + O(t^2) = G_r(t) \cdot y_t + O(t^2) \\ &\leq G_r(t) \cdot y_t + \delta(t)\|y_t\|_1 = (G_r(t) + \Delta_r(y_t)) \cdot y_t. \end{aligned}$$

□

By 4.4 and 4.1, for t in $(-\epsilon, \epsilon)$, the program

$$\max_{y_t \in H} (F(t) + \Delta) \cdot y_t \text{ subject to } (G_r + \Delta_r) \cdot y_t$$

is uniquely maximized at $y_t = 0$ for any choice of Δ , Δ_r in the $2\delta(t)$ cube $Q(t)$. Combined with 4.5, there is an ϵ_t neighborhood of 0 where $f(y_t + te_1)$ is less than $f(te_1)$ on $\cup_{\Delta_r \in Q(t)} \{y_t : (G_r + \Delta_r) \cdot y_t \geq 0, r \in I, y_t \in H\}$, which contains the feasible set $\{y_t : g_r(y_t + te_1) \geq 0, r \in I, y_t \in H\}$. Therefore the nonlinear programs $f(y_t + te_1)$ subject to $g_r(y_t + te_1) \geq 0$, $y_t \in H$, which are parameterized by t in $(-\epsilon, \epsilon)$, have local maxima at $y_t = 0$. This gives the following:

Theorem 4.1. *Given 4.1, a fixed t in $(-\epsilon, \epsilon)$ and choosing Δ and Δ_r as in 4.5, for x satisfying $g_r(x) \geq 0$ for all r in I and $y_t = x - te_1$ in H , there exist linear programs⁵*

$$\max_{y_t \in H} (F(t) + \Delta(y_t)) \cdot y_t \text{ subject to } (G_r(t) + \Delta_r(y_t)) \cdot y_t \geq 0$$

that give solutions to the nonlinear programs

$$\max_{x \in H + te_1} f(x) \text{ subject to } g_r(x) \geq 0$$

in an ϵ_t neighborhood of te_1 in $H + te_1$.

□

By choice of a sufficiently small ϵ and a minimal⁶ non-zero ϵ_t , 4.1 gives an open neighborhood of 0 in which the maximum value of the original nonlinear program occurs on E . The assumptions for the first and second t -derivatives at 0 shows 0 to be a local maximum for the nonlinear program

$$\max_{x \in \mathbb{R}^n} f(x) \text{ subject to } g_r(x) \geq 0.$$

Theorem 4.2. *A nonlinear program satisfying 4.1 has an isolated local maximum at 0 with $f(0) = 0$.*

□

⁵These programs may depend on a choice of $y_t \in H$, but $f(x)$ is always less than $f(te_1)$ by 4.5.

⁶This exists by a compactness argument.

References

- [1] P Brass, W Moser, and J Pach. *Research Problems in Discrete Geometry*, volume 1. Springer, 2005.
- [2] J H Conway, C Goodman-Strauss, and N Sloane. Recent progress in sphere packing. *Current Developments in Mathematics*, pages 37–76, 1999.
- [3] L Fejes Tóth. Some packing and covering theorems. *Acta Sci. Math. Szeged*, 12(Leopoldo Fejer et Frederico Riesz LXX annos natis dedicatus, Pars A):62–67, 1950.
- [4] S Gravel, V Elser, and Y Kallus. Upper bound on the packing density of regular tetrahedra and octahedra. *Discrete & Computational Geometry*, 46(4):799–818, 2011.
- [5] J B Keiper. Interval arithmetic in mathematica. *Mathematica Journal*, 5(2):66–71, 1995.
- [6] G Kuperberg and W Kuperberg. Double-lattice packings of convex bodies in the plane. *Discrete & Computational Geometry*, 5(1):389–397, 1990.
- [7] F Mário de Oliveira Filho and F Vallentin. Computing upper bounds for the packing density of congruent copies of a convex body I. *arXiv preprint arXiv:1308.4893*, 2013.
- [8] Wolfram Research. *Mathematica 10.0*. Wolfram Research, Inc., Champaign, Illinois, 2014.