

# An upper bound on packing density for circular cylinders with high aspect ratio.

Wöden Kusner  
University of Pittsburgh

March 17, 2013

# Abstract

In the early nineties, A. Bezdek and W. Kuperberg used a relatively simple to argument to show a surprising result: The maximum packing density of circular cylinders of infinite length in three-dimensional Euclidian space is exactly  $\pi/\sqrt{12}$ , the planar packing density of the circle. We modify this result to find a bound on the packing density of finite length circular cylinders. In fact, the maximum density for a capped unit radius cylinder of length  $t$  is bounded above by  $\pi/\sqrt{12} + 10/t$ .

# Circle Packing/Comments

## Results in Circle Packing

### ► Toth

A packing of a region  $R \subset \mathbb{R}^n$  by objects  $C_i$  is a family  $\mathfrak{C} = \{C_i\}_{i \in I}$  in  $\mathbb{R}^n$  with disjoint interiors.

The *upper density*  $\rho^+$  of a packing  $\mathfrak{C}$  in  $\mathbb{R}^n$  will be defined as

$$\rho^+(\mathfrak{C}) = \limsup_{r \rightarrow \infty} \frac{\text{Vol}(C_i \cap B_0(r))}{\text{Vol}(B_0(r))}$$

where  $B_0(r)$  is the ball of radius  $r$ , centered at 0.

# Circle Packing/Comments

## Results in Circle Packing

- ▶ Voronoi Cells

- ▶ Toth

A packing of a region  $R \subset \mathbb{R}^n$  by objects  $C_i$  is a family  $\mathfrak{C} = \{C_i\}_{i \in I}$  in  $\mathbb{R}^n$  with disjoint interiors.

The *upper density*  $\rho^+$  of a packing  $\mathfrak{C}$  in  $\mathbb{R}^n$  will be defined as

$$\rho^+(\mathfrak{C}) = \limsup_{r \rightarrow \infty} \frac{\text{Vol}(C_i \cap B_0(r))}{\text{Vol}(B_0(r))}$$

where  $B_0(r)$  is the ball of radius  $r$ , centered at 0.

# Circle Packing/Comments

## Results in Circle Packing

- ▶ Voronoi Cells
- ▶ Volumes and Densities
- ▶ Toth

A packing of a region  $R \subset \mathbb{R}^n$  by objects  $C_i$  is a family  $\mathfrak{C} = \{C_i\}_{i \in I}$  in  $\mathbb{R}^n$  with disjoint interiors.

The *upper density*  $\rho^+$  of a packing  $\mathfrak{C}$  in  $\mathbb{R}^n$  will be defined as

$$\rho^+(\mathfrak{C}) = \limsup_{r \rightarrow \infty} \frac{\text{Vol}(C_i \cap B_0(r))}{\text{Vol}(B_0(r))}$$

where  $B_0(r)$  is the ball of radius  $r$ , centered at 0.

# Review of Bezdek and Kuperberg

Lower bound in density:

- ▶ Obvious construction.

Upper bound on density:

# Review of Bezdek and Kuperberg

Lower bound in density:

- ▶ Obvious construction.

Upper bound on density:

- ▶ Fix a packing of Euclidian 3-space with bi-infinite cylinders.

# Review of Bezdek and Kuperberg

Lower bound in density:

- ▶ Obvious construction.

Upper bound on density:

- ▶ Fix a packing of Euclidian 3-space with bi-infinite cylinders.
- ▶ Decompose 3-space into "Voronoi" cells associated with said cylinders. The Voronoi cell of a cylinder is the region closer to that particular cylinder.



# Review of Bezdek and Kuperberg

Lower bound in density:

- ▶ Obvious construction.

Upper bound on density:

- ▶ Fix a packing of Euclidian 3-space with bi-infinite cylinders.
- ▶ Decompose 3-space into "Voronoi" cells associated with said cylinders. The Voronoi cell of a cylinder is the region closer to that particular cylinder.
- ▶ Slice the Voronoi cells perpendicular to the axis of associated cylinder.

# Review of Bezdek and Kuperberg

Lower bound in density:

- ▶ Obvious construction.

Upper bound on density:

- ▶ Fix a packing of Euclidian 3-space with bi-infinite cylinders.
- ▶ Decompose 3-space into "Voronoi" cells associated with said cylinders. The Voronoi cell of a cylinder is the region closer to that particular cylinder.
- ▶ Slice the Voronoi cells perpendicular to the axis of associated cylinder.
- ▶ Show the area of each slice of a Voronoi cell is large.
- ▶ \*Integrate\*

## Showing slices of Voronoi cells are large

- ▶ The slices are "parabolic polygons."

# Showing slices of Voronoi cells are large

- ▶ The slices are "parabolic polygons."
- ▶ Satisfy certain conditions that allow a truncation and replacement to a nicer object.

# Showing slices of Voronoi cells are large

- ▶ The slices are "parabolic polygons."
- ▶ Satisfy certain conditions that allow a truncation and replacement to a nicer object.
- ▶ Explicit computation of the area bound of such objects.

# Modifications for Finite Cylinders

- ▶ The Voronoi cell of a finite cylinder doesn't need to contain the cylinder.
- ▶ Solution: We do know that the area is less than one. (was problem 1.) We can integrate areas of slices of  
Also need to work in a bounded area/  
So we have a series of inequalities



# Modifications for Finite Cylinders

- ▶ The Voronoi cell of a finite cylinder doesn't need to contain the cylinder.
  - ▶ Solution: Cap the ends of the cylinder.
  - ▶ Voronoi slices are no longer parabolic polygons
- 
- ▶ Solution: We do know that the area is less than one. (was problem 1.) We can integrate areas of slices of  
Also need to work in a bounded area/  
So we have a series of inequalities



# Modifications for Finite Cylinders

- ▶ The Voronoi cell of a finite cylinder doesn't need to contain the cylinder.
  - ▶ Solution: Cap the ends of the cylinder.
  - ▶ Voronoi slices are no longer parabolic polygons
  - ▶ Theorem: Slices that are far away from the ends of ANY CYLINDER satisfy conditions that allow the same truncation and replacements done by Bezdek and Kuperberg.
- 
- ▶ Solution: We do know that the area is less than one. (was problem 1.) We can integrate areas of slices of  
Also need to work in a bounded area/  
So we have a series of inequalities

# Modifications for Finite Cylinders

- ▶ The Voronoi cell of a finite cylinder doesn't need to contain the cylinder.
- ▶ Solution: Cap the ends of the cylinder.
- ▶ Voronoi slices are no longer parabolic polygons
- ▶ Theorem: Slices that are far away from the ends of ANY CYLINDER satisfy conditions that allow the same truncation and replacements done by Bezdek and Kuperberg.
- ▶ We now don't know the area of some cells, also, the caps contribute volume.
- ▶ Solution: We do know that the area is less than one. (was problem 1.) We can integrate areas of slices of  
Also need to work in a bounded area/  
So we have a series of inequalities

# Modifications for Finite Cylinders

- ▶ The Voronoi cell of a finite cylinder doesn't need to contain the cylinder.
- ▶ Solution: Cap the ends of the cylinder.
- ▶ Voronoi slices are no longer parabolic polygons
- ▶ Theorem: Slices that are far away from the ends of ANY CYLINDER satisfy conditions that allow the same truncation and replacements done by Bezdek and Kuperberg.
- ▶ We now don't know the area of some cells, also, the caps contribute volume.
- ▶ Solution: We do know that the area is less than one. (was problem 1.) We can integrate areas of slices of  
Also need to work in a bounded area/  
So we have a series of inequalities

# Main Theorem

Fix  $t_0 = \frac{4}{3}(\frac{4}{\sqrt{3}} + 1)^3 = 48.3266786\dots$

## Theorem

*Fix  $t \geq 2t_0$ . Fix  $B_0(R)$ , a ball of radius  $R$  centered at 0. Fix a packing  $\mathfrak{C}$  of  $B_0(R)$  by capped  $t$ -cylinders completely contained in  $B_0(R - 2/\sqrt{3})$ . Then*

$$\rho(\mathfrak{C}) \leq \left( \frac{t + \frac{4}{3}}{\frac{\sqrt{12}}{\pi}(t - 2t_0) + (2t_0) + \frac{4}{3}} \right).$$

# Heuristic

We can expand and find a dominating hyperbola, giving

$$\rho(\mathfrak{C}) \leq \pi/\sqrt{12} + 10/t$$

for a packing  $\mathfrak{C}$  given by congruent capped cylinders.

# Quick Corollaries

- ▶ Upper bound on the packing density cylinders/spherocylinders.

## Quick Corollaries

- ▶ Upper bound on the packing density cylinders/spherocylinders.
- ▶ Strict result for ray cylinders.

# Quick Corollaries

- ▶ Upper bound on the packing density cylinders/spherocylinders.
- ▶ Strict result for ray cylinders.
- ▶ Bounds for packings of non-congruent unit radius cylinders.