The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

Iowa State University

December 2, 2014

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

i ne problem

Simulated data

The contenders
edgeR
ShrinkBayes

The contest
Constructing ROC curves
ROC results
FDR control

#### The problem

The workflow

Simulated data

#### The contenders

edgeR

ShrinkBayes

baySec

#### The contest

Constructing ROC curves

ROC results

FDR control procedures

FDR control results

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

#### The problem

i ne worktio

Simulated data

The contenders

edgeR ShrinkBayes

he contest

Constructing ROC curves ROC results FDR control procedures

### Mock heterosis data

		-	Parei	nt (1)	)		Pare	nt (2)	)		Hybr	id (3	)	Truth
HPH (	Feature 1	3	4	2	1	0	0	1	0	700	900	825	860	1
HPH (	Feature 2	0	1	1	0	2	7	5	18	50	501	400	90	1
	Feature 3	100	225	0	15	300	106	200	400	70	279	100	123	0
LPH (	Feature 4	893	400	760	901	1000	513	760	580	5	5	6	7	1
	Feature 25000	10	13	6	4	902	912	999	825	819	761	800	465	0

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

he contenders dgeR BrinkBayes

The contest
Constructing ROC curves

orves OC results DR control rocedures DR control result

The problem

The workflow

Simulated data

The contenders

edgeR

ShrinkBayes

baySec

The contest

Constructing ROC curves

ROC results

FDR control procedures

FDR control results

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

The workflow

Simulated data

Jiiilalated data

edgeR ShrinkBayes

baySeq The contest Constructing RO

Constructing ROC curves
ROC results
FDR control procedures

#### Simulation workflow

- Simulate 30 datasets:
  - 10 datasets with 4 samples (libraries, columns, etc.) per group
  - ▶ 10 with 8 per group
  - ▶ 10 with 16 per group
- For each simulated dataset, test for heterosis with
  - empirical Bayes with STAN (Eric's method)
  - edgeR
  - baySeq
  - ► ShrinkBayes
- Compare methods with ROC curves
- Check FDR control

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

The workflow

Simulated data

The contenders
edgeR
ShrinkBayes

he contest

Constructing RO curves ROC results FDR control procedures

The problem

The workflow

#### Simulated data

The contenders

edgeF

ShrinkBayes

baySed

#### The contest

Constructing ROC curves

ROC results

FDR control procedures

FDR control results

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

THE WOLKING

Simulated data

The contenders

edgeR ShrinkBayes baySeq

aySeq he contest

Constructing ROC curves
ROC results
FDR control procedures

# Apply edgeR to real data to get simulation parameters

Normalization factors



Main effects and dispersions

Parent (1)	Parent (2)	Hybrid (3)	Dispersion
$\mu_{1,1}$	$\mu_{1,2}$	$\mu_{1,3}$	$\psi_1$
$\mu_{2,1}$	$\mu_{2,2}$	$\mu_{2,3}$	$\psi_2$
$\mu_{27888,1}$	$\mu_{27888,2}$	$\mu_{27888,3}$	$\psi_{27888}$

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

The working

Simulated data

he contenders edgeR ShrinkBayes

he contest

urves OC results DR control rocedures DR control resu

# Generating the data

▶ Does feature *f* truly have heterosis?

$$\operatorname{truth}_f = I(\mu_{f,3} > \max(\mu_{f,1}, \mu_{f,2}) \text{ or } \mu_{f,3} < \min(\mu_{f,1}, \mu_{f,2}))$$

► For a dataset with 4 libraries per group,

$$y_{f,i} \stackrel{\text{iid}}{\sim} NB \left( \exp \left( c_i + \mu_{f,t(i)} \right), \ \psi_f \right)$$

- $\blacktriangleright$  t(i) is the group of library i.
- ▶ Resimulate to increase the number of libraries per group.
- Remove extremely low-count features.
- ► Take a random subset of 25000 features from the remaining ones.

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

Simulated data

The contenders edgeR ShrinkBayes

The contest
Constructing RO

Constructing ROC curves
ROC results
FDR control procedures
FDR control results

# Mock example data with 4 samples per treatment group

			Pare	nt (1)	)		Pare	nt (2)	)	ı	Hybr	id (3	)	Truth
HPH (	Feature 1	3	4	2	1	0	0	1	0	700	900	825	860	1
HPH (	Feature 2	0	1	1	0	2	7	5	18	50	501	400	90	1
	Feature 3	100	225	0	15	300	106	200	400	70	279	100	123	0
LPH (	Feature 4	893	400	760	901	1000	513	760	580	5	5	6	7	1
	Feature 25000	10	13	6	4	902	912	999	825	819	761	800	465	0

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

HE WOLKHOW

Simulated data

edgeR ShrinkBayes DaySeq

The contest Constructing RC curves

urves ROC results EDR control rocedures EDR control result:

#### The contenders

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

#### The contenders

- ▶ Fit a loglinear model to estimate main effects  $\mu_{f,t}$ 
  - Feature f = 1, ..., 25000
  - ▶ Treatment group t = 1 (parent), 2 (parent), 3 (hybrid)
- Likelihood ratio tests to get p-values  $p_{f,1}$ ,  $p_{f,2}$

$$H_{0,1}: \mu_{f,3} = \mu_{f,1}$$
  $H_{a,1}: \mu_{f,3} \neq \mu_{f,1}$   
 $H_{0,2}: \mu_{f,3} = \mu_{f,2}$   $H_{a,2}: \mu_{f,3} \neq \mu_{f,2}$ 

Final p-value:

$$p_{f, \text{edgeR}} = \begin{cases} \frac{p_{f,1}}{2} & \widehat{\mu}_{f,3} < \widehat{\mu}_{f,1} \leq \widehat{\mu}_{f,2} \text{ or } \widehat{\mu}_{f,3} > \widehat{\mu}_{f,1} \geq \widehat{\mu}_{f,2} \\ p_{f,2}/2 & \widehat{\mu}_{f,3} < \widehat{\mu}_{f,2} \leq \widehat{\mu}_{f,1} \text{ or } \widehat{\mu}_{f,3} > \widehat{\mu}_{f,2} \geq \widehat{\mu}_{f,1} \\ 1 & \widehat{\mu}_{f,1} \leq \widehat{\mu}_{f,3} \leq \widehat{\mu}_{f,2} \text{ or } \widehat{\mu}_{f,2} \leq \widehat{\mu}_{f,3} \leq \widehat{\mu}_{f,1} \end{cases}$$

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

I he problem

ormulated data

The contenders
edgeR
ShrinkBayes

The contest

Constructing ROC curves

ROC results

FDR control

- ▶ Built on inla (integrated nested Laplace approximation).
- empirical Bayes with a zero-inflated NB likelihood and normal priors.
- ▶ I reparameterize

$$\begin{split} \phi_f &= \frac{\mu_{f,1} + \mu_{f,2}}{2} \qquad \text{(parental mean)} \\ \alpha_f &= \frac{\mu_{f,2} - \mu_{f,1}}{2} \qquad \text{(half parental difference)} \\ \delta_f &= \mu_{f,3} - \frac{\mu_{f,1} + \mu_{f,2}}{2} \qquad \text{(hybrid effect)} \end{split}$$

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

The contenders

edgeR

ShrinkBayes

The contest
Constructing ROC curves
ROC results
FDR control

$$\alpha_f = \text{ half parental difference}$$
 $\delta_f = \text{ hybrid effect}$ 

Use contrasts to calculate final posterior probabilities of heterosis:

$$p_{f,\mathtt{ShrinkBayes}} = \begin{cases} 0 & \left| \widehat{\delta}_f \right| < \left| \widehat{\alpha}_f \right|, \, \mathtt{o.w.:} \\ P(\delta_f + \alpha_f > 0 \mid \mathtt{data}) & \widehat{\delta}_f > -\widehat{\alpha}_f \\ P(\delta_f - \alpha_f > 0 \mid \mathtt{data}) & \widehat{\delta}_f > \widehat{\alpha}_f \\ P(\delta_f - \alpha_f < 0 \mid \mathtt{data}) & \widehat{\delta}_f < \widehat{\alpha}_f \\ P(\delta_f + \alpha_f < 0 \mid \mathtt{data}) & \widehat{\delta}_f < -\widehat{\alpha}_f \end{cases}$$

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

. . .

The contenders

edgeR

ShrinkBayes

edgeR ShrinkBayes baySeq

The contest

Constructing ROC curves

ROC results

FDR control procedures

FDR control results

- ▶ Estimate main effects  $\mu_{f,t}$  using edgeR.
- Calculate the posterior probability that each feature satisfies:

Model	Constraint
$M_1$	All $\mu_{f,t}$ 's equal
$M_2$	$\mu_{f,1} = \mu_{f,2}$
$M_3$	$\mu_{f,1} = \mu_{f,3}$
$M_4$	$\mu_{f,2} = \mu_{f,3}$
$M_5$	All $\mu_{f,t}$ 's distinct

► Final posterior probabilities of heterosis:

$$p_{f, \text{baySeq}} = \begin{cases} 0 & \widehat{\mu}_{f, 1} \leq \widehat{\mu}_{f, 3} \leq \widehat{\mu}_{f, 2} \text{ or} \\ & \widehat{\mu}_{f, 2} \leq \widehat{\mu}_{f, 3} \leq \widehat{\mu}_{f, 1} \end{cases}$$

$$P(M_2 \mid \text{data}) + P(M_5 \mid \text{data}) & \text{otherwise}$$

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

i ne problem

The contenders

edgeR ShrinkBayes baySeq

The contest

Constructing ROC curves

ROC results

FDR control procedures

#### The contest

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The contest

# Calculating false positive rate (FPR) and true positive rate (TPR)

- ROC = Receiver Operating Characteristic
- N<sub>true</sub> heterosis features, N<sub>false</sub> null features.
- Results of testing each feature for heterosis
  - For edgeR, the ranks are p-values.
  - For the other methods, take p(no heterosis | data) for each feature.

rank	0.802	0.935	0.539	0.001		0.500	0.603
truth	0	0	1	1		1	0

Sort by rank.

rank	0.000	0.001	1	l	l .	1	l
truth	1	1	0	1		0	0

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

Constructing ROC

# Calculating false positive rate (FPR) and true positive rate (TPR)

▶ In practice, we would declare the lowest-rank features to have heterosis.

rank	0.000	0.001	0.005	0.006	 0.901	1.000
truth	1	1	0	1	 0	0

With 2 heterosis genes and 1 null gene, say,

$$FPR = \frac{1}{N_{false}}$$
  $TPR = \frac{2}{N_{true}}$ 

 Repeat for multiple cutoffs to get multiple (FPR, TPR) pairs.


The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric

edgeR

Constructing ROC

# Example ROC curves



The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

Simulated data

edgeR ShrinkBayes oaySeq


The contest

Constructing ROC curves

ROC results

FDR control

#### Areas under ROC curves



The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

....

Simulated data


The contenders
edgeR
ShrinkBayes

The contest

Constructing ROC

ROC results
FDR control
procedures

#### Areas under ROC curves



The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

ROC results

- Let  $p_{(f)}$  (f = 1, ..., 25000) be the ordered ranks
  - For edgeR, the ranks are now q-values.
  - For the other methods, take p(no heterosis | data) for each feature.
- ▶ Let  $I_{(f)} = I$ (feature of  $p_{(f)}$  is not heterosis)
- ► Method 1:

$$\overline{p}_m = \frac{1}{m} \sum_{f=1}^m p_{(f)} \qquad \overline{l}_m = \frac{1}{m} \sum_{f=1}^m l_{(f)}$$

- ▶ To control FDR, we need  $\overline{I}_m \leq \overline{p}_m$ .
- ▶ Plot  $\overline{I}_m \overline{p}_m$  vs  $\overline{p}_m$  for multiple m.
- ► Method 2:

$$\overline{p}_m = \frac{1}{100} \sum_{f=m}^{m+100} p_{(f)} \qquad \overline{I}_m = \frac{1}{100} \sum_{f=m}^{m+100} I_{(f)}$$

The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

Cianalana da Jan

The contenders
edgeR
ShrinkBayes


The contest

Constructing ROC curves

ROC results

FDR control procedures

# Method 1: stan\_laplace



The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

THE WOLKHOW


Simulated data

The contenders
edgeR
ShrinkBayes

The contest
Constructing ROC curves
ROC results
FDR control

FDR control results

# Method 2: stan\_laplace



The heterosis problem: a comparison of Eric's method with edgeR, baySeq, and ShrinkBayes

Will Landau, Eric Mittman

The problem

THE WORKHOW

Simulated data

The contenders
edgeR
ShrinkBayes
baySeq

The contest
Constructing ROC curves
ROC results
FDR control procedures