


# Lecture 05: Probability and Statistical Inference

Bill Perry


## Lecture 4: Review

- Introduction to histograms or frequency distributions
- Probability Distribution Functions (PDF)
- Descriptive Statistics
  - Center - mean, median, mode
  - Spread - range, variance, standard deviation



## Lecture 4: Review - Statistical Concepts

- Introduction to histograms or frequency distributions
- Probability Distribution Functions (PDF)
- Descriptive Statistics
  - Center - mean, median, mode
  - Spread - range, variance, standard deviation



## Lecture 4: Review - Summary Statistics


- Introduction to histograms or frequency distributions
- Probability Distribution Functions (PDF)
- Descriptive Statistics
  - Center - mean, median, mode
  - Spread - range, variance, standard deviation

| lake | mean_length | sd_length | se_length | count |
|------|-------------|-----------|-----------|-------|
| I3   | 265.6061    | 28.30378  | 3.483954  | 66    |
| I8   | 362.5980    | 52.33901  | 5.182334  | 102   |

## Lecture 5: Probability and Statistical Inference

The goals for today

- Statistical inference fundamentals
- Hypothesis testing principles
- T Distributions
- One sample T Tests
- Two sample T Test



## Lecture 5: Confidence intervals

In the more typical case DON'T know the population  $\sigma$  or standard deviation

- estimate it from the samples
- and when sample size is  $< \sim 30$ )
- can't use the standard normal (z) distribution

*Instead, we use Student's t distribution*




## Lecture 5: Understanding t-distribution

When sample sizes are small, the **t-distribution** is more appropriate than the normal distribution.

- Similar to normal distribution but with heavier tails
- Shape depends on **degrees of freedom** ( $df = n-1$ )
- With large  $df$  ( $> 30$ ), approaches the normal distribution
- Used for:

- Small sample sizes
- When population standard deviation is unknown
- Calculating confidence intervals
- Conducting t-tests

### Comparison of Normal and t-distributions



## Lecture 5: t-distribution Properties

When sample sizes are small, the **t-distribution** is more appropriate than the normal distribution.

- Similar to normal distribution (1.96 = 2.5% tails) but with heavier tails
- Shape depends on **degrees of freedom** ( $df = n-1$ )
- With large  $df$  ( $>30$ ), approaches the normal distribution
- Used for:
  - Small sample sizes
  - When population standard deviation is unknown
  - Calculating confidence intervals
  - Conducting t-tests

| df   | 0.000 | 1.000 | 1.376 | 1.963 | 3.078 | 6.314 | 12.71 | 31.82 | 63.66 | 318.31 | 636.62 |
|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|--------|
| 1    | 0.000 | 0.816 | 1.061 | 1.386 | 1.886 | 2.920 | 4.303 | 6.965 | 9.925 | 22.327 | 31.599 |
| 2    | 0.000 | 0.765 | 0.978 | 1.250 | 1.638 | 2.353 | 3.182 | 4.541 | 5.841 | 10.215 | 12.924 |
| 3    | 0.000 | 0.741 | 0.941 | 1.190 | 1.533 | 2.132 | 2.776 | 3.747 | 4.604 | 7.173  | 8.610  |
| 4    | 0.000 | 0.727 | 0.920 | 1.156 | 1.476 | 2.015 | 2.571 | 3.365 | 4.032 | 5.893  | 6.869  |
| 5    | 0.000 | 0.718 | 0.906 | 1.134 | 1.440 | 1.943 | 2.447 | 3.143 | 3.707 | 5.208  | 5.959  |
| 6    | 0.000 | 0.711 | 0.896 | 1.119 | 1.415 | 1.895 | 2.365 | 2.998 | 3.499 | 4.785  | 5.408  |
| 7    | 0.000 | 0.706 | 0.889 | 1.108 | 1.397 | 1.860 | 2.306 | 2.896 | 3.355 | 4.501  | 5.041  |
| 8    | 0.000 | 0.703 | 0.883 | 1.100 | 1.383 | 1.833 | 2.262 | 2.821 | 3.250 | 4.297  | 4.781  |
| 9    | 0.000 | 0.700 | 0.879 | 1.093 | 1.372 | 1.812 | 2.228 | 2.784 | 3.169 | 4.144  | 4.587  |
| 10   | 0.000 | 0.697 | 0.876 | 1.088 | 1.363 | 1.796 | 2.201 | 2.718 | 3.106 | 4.025  | 4.437  |
| 11   | 0.000 | 0.695 | 0.873 | 1.083 | 1.356 | 1.782 | 2.179 | 2.681 | 3.055 | 3.930  | 4.318  |
| 12   | 0.000 | 0.694 | 0.870 | 1.079 | 1.350 | 1.771 | 2.160 | 2.650 | 3.012 | 3.852  | 4.221  |
| 13   | 0.000 | 0.692 | 0.868 | 1.076 | 1.345 | 1.761 | 2.145 | 2.624 | 2.977 | 3.787  | 4.140  |
| 14   | 0.000 | 0.691 | 0.866 | 1.074 | 1.341 | 1.753 | 2.131 | 2.602 | 2.947 | 3.733  | 4.073  |
| 15   | 0.000 | 0.690 | 0.865 | 1.071 | 1.337 | 1.746 | 2.120 | 2.583 | 2.921 | 3.686  | 4.015  |
| 16   | 0.000 | 0.689 | 0.863 | 1.069 | 1.333 | 1.740 | 2.110 | 2.567 | 2.898 | 3.646  | 3.965  |
| 17   | 0.000 | 0.688 | 0.862 | 1.067 | 1.330 | 1.734 | 2.101 | 2.552 | 2.878 | 3.610  | 3.922  |
| 18   | 0.000 | 0.688 | 0.861 | 1.066 | 1.328 | 1.729 | 2.093 | 2.539 | 2.861 | 3.579  | 3.883  |
| 19   | 0.000 | 0.687 | 0.860 | 1.064 | 1.325 | 1.725 | 2.086 | 2.528 | 2.845 | 3.552  | 3.850  |
| 20   | 0.000 | 0.686 | 0.859 | 1.063 | 1.323 | 1.721 | 2.080 | 2.518 | 2.831 | 3.527  | 3.819  |
| 21   | 0.000 | 0.686 | 0.858 | 1.061 | 1.321 | 1.717 | 2.074 | 2.508 | 2.819 | 3.505  | 3.792  |
| 22   | 0.000 | 0.685 | 0.858 | 1.060 | 1.319 | 1.714 | 2.069 | 2.500 | 2.807 | 3.485  | 3.768  |
| 23   | 0.000 | 0.685 | 0.857 | 1.059 | 1.318 | 1.711 | 2.064 | 2.492 | 2.797 | 3.467  | 3.745  |
| 24   | 0.000 | 0.684 | 0.856 | 1.058 | 1.316 | 1.708 | 2.060 | 2.485 | 2.787 | 3.450  | 3.725  |
| 25   | 0.000 | 0.684 | 0.856 | 1.058 | 1.315 | 1.706 | 2.056 | 2.479 | 2.779 | 3.435  | 3.707  |
| 26   | 0.000 | 0.684 | 0.855 | 1.057 | 1.314 | 1.703 | 2.052 | 2.473 | 2.771 | 3.421  | 3.690  |
| 27   | 0.000 | 0.683 | 0.855 | 1.056 | 1.313 | 1.701 | 2.048 | 2.467 | 2.763 | 3.408  | 3.674  |
| 28   | 0.000 | 0.683 | 0.854 | 1.055 | 1.311 | 1.699 | 2.045 | 2.462 | 2.756 | 3.396  | 3.659  |
| 29   | 0.000 | 0.683 | 0.854 | 1.055 | 1.310 | 1.697 | 2.042 | 2.457 | 2.750 | 3.385  | 3.646  |
| 30   | 0.000 | 0.681 | 0.851 | 1.050 | 1.303 | 1.684 | 2.021 | 2.423 | 2.704 | 3.307  | 3.551  |
| 40   | 0.000 | 0.679 | 0.848 | 1.045 | 1.296 | 1.671 | 2.000 | 2.390 | 2.660 | 3.232  | 3.460  |
| 60   | 0.000 | 0.678 | 0.846 | 1.043 | 1.292 | 1.664 | 1.990 | 2.374 | 2.639 | 3.195  | 3.416  |
| 80   | 0.000 | 0.677 | 0.845 | 1.042 | 1.290 | 1.660 | 1.984 | 2.364 | 2.626 | 3.174  | 3.390  |
| 100  | 0.000 | 0.675 | 0.842 | 1.037 | 1.282 | 1.646 | 1.962 | 2.330 | 2.581 | 3.098  | 3.300  |
| 1000 | 0.000 | 0.674 | 0.842 | 1.036 | 1.282 | 1.645 | 1.960 | 2.326 | 2.576 | 3.090  | 3.291  |

## Practice Exercise 4: Using the t-distribution

## 🔗 Practice Exercise 4: Using the t-distribution

Let's compare confidence intervals using the normal approximation (z) versus the t-distribution for our fish data.

```
# Calculate CI using both z and t distributions for a smaller subset
small_sample <- grayling_df %>%
  filter(lake == "I3") %>%
  slice_sample(n = 10)

# Calculate statistics
sample_mean <- mean(small_sample$length_mm)
sample_sd <- sd(small_sample$length_mm)
sample_n <- nrow(small_sample)
sample_se <- sample_sd / sqrt(sample_n)

# Calculate confidence intervals
z_ci_lower <- sample_mean - 1.96 * sample_se
z_ci_upper <- sample_mean + 1.96 * sample_se

# For t-distribution, get critical value for 95% CI with df = n-1
t_crit <- qt(0.975, df = sample_n - 1)
t_ci_lower <- sample_mean - t_crit * sample_se
t_ci_upper <- sample_mean + t_crit * sample_se

# Display results
cat("Mean:", round(sample_mean, 1), "mm\n")
```

Mean: 252.8 mm

```
cat("Standard deviation:", round(sample_sd, 2), "mm\n")
```

Standard deviation: 31.59 mm

```
cat("Standard error:", round(sample_se, 2), "mm\n")
```

Standard error: 9.99 mm

```
cat("95% CI using z:", round(z_ci_lower, 1), "to", round(z_ci_upper, 1), "mm\n")
```

95% CI using z: 233.2 to 272.4 mm

```
cat("95% CI using t:", round(t_ci_lower, 1), "to", round(t_ci_upper, 1), "mm\n")
```

95% CI using t: 230.2 to 275.4 mm

```
cat("t critical value:", round(t_crit, 3), "vs z critical value: 1.96\n")
```

t critical value: 2.262 vs z critical value: 1.96

# Student's t-distribution Formula

To calculate CI for sample from “unknown” population:

$$CI = \bar{y} \pm t \cdot \frac{s}{\sqrt{n}}$$

Where:

- $\bar{y}$  is sample mean
- $n$  is sample size
- $s$  is sample standard deviation
- $t$  t-value corresponding the probability of the CI
- $t$  in t-table for different degrees of freedom (n-1)

| two-tails | 1.00  | 0.50  | 0.40  | 0.30  | 0.20  | 0.10  | 0.05  | 0.02  | 0.01  | 0.002  | 0.001  |
|-----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|--------|
| df        |       |       |       |       |       |       |       |       |       |        |        |
| 1         | 0.000 | 1.000 | 1.376 | 1.963 | 3.078 | 6.314 | 12.71 | 31.82 | 63.66 | 318.31 | 636.62 |
| 2         | 0.000 | 0.816 | 1.061 | 1.386 | 1.886 | 2.920 | 4.303 | 6.965 | 9.925 | 22.327 | 31.599 |
| 3         | 0.000 | 0.765 | 0.978 | 1.250 | 1.638 | 2.353 | 3.182 | 4.541 | 5.841 | 10.215 | 12.924 |
| 4         | 0.000 | 0.741 | 0.941 | 1.190 | 1.533 | 2.132 | 2.776 | 3.747 | 4.604 | 7.173  | 8.610  |
| 5         | 0.000 | 0.727 | 0.920 | 1.156 | 1.476 | 2.015 | 2.571 | 3.365 | 4.032 | 5.893  | 6.869  |
| 6         | 0.000 | 0.718 | 0.906 | 1.134 | 1.440 | 1.943 | 2.447 | 3.143 | 3.707 | 5.208  | 5.959  |
| 7         | 0.000 | 0.711 | 0.896 | 1.119 | 1.415 | 1.895 | 2.365 | 2.998 | 3.499 | 4.785  | 5.408  |
| 8         | 0.000 | 0.706 | 0.889 | 1.108 | 1.397 | 1.860 | 2.306 | 2.896 | 3.355 | 4.501  | 5.041  |
| 9         | 0.000 | 0.703 | 0.883 | 1.100 | 1.383 | 1.833 | 2.262 | 2.821 | 3.250 | 4.297  | 4.781  |
| 10        | 0.000 | 0.700 | 0.879 | 1.093 | 1.372 | 1.812 | 2.228 | 2.764 | 3.169 | 4.144  | 4.587  |
| 11        | 0.000 | 0.697 | 0.876 | 1.088 | 1.363 | 1.796 | 2.201 | 2.718 | 3.106 | 4.025  | 4.437  |
| 12        | 0.000 | 0.695 | 0.873 | 1.083 | 1.356 | 1.782 | 2.179 | 2.681 | 3.055 | 3.930  | 4.318  |
| 13        | 0.000 | 0.694 | 0.870 | 1.079 | 1.350 | 1.771 | 2.160 | 2.650 | 3.012 | 3.852  | 4.221  |
| 14        | 0.000 | 0.692 | 0.868 | 1.076 | 1.345 | 1.761 | 2.145 | 2.624 | 2.977 | 3.787  | 4.140  |
| 15        | 0.000 | 0.691 | 0.866 | 1.074 | 1.341 | 1.753 | 2.131 | 2.602 | 2.947 | 3.733  | 4.073  |
| 16        | 0.000 | 0.690 | 0.865 | 1.071 | 1.337 | 1.746 | 2.120 | 2.583 | 2.921 | 3.686  | 4.015  |
| 17        | 0.000 | 0.689 | 0.863 | 1.069 | 1.333 | 1.740 | 2.110 | 2.567 | 2.898 | 3.646  | 3.965  |
| 18        | 0.000 | 0.688 | 0.862 | 1.067 | 1.330 | 1.734 | 2.101 | 2.552 | 2.878 | 3.610  | 3.922  |
| 19        | 0.000 | 0.688 | 0.861 | 1.066 | 1.328 | 1.729 | 2.093 | 2.539 | 2.861 | 3.579  | 3.883  |
| 20        | 0.000 | 0.687 | 0.860 | 1.064 | 1.325 | 1.725 | 2.086 | 2.528 | 2.845 | 3.552  | 3.850  |
| 21        | 0.000 | 0.686 | 0.859 | 1.063 | 1.323 | 1.721 | 2.080 | 2.518 | 2.831 | 3.527  | 3.819  |
| 22        | 0.000 | 0.686 | 0.858 | 1.061 | 1.321 | 1.717 | 2.074 | 2.508 | 2.819 | 3.505  | 3.792  |
| 23        | 0.000 | 0.685 | 0.858 | 1.060 | 1.319 | 1.714 | 2.069 | 2.500 | 2.807 | 3.485  | 3.768  |
| 24        | 0.000 | 0.685 | 0.857 | 1.059 | 1.318 | 1.711 | 2.064 | 2.492 | 2.797 | 3.467  | 3.745  |
| 25        | 0.000 | 0.684 | 0.856 | 1.058 | 1.316 | 1.708 | 2.060 | 2.485 | 2.787 | 3.450  | 3.725  |
| 26        | 0.000 | 0.684 | 0.856 | 1.058 | 1.315 | 1.706 | 2.056 | 2.479 | 2.779 | 3.435  | 3.707  |
| 27        | 0.000 | 0.684 | 0.855 | 1.057 | 1.314 | 1.703 | 2.052 | 2.473 | 2.771 | 3.421  | 3.690  |
| 28        | 0.000 | 0.683 | 0.855 | 1.056 | 1.313 | 1.701 | 2.048 | 2.467 | 2.763 | 3.408  | 3.674  |
| 29        | 0.000 | 0.683 | 0.854 | 1.055 | 1.311 | 1.699 | 2.045 | 2.462 | 2.756 | 3.396  | 3.659  |
| 30        | 0.000 | 0.683 | 0.854 | 1.055 | 1.310 | 1.697 | 2.042 | 2.457 | 2.750 | 3.385  | 3.646  |
| 40        | 0.000 | 0.681 | 0.851 | 1.050 | 1.303 | 1.684 | 2.021 | 2.423 | 2.704 | 3.307  | 3.551  |
| 60        | 0.000 | 0.679 | 0.848 | 1.045 | 1.296 | 1.671 | 2.000 | 2.390 | 2.660 | 3.232  | 3.460  |
| 80        | 0.000 | 0.678 | 0.846 | 1.043 | 1.292 | 1.664 | 1.990 | 2.374 | 2.639 | 3.195  | 3.416  |
| 100       | 0.000 | 0.677 | 0.845 | 1.042 | 1.290 | 1.660 | 1.984 | 2.364 | 2.626 | 3.174  | 3.390  |
| 1000      | 0.000 | 0.675 | 0.842 | 1.037 | 1.282 | 1.646 | 1.962 | 2.330 | 2.581 | 3.098  | 3.300  |
| Z         | 0.000 | 0.674 | 0.842 | 1.036 | 1.282 | 1.645 | 1.960 | 2.326 | 2.576 | 3.090  | 3.291  |

## Lecture 5: Student's t-distribution Table

Here is a t-table


- Values of  $t$  that correspond to probabilities
- Probabilities listed along top
- Sample  $df$ s are listed in the left-most column
- Probabilities are given for one-tailed and two-tailed “questions”

| two-tails | 1.00  | 0.50  | 0.40  | 0.30  | 0.20  | 0.10  | 0.05  | 0.02  | 0.01  | 0.002  | 0.001  |
|-----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|--------|
| df        |       |       |       |       |       |       |       |       |       |        |        |
| 1         | 0.000 | 1.000 | 1.376 | 1.963 | 3.078 | 6.314 | 12.71 | 31.82 | 63.66 | 318.31 | 636.62 |
| 2         | 0.000 | 0.816 | 1.061 | 1.386 | 1.886 | 2.920 | 4.303 | 6.965 | 9.925 | 22.327 | 31.599 |
| 3         | 0.000 | 0.765 | 0.978 | 1.250 | 1.638 | 2.353 | 3.182 | 4.541 | 5.841 | 10.215 | 12.924 |
| 4         | 0.000 | 0.741 | 0.941 | 1.190 | 1.533 | 2.132 | 2.776 | 3.747 | 4.604 | 7.173  | 8.610  |
| 5         | 0.000 | 0.727 | 0.920 | 1.156 | 1.476 | 2.015 | 2.571 | 3.365 | 4.032 | 5.893  | 6.869  |
| 6         | 0.000 | 0.718 | 0.906 | 1.134 | 1.440 | 1.943 | 2.447 | 3.143 | 3.707 | 5.208  | 5.959  |
| 7         | 0.000 | 0.711 | 0.896 | 1.119 | 1.415 | 1.895 | 2.365 | 2.998 | 3.499 | 4.785  | 5.408  |
| 8         | 0.000 | 0.706 | 0.889 | 1.108 | 1.397 | 1.860 | 2.306 | 2.896 | 3.355 | 4.501  | 5.041  |
| 9         | 0.000 | 0.703 | 0.883 | 1.100 | 1.383 | 1.833 | 2.262 | 2.821 | 3.250 | 4.297  | 4.781  |
| 10        | 0.000 | 0.700 | 0.879 | 1.093 | 1.372 | 1.812 | 2.228 | 2.764 | 3.169 | 4.144  | 4.587  |
| 11        | 0.000 | 0.697 | 0.876 | 1.088 | 1.363 | 1.796 | 2.201 | 2.718 | 3.106 | 4.025  | 4.437  |
| 12        | 0.000 | 0.695 | 0.873 | 1.083 | 1.356 | 1.782 | 2.179 | 2.681 | 3.055 | 3.930  | 4.318  |
| 13        | 0.000 | 0.694 | 0.870 | 1.079 | 1.350 | 1.771 | 2.160 | 2.650 | 3.012 | 3.852  | 4.221  |
| 14        | 0.000 | 0.692 | 0.868 | 1.076 | 1.345 | 1.761 | 2.145 | 2.624 | 2.977 | 3.787  | 4.140  |
| 15        | 0.000 | 0.691 | 0.866 | 1.074 | 1.341 | 1.753 | 2.131 | 2.602 | 2.947 | 3.733  | 4.073  |
| 16        | 0.000 | 0.690 | 0.865 | 1.071 | 1.337 | 1.746 | 2.120 | 2.583 | 2.921 | 3.686  | 4.015  |
| 17        | 0.000 | 0.689 | 0.863 | 1.069 | 1.333 | 1.740 | 2.110 | 2.567 | 2.898 | 3.646  | 3.965  |
| 18        | 0.000 | 0.688 | 0.862 | 1.067 | 1.330 | 1.734 | 2.101 | 2.552 | 2.878 | 3.610  | 3.922  |
| 19        | 0.000 | 0.688 | 0.861 | 1.066 | 1.328 | 1.729 | 2.093 | 2.539 | 2.861 | 3.579  | 3.883  |
| 20        | 0.000 | 0.687 | 0.860 | 1.064 | 1.325 | 1.725 | 2.086 | 2.528 | 2.845 | 3.552  | 3.850  |
| 21        | 0.000 | 0.686 | 0.859 | 1.063 | 1.323 | 1.721 | 2.080 | 2.518 | 2.831 | 3.527  | 3.819  |
| 22        | 0.000 | 0.686 | 0.858 | 1.061 | 1.321 | 1.717 | 2.074 | 2.508 | 2.819 | 3.505  | 3.792  |
| 23        | 0.000 | 0.685 | 0.858 | 1.060 | 1.319 | 1.714 | 2.069 | 2.500 | 2.807 | 3.485  | 3.768  |
| 24        | 0.000 | 0.685 | 0.857 | 1.059 | 1.318 | 1.711 | 2.064 | 2.492 | 2.797 | 3.467  | 3.745  |
| 25        | 0.000 | 0.684 | 0.856 | 1.058 | 1.316 | 1.708 | 2.060 | 2.485 | 2.787 | 3.450  | 3.725  |
| 26        | 0.000 | 0.684 | 0.856 | 1.058 | 1.315 | 1.706 | 2.056 | 2.479 | 2.779 | 3.435  | 3.707  |
| 27        | 0.000 | 0.684 | 0.855 | 1.057 | 1.314 | 1.703 | 2.052 | 2.473 | 2.771 | 3.421  | 3.690  |
| 28        | 0.000 | 0.683 | 0.855 | 1.056 | 1.313 | 1.701 | 2.048 | 2.467 | 2.763 | 3.408  | 3.674  |
| 29        | 0.000 | 0.683 | 0.854 | 1.055 | 1.311 | 1.699 | 2.045 | 2.462 | 2.756 | 3.396  | 3.659  |
| 30        | 0.000 | 0.683 | 0.854 | 1.055 | 1.310 | 1.697 | 2.042 | 2.457 | 2.750 | 3.385  | 3.646  |
| 40        | 0.000 | 0.681 | 0.851 | 1.050 | 1.303 | 1.684 | 2.021 | 2.423 | 2.704 | 3.307  | 3.551  |
| 60        | 0.000 | 0.679 | 0.848 | 1.045 | 1.296 | 1.671 | 2.000 | 2.390 | 2.660 | 3.232  | 3.460  |
| 80        | 0.000 | 0.678 | 0.846 | 1.043 | 1.292 | 1.664 | 1.990 | 2.374 | 2.639 | 3.195  | 3.416  |
| 100       | 0.000 | 0.677 | 0.845 | 1.042 | 1.290 | 1.660 | 1.984 | 2.364 | 2.626 | 3.174  | 3.390  |
| 1000      | 0.000 | 0.675 | 0.842 | 1.037 | 1.282 | 1.646 | 1.962 | 2.330 | 2.581 | 3.098  | 3.300  |
| Z         | 0.000 | 0.674 | 0.842 | 1.036 | 1.282 | 1.645 | 1.960 | 2.326 | 2.576 | 3.090  | 3.291  |

## Lecture 5: One-tailed Questions

One-tailed questions: area of distribution left or (right) of a certain value

- $n=20$  ( $df=19$ ) - 90% of the observations found left
- $t = 1.328$  (10% are outside)




|    |       |       |       |       |       |       |       |       |       |       |       |
|----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 4  | 0.000 | 0.741 | 0.941 | 1.190 | 1.533 | 2.132 | 2.776 | 3.747 | 4.604 | 7.173 | 8.610 |
| 5  | 0.000 | 0.727 | 0.920 | 1.156 | 1.476 | 2.015 | 2.571 | 3.365 | 4.032 | 5.893 | 6.869 |
| 6  | 0.000 | 0.718 | 0.906 | 1.134 | 1.440 | 1.943 | 2.447 | 3.143 | 3.707 | 5.208 | 5.959 |
| 7  | 0.000 | 0.711 | 0.896 | 1.119 | 1.415 | 1.895 | 2.365 | 2.998 | 3.499 | 4.785 | 5.408 |
| 8  | 0.000 | 0.706 | 0.889 | 1.108 | 1.397 | 1.860 | 2.306 | 2.896 | 3.355 | 4.501 | 5.041 |
| 9  | 0.000 | 0.703 | 0.883 | 1.100 | 1.383 | 1.833 | 2.262 | 2.821 | 3.250 | 4.297 | 4.781 |
| 10 | 0.000 | 0.700 | 0.879 | 1.093 | 1.372 | 1.812 | 2.228 | 2.764 | 3.169 | 4.144 | 4.587 |
| 11 | 0.000 | 0.697 | 0.876 | 1.088 | 1.363 | 1.796 | 2.201 | 2.718 | 3.106 | 4.025 | 4.437 |
| 12 | 0.000 | 0.695 | 0.873 | 1.083 | 1.356 | 1.782 | 2.179 | 2.681 | 3.055 | 3.930 | 4.318 |
| 13 | 0.000 | 0.694 | 0.870 | 1.079 | 1.350 | 1.771 | 2.160 | 2.650 | 3.012 | 3.852 | 4.221 |
| 14 | 0.000 | 0.692 | 0.868 | 1.076 | 1.345 | 1.761 | 2.145 | 2.624 | 2.977 | 3.787 | 4.140 |
| 15 | 0.000 | 0.691 | 0.866 | 1.074 | 1.341 | 1.753 | 2.131 | 2.602 | 2.947 | 3.733 | 4.073 |
| 16 | 0.000 | 0.690 | 0.865 | 1.071 | 1.337 | 1.746 | 2.120 | 2.583 | 2.921 | 3.686 | 4.015 |
| 17 | 0.000 | 0.689 | 0.863 | 1.069 | 1.333 | 1.740 | 2.110 | 2.567 | 2.898 | 3.646 | 3.965 |
| 18 | 0.000 | 0.688 | 0.862 | 1.067 | 1.330 | 1.734 | 2.101 | 2.552 | 2.878 | 3.610 | 3.922 |
| 19 | 0.000 | 0.688 | 0.861 | 1.066 | 1.328 | 1.729 | 2.093 | 2.539 | 2.861 | 3.579 | 3.883 |
| 20 | 0.000 | 0.687 | 0.860 | 1.064 | 1.325 | 1.725 | 2.086 | 2.528 | 2.845 | 3.552 | 3.850 |
| 21 | 0.000 | 0.686 | 0.859 | 1.063 | 1.323 | 1.721 | 2.080 | 2.518 | 2.831 | 3.527 | 3.819 |
| 22 | 0.000 | 0.686 | 0.858 | 1.061 | 1.321 | 1.717 | 2.074 | 2.508 | 2.819 | 3.505 | 3.792 |
| 23 | 0.000 | 0.685 | 0.858 | 1.060 | 1.319 | 1.714 | 2.069 | 2.500 | 2.807 | 3.485 | 3.768 |
| 24 | 0.000 | 0.685 | 0.857 | 1.059 | 1.318 | 1.711 | 2.064 | 2.492 | 2.797 | 3.467 | 3.745 |
| 25 | 0.000 | 0.684 | 0.856 | 1.058 | 1.316 | 1.708 | 2.060 | 2.485 | 2.787 | 3.450 | 3.725 |
| 26 | 0.000 | 0.684 | 0.856 | 1.058 | 1.315 | 1.706 | 2.056 | 2.479 | 2.779 | 3.435 | 3.707 |
| 27 | 0.000 | 0.684 | 0.855 | 1.057 | 1.314 | 1.703 | 2.052 | 2.473 | 2.771 | 3.421 | 3.690 |
| 28 | 0.000 | 0.683 | 0.855 | 1.056 | 1.313 | 1.701 | 2.048 | 2.467 | 2.763 | 3.408 | 3.674 |
| 29 | 0.000 | 0.683 | 0.854 | 1.055 | 1.311 | 1.699 | 2.045 | 2.462 | 2.756 | 3.396 | 3.659 |
| 30 | 0.000 | 0.683 | 0.854 | 1.055 | 1.310 | 1.697 | 2.042 | 2.457 | 2.750 | 3.385 | 3.646 |
| 40 | 0.000 | 0.681 | 0.851 | 1.050 | 1.303 | 1.684 | 2.021 | 2.423 | 2.704 | 3.307 | 3.551 |
| 60 | 0.000 | 0.679 | 0.848 | 1.045 | 1.296 | 1.671 | 2.000 | 2.390 | 2.660 | 3.232 | 3.460 |

## Lecture 5: Two-tailed Questions

Two-tailed questions refer to area between certain values

- $n = 20$  ( $df=19$ ), 90% of the observations are between
- $t=-1.729$  and  $t=1.729$  (10% are outside)



|    |       |       |       |       |       |       |       |       |       |       |       |
|----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 5  | 0.000 | 0.727 | 0.920 | 1.156 | 1.476 | 2.015 | 2.571 | 3.365 | 4.032 | 5.893 | 6.869 |
| 6  | 0.000 | 0.718 | 0.906 | 1.134 | 1.440 | 1.943 | 2.447 | 3.143 | 3.707 | 5.208 | 5.959 |
| 7  | 0.000 | 0.711 | 0.896 | 1.119 | 1.415 | 1.895 | 2.365 | 2.998 | 3.499 | 4.785 | 5.408 |
| 8  | 0.000 | 0.706 | 0.889 | 1.108 | 1.397 | 1.860 | 2.306 | 2.896 | 3.355 | 4.501 | 5.041 |
| 9  | 0.000 | 0.703 | 0.883 | 1.100 | 1.383 | 1.833 | 2.262 | 2.821 | 3.250 | 4.297 | 4.781 |
| 10 | 0.000 | 0.700 | 0.879 | 1.093 | 1.372 | 1.812 | 2.228 | 2.764 | 3.169 | 4.144 | 4.587 |
| 11 | 0.000 | 0.697 | 0.876 | 1.088 | 1.363 | 1.796 | 2.201 | 2.718 | 3.106 | 4.025 | 4.437 |
| 12 | 0.000 | 0.695 | 0.873 | 1.083 | 1.356 | 1.782 | 2.179 | 2.681 | 3.055 | 3.930 | 4.318 |
| 13 | 0.000 | 0.694 | 0.870 | 1.079 | 1.350 | 1.771 | 2.160 | 2.650 | 3.012 | 3.852 | 4.221 |
| 14 | 0.000 | 0.692 | 0.868 | 1.076 | 1.345 | 1.761 | 2.145 | 2.624 | 2.977 | 3.787 | 4.140 |
| 15 | 0.000 | 0.691 | 0.866 | 1.074 | 1.341 | 1.753 | 2.131 | 2.602 | 2.947 | 3.733 | 4.073 |
| 16 | 0.000 | 0.690 | 0.865 | 1.071 | 1.337 | 1.746 | 2.120 | 2.583 | 2.921 | 3.686 | 4.015 |
| 17 | 0.000 | 0.689 | 0.863 | 1.069 | 1.333 | 1.740 | 2.110 | 2.567 | 2.898 | 3.646 | 3.965 |
| 18 | 0.000 | 0.688 | 0.862 | 1.067 | 1.330 | 1.734 | 2.101 | 2.552 | 2.878 | 3.610 | 3.922 |
| 19 | 0.000 | 0.688 | 0.861 | 1.066 | 1.328 | 1.729 | 2.093 | 2.539 | 2.861 | 3.579 | 3.883 |
| 20 | 0.000 | 0.687 | 0.860 | 1.064 | 1.325 | 1.725 | 2.086 | 2.528 | 2.845 | 3.552 | 3.850 |
| 21 | 0.000 | 0.686 | 0.859 | 1.063 | 1.323 | 1.721 | 2.080 | 2.518 | 2.831 | 3.527 | 3.819 |
| 22 | 0.000 | 0.686 | 0.858 | 1.061 | 1.321 | 1.717 | 2.074 | 2.508 | 2.819 | 3.505 | 3.792 |
| 23 | 0.000 | 0.685 | 0.858 | 1.060 | 1.319 | 1.714 | 2.069 | 2.500 | 2.807 | 3.485 | 3.768 |
| 24 | 0.000 | 0.685 | 0.857 | 1.059 | 1.318 | 1.711 | 2.064 | 2.492 | 2.797 | 3.467 | 3.745 |
| 25 | 0.000 | 0.684 | 0.856 | 1.058 | 1.316 | 1.708 | 2.060 | 2.485 | 2.787 | 3.450 | 3.725 |
| 26 | 0.000 | 0.684 | 0.856 | 1.058 | 1.315 | 1.706 | 2.056 | 2.479 | 2.779 | 3.435 | 3.707 |
| 27 | 0.000 | 0.684 | 0.855 | 1.057 | 1.314 | 1.703 | 2.052 | 2.473 | 2.771 | 3.421 | 3.690 |
| 28 | 0.000 | 0.683 | 0.855 | 1.056 | 1.313 | 1.701 | 2.048 | 2.467 | 2.763 | 3.408 | 3.674 |
| 29 | 0.000 | 0.683 | 0.854 | 1.055 | 1.311 | 1.699 | 2.045 | 2.462 | 2.756 | 3.396 | 3.659 |
| 30 | 0.000 | 0.683 | 0.854 | 1.055 | 1.310 | 1.697 | 2.042 | 2.457 | 2.750 | 3.385 | 3.646 |
| 40 | 0.000 | 0.681 | 0.851 | 1.050 | 1.303 | 1.684 | 2.021 | 2.423 | 2.704 | 3.307 | 3.551 |
| 60 | 0.000 | 0.679 | 0.848 | 1.045 | 1.296 | 1.671 | 2.000 | 2.390 | 2.660 | 3.232 | 3.460 |

## Lecture 5: Calculating CI Example

Let's calculate CIs again:

Use two-sided test



- $CI = \bar{y} \pm t \cdot \frac{s}{\sqrt{n}}$
- 95% CI Sample A: =  $272.8 \pm 2.262 \cdot (37.81/(9^{.5})) = 1.650788$
- The 95% CI is between 244.3 and 301.3
- “The 95% CI for the population mean from sample A is  $272.8 \pm 28.5$ ”

| two-tails | 1.00  | 0.50  | 0.40  | 0.30  | 0.20  | 0.10  | 0.05  | 0.02  | 0.01  | 0.002  | 0.001  |
|-----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|--------|
| df        |       |       |       |       |       |       |       |       |       |        |        |
| 1         | 0.000 | 1.000 | 1.376 | 1.963 | 3.078 | 6.314 | 12.71 | 31.82 | 63.66 | 318.31 | 636.62 |
| 2         | 0.000 | 0.816 | 1.061 | 1.386 | 1.886 | 2.920 | 4.303 | 6.965 | 9.925 | 22.327 | 31.599 |
| 3         | 0.000 | 0.765 | 0.978 | 1.250 | 1.638 | 2.353 | 3.182 | 4.541 | 5.841 | 10.215 | 12.924 |
| 4         | 0.000 | 0.741 | 0.941 | 1.190 | 1.533 | 2.132 | 2.776 | 3.747 | 4.604 | 7.173  | 8.610  |
| 5         | 0.000 | 0.727 | 0.920 | 1.156 | 1.476 | 2.015 | 2.571 | 3.365 | 4.032 | 5.893  | 6.869  |
| 6         | 0.000 | 0.718 | 0.906 | 1.134 | 1.440 | 1.943 | 2.447 | 3.143 | 3.707 | 5.208  | 5.959  |
| 7         | 0.000 | 0.711 | 0.896 | 1.119 | 1.415 | 1.895 | 2.365 | 2.998 | 3.499 | 4.785  | 5.408  |
| 8         | 0.000 | 0.706 | 0.889 | 1.108 | 1.397 | 1.860 | 2.306 | 2.896 | 3.355 | 4.501  | 5.041  |
| 9         | 0.000 | 0.703 | 0.883 | 1.100 | 1.383 | 1.833 | 2.262 | 2.821 | 3.250 | 4.297  | 4.781  |
| 10        | 0.000 | 0.700 | 0.879 | 1.093 | 1.372 | 1.812 | 2.228 | 2.764 | 3.169 | 4.144  | 4.587  |
| 11        | 0.000 | 0.697 | 0.876 | 1.088 | 1.363 | 1.796 | 2.201 | 2.718 | 3.106 | 4.025  | 4.437  |
| 12        | 0.000 | 0.695 | 0.873 | 1.083 | 1.356 | 1.782 | 2.179 | 2.681 | 3.055 | 3.930  | 4.318  |
| 13        | 0.000 | 0.694 | 0.870 | 1.079 | 1.350 | 1.771 | 2.160 | 2.650 | 3.012 | 3.852  | 4.221  |
| 14        | 0.000 | 0.692 | 0.868 | 1.076 | 1.345 | 1.761 | 2.145 | 2.624 | 2.977 | 3.787  | 4.140  |
| 15        | 0.000 | 0.691 | 0.866 | 1.074 | 1.341 | 1.753 | 2.131 | 2.602 | 2.947 | 3.733  | 4.073  |
| 16        | 0.000 | 0.690 | 0.865 | 1.071 | 1.337 | 1.746 | 2.120 | 2.583 | 2.921 | 3.686  | 4.015  |
| 17        | 0.000 | 0.689 | 0.863 | 1.069 | 1.333 | 1.740 | 2.110 | 2.567 | 2.898 | 3.646  | 3.965  |
| 18        | 0.000 | 0.688 | 0.862 | 1.067 | 1.330 | 1.734 | 2.101 | 2.552 | 2.878 | 3.610  | 3.922  |
| 19        | 0.000 | 0.688 | 0.861 | 1.066 | 1.328 | 1.729 | 2.093 | 2.539 | 2.861 | 3.579  | 3.883  |
| 20        | 0.000 | 0.687 | 0.860 | 1.064 | 1.325 | 1.725 | 2.086 | 2.528 | 2.845 | 3.552  | 3.850  |
| 21        | 0.000 | 0.686 | 0.859 | 1.063 | 1.323 | 1.721 | 2.080 | 2.518 | 2.831 | 3.527  | 3.819  |
| 22        | 0.000 | 0.686 | 0.858 | 1.061 | 1.321 | 1.717 | 2.074 | 2.508 | 2.819 | 3.505  | 3.792  |
| 23        | 0.000 | 0.685 | 0.858 | 1.060 | 1.319 | 1.714 | 2.069 | 2.500 | 2.807 | 3.485  | 3.768  |
| 24        | 0.000 | 0.685 | 0.857 | 1.059 | 1.318 | 1.711 | 2.064 | 2.492 | 2.797 | 3.467  | 3.745  |
| 25        | 0.000 | 0.684 | 0.856 | 1.058 | 1.316 | 1.708 | 2.060 | 2.485 | 2.787 | 3.450  | 3.725  |
| 26        | 0.000 | 0.684 | 0.856 | 1.058 | 1.315 | 1.706 | 2.056 | 2.479 | 2.779 | 3.435  | 3.707  |
| 27        | 0.000 | 0.684 | 0.855 | 1.057 | 1.314 | 1.703 | 2.052 | 2.473 | 2.771 | 3.421  | 3.690  |
| 28        | 0.000 | 0.683 | 0.855 | 1.056 | 1.313 | 1.701 | 2.048 | 2.467 | 2.763 | 3.408  | 3.674  |
| 29        | 0.000 | 0.683 | 0.854 | 1.055 | 1.311 | 1.699 | 2.045 | 2.462 | 2.756 | 3.396  | 3.659  |
| 30        | 0.000 | 0.683 | 0.854 | 1.055 | 1.310 | 1.697 | 2.042 | 2.457 | 2.750 | 3.385  | 3.646  |
| 40        | 0.000 | 0.681 | 0.851 | 1.050 | 1.303 | 1.684 | 2.021 | 2.423 | 2.704 | 3.307  | 3.551  |
| 50        | 0.000 | 0.679 | 0.848 | 1.045 | 1.296 | 1.671 | 2.000 | 2.390 | 2.660 | 3.232  | 3.460  |
| 60        | 0.000 | 0.678 | 0.846 | 1.043 | 1.292 | 1.664 | 1.990 | 2.374 | 2.639 | 3.195  | 3.416  |
| 80        | 0.000 | 0.677 | 0.845 | 1.042 | 1.290 | 1.660 | 1.984 | 2.364 | 2.626 | 3.174  | 3.390  |
| 100       | 0.000 | 0.677 | 0.845 | 1.042 | 1.290 | 1.660 | 1.984 | 2.364 | 2.626 | 3.174  | 3.390  |
| 1000      | 0.000 | 0.675 | 0.842 | 1.037 | 1.282 | 1.646 | 1.962 | 2.330 | 2.581 | 3.098  | 3.300  |
| Z         | 0.000 | 0.674 | 0.842 | 1.036 | 1.282 | 1.645 | 1.960 | 2.326 | 2.576 | 3.090  | 3.291  |

## Lecture 5: Applications of t-distribution

So:

- Can assess confidence that population mean is within a certain range
- Can use t distribution to ask questions like:
  - “What is probability of getting sample with mean =  $\bar{y}$  from population with mean =  $\mu$ ?” (1 sample t-test)
  - “What is the probability that two samples came from same population?” (2 sample t-test)

## Lecture 5: Single Sample T-Test

We want to test if the mean fish length in I3 differs from 240mm.

**Activity: Define hypotheses and identify assumptions**

$H_0: \mu = 240$  (The mean fish length in I3 is 240mm)

$H_1: \mu \neq 240$  (The mean fish length in I3 is not 240mm)

### Assumptions for t-test:


1. Data is normally distributed
2. Observations are independent
3. No significant outliers

## Assumptions in R - qqplots from car

```
# Filter for just windward side needles

# YOUR TASK: Test normality of windward pine needle lengths
# QQ Plot
qqPlot(i3_df$length_mm,
       main = "QQ Plot for length of Grayling",
       ylab = "Sample Quantiles")
```

## QQ Plot for length of Grayling



```
[1] 53 35
```

## Statistical Test of Normality

### Shapiro-Wilk test


```
# Shapiro-Wilk test
shapiro_test <- shapiro.test(i3_df$length_mm)
print(shapiro_test)
```

Shapiro-Wilk normality test

data: i3\_df\$length\_mm  
W = 0.91051, p-value = 0.0001623

## Checking for Outliers

```
# Check for outliers using boxplot
# YOUR CODE HERE
i3_df %>% ggplot(aes(lake, length_mm))+geom_boxplot()
```



## Practice Exercise 1: One-Sample t-Test

## 💡 Practice Exercise 1: One-Sample t-Test

Let's perform a one-sample t-test to determine if the mean fish length in I3 Lake differs from 240 mm:

```
# what is the mean
i3_mean <- mean(i3_df$length_mm, na.rm=TRUE)
cat("Mean:", round(i3_mean, 1), "mm\n")
```

```
Mean: 265.6 mm
```

```
# Perform a one-sample t-test
t_test_result <- t.test(i3_df$length_mm, mu = 240)

# View the test results
t_test_result
```

### One Sample t-test

```
data: i3_df$length_mm
t = 7.3497, df = 65, p-value = 4.17e-10
alternative hypothesis: true mean is not equal to 240
95 percent confidence interval:
 258.6481 272.5640
sample estimates:
mean of x
 265.6061
```

Interpret this test result by answering these questions:

1. What was the null hypothesis?
2. What was the alternative hypothesis?
3. What does the p-value tell us?
4. Should we reject or fail to reject the null hypothesis at  $\alpha = 0.05$ ?
5. What is the practical interpretation of this result for fish biologists?

## Lecture 5: Hypothesis Testing Framework

Hypothesis testing is a systematic way to evaluate research questions using data.


### Key components:

1. **Null hypothesis ( $H_0$ ):** Typically assumes “no effect” or “no difference”
2. **Alternative hypothesis ( $H_a$ ):** The claim we’re trying to support
3. **Statistical test:** Method for evaluating evidence against  $H_0$
4. **P-value:** Probability of observing our results (or more extreme) if  $H_0$  is true
5. **Significance level ( $\alpha$ ):** Threshold for rejecting  $H_0$ , typically 0.05

**Decision rule:** Reject  $H_0$  if  $p\text{-value} < \alpha \Rightarrow p < 0.05$

## One-Sample t-Test

$H_0: \mu = 240$  vs  $H_1: \mu \neq 240$  ( $\alpha = 0.05$ ,  $df = 19$ )



## Lecture 5: Hypothesis Testing - Original Scale

Hypothesis testing is a systematic way to evaluate research questions using data.


### Key components:

1. **Null hypothesis ( $H_0$ ):** Typically assumes “no effect” or “no difference”
2. **Alternative hypothesis ( $H_a$ ):** The claim we’re trying to support
3. **Statistical test:** Method for evaluating evidence against  $H_0$
4. **P-value:** Probability of observing our results (or more extreme) if  $H_0$  is true
5. **Significance level ( $\alpha$ ):** Threshold for rejecting  $H_0$ , typically 0.05

**Decision rule:** Reject  $H_0$  if p-value  $< \alpha$

## One-Sample t-Test in Original Scale

Testing  $H_0: \mu = 240$  ( $\alpha = 0.05$ ,  $df = 19$ )



## Lecture 5: Interpreting One-Sample T-Test Results

**Activity: Interpret the t-test results**

- What does the p-value tell us?
- Should we reject or fail to reject the null hypothesis?

**How to report this result in a scientific paper:**

“A two-tailed, one-sample t-test at  $\alpha=0.05$  showed that the mean pine needle length on the windward side (... mm, SD = ...) [was/was not] significantly different from the expected 55 mm,  $t(\dots) = \dots$ ,  $p = \dots$ ”


## Lecture 5: Two Sample T-Tests Introduction

For example

- what is probability that population X is the same as population Y?

How would you assess this question using what we learned?

This is what we will do with the pine needles...



## Lecture 5: Comparing Two Samples


For example

- what is probability that population X is the same as population Y?

How would you assess this question using what we learned?

```
# Now create a boxplot to visualize the difference in fish lengths between these lakes:
pine_df <- read_csv("data/pine_needles.csv")

# Create a boxplot comparing the two lakes
pine_wind_plot <- pine_df %>%
  ggplot(aes(x = wind, y = length_mm, fill = wind)) +
  geom_boxplot() +
  labs(title = "Pine Needle Lengths by Wind Exposure",
       x = "Position",
       y = "Length (mm)",
       fill = "Wind Position") +
  scale_fill_manual(values = c("lee" = "forestgreen", "wind" = "skyblue"),
                   labels = c("lee" = "Leeward", "wind" = "Windward"))
pine_wind_plot
```



```
# Based on the t-test results and the boxplot
#
# what can you conclude about the fish populations in these two lakes?
```

## Practice Exercise 2: Formulating Hypotheses

### 💡 Practice Exercise 2: Formulating Hypotheses

For the following research questions about pine needles write the null and alternative hypotheses:

1. Are needles on the lee side longer than the needles on the windy side?

What are the hypotheses?

Ho =

Ha =

## Lecture 5: Two-Sample T-Test Framework

Now, let's compare pine needle lengths between windward and leeward sides of trees.

Question: **Is there a significant difference in needle length between the windward and leeward sides?**

This requires a two-sample t-test.

Two-sample t-test compares means from two independent groups.

$$t = \frac{\bar{x}_1 - \bar{x}_2}{S_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

where:

- $\bar{x}_1$  and  $\bar{x}_2$ : These represent the sample means of the two groups you're comparing



- $s_p^2$ : This is the pooled variance, calculated as:  $s_p^2 = [(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2] / (n_1 + n_2 - 2)$ , where  $s_1^2$  and  $s_2^2$  are the sample variances of the two groups.
- $n_1$  and  $n_2$ : These are the sample sizes of the two groups.
- $\sqrt{(1/n_1 + 1/n_2)}$ : This represents the pooled standard error.

$$t = \frac{SIGNAL}{NOISE}$$

## Practice Exercise 3: Summary Statistics

### 💡 Practice Exercise 3: Calculate summary statistics grouped by wind exposure

Before conducting the test, we need to understand the data for each group.

1. You need this and the graph to see what is going on ....


```
group_summary <- pine_df %>%
  group_by(wind) %>%
  summarize(
    mean_length = mean(length_mm),
    sd_length = sd(length_mm),
    n = n(),
    se_length = sd_length / sqrt(n)
  )

print(group_summary)
```

```
# A tibble: 2 × 5
  wind mean_length sd_length    n se_length
<chr>      <dbl>      <dbl> <int>    <dbl>
1 lee         20.4         2.45    24     0.500
2 wind        14.9         1.91    24     0.390
```

## Visualizing Group Differences

```
# Create a boxplot comparing the two sides
pine_wind_plot
```



## Practice Exercise 4: Effect Size

### 💡 Practice Exercise 4: Effect size

We could also look at the difference in means... some cool code here

```
# Assuming your dataframe is called df
group_summary %>%
  summarize(difference = mean_length[wind == "wind"] - mean_length[wind == "lee"])
```


```
# A tibble: 1 × 1
  difference
  <dbl>
1      -5.5
```

## Practice Exercise 5: ggplot Summary Statistics

## 💡 Practice Exercise 5: Using GGPLOT to get summary stats

GGplot also has code to make the mean and standard error plots we are interested in along with a lot of others

```
# Assuming your dataframe is called df
pine_mean_se_plot <- ggplot(pine_df, aes(x = wind, y = length_mm, color = wind)) +
  stat_summary(fun = mean, geom = "point") +
  stat_summary(fun.data = mean_se, geom = "errorbar", width = 0.2) +
  labs(title = "Mean Pine Needle Length by Wind Exposure",
       x = "Wind Exposure",
       y = "Mean Length (mm)") +
  coord_cartesian(ylim = c(0,25)) +
  scale_color_manual(values = c("lee" = "forestgreen", "wind" = "skyblue"),
                    labels = c("lee" = "Leeward", "wind" = "Windward")) +
  theme_classic()
pine_mean_se_plot
```



## Lecture 5: Testing Assumptions for Two-Sample T-Test

For a two-sample t-test, we need to check:

1. Normality within each group
2. Equal variances between groups (for standard t-test)
3. Independent observations

If assumptions are violated:

- Welch's t-test (unequal variances)
- Non-parametric alternatives (Mann-Whitney U test)


## Practice Exercise 6: Creating Group Data

## 💡 Practice Exercise 6: Test normality of windward pine needle lengths

qqplots

Note you need to test each groups separately...

```
# Assuming your dataframe is called df  
pine_mean_se_plot
```



## Practice Exercise 7: Separate Group Data

## 💡 Practice Exercise 7: Test normality of windward pine needle lengths

qqplots

Note you need to test each groups separately...

```
# how do you make separate dataframes to do this on?
# Separate data by groups
windward_data <- pine_df %>% filter(wind == "wind")
leeward_data <- pine_df %>% filter(wind == "lee")
head(leeward_data)
```

```
# A tibble: 6 × 6
  date      group      n_s  wind  tree_no length_mm
<chr>   <chr>    <chr> <chr>   <dbl>     <dbl>
1 3/20/25 cephalopods n    lee      1         20
2 3/20/25 cephalopods n    lee      1         21
3 3/20/25 cephalopods n    lee      1         23
4 3/20/25 cephalopods n    lee      1         25
5 3/20/25 cephalopods n    lee      1         21
6 3/20/25 cephalopods n    lee      1         16
```

## Practice Exercise 8: QQ Plot for Windward Data


## 💡 Practice Exercise 8: Test normality of windward pine needle lengths

qqplots

Note you need to test each groups separately...

```
# QQ Plot for windward group
qqPlot(windward_data$length_mm,
       main = "QQ Plot for Windward Pine Needles",
       ylab = "Sample Quantiles")
```

### QQ Plot for Windward Pine Needles



```
[1] 21 22
```

## Practice Exercise 9: Shapiro-Wilk Test

### 💡 Practice Exercise 9: Test normality of windward pine needle lengths

Shapiro-Wilk test

Note you need to test each groups separately...

```
# Shapiro-Wilk test for windward group
shapiro_windward <- shapiro.test(windward_data$length_mm)
print("Shapiro-Wilk test for windward data:")
```

```
[1] "Shapiro-Wilk test for windward data:"
```

```
print(shapiro_windward)
```

Shapiro-Wilk normality test

```
data:  windward_data$length_mm
W = 0.96062, p-value = 0.451
```

## Practice Exercise 10: QQ Plot for Leeward Data


## 💡 Practice Exercise 10: Test normality of leeward pine needle lengths

qqplots

Note you need to test each groups separately...

```
# You can also test the leeward group
# QQ Plot for leeward group
qqPlot(leeward_data$length_mm,
       main = "QQ Plot for Leeward Pine Needles",
       ylab = "Sample Quantiles")
```

### QQ Plot for Leeward Pine Needles



```
[1] 4 16
```

## Practice Exercise 11: Shapiro-Wilk for Leeward



### 💡 Practice Exercise 11: Test normality of leeward pine needle lengths

Shapiro-Wilk test

Note you need to test each groups separately...

```
# Shapiro-Wilk test for leeward group
shapiro_lee <- shapiro.test(leeward_data$length_mm)
print("Shapiro-Wilk test for leeward data:")
```

```
[1] "Shapiro-Wilk test for leeward data:"
```

```
print(shapiro_lee)
```

Shapiro-Wilk normality test

```
data: leeward_data$length_mm
W = 0.95477, p-value = 0.3425
```

## Practice Exercise 12: Combined Normality Test

### 💡 Practice Exercise 12: Test Normality at one time

There are always a lot of ways to do this in R

```
# there are always two ways
# Test for normality using Shapiro-Wilk test for each wind group
# All in one pipeline using tidyverse approach
normality_results <- pine_df %>%
  group_by(wind) %>%
  summarize(
    shapiro_stat = shapiro.test(length_mm)$statistic,
    shapiro_p_value = shapiro.test(length_mm)$p.value,
    normal_distribution = if_else(shapiro_p_value > 0.05, "Normal", "Non-normal")
  )

# Print the results
print(normality_results)
```

```
# A tibble: 2 × 4
  wind shapiro_stat shapiro_p_value normal_distribution
<chr>    <dbl>         <dbl> <chr>
1 lee      0.955         0.343 Normal
2 wind     0.961         0.451 Normal
```

## Practice Exercise 13: Test Equal Variances

### 💡 Practice Exercise 13: Test equal variances

Levenes test can be done on the original dataframe

```
# Method 1: Using car package's leveneTest
# This is often preferred as it's more robust to departures from normality
levene_result <- leveneTest(length_mm ~ wind, data = pine_df)
print("Levene's Test for Homogeneity of Variance:")
```

```
[1] "Levene's Test for Homogeneity of Variance:"
```

```
print(levene_result)
```

```
Levene's Test for Homogeneity of Variance (center = median)
      Df F value Pr(>F)
group  1  1.2004 0.2789
      46
```

## Lecture 5: Conducting the Two-Sample T-Test

Now we can compare the mean pine needle lengths between windward and leeward sides.

Ho:  $\mu_1 = \mu_2$  (The mean needle lengths are equal)

Ha:  $\mu_1 \neq \mu_2$  (The mean needle lengths are different)

Deciding between:

- Standard t-test (equal variances)
- Welch's t-test (unequal variances)

**Note the Levenes Test should be NOT SIGNIFICANT - What is the null hypothesis**

```
Levene's Test for Homogeneity of Variance (center = median)
      Df F value Pr(>F)
group  1  1.2004 0.2789
      46
```

## Lecture 5: Running the Two-Sample T-Test

Now we can do a two sample TTEST

Calculate t-statistic manually (optional)

YOUR CODE HERE:

```
t = (mean1 - mean2) / sqrt((s1^2/n1) + (s2^2/n2))
```



Tip

```
# YOUR TASK: Conduct a two-sample t-test
# Use var.equal=TRUE for standard t-test or var.equal=FALSE for Welch's t-test

# Standard t-test (if variances are equal)
t_test_result <- t.test(length_mm ~ wind, data = pine_df, var.equal = TRUE)
print("Standard two-sample t-test:")
```

```
[1] "Standard two-sample t-test:"
```

```
print(t_test_result)
```

Two Sample t-test

```
data: length_mm by wind
t = 8.6792, df = 46, p-value = 3.01e-11
alternative hypothesis: true difference in means between group lee and group wind is not
equal to 0
95 percent confidence interval:
 4.224437 6.775563
sample estimates:
 mean in group lee mean in group wind
      20.41667      14.91667
```

```
# Welch's t-test (if variances are unequal)
# YOUR CODE HERE
```

## Lecture 5: Interpreting Two-Sample T-Test Results

### Interpret the results of the two-sample t-test


What can we conclude about the needle lengths on windward vs. leeward sides?

#### How to report this result in a scientific paper:

“A two-tailed, two-sample t-test at  $\alpha=0.05$  showed [a significant/no significant] difference in needle length between windward ( $M = \dots$ ,  $SD = \dots$ ) and leeward ( $M = \dots$ ,  $SD = \dots$ ) sides of pine trees,  $t(\dots) = \dots$ ,  $p = \dots$ ”

## T-Distribution with Observed T-Value

Two-Sample T-Test for Pine Needle Length (df = 46)



## Lecture 5: Visualizing the Results

### Interpret the results of the two-sample t-test


What can we conclude about the needle lengths on windward vs. leeward sides?

### How to report this result in a scientific paper:

“A two-tailed, two-sample t-test at  $\alpha=0.05$  showed [a significant/no significant] difference in needle length between windward (M = ..., SD = ...) and leeward (M = ..., SD = ...) sides of pine trees,  $t(\dots) = \dots$ ,  $p = \dots$ ”

## Density Plot of Pine Needle Lengths by Wind Exposure

Mean Difference = 5.5mm ( $t = 8.68$ ,  $p < 0.001$ )



## Lecture 5: Assumptions of Parametric Tests

### Common assumptions for t-tests:

1. Normality: Data comes from normally distributed populations
2. Equal variances (for two-sample tests)
3. Independence: Observations are independent
4. No outliers: Extreme values can influence results

What can we do if our data violates these assumptions?

Alternatives when assumptions are violated:

- Data transformation (log, square root, etc.)
- Non-parametric tests
- Robust statistical methods

## Lecture 5: Summary and Conclusions

In this activity, we've:

1. Formulated hypotheses about pine needle length
2. Tested assumptions for parametric tests

3. Conducted one-sample and two-sample t-tests
4. Visualized data using appropriate methods
5. Learned how to interpret and report t-test results

**Key takeaways:**

- Always check assumptions before conducting tests
- Visualize your data to understand patterns
- Report results comprehensively
- Consider alternatives when assumptions are violated