


Lecture 06

Bill Perry

Lecture 5: Review

Covered


- Statistical inference fundamentals
- Hypothesis testing principles
- T Distributions
- One sample T Tests
- Two sample T Tests
- Paired T Tests



Lecture 6: Overview

The objectives:

- Brief review
- Hypothesis tests for two populations
- Assumptions of parametric tests
- Power of Parametric Tests
- Introduction to Non-Parametric Tests



Lecture 6: Statistical hypothesis testing

- Major goal of statistics:
 - inferences about populations from samples...
 - assign degree of confidence to inferences
 - Statistical hypothesis testing:
 - formalized approach to inference
 - Hypotheses ask whether samples come from populations with certain properties
 - Often interested in questions about population means
 - but other questions are of interest



Lecture 6: Hypothesis Components

Useful hypotheses:

- Rely on specifying
 - H_0 is the hypothesis of “no effect”
 - two samples from population with same mean
 - sample is from population of mean = X
 - H_a (research or alternate hypothesis)
 - is the opposite of the H_0
 - or predicts that there is an effect of x on y
 - *but does NOT suggest a direction which is a prediction*



Lecture 6: Hypothesis Examples

Together H_0 and H_a encompass all possible outcomes:

- $H_0: \mu = 0$, $H_a: \mu \neq 0$
 - mean equals 0 or mean does not equal 0
- $H_0: \mu = 35$, $H_a: \mu \neq 35$
 - mean equals 35 or mean does not equal 35
- $H_0: \mu_1 = \mu_2$, $H_a: \mu_1 \neq \mu_2$
 - mean of population 1 equals mean of population 2 or it does not
- $H_0: \mu > 0$, $H_a: \mu \leq 0$
 - can be directional mean is greater than 0 or mean is not equal or less than 0
 - this becomes a one sided test as it predicts only one direction



Lecture 6: Statistical Testing Framework

Statistical tests assess likelihood of the null hypothesis being true

- If the H_0 is likely false, then H_a assumed to be correct
- More precisely:
 - the long run probability of obtaining sample value (or more extreme one) if the null hypothesis is true
 - $p(\text{data}|H_0)$ - the probability of observing the data given that the null hypothesis H_0 is true



Lecture 6: Understanding P-values

Hypothesis tests

- Expressed as p-value (0-never to 1-always)
- Interpret p-value as:
 - probability of obtaining sample value of statistic (or more extreme one) if H_0 is true
- High p-value:
 - high probability of obtaining sample statistic under H_0
 - if the null hypothesis (H_0) were true, you would frequently observe data similar to your sample statistic
 - your observed results are quite compatible with what the null hypothesis predicts
- Low p-value: low probability of obtaining sample statistic under H_0
 - if the null hypothesis (H_0) were true, you would rarely observe data similar to or more extreme than your sample statistic
 - Your results are unusual under the null hypothesis, suggesting that either you've witnessed a rare event or the null hypothesis may be incorrect



Lecture 6: P-value Interpretation

Statistical test results:

- $p = 0.3$ means that if I repeated the study 100 times, I would get this (or more extreme) result due to chance **30 times**

- $p = 0.03$ means that if I repeated the study 100 times, I would get this (or more extreme) result due to chance **3 times**

Which p -value suggests H_0 likely false?

At what point reject H_0 ?

$p < 0.05$ conventional “significance threshold” (α = alpha or p value)

$p < 0.05$ means: if H_0 is true and we repeated the study 100 times

- we would get this (or more extreme) result less than 5 times due to chance

Lecture 6: Significance Levels and Interpretation

Statistical test results:

- **α is the rate at which we will reject a true null hypothesis (Type I error rate)**
- Lowering α will lower likelihood of incorrectly rejecting a true null hypothesis (e.g., 0.01, 0.001)
- *Both H_s and α are specified BEFORE collection of data and analysis*

Traditionally $\alpha=0.05$ is used as a cut off for rejecting null hypothesis

There is nothing magical about 0.05 - actual p -values need to be reported - also need to decide prior to study

p-value range	Interpretation
$P > 0.10$	No evidence against H_0 - data appear consistent with H_0
$0.05 < P < 0.10$	Weak evidence against the H_0 in favor of H_a
$0.01 < P < 0.05$	Moderate evidence against H_0 in favor of H_a
$0.001 < P < 0.01$	Strong evidence against H_0 in favor of H_a
$P < 0.001$	Very strong evidence against H_0 in favor of H_a


Lecture 6: Understanding P-values Visually

A **p -value** is the probability of observing the sample result (or something more extreme) if the null hypothesis is true.

- **Common interpretations:**
 - $p < 0.05$: Strong evidence against H_0
 - $0.05 \leq p < 0.10$: Moderate evidence against H_0
 - $p \geq 0.10$: Insufficient evidence against H_0
- **Common misinterpretations:**
 - p -value is NOT the probability that H_0 is true
 - p -value is NOT the probability that results occurred by chance
 - Statistical significance $\neq$ practical significance
- Note that there is a difference in how to state the hypotheses
 - one sample T-TEST
 - two sample T-TEST

df = 20 , $\alpha = 0.05$ (0.025 in each tail)

df = 20 , $\alpha = 0.05$ (0.025 in each tail)



Lecture 6: Historical Context

end to hyped claims and the dismissal of possibly crucial effects.


Valentin Arrheim¹, Sander Greenland & Blake McShane

Lecture 6: Fisher's Perspective

Fisher:

p-value as informal measure of discrepancy between data and H_0

“If p is between 0.1 and 0.9 there is certainly no reason to suspect the hypothesis tested. If it is below 0.02 it is strongly indicated that the hypothesis fails to account for the whole of the facts. We shall not often be astray if we draw a conventional line at .05 ...”



Ronald Fisher: 1890-1962


Decision errors

- Even good studies can reach incorrect conclusions
- “Decision errors”
- Two types of decision errors
- Want to know probability of making these errors

		Statistical Conclusion	
		Reject H_0	Fail to Reject
Effect	Effect	Correct Decision Effect detected ; Effect exists	“FALSE NEGATIVE” Type II Error Effect not detected Effect exists
Effect	No effect	“FALSE POSITIVE” Type I Error Effect detected; none exists	Correct Decision No effect detected None exists

Type I and Type II Errors - Concept

- Type I error rate
 - α : wrongly reject H_0 when it's true
 - $\alpha = 0.05$ means a type I error rate of 5%
- Type II error rate, β
 - wrongly fail to reject H_0 when it's false
- **Power = $1 - \beta$: probability of correctly rejecting H_0 when H_1 is true**
- Inverse relationship between type I and type II error - but not straightforward
- Result of chance - sample not representative of population
- Which type of error is more dangerous?



the dotted line is the $\alpha = 0.05$

Lecture 6: Type I and Type II Errors - Details

When making decisions based on hypothesis tests, two types of errors can occur:

Type I Error (False Positive)

- Rejecting H_0 when it's actually true
- Probability = α (significance level)
- "Finding an effect that isn't real"

Type II Error (False Negative)

- Failing to reject H_0 when it's actually false
- Probability = β - "Missing an effect that is real"

Statistical Power = $1 - \beta$

- Probability of correctly rejecting a false H_0
- Increases with:
 - Larger sample size
 - Larger effect size
 - Lower variability
 - Higher α level

The farther apart the means or lower variance the lower the beta error is ~ have higher power. - CLICK TO NEXT AS SAME TEXT

Type I error
(false positive)



Type II error
(false negative)



Lecture 6: Type I and Type II Errors - Visualization

When making decisions based on hypothesis tests, two types of errors can occur:

Type I Error (False Positive)

- Rejecting H_0 when it's actually true
- Probability = α (significance level)
- "Finding an effect that isn't real"

Type II Error (False Negative)


- Failing to reject H_0 when it's actually false
- Probability = β - "Missing an effect that is real"

Statistical Power = $1 - \beta$

- Probability of correctly rejecting a false H_0
- Increases with:
 - Larger sample size
 - Larger effect size
 - Lower variability
 - Higher α level

The farther apart the means or lower variance the lower the beta error is ~ have higher power.

Type I and Type II Errors



Practice Exercise: Interpreting Errors and Power

🔗 Practice Exercise 6: Interpreting P-values and Errors

Given the following scenarios, identify whether a Type I or Type II error might have occurred:

1. A researcher concludes that island mice size is larger, when in fact it is not.
2. A study fails to detect a real difference in mouse size on islands when there is and concludes there is no effect.
3. Let's calculate the power of our t-test to detect a 5 mm difference in mass sampling_sites:
 - pooled standard deviation
 - This is the combined standard deviation of both groups weighted by respective degrees of freedom.
 - Cohen's d
 - standardized difference between means - here assuming a difference of 1 units (g)
 - $\delta = 0.423$: The standardized effect size (Cohen's d)

```
library(car)
library(patchwork)
library(tidyverse)
library(readxl)

m_df <- read_csv("data/mice_weights.csv") %>%
  filter(!is.na(mass_g))

sidney_df <- m_df %>% filter(sampling_site=="Sidney Island")
vancouver_df <- m_df %>% filter(sampling_site=="Vancouver")
# Calculate power for detecting a 30 mm difference

n1 <- nrow(sidney_df)
n2 <- nrow(vancouver_df)
sd_pooled <- sqrt((var(sidney_df$mass_g) * (n1-1) +
  var(vancouver_df$mass_g) * (n2-1)) /
  (n1 + n2 - 2))

# Calculate power
effect_size <- 1 / sd_pooled # Cohen's d
df <- n1 + n2 - 2
alpha <- 0.05
power <- power.t.test(n = min(n1, n2),
  delta = effect_size,
  sd = 1, # Using standardized effect size
  sig.level = alpha,
  type = "two.sample",
  alternative = "two.sided")

# Display results
power
```

Two-sample t test power calculation

```
n = 28
delta = 0.4231154
sd = 1
sig.level = 0.05
power = 0.3427604
```

alternative = two.sided

What if we calculated power for the pine needles you measured?

```
pine_switch_df <- read_excel("data/class_pine needle length switched.xlsx")
ps_df <- pine_switch_df %>%
  group_by(group, tree_no, tree_char, side) %>%
  summarise(length_mm = mean(length_mm, na.rm=TRUE))

ps_shady_df <- ps_df %>%
  filter(side == "shady")

ps_sunny_df <- ps_df %>%
  filter(side == "sunny")

pine_alpha <- 0.05

# Calculate power for detecting a 30 mm difference

pine_n1 <- nrow(ps_shady_df) # FIX: was using sidney_df instead of ps_shady_df
pine_n2 <- nrow(ps_sunny_df) # FIX: was using vancouver_df instead of ps_sunny_df

pine_sd_pooled <- sqrt((var(ps_shady_df$length_mm) * (pine_n1-1) +
  var(ps_sunny_df$length_mm) * (pine_n2-1)) /
  (pine_n1 + pine_n2 - 2)) # FIX: was using n1, n2 instead of pine_n1,
pine_n2

cat("Pooled SD:", round(pine_sd_pooled, 2), "mm\n\n")
```

Pooled SD: 2.58 mm

```
# More realistic effect sizes to test:
observed_diff <- abs(mean(ps_shady_df$length_mm) - mean(ps_sunny_df$length_mm))
pine_effect_size_observed <- observed_diff / pine_sd_pooled
cat("Effect size for observed", round(observed_diff, 2), "mm difference: Cohen's d =",
  round(pine_effect_size_observed, 2), "\n")
```

Effect size for observed 1.45 mm difference: Cohen's d = 0.56

```
# 2. Power for the observed difference
pine_power_observed <- power.t.test(n = min(pine_n1, pine_n2),
  delta = pine_effect_size_observed,
  sd = 1,
  sig.level = pine_alpha,
  type = "two.sample",
  alternative = "two.sided")

cat("Power for observed", round(observed_diff, 2), "mm difference:",
  round(pine_power_observed$power, 3), "\n")
```

Power for observed 1.45 mm difference: 0.182

What is Power

Statistical power represents the probability of detecting a true effect (rejecting the null hypothesis when it is false). In this case, with a power of 97%, there's a 97% chance of detecting a true difference of X units between the means of the two groups if such a difference actually exists.

A power analysis like this is typically done for one of these purposes:

1. Before data collection to determine required sample size
2. After a study to evaluate if the sample size was adequate
3. To determine the minimum detectable effect size with the given sample

With 97% power, this test has excellent ability to detect the specified effect size. Generally, **80% power is considered acceptable**, so 97% indicates a very well-powered study for detecting a difference of XXg between the groups.

Lecture 6: Error Bars and Their Interpretation

Error bars are graphical representations of the variability of data that show:

- The **precision** of a measurement
- The **uncertainty** around an estimate
- A **confidence interval** for a parameter

Common types of error bars:


1. **Standard Error (SE)**: Shows precision of the mean
2. **Standard Deviation (SD)**: Shows variability in the data
3. **Confidence Interval (CI)**: Shows plausible range for parameter

When interpreting graphs:

- Always check what the error bars represent
- Non-overlapping 95% CI bars suggest statistically significant differences
- Error bars help assess both statistical and practical significance

Different Types of Error Bars

Comparing SD, SE, and 95% CI



Lecture 6: Sampling and Pseudoreplication

Pseudoreplication occurs when measurements that are not independent are analyzed as if they were independent.


- A critical consideration in experimental design
- Results in underestimated standard errors and confidence intervals
- Leads to inflated Type I error rates (false positives)

Examples of pseudoreplication:

- Measuring the same individual multiple times
- Treating multiple fish from the same tank as independent
- Using multiple data points from a single site

How to avoid pseudoreplication:

- Identify the true experimental unit
- Use appropriate statistical techniques (e.g., mixed models)
- Be clear about the level of replication



Lecture 6: Summary and Key Takeaways

Key concepts covered:

1. **P-values** measure evidence against the null hypothesis
 - Not the probability that H_0 is true
 - Should be interpreted in context with effect size
2. **Hypothesis testing** provides a framework for making decisions
 - Null and alternative hypotheses must be specified beforehand
 - α level determines Type I error rate
3. **Type I and Type II errors** represent different kinds of mistakes
 - Type I (α): False positive - rejecting true H_0
 - Type II (β): False negative - failing to reject false H_0
 - Statistical power = $1 - \beta$

4. **Error bars** communicate uncertainty in different ways
 - Always check what type of error bar is shown
 - CI bars help assess statistical significance
5. **Pseudoreplication** inflates significance
 - Identify true experimental units
 - Account for non-independence in analysis