

Fibonacci Polynomials & *Lights Out*

William Boyles *

August 6, 2026

Abstract

Consider a game played on a simple graph $G = (V, E)$ where each vertex consists of clickable light. A move is clicking any $v \in V$, which toggles the on/off state of v and its neighbors. Given an initial configuration, one wins the game by finding a sequence of moves that turns off all of the lights. This game was commercially available from Tiger Electronics as *Lights Out* for G a 5×5 grid.

Let $\Phi_G : P(V) \rightarrow P(V)$ be the function which maps the vertices clicked to the vertices that change state. For G and $n \times n$ grid, let $d(n) = \dim(\ker \Phi_G)$. We show several that many values of $d(n)$ never occur. We show more general formulas for the value of $d(n)$, including a complete characterization when $n = 2^a p^b - 1$ for $a, b \geq 0$ and p not a Wieferich prime.

1 Fibonacci Polynomials

Definition 1.1 (Fibonacci Polynomial). *Let $F_n(x)$ be the polynomial in the ring $\mathbb{F}_2[x]$ defined recursively by*

$$F_n(x) = \begin{cases} 0 & n = 0 \\ 1 & n = 1 \\ xF_{n-1}(x) + F_{n-2}(x) & \text{otherwise} \end{cases}.$$

These polynomials are called "Fibonacci" because when defined over the real numbers, $F_n(1)$ is the n th Fibonacci number.

In $\mathbb{F}_2[x]$, the coefficients of $F_n(x)$ are either 1 or 0, so we can visualize them by coloring squares black or white to represent the coefficients. For example,

$$F_6(x) = 1x^5 + 0x^4 + 0x^3 + 0x^2 + 1x + 0 = \blacksquare \square \square \square \blacksquare \square.$$

Plotting $F_1(x)$ through $F_{128}(x)$ in the same way, aligning terms of the same degree, we see a Sierpinski triangle.

*Independent (williamboyles22900@gmail.com)




Figure 1: $F_1(x)$ through $F_{128}(x)$, right-aligned.

Using a well-known connection between the Sierpinski triangle and the parity of binomial coefficients, we may see that

$$F_n(x) = \sum_{i=0}^{n-1} \binom{n+i}{2i+1} x^i \pmod{2}.$$

The above construction and formula was also shown by Sutner [10]. Sutner also showed the following relationship between square *Lights Out* grids and Fibonacci Polynomials.

Lemma 1.1 (Sutner). *Let $G = (V, E)$ be an $n \times m$ grid graph. Let $\Phi_G : P(V) \rightarrow P(V)$ be the function that maps which lights are clicked to which lights change state. Then*

$$\dim(\ker \Phi_G) = \deg(\gcd(F_{n+1}(x), F_{m+1}(x+1))) = \deg(\gcd(F_{n+1}(x+1), F_{m+1}(x))).$$

Proof. See [10] theorem 5.1. ■

Sutner and others also shows several divisibility rules for these polynomials

Lemma 1.2 (Hunziker, Machiavelo, and Park). *A polynomial $\tau(x)$ in $\mathbb{F}_2[x]$ divides both $F_n(x)$ and $F_m(x)$ if and only if it divides $F_{\gcd(m,n)}(x)$. In particular,*

$$\gcd(F_m(x), F_n(x)) = F_{\gcd(m,n)}(x).$$

Proof. See [5] proposition 2.2, [10] proposition 2.1, and [6]. ■

Corollary 1.2.1. *The following are true:*

- (i) x divides $F_n(x)$ if and only if $n \equiv 0 \pmod{2}$.
- (ii) $x + 1$ divides $F_n(x + 1)$ if and only if $n \equiv 0 \pmod{2}$.
- (iii) $x + 1$ divides $F_n(x)$ if and only if $n \equiv 0 \pmod{3}$.
- (iv) x divides $F_n(x + 1)$ if and only if $n \equiv 0 \pmod{3}$.

Proof. Notice,

- (i) We find that $f_2(x) = x$ is the smallest Fibonacci polynomial divisible by x , so we apply lemma 1.2 to get the desired result.
 - (ii) Follows from (i) by substituting $x + 1$ for x .
 - (iii) We find that $f_3(x) = (x + 1)^2$ is the smallest Fibonacci polynomial divisible by $x + 1$, so we apply lemma 1.2 to get the desired result.
 - (iv) Follows from (iii) by substituting $x + 1$ for x .
-

Lemma 1.3 (Hunziker, Machiavelo, and Park). *If n is odd, then $F_n(x)$ is the square of a square-free polynomial.*

Proof. See [5] proposition 2.11 and [10] Theorem 2.1 claim 2. ■

Corollary 1.3.1. *If n is odd, then every root of $F_n(x)$ has multiplicity 2.*

Proof. Let n be odd and α a root of $F_n(x)$. By 1.3, there exists $\tau_n \in \mathbb{F}_2[x]$ such that

$$F_n(x) = \tau_n(x)^2,$$

and $\tau(x)$ is square-free. Since α is a root of $F_n(x)$, it must also be a root of $\tau_n(x)$ and thus has multiplicity 2. ■

We can now prove some of our own facts about these polynomials and their GCDs, particularly on square grids.

Lemma 1.4. *Let $d(n) = \dim(\ker \Phi_G)$ for G an $n \times n$ grid graph. Then $d(n)$ is always even.*

Proof. Let

$$G_n(x) = \gcd(F_n(x), F_n(x + 1)).$$

By lemma 1.1, we know that $d(n) = \deg G_{n+1}(x)$ and $G_n(x) = G_n(x + 1)$.

$y = x(x + 1)$ is also unchanged by this $x \mapsto x + 1$ translation. It is well-known that every polynomial $P(x)$ in $\mathbb{F}_2[x]$ can be uniquely written as

$$P(x) = A(y) + xB(y) \quad A, B \in \mathbb{F}_2[y].$$

We can then determine exactly when P is also unchanged by translation.

$$\begin{aligned} P(x+1) &= P(x) \\ A(y) + (x+1)B(y) &= A(y) + xB(y) \\ B(y) &= 0. \end{aligned}$$

In other words, $P(x)$ is unchanged by translation when P is a polynomial in $x^2 + x$.

Since G_n is also unchanged by translation, there exists $H_n \in \mathbb{F}_2[x]$ such that

$$G_n(x) = H_n(x^2 + x).$$

So,

$$d(n) = \deg G_{n+1}(x) = 2 \deg H_{n+1}(x)$$

and thus $d(n)$ is even. ■

Corollary 1.4.1. *If n is even, then $d(n)$ is divisible by 4.*

Proof. If n is even, then $n+1$ is odd. Then by 1.3, $F_{n+1}(x)$ and $F_{n+1}(x+1)$ are squares of square-free polynomials $\tau(x)$ and $\tau(x+1)$.

$$\begin{aligned} G_{n+1}(x) &= \gcd(F_{n+1}(x), F_{n+1}(x+1)) \\ &= \gcd(\tau_{n+1}(x)^2, \tau_{n+1}(x+1)^2) \\ &= \gcd(\tau_{n+1}(x), \tau_{n+1}(x+1))^2 \\ \sqrt{G_{n+1}(x)} &= \gcd(\tau_{n+1}(x), \tau_{n+1}(x+1)) \end{aligned}$$

We can see from the right-hand side of the equation that this function is also unaffected by translation. So, by lemma 1.4, there exists $H_{n+1} \in \mathbb{F}_2[x]$ such that

$$\begin{aligned} \sqrt{G_{n+1}(x)} &= H_{n+1}(x^2 + x) \\ G_{n+1}(x) &= H_{n+1}(x^2 + x)^2. \end{aligned}$$

So,

$$d(n) = \deg G_{n+1}(x) = 4 \deg H_{n+1}(x),$$

as desired. ■

Note that the implication does not hold in the other direction: $d(n)$ also be divisible by 4 when n is odd. For example, $d(9) = 8$.

Lemma 1.5. *Let $x = t + t^{-1}$. Then for $t \neq 0$,*

$$xF_n(x) = t^n + t^{-n}.$$

Proof. Define $G_n(t) = t^{n+1} + t^{-(n+1)}$. Then

$$\begin{aligned} xG_{n-1}(t) + G_{n-2}(t) &= (t + t^{-1})(t^n + t^{-n}) + (t^{n-1} + t^{-(n-1)}) \\ &= t^{n+1} + t^{-(n-1)} + t^{n-1} + t^{-(n+1)} + t^{n-1} + t^{-(n-1)} \\ &= t^{n+1} + t^{-(n+1)} \\ &= G_n. \end{aligned}$$

Thus we see that $G_n(t)$ satisfies the same recurrence relation as $F_n(x)$. $xF_n(x)$ also satisfies the same recurrence relation.

$$\begin{aligned} xF_n(x) &= x(xF_{n-1}(x) + F_{n-2}(x)) \\ &= x(xF_{n-1}(x)) + xF_{n-2}(x). \end{aligned}$$

Now we compute $xF_1(x)$ and $xF_2(x)$ in terms of t and G :

$$\begin{aligned} xF_1(x) &= x \\ &= t + t^{-1} \\ &= G_0(t), \end{aligned}$$

and

$$\begin{aligned} xF_2(x) &= x^2 \\ &= (t + t^{-1})^2 \\ &= t^2 + t^{-2} \\ &= G_1(t). \end{aligned}$$

Since the first two terms match and they satisfy the same recurrence relation then we've shown by induction that for $t \neq 0$,

$$xF_n(x) = G_{n-1}(t) = t^n + t^{-n}.$$

■

This was also proven in [5], who also prove the following identities.

Lemma 1.6 (Hunziker, Machiavelo, and Park). *Let $n = 2^k b$ where $b, k \in \mathbb{N}$. Then*

$$F_n(x) = x^{2^k - 1} F_b(x)^{2^k}.$$

Proof. See [5] lemma 2.6. ■

Lemma 1.7. *The distinct roots of $F_n(x)$ are $\{\zeta + \zeta^{-1} : \zeta^n = 1, \zeta \neq 1\}$, plus 0 when n is even.*

Proof. Notice that $F_n(0) = F_{n-2}(0)$, so

$$F_n(0) = \begin{cases} 1 & n \text{ odd} \\ 0 & n \text{ even} \end{cases}.$$

Thus 0 is a root of $F_n(x)$ whenever n is even.

Let α be a non-zero root of F_n . Choose ζ satisfying

$$\zeta^2 + \alpha\zeta + 1 = 0.$$

Notice that ζ cannot be 0. Dividing by ζ ,

$$\begin{aligned} \zeta + \alpha + \zeta^{-1} &= 0 \\ \alpha &= \zeta + \zeta^{-1}. \end{aligned}$$

By, Lemma 1.5,

$$\begin{aligned} 0 &= \alpha F_n(\alpha) \\ &= \zeta^n + \zeta^{-n} \end{aligned}$$

Since a field has no nonzero nilpotents, we may multiply by ζ^n .

$$\begin{aligned} 0 &= \zeta^{2n} + 1 \\ &= (\zeta^n + 1)^2, \end{aligned}$$

and

$$\begin{aligned} 0 &= \zeta^n + 1 \\ \zeta^n &= 1. \end{aligned}$$

Thus the order of ζ divides n . Notice too that $\zeta \neq 1$; otherwise, $\alpha = 1 + 1^{-1} = 0$, which we already showed as not possible.

Conversely, if $\zeta^n = 1$ and $\zeta \neq 1$, then let $\alpha = \zeta + \zeta^{-1}$. Then we see $\alpha \neq 0$ and

$$\alpha F_n(\alpha) = \zeta^n + \zeta^{-n} = 1 + 1^{-1} = 0.$$

Thus, α is a root of F_n . ■

2 Even Parity Covers

$\ker \Phi_G$ may also be understood via parity covers and dominating sets.

Definition 2.1. *An even dominating set of a graph $G = (V, E)$ is a subset $V' \subseteq V$ such that every vertex contains an even number of vertices from V' in its neighborhood.*

Odd dominating sets can be analogously defined. Sutner specifically studied odd dominating sets as the solution to *Lights Out* when all lights are initially on. He showed that such an odd dominating set always exists for any finite graph [9].

Lemma 2.1. *$V' \in \ker \Phi_G$ if and only if V' is an even dominating set of G .*

Proof. Let $V' \in \ker \Phi_G$. Let $N(v)$ be the neighborhood of vertex v : v and its immediate neighbors. Then for every $v \in V$,

$$|N(v) \cap V'| \text{ is even.}$$

Thus, V' is an even dominating set of G . We can apply the same steps in reverse to show the converse. ■

Lemma 2.2. *Φ_G is linear with respect to symmetric difference. That is, for all $V_1, V_2 \subseteq V$,*

$$\Phi_G(V_1 \triangle V_2) = \Phi_G(V_1) \triangle \Phi_G(V_2).$$

Proof. Notice that for any $v \in V$,

$$|N(v) \cap (V_1 \triangle V_2)| \equiv |\{N(v) \cap V_1\}| + |\{N(v) \cap V_2\}| \pmod{2}.$$

So, we can write

$$\begin{aligned} \Phi_G(V_1 \triangle V_2) &= \{v \in V : |N(v) \cap (V_1 \triangle V_2)| \text{ is odd}\} \\ &= (\{v \in V : |N(v) \cap V_1| \text{ is odd}\} \cup \{v \in V : |N(v) \cap V_2| \text{ is odd}\}) \setminus \{v \in V : |N(v) \cap (V_1 \cap V_2)| \text{ is odd}\} = \Phi_G(V_1) \triangle \Phi_G(V_2) \end{aligned}$$

as desired. ■

Corollary 2.2.1. *If V_1 and V_2 have the same image under Φ_G if and only if $V_1 \triangle V_2 \in \ker \Phi_G$.*

Proof. Assume that V_1 and V_2 have the same image under Φ_G . Then by lemma 2.2,

$$\begin{aligned} \Phi_G(V_1 \triangle V_2) &= \Phi_G(V_1) \triangle \Phi_G(V_2) \\ &= \Phi_G(V_1) \triangle \Phi_G(V_1) \\ &= \emptyset. \end{aligned}$$

Thus, $V_1 \triangle V_2 \in \ker \Phi_G$.

Now assume $V_1 \triangle V_2 \in \ker \Phi_G$. Then by lemma 2.2,

$$\begin{aligned}\Phi_G(V_1) &= \Phi_G(V_1) \triangle \emptyset \\ &= \Phi_G(V_1) \triangle \Phi_G(V_1 \triangle V_2) \\ &= \Phi_G(V_1) \triangle \Phi_G(V_1) \triangle \Phi_G(V_2) \\ &= \Phi_G(V_2).\end{aligned}$$

Thus V_1 and V_2 have the same image under Φ_G . ■

We may now use our geometric understanding of $\ker \Phi_G$ as even parity covers to construct a lower bound on $d(n)$.

Theorem 2.1. *For all $n, k \in \mathbb{N}$,*

$$d(nk - 1) \geq d(n - 1).$$

Proof. Let G_L be an $nk - 1 \times nk - 1$ grid and G_ℓ be an $n - 1 \times n - 1$ grid. Since $nk - 1 = k(n - 1) + (k - 1)$, an G may be broken into a $k \times k$ grid of $n - 1 \times n - 1$ grids, each separated by a horizontal or vertical strip of size 1.

Let Q be an even dominating set of G_ℓ . Let R be Q tiled k times horizontally and vertically, with size 1 separators between each tile, onto G_L where each tile is the reflection across its neighboring separators, as shown below.




Figure 2: Pattern Q tiled 5 times.

Consider a vertex v in G_L . There are 3 cases.

1. v is within a tile. Then $|N(v) \cap R|$ is even because Q being is an even parity cover.
2. v is on a separator between two tiles. Then only two elements in $N(v)$, the vertices within a tile, are also in R . Because neighboring tiles are reflections of each other across separators, these two vertices are either both in R or both not in R . In either case, $|N(v) \cap R|$ is even.

3. v is in the intersection of two separators. Then all of $N(v)$ is in a separator and thus not in R , so $|N(v) \cap R| = 0$, which is even.

Since for every vertex v of G_L , $|N(v) \cap R|$ is even, so R is an even dominating set of G_L . Thus, for every even dominating set of G_ℓ , there exists an even dominating set of G_L obtained through tiling. This tiling function is also injective because we can recover Q by restricting R to just the top left tile. Thus,

$$d(nk - 1) \geq d(n - 1),$$

as desired. ■

Sutner shows a similar result [8]. There is also a simple proof in terms of Fibonacci polynomials that also provides a divisibility relationship.

Proof. Since n divides nk , lemma 1.2 tells us $F_n(x)$ divides $F_{nk}(x)$. So,

$$\gcd(F_n(x), F_n(x+1)) \mid \gcd(F_{nk}(x), F_{nk}(x+1)).$$

So we may write

$$\gcd(F_{nk}(x), F_{nk}(x+1)) = \gcd(F_n(x), F_n(x+1))Q(x),$$

for some $Q(x) \in \mathbb{F}_2[x]$. Taking the degrees, we obtain $d(nk - 1) = d(n - 1) + \deg Q(x)$, and so the desired inequality follows. ■

Note that although the smaller GCD polynomial divides the larger, it's not the case that $d(n - 1)$ divides $d(nk - 1)$. For example, if $n = 5$ and $k = 6$, then $d(n - 1) = d(4) = 4$, which does not divide $d(nk - 1) = d(29) = 10$.

3 Number Theory

We may also understand $d(n)$ and the Fibonacci polynomials with some results from number theory. Many of the results in this section were first discussed by the author in [2] and [1]. However, this writing restates several proofs for clarity and to correct some small mistakes in those earlier proofs.

Lemma 3.1 (Ore). *Let a, b, c , and d be integers. Let (a, b) denote $\gcd(a, b)$. Then*

$$(ab, cd) = (a, c)(b, d) \left(\frac{a}{(a, c)}, \frac{d}{(b, d)} \right) \left(\frac{c}{(a, c)}, \frac{b}{(b, d)} \right).$$

Proof. See [11]. ■

Although Ore's result deals specifically with integers, both the integers and $\mathbb{F}_2[x]$ are Euclidean domains, so the result will still hold for our purposes.

Theorem 3.1. *For all $n \in \mathbb{N}$,*

$$d(2n + 1) = 2d(n) + \delta_n, \quad \delta_n = \begin{cases} 2 & 3 \mid n + 1 \\ 0 & \text{otherwise} \end{cases}.$$

Proof. Let (a, b) denote $\gcd(a, b)$. Applying lemmas 1.6 and 3.1,

$$\begin{aligned}
d(2n+1) &= \deg(F_{2n+2}(x), F_{2n+2}(x+1)) \\
&= \deg(xF_{n+1}(x)^2, (x+1)F_{n+1}(x+1)^2) \\
&= \deg(x, x+1) (F_{n+1}(x)^2, F_{n+1}(x+1)^2) \left(\frac{x+1}{(x, x+1)}, \frac{F_{n+1}(x)^2}{(F_{n+1}(x)^2, F_{n+1}(x+1)^2)} \right) \left(\frac{x}{(x, x+1)}, \frac{F_{n+1}(x+1)^2}{(F_{n+1}(x)^2, F_{n+1}(x+1)^2)} \right) \\
&= \deg(F_{n+1}(x), F_{n+1}(x+1))^2 \left(x+1, \frac{F_{n+1}(x)^2}{(F_{n+1}(x), F_{n+1}(x+1))^2} \right) \left(x, \frac{F_{n+1}(x+1)^2}{(F_{n+1}(x), F_{n+1}(x+1))^2} \right) \\
&= 2d(n) + 2 \deg \left(x, \frac{F_{n+1}(x+1)^2}{(F_{n+1}(x), F_{n+1}(x+1))^2} \right)
\end{aligned}$$

Thus, we have that

$$\delta_n = 2 \deg \left(x, \frac{F_{n+1}(x+1)^2}{(F_{n+1}(x), F_{n+1}(x+1))^2} \right).$$

So, δ_n is 2 exactly when $F_{n+1}(x)$ can be divided without remainder by $x+1$ more times than $F_{n+1}(x+1)$. When $n+1$ is not divisible by 3, corollary 1.2.1 tells us that $F_{n+1}(x)$ is not divisible by $x+1$, so $\delta_n = 0$. When $n+1$ is divisible by 3, let $n+1 = b2^k$ for $b, k \geq 0$ and b odd. Then b must be divisible by 3. Applying Lemma 1.6,

$$F_{n+1}(x) = x^{2^k-1} F_b(x)^{2^k} \quad F_{n+1}(x+1) = (x+1)^{2^k-1} F_b(x+1)^{2^k}.$$

Since b is odd, lemma 1.3 tells us $F_b(x)$ is the square of a square-free polynomial. Since b is a multiple of 3, corollary 1.2.1 tells us that $x+1$ divides $F_b(x)$, but $x+1$ does not divide $F_b(x+1)$. So,

$$F_{n+1}(x) = x^{2^k-1} (x+1)^{2^{k+1}} g(x)^{2^{k+1}} \quad F_{n+1}(x+1) = x^{2^{k+1}} (x+1)^{2^k-1} g(x+1)^{2^{k+1}},$$

for some $g(x) \in \mathbb{F}_2[x]$, where $g(x)$ and $g(x+1)$ are both square-free and divisible by neither x nor $x+1$. So, we see that $F_{n+1}(x)$ can be divided without remainder by $x+1$ more times than $F_{n+1}(x+1)$, so $\delta_n = 2$. ■

This result was first proven by Boyles in [2] and resolves a conjecture of Sutner's [9].

We can repeatedly apply theorem 3.1 to obtain explicit formulas for $d(n)$.

Corollary 3.1.1. *Write n as $n+1 = 2^s b$ for $s \geq 0$ and b odd. Then*

$$d(n) = 2^s d(b-1) + \begin{cases} 2(2^s - 1) & 3 \mid b \\ 0 & \text{otherwise} \end{cases}.$$

Proof. The result is a tautology for even n ($s = 0$ and $n+1 = b$), so we may continue assuming n is odd.

By induction on s . For $s = 0$,

$$\begin{aligned}
n+1 &= 2^0 b \\
n &= b-1 \\
d(n) &= d(b-1) \\
&= 2^0 d(b-1) + \begin{cases} 2(2^0 - 1) & 3 \mid b \\ 0 & \text{otherwise} \end{cases},
\end{aligned}$$

as desired. Assume that our claim holds for $s \geq 0$. Then for $n + 1 = 2^{s+1}b$,

$$\begin{aligned}
d(n) &= d(2^{s+1}b - 1) \\
&= d(2(2^s b - 1) + 1) \\
&= 2d(2^s b - 1) + \begin{cases} 2 & 3 \mid 2^s b \\ 0 & \text{otherwise} \end{cases} \\
&= 2 \left(2^s d(b - 1) + \begin{cases} 2(2^s - 1) & 3 \mid b \\ 0 & \text{otherwise} \end{cases} \right) + \begin{cases} 2 & 3 \mid 2^s b \\ 0 & \text{otherwise} \end{cases} \\
&= 2^{s+1}d(b - 1) + \begin{cases} 2(2^{s+1} - 1) & 3 \mid b \\ 0 & \text{otherwise} \end{cases} + \begin{cases} 2 & 3 \mid b \\ 0 & \text{otherwise} \end{cases} \\
&= 2^{s+1}d(b - 1) + \begin{cases} 2(2^{s+1} - 1) & 3 \mid b \\ 0 & \text{otherwise} \end{cases},
\end{aligned}$$

as desired. ■

We can also use theorem 3.1 to narrow down more when $d(n) = 2$.

Corollary 3.1.2. *If $d(n) = 2$, then $n \equiv -1 \pmod{6}$.*

Proof. Since 2 is not divisible by 4, corollary 1.4.1 tells us that n must be odd. So, we can write $n = 2^k b - 1$ for b odd and $k \geq 1$. Using corollary 3.1.1, we can define

$$D_i = d(2^i b - 1) = \begin{cases} 2^i d(b - 1) + 2(2^i - 1) & 3 \mid b \\ 2^i d(b - 1) & \text{otherwise} \end{cases}.$$

If $3 \nmid b$, then we must have $D_k = 2 = 2^k d(b - 1)$. However, corollary 1.4.1 tells us that $d(b - 1)$ is divisible by 4, which cannot be the case.

If $3 \mid b$, then we must have $D_k = 2 = 2^k d(b - 1) + 2(2^k - 1)$. If $k \geq 2$, then this expression is greater than 2, which cannot be the case. Thus, we must have $k = 1$ and $D_1 = 2 = 2d(b - 1) + 2$, which simplifies to $d(b - 1) = 0$. Thus we may write $b = 3j$.

Solving for n , we find

$$\begin{aligned}
n &= 2^k b - 1 \\
&= 2b - 1 \\
&= 6j - 1,
\end{aligned}$$

so $n \equiv -1 \pmod{6}$, as desired. ■

Definition 3.1. *Let P be a polynomial in $\mathbb{F}_2[x]$. Then*

$$\text{rank}(P) = \min \{r > 0 : P(x) \mid F_r(x)\}.$$

Hunziker, Machiavelo, and Park proved that every polynomial divides some $F_r(x)$ and therefore has a finite rank [5] (see Corollary 2.8).

We now show a relationship between divisibility of polynomials and divisibility of their ranks.

Lemma 3.2.

$$P(x) \mid F_m(x) \iff \text{rank}(P) \mid m.$$

Proof. Let $r = \text{rank}(P)$.

Assume that $P(x)$ divides $F_m(x)$. $P(x)$ must also divide $F_r(x)$. So by lemma 1.2,

$$\begin{aligned} P(x) &| \gcd(F_r(x), F_m(x)) \\ P(x) &| F_{\gcd(r, m)}(x). \end{aligned}$$

Since r is minimal, $\gcd(r, m) = r$, and so $r \mid m$.

Now assume that $r \mid m$. Then $\gcd(r, m) = r$. By lemma 1.2, $F_r(x) \mid F_m(x)$, and so $P(x) \mid F_m(x)$. ■

We can now use this recurrence result to prove certain even values of $d(n)$ are impossible.

Theorem 3.2. *The smallest even k such that for all $n \in \mathbb{N}$, $d(n) \neq k$ is $k = 26$.*

Proof. Recall from lemma 1.4 that $d(n)$ is always even. Through calculation, we can find that

$$\begin{array}{llllll} d(5) = 2 & d(4) = 4 & d(11) = 6 & d(16) = 8 & d(29) = 10 & d(84) = 12 \\ d(23) = 14 & d(19) = 16 & d(101) = 18 & d(30) = 20 & d(59) = 22 & d(62) = 24. \end{array}$$

We will show than $d(n) = 26$ is impossible.

Assume for the sake of contradiction that there exists $n \in \mathbb{N}$ such that $d(n) = 26$. Since 26 is not divisible by 4, corollary 1.4.1 that n must be odd. So, we may write $n = 2m + 1$. By theorem 3.1,

$$26 = d(n) = d(2m + 1) = 2d(m) + \begin{cases} 2 & 3 \mid m + 1 \\ 0 & \text{otherwise} \end{cases}.$$

Since lemma 1.4 tells us that $d(m) = 13$ is not possible, we must have $d(m) = 12$ and $m + 1$ is divisible by 3. Let $m + 1 = 2^s b$ for odd b and $s \geq 0$. By corollary 3.1.1,

$$d(m) = 2^s d(b - 1) + 2(2^s - 1) = 12.$$

The only possible integer solutions here are $d(b - 1) = 12$ and $s = 0$ or $d(b - 1) = 5$ and $s = 1$. Since lemma 1.4 tells us that $d(b - 1) = 5$ is impossible, the only remaining possibility is that $d(m) = d(b - 1) = 12$ and $s = 0$. Thus, m must be even.

We know from lemma 1.1 that

$$d(m) = \deg(\gcd(F_{m+1}(x), F_{m+1}(x + 1))).$$

Since $m + 1$ is odd lemma 1.3 tells us that there exists some square-free $R_{m+1}(x)$ such that

$$F_{m+1}(x) = R_{m+1}(x)^2.$$

Define $H_{m+1}(x)$

$$H_{m+1}(x) = \gcd(R_{m+1}(x), R_{m+1}(x + 1)) \quad \gcd(F_{m+1}(x), F_{m+1}(x + 1)) = H(x)^2.$$

Since $R_{m+1}(x)$ is square-free, then $H_{m+1}(x)$ is too. Since $d(m) = 12$, $\deg H_{m+1}(x) = 6$. Notice too that $H_{m+1}(x) = H_{m+1}(x + 1)$, so our work in lemma 1.5 tells us that for $y = x^2 + x$, there exists square-free a cubic polynomial $h_{m+1}(y) \in \mathbb{F}_2[y]$ such that

$$H_{m+1}(x) = h_{m+1}(y).$$

There are only three possible square-free cubic polynomials in $\mathbb{F}_2[y]$ that are also 1 at $y = 0$:

1. $C_1(y) = y^3 + 1$

2. $C_2(y) = y^3 + y + 1$
3. $C_3(y) = y^3 + y^2 + 1$.

We will handle these case-by-case.

Calculating the rank of our three candidates for $h_{m+1}(x)$,

$$\begin{aligned}\text{rank}((x^2 + x)^3 + 1) &= 85 \\ \text{rank}((x^2 + x)^3 + (x^2 + x) + 1) &= 63 \\ \text{rank}((x^2 + x)^3 + (x^2 + x)^2 + 1) &= 63\end{aligned}$$

By lemma 3.2, if either rank-63 polynomials C_2 or C_3 divides $R_{m+1}(x)$, then 63 divides $m + 1$. However, both polynomials would divide $R_{m+1}(x)$. C_2 and C_3 are coprime and each of degree 6 in x , so $\deg H_{m+1}(x) = \deg C_2(x) + \deg C_3(x) = 12$, contradicting our assumption that the degree was 6. If C_1 divides R_{m+1} , then 85 divides $m + 1$. Combining this with the fact that 3 divides $m + 1$ we showed earlier, we see that 255 must divide $m + 1$.

Define

$$P(x) = x^4 + x^3 + 1 \quad Q(x) = x^4 + x^3 + x^2 + x + 1.$$

P and Q are closed under translation ($P(x + 1) = Q(x)$), coprime to each other, and coprime with C_1 . The ranks of $P(x)$ and $Q(x)$ are 15 and 17 respectively. By lemma 3.2, since both 15 and 17 divide 255, and 255 divides $m + 1$, then both $P(x)$ and $Q(x)$ must divide $H_{m+1}(x)$. Therefore,

$$\deg H_{m+1}(x) \geq \deg C_1(x) + \deg P(x) + \deg Q(x) = 14,$$

which contradicts our assumption that the degree was 6.

Thus, there is no $3 \mid m + 1$ such that $d(m) = 12$ and therefore no n such that $d(n) = 26$. ■

Now that we know that one impossible even value of $d(n)$ exists, we can show an entire family of impossible even values also exists.

Corollary 3.2.1. *Suppose $a \in \mathbb{N}$ is even and an impossible value of $d(n)$. Then $2a + 2$ is also an impossible value of $d(n)$.*

Proof. Suppose for the sake of contradiction that there exists n such that $d(n) = 2a + 2$. Since $2a + 2$ is not divisible by 4, then corollary 1.4.1 tells us that n must be odd, so we may write $n = 2m + 1$. So, by theorem 3.1,

$$2a + 2 = d(n) = d(2m + 1) = 2d(m) + \delta_m \quad \delta_m = \begin{cases} 2 & 3 \mid m + 1 \\ 0 & \text{otherwise} \end{cases}.$$

If $\delta_m = 0$, then $2d(m)$ is divisible by 4, which cannot be the case since we know the value is $2a + 2$. If $\delta_m = 2$, then $d(m) = a$, contradicting the impossibility of a as an output of d . ■

Expanding the recurrence, if a is an impossible value, then $2^t(a + 2) - 2$ is also impossible for all $t \geq 0$. In other words, 26, 54, 110, 222, 446, ... are also all impossible values of $d(n)$.

4 Finite Fields

Although we have already been working in $\mathbb{F}_2$, we will now more directly use results from the study of finite fields.

Lemma 4.1. *Let p be an odd prime. Define $h = \text{ord}_p(2)$ and $c = v_p(2^h - 1)$, where $v_p(a)$ is the p -adic valuation of a . Then for every $j \geq 1$,*

$$\text{ord}_{p^j}(2) = hp^{\max(0, j-c)}.$$

Proof. Suppose $2^r \equiv 1 \pmod{p^j}$. Then h divides r , so we may write $r = ht$. By the Lifting the Exponent Lemma,

$$\begin{aligned} v_p(2^r - 1) &= v_p(2^{ht} - 1) \\ &= v_p(2^h - 1) + v_p(t) \\ &= c + v_p(t). \end{aligned}$$

So, $p^j \mid 2^{ht} - 1$ exactly when $c + v_p(t) \geq j$. The smallest possible t satisfying this inequality is $t = p^{\max(0, j-c)}$, so

$$\text{ord}_{p^j}(2) = hp^{\max(0, j-c)},$$

as desired. ■

Definition 4.1. *A prime number p is a Wieferich prime when*

$$p^2 \mid 2^{p-1} - 1.$$

Although infinitely any Wieferich primes are conjectured to exist, the only ones known to date are 1093 and 3511 [4], and any larger ones must be greater than 2^{64} [3].

We will show an equivalent definition of Wieferich primes.

Lemma 4.2. *A odd prime number p is a Wieferich prime if and only if*

$$p^2 \mid 2^{\text{ord}_p(2)} - 1.$$

Proof. Let $n = \text{ord}_p(2)$. Then $p - 1 = kn$ for some natural number k , and

$$2^{p-1} - 1 = 2^{kn} - 1.$$

So, by the Lifting the Exponent lemma,

$$v_p(2^{kn} - 1) = v_p(2^n - 1) + v_p(k).$$

However, $k = (p - 1)/n$ is smaller than p and thus not divisible by p , so $v_p(k) = 0$. Thus,

$$v_p(2^{p-1} - 1) = v_p(2^n - 1).$$

So, $p^2 \mid 2^{p-1} - 1$ if and only if $p^2 \mid 2^{\text{ord}_p(2)} - 1$, as desired. ■

Corollary 4.2.1. *Let p be an odd prime. If p isn't a Wieferich prime, then*

$$\text{ord}_{p^j}(2) = \text{ord}_p(2) p^{j-1}.$$

Proof. Lemma 4.1 tells us that

$$\text{ord}_{p^j}(2) = \text{ord}_p(2) p^{\max(0, j-c)}, \quad c = v_p(2^{\text{ord}_p(2)} - 1).$$

Thus it is sufficient to show that $c = 1$. By definition of $\text{ord}_p(2)$, $c \geq 1$. p is not a Wieferich prime, so by lemma 4.2, for all $t \geq 2$,

$$p^t \nmid 2^{\text{ord}_p(2)} - 1.$$

Thus $c < 2$ and therefore must be exactly 1, as desired. ■

Definition 4.2. Let b be coprime to n . Then “signed order” of b modulo n is

$$\rho_b(n) = \min \{r > 0 : b^r \equiv \pm 1 \pmod{n}\}.$$

This is also sometimes called the “index” of ± 1 relative to b , written $\text{ind}_b(\pm 1)$.

Lemma 4.3. Let p be an odd prime, $j \geq 1$, and $H_j = \text{ord}_{p^j}(2)$. Then

$$\rho_2(p^j) = \begin{cases} H_j/2 & H_j \text{ is even} \\ H_j & H_j \text{ is odd} \end{cases}.$$

Moreover, when H_j is even,

$$2^{H_j/2} \equiv -1 \pmod{p^j}.$$

Proof. Assume H_j is even. Then by definition of H_j ,

$$\left(2^{H_j/2}\right)^2 \equiv 1 \pmod{p^j}.$$

Thus the only solutions are $2^{H_j/2} \equiv \pm 1 \pmod{p^j}$. However, 1 cannot be a solution; otherwise, H_j would not be minimal. So,

$$2^{H_j/2} \equiv -1 \pmod{p^j}.$$

Notice that if $r \leq H_j/2$, and $2^r \equiv -1 \pmod{p^j}$, then $2^{2r} \equiv 1 \pmod{p^j}$; since H_j is minimal, $H_j \leq 2r$. Thus to satisfy both constraints we must have $H_j/2 = r$.

Now assume H_j is odd. Notice that by definition,

$$H_j = \text{ord}_{p^j}(2) = \min \{h > 0 : 2^h \equiv 1 \pmod{p^j}\},$$

so smaller solutions can only be of the form $2^r \equiv -1 \pmod{p^j}$. Then $2^{2r} \equiv 1 \pmod{p^j}$, which means $H_j \mid 2r$, and since H_j is odd, $H_j \mid r$. Thus $H_j \leq r$, so H_j is minimal, as desired. ■

We can use this notion of a signed order to find the extension degree of extensions of $\mathbb{F}_2$.

Lemma 4.4. Let $M > 1$ be odd; let ζ have exact order M ; set $\alpha = \zeta + \zeta^{-1}$. Then

$$[\mathbb{F}_2(\alpha) : \mathbb{F}_2] = \rho_2(M).$$

Proof. A standard finite field fact says

$$[\mathbb{F}_2(\alpha) : \mathbb{F}_2] = \min \{r > 0 : \alpha^{2^r} = \alpha\}.$$

By definition of α , $\alpha^{2^r} = \zeta^{2^r} + \zeta^{-2^r}$, so $\alpha^{2^r} = \alpha$ is equivalent to $\zeta^{2^r} + \zeta^{-2^r} = \zeta + \zeta^{-1}$.

For any non-zero $a, b \in \overline{\mathbb{F}_2}$, $a + a^{-1} = b + b^{-1} \iff (a+b)(ab+1) = 0$. So, either $a = b$ or $a = b^{-1}$. Taking $a = \zeta^{2^r}$ and $b = \zeta$, we obtain

$$\zeta^{2^r} = \zeta \quad \text{or} \quad \zeta^{2^r} = \zeta^{-1}.$$

Since ζ has exact order M , we can equivalently write

$$2^r \equiv 1 \pmod{M} \quad \text{or} \quad 2^r \equiv -1 \pmod{M}.$$

So, $\alpha^{2^r} = \alpha \iff 2^r \equiv \pm 1 \pmod{M}$, and

$$[\mathbb{F}_2(\alpha) : \mathbb{F}_2] = \min \{r > 0 : \alpha^{2^r} = \alpha\} = \rho_2(M),$$

as desired. ■

Lemma 4.5. *Let a be algebraic over $\mathbb{F}_q$. Then for every $z \in \mathbb{F}_q(a)$,*

$$\mathrm{Tr}_{\mathbb{F}_q(a)/\mathbb{F}_q}(z) = \sum_{r=0}^{[\mathbb{F}_q(a):\mathbb{F}_q]-1} z^{q^r}.$$

Proof. See [7]. ■

Corollary 4.5.1.

$$\mathrm{Tr}_{\mathbb{F}_q(a)/\mathbb{F}_q}(1) = [\mathbb{F}_q(a) : \mathbb{F}_q].$$

Proof. Immediate from evaluating lemma 4.5 at $z = 1$. ■

Lemma 4.6. *Let $M > 1$ be odd, and let ζ have exact order M . Let $\alpha = \zeta + \zeta^{-1}$. Define*

$$G_M = \{\pm 2^r \pmod{M} : r \geq 0\} \subseteq (\mathbb{Z}/M\mathbb{Z})^\times.$$

Then

$$\mathrm{Tr}_{\mathbb{F}_2(\alpha)/\mathbb{F}_2}(\alpha) = \sum_{u \in G_M} \zeta^u.$$

Proof. Lemma 4.4 showed that

$$[\mathbb{F}_2(\alpha) : \mathbb{F}_2] = \rho_2(M).$$

So, by lemma 4.5,

$$\mathrm{Tr}_{\mathbb{F}_2(\alpha)/\mathbb{F}_2}(\alpha) = \sum_{r=0}^{\rho_2(M)-1} \zeta^{2^r} + \zeta^{-2^r}.$$

By the definition of $\rho_2(M)$, the residues $2^0, -2^0, 2^1, -2^1, \dots, 2^{\rho_2(M)-1}, -2^{\rho_2(M)-1}$ are all distinct modulo M . Otherwise, we would obtain $2^r \equiv \pm 2^s \pmod{M}$, or equivalently, $2^{r-s} \equiv \pm 1 \pmod{M}$ for some $0 \leq s < r < d$, which contradicts $\rho_2(M)$ being minimal.

The residues are exactly the elements of G_M , so

$$\mathrm{Tr}_{\mathbb{F}_2(\alpha)/\mathbb{F}_2}(\alpha) = \sum_{u \in G_M} \zeta^u,$$

as desired. ■

Corollary 4.6.1. *If $G_M = (\mathbb{Z}/M\mathbb{Z})^\times$, then*

$$\mathrm{Tr}_{\mathbb{F}_2(\alpha)/\mathbb{F}_2}(\alpha) = \mu(M) \pmod{2},$$

where μ is the number-theoretic Möbius function.

Proof. From lemma 4.6, we know that

$$\mathrm{Tr}_{\mathbb{F}_2(\alpha)/\mathbb{F}_2}(\alpha) = \sum_{u \in G_M} \zeta^u.$$

When $G_M = (\mathbb{Z}/M\mathbb{Z})^\times$, we also have

$$\mathrm{Tr}_{\mathbb{F}_2(\alpha)/\mathbb{F}_2}(\alpha) = \sum_{0 \leq u < M, \gcd(u, M)=1} \zeta^u = \mu(M) \pmod{2},$$

by Ramanujan's sum identity of roots of unity, as desired. ■

Now we may combine together many of our partial results to state a more significant result.

Theorem 4.1. *If p is not a Wieferich prime, then for all $k \geq 1$,*

$$d(p^k - 1) = d(p - 1).$$

Proof. Let p be a non-Wieferich prime. The case for $p = 2$ was shown in [9], so we will consider the case when p is odd. For $1 \leq j \leq k$, let

$$R_j = \{\zeta + \zeta^{-1} : \text{ord}(\zeta) = p^j\}.$$

By lemma 1.7,

$$\text{roots } F_{p^k} = \bigcup_{j=1}^k R_j. \quad (1)$$

Furthermore, since p^k is odd, corollary 1.3.1 and lemma 1.3 tell us that F_{p^k} is the square of a square-free polynomial. In other words, F_{p^k} has $(p^k - 1)/2$ distinct roots. Let $h = \text{ord}_p(2)$. Since p is not Wieferich, corollary 4.2.1 tells us that

$$\text{ord}_{p^j}(2) = \text{ord}_p(2) p^{j-1},$$

and lemma 4.3 tells us that

$$\rho_2(p^j) = \rho_2(p) p^{j-1}.$$

So, by lemma 4.4, if $\alpha \in R_j$,

$$[\mathbb{F}_2(\alpha) : \mathbb{F}_2] = \rho_2(p) p^{j-1}.$$

Suppose both α and $\alpha + 1$ are roots of F_{p^k} with $\alpha \in R_j$ and $\alpha + 1 \in R_\ell$. Since $\mathbb{F}_2(\alpha) = \mathbb{F}_2(\alpha + 1)$, that is they are the same field with the same degree, α and $\alpha + 1$ must both be members of the same R_j and so $j = \ell$.

Suppose for the sake of contradiction that α and $\alpha + 1$ are both members of R_j for some $j \geq 2$. Set $M = p^j$ and $m = p^{j-1}$ and choose a primitive M th root ζ such that $\alpha = \zeta + \zeta^{-1}$. Since $\alpha + 1$ is also in R_j , there must exist s such that $\gcd(s, p) = 1$ and $\alpha + 1 = \zeta^s + \zeta^{-s}$. Then

$$1 + \alpha + (\alpha + 1) = 1 + \zeta + \zeta^{-1} + \zeta^s + \zeta^{-s} = 0.$$

Equivalently, ζ is a root of

$$P_s(x) = 1 + x + x^{M-1} + x^s + x^{M-s}.$$

Notice that ζ^m is a primitive p th root. Let $f(x)$ be the minimal polynomial of ζ^m over $\mathbb{F}_2$. The degree of $f(x)$ is $\text{ord}_p(2)$. Since $f(\zeta^m) = 0$, ζ is a root of $f(x^m)$. The degree of $f(x^m)$ is $\text{ord}_p(2) m = \text{ord}_{p^j}(2)$. This is also the degree of the minimal polynomial of ζ and so

$$f(x^m) \mid P_s(x).$$

So, every polynomial can be decomposed by exponents modulo m :

$$P_s(x) = \sum_{a=0}^{m-1} x^a P_a(x^m). \quad (2)$$

Since $f(x^m) \mid P_s(x)$, each $P_a(y)$ must be divisible by $f(y)$. However, only exponent 0 among 0, 1, $M - 1$, s , $M - s$ is divisible by m , so $P_0(y) = 1$, a contradiction. Instead, we must conclude that no common roots of $F_{p^k}(x)$ and $F_{p^k}(x + 1)$ are in R_j for $j \geq 2$.

So the only common roots of $F_{p^k}(x)$ and $F_{p^k}(x + 1)$ must come from R_1 and so we must conclude

$$d(p^k - 1) = d(p - 1).$$

■

We can combine this result with our recurrence for $d(2n+1)$.

Corollary 4.6.2. *Let p be an odd non-Wieferich prime. Then for all $n, m \geq 0$,*

$$d(2^n p^m - 1) = \begin{cases} 0 & m = 0 \\ 2^n d(p-1) & m \geq 1, p \neq 3 \\ 2^{n+1} - 2 & m \geq 1, p = 3 \end{cases}.$$

Proof. The $m = 0$ case reduces to what Sutner already showed [9]. For the other cases, we first apply corollary 3.1.1 to obtain

$$d(2^n p^m - 1) = \begin{cases} 0 & m = 0 \\ 2^n d(p^m - 1) & m \geq 1, p \neq 3 \\ 2^{n+1} - 2 & m \geq 1, p = 3 \end{cases}.$$

We then apply theorem 4.1 to change $d(p^m - 1)$ to $d(p - 1)$. ■

Corollary 4.6.3. *For all $n, m \geq 1$, the following relationships hold*

1. $d(3^n - 1) = 0$
2. $d(5^n - 1) = 4$
3. $d(7^n - 1) = 0$
4. $d(17^n - 1) = 8$
5. $d(31^n - 1) = 20$
6. $d(127^n - 1) = 56$
7. $d(257^n - 1) = 144$
8. $d(683^n - 1) = 132$
9. $d(8191^n - 1) = 4004$
10. $d(2^n 3^m - 1) = 2^{n+1} - 2$
11. $d(2^n 5^m - 1) = 2^{n+2}$
12. $d(2^n 7^m - 1) = 0$
13. $d(2^n 17^m - 1) = 2^{n+3}$
14. $d(2^n 31^m - 1) = 2^{n+2} \cdot 5$
15. $d(2^n 127^m - 1) = 2^{n+3} \cdot 7$
16. $d(2^n 257^m - 1) = 2^{n+4} \cdot 9$
17. $d(2^n 683^m - 1) = 2^{n+2} \cdot 33$
18. $d(2^n 8191^m - 1) = 2^{n+2} \cdot 7 \cdot 11 \cdot 13$

You may have noticed that for many sizes, we have $d(p-1) = 0$. We can actually classify some p for which this is the case.

Theorem 4.2. *Let p be a prime one less than a multiple of 4 where*

$$\{\pm 2^r \pmod{p} : r \geq 0\} = (\mathbb{Z}/p\mathbb{Z})^\times.$$

Then $d(p-1) = 0$.

Proof. Assume for the sake of contradiction that $d(p-1) \neq 0$. Then $F_p(x)$ and $F_p(x+1)$ have a common factor and thus a common root α , so

$$F_p(\alpha) = F_p(\alpha+1) = 0.$$

By lemma 1.7, there must exist non-trivial p th roots of unity ζ and η such that

$$\alpha = \zeta + \zeta^{-1} \quad \alpha + 1 = \eta + \eta^{-1}.$$

By lemmas 4.4 and 4.3,

$$[\mathbb{F}_2(\alpha) : \mathbb{F}_2] = [\mathbb{F}_2(\alpha+1) : \mathbb{F}_2] = \rho_2(p) = \frac{p-1}{2}.$$

Since $p \equiv 3 \pmod{4}$, this value is odd. Applying the linearity of the trace and corollary 4.5.1,

$$\mathrm{Tr}_{\mathbb{F}_2(\alpha)/\mathbb{F}_2}(\alpha+1) = \mathrm{Tr}_{\mathbb{F}_2(\alpha)/\mathbb{F}_2}(\alpha) + \mathrm{Tr}_{\mathbb{F}_2(\alpha)/\mathbb{F}_2}(1) = \mathrm{Tr}_{\mathbb{F}_2(\alpha)/\mathbb{F}_2}(\alpha) + 1 = 0.$$

However, we can see that $\mathbb{F}_2(\alpha+1) = \mathbb{F}_2(\alpha)$ (α and $\alpha+1$ can be obtained from each other by adding 1), and so by corollary 4.6.1,

$$\mathrm{Tr}_{\mathbb{F}_2(\alpha+1)/\mathbb{F}_2}(\alpha+1) = \mathrm{Tr}_{\mathbb{F}_2(\alpha)/\mathbb{F}_2}(\alpha) = \mu(p) \pmod{2} = 1.$$

Thus we have reached a contradiction. ■

References

- [1] William Boyles, *Most clicks problem in lights out*, 2022, arXiv:2201.03452 [math.CO].
- [2] ———, *Resolution to sutner’s conjecture*, 2022, arXiv:2202.09878 [math.CO].
- [3] PrimeGrid Collaboration, *Ww statistics*, PrimeGrid, 2026, Online resource.
- [4] Richard Crandall, Karl Dilcher, and Carl Pomerance, *A search for wieferich and wilson primes*, Mathematics of Computation **66** (1997), no. 217, 433–449.
- [5] Markus Hunziker, António Machiavelo, and Jihun Park, *Chebyshev polynomials over finite fields and reversibility of σ -automata on square grids*, Theoretical Computer Science **320** (2004), no. 2, 465–483.
- [6] Donald Knuth, *The art of computer programming, volume 4A: Combinatorial algorithms, part 1*, Addison-Wesley, 2011, Section 7.1.3, Exercises 190-194.
- [7] Rudolf Lidl and Harald Niederreiter, *Finite fields*, 2nd ed., p. 54, Cambridge University Press, 1997.
- [8] Klaus Sutner, *On σ -automata*, Complex Systems **2** (1988), no. 1, 1–28.
- [9] ———, *Linear cellular automata and the Garden-of-Eden*, The Mathematical Intelligencer **11** (1989), no. 2, 49–53.
- [10] ———, *σ -automata and chebyshev-polynomials*, Theoretical Computer Science **230** (2000), no. 1, 49–73.
- [11] Øystein Ore, *Number theory and its history*, p. 109, McGraw-Hill, 1948, Section 5-4, Problem 2.