

Fibonacci Polynomials & *Lights Out*

William Boyles *

August 2, 2026

Abstract

Consider a game played on a simple graph $G = (V, E)$ where each vertex consists of clickable light. A move is clicking any $v \in V$, which toggles the on/off state of v and its neighbors. Given an initial configuration, one wins the game by finding a sequence of moves that turns off all of the lights. This game was commercially available from Tiger Electronics as *Lights Out* for G a 5×5 grid. We explore this game for an $n \times n$ grid and show several new results.

1 Fibonacci Polynomials

Definition 1.1 (Fibonacci Polynomial). *Let $F_n(x)$ be the polynomial in the ring $\mathbb{F}_2[x]$ defined recursively by*

$$F_n(x) = \begin{cases} 0 & n = 0 \\ 1 & n = 1 \\ xF_{n-1}(x) + F_{n-2}(x) & \text{otherwise} \end{cases}.$$

These polynomials are called "Fibonacci" because when defined over the real numbers, $F_n(1)$ is the n th Fibonacci number.

In $\mathbb{F}_2[x]$, the coefficients of $F_n(x)$ are either 1 or 0, so we can visualize them by coloring squares black or white to represent the coefficients. For example,

$$F_6(x) = 1x^5 + 0x^4 + 0x^3 + 0x^2 + 1x + 0 = \blacksquare \square \square \square \blacksquare \square.$$

Plotting $F_1(x)$ through $F_{128}(x)$ in the same way, aligning terms of the same degree, we see a Sierpinski triangle.

*Independent (williamboyles22900@gmail.com)




Figure 1: $F_1(x)$ through $F_{128}(x)$, right-aligned.

Using a well-known connection between the Sierpinski triangle and the parity of binomial coefficients, we may see that

$$F_n(x) = \sum_{i=0}^{n-1} \binom{n+i}{2i+1} x^i \pmod{2}.$$

The above construction and formula was also shown by Sutner [5]. Sutner also showed the following relationship between square *Lights Out* grids and Fibonacci Polynomials.

Lemma 1.1 (Sutner). *Let $G = (V, E)$ be an $n \times m$ grid graph. Let $\Phi_G : P(V) \rightarrow P(V)$ be the function that maps which lights are clicked to which lights change state. Then*

$$\dim(\ker \Phi_G) = \deg(\gcd(F_{n+1}(x), F_{m+1}(x+1))) = \deg(\gcd(F_{n+1}(x+1), F_{m+1}(x))).$$

Proof. See [5] theorem 5.1. ■

Sutner and others also shows several divisibility rules for these polynomials

Lemma 1.2 (Hunziker, Machiavelo, and Park). *A polynomial $\tau(x)$ in $\mathbb{F}_2[x]$ divides both $F_n(x)$ and $F_m(x)$ if and only if it divides $F_{\gcd(m,n)}(x)$. In particular,*

$$\gcd(F_m(x), F_n(x)) = F_{\gcd(m,n)}(x).$$

Proof. See [1] proposition 2.2, [5] proposition 2.1, and [2]. ■

Corollary 1.2.1. *The following are true:*

- (i) x divides $f_n(x)$ if and only if $n \equiv 0 \pmod{2}$.
- (ii) $x + 1$ divides $f_n(x + 1)$ if and only if $n \equiv 0 \pmod{2}$.
- (iii) $x + 1$ divides $f_n(x)$ if and only if $n \equiv 0 \pmod{3}$.
- (iv) x divides $f_n(x + 1)$ if and only if $n \equiv 0 \pmod{3}$.

Proof. Notice,

- (i) We find that $f_2(x) = x$ is the smallest Fibonacci polynomial divisible by x , so we apply lemma 1.2 to get the desired result.
 - (ii) Follows from (i) by substituting $x + 1$ for x .
 - (iii) We find that $f_3(x) = (x + 1)^2$ is the smallest Fibonacci polynomial divisible by $x + 1$, so we apply lemma 1.2 to get the desired result.
 - (iv) Follows from (iii) by substituting $x + 1$ for x .
-

Lemma 1.3 (Hunziker, Machiavelo, and Park). *If n is odd, then $F_n(x)$ is the square of a square-free polynomial.*

Proof. See [1] proposition 2.11 and [5] Theorem 2.1 claim 2. ■

Corollary 1.3.1. *If n is odd, then every root of $F_n(x)$ has multiplicity 2.*

Proof. Let n be odd and α a root of $F_n(x)$. By 1.3, there exists $\tau_n \in \mathbb{F}_2[x]$ such that

$$F_n(x) = \tau_n(x)^2,$$

and $\tau(x)$ is square-free. Since α is a root of $F_n(x)$, it must also be a root of $\tau_n(x)$ and thus has multiplicity 2. ■

We can now prove some of our own facts about these polynomials and their GCDs, particularly on square grids.

Lemma 1.4. *Let $d(n) = \dim(\ker \Phi_G)$ for G an $n \times n$ grid graph. Then $d(n)$ is always even.*

Proof. Let

$$G_n(x) = \gcd(F_n(x), F_n(x + 1)).$$

By lemma 1.1, we know that $d(n) = \deg G_{n+1}(x)$ and $G_n(x) = G_n(x + 1)$.

$y = x(x + 1)$ is also unchanged by this $x \mapsto x + 1$ translation. It is well-known that every polynomial $P(x)$ in $\mathbb{F}_2[x]$ can be uniquely written as

$$P(x) = A(y) + xB(y) \quad A, B \in \mathbb{F}_2[y].$$

We can then determine exactly when P is also unchanged by translation.

$$\begin{aligned} P(x+1) &= P(x) \\ A(y) + (x+1)B(y) &= A(y) + xB(y) \\ B(y) &= 0. \end{aligned}$$

In other words, $P(x)$ is unchanged by translation when P is a polynomial in $x^2 + x$.

Since G_n is also unchanged by translation, there exists $H_n \in \mathbb{F}_2[x]$ such that

$$G_n(x) = H_n(x^2 + x).$$

So,

$$d(n) = \deg G_{n+1}(x) = 2 \deg H_{n+1}(x)$$

and thus $d(n)$ is even. ■

Corollary 1.4.1. *If n is even, then $d(n)$ is divisible by 4.*

Proof. If n is even, then $n+1$ is odd. Then by 1.3, $F_{n+1}(x)$ and $F_{n+1}(x+1)$ are squares of square-free polynomials $\tau(x)$ and $\tau(x+1)$.

$$\begin{aligned} G_{n+1}(x) &= \gcd(F_{n+1}(x), F_{n+1}(x+1)) \\ &= \gcd(\tau_{n+1}(x)^2, \tau_{n+1}(x+1)^2) \\ &= \gcd(\tau_{n+1}(x), \tau_{n+1}(x+1))^2 \\ \sqrt{G_{n+1}(x)} &= \gcd(\tau_{n+1}(x), \tau_{n+1}(x+1)) \end{aligned}$$

We can see from the right-hand side of the equation that this function is also unaffected by translation. So, by lemma 1.4, there exists $H_{n+1} \in \mathbb{F}_2[x]$ such that

$$\begin{aligned} \sqrt{G_{n+1}(x)} &= H_{n+1}(x^2 + x) \\ G_{n+1}(x) &= H_{n+1}(x^2 + x)^2. \end{aligned}$$

So,

$$d(n) = \deg G_{n+1}(x) = 4 \deg H_{n+1}(x),$$

as desired. ■

Note that the implication does not hold in the other direction: $d(n)$ also be divisible by 4 when n is odd. For example, $d(9) = 8$.

Lemma 1.5. *Let $x = t + t^{-1}$. Then for $t \neq 0$,*

$$xF_n(x) = t^n + t^{-n}.$$

Proof. Define $G_n(t) = t^{n+1} + t^{-(n+1)}$. Then

$$\begin{aligned} xG_{n-1}(t) + G_{n-2}(t) &= (t + t^{-1})(t^n + t^{-n}) + (t^{n-1} + t^{-(n-1)}) \\ &= t^{n+1} + t^{-(n-1)} + t^{n-1} + t^{-(n+1)} + t^{n-1} + t^{-(n-1)} \\ &= t^{n+1} + t^{-(n+1)} \\ &= G_n. \end{aligned}$$

Thus we see that $G_n(t)$ satisfies the same recurrence relation as $F_n(x)$. $xF_n(x)$ also satisfies the same recurrence relation.

$$\begin{aligned} xF_n(x) &= x(xF_{n-1}(x) + F_{n-2}(x)) \\ &= x(xF_{n-1}(x)) + xF_{n-2}(x). \end{aligned}$$

Now we compute $xF_1(x)$ and $xF_2(x)$ in terms of t and G :

$$\begin{aligned} xF_1(x) &= x \\ &= t + t^{-1} \\ &= G_0(t), \end{aligned}$$

and

$$\begin{aligned} xF_2(x) &= x^2 \\ &= (t + t^{-1})^2 \\ &= t^2 + t^{-2} \\ &= G_1(t). \end{aligned}$$

Since the first two terms match and they satisfy the same recurrence relation then we've shown by induction that for $t \neq 0$,

$$xF_n(x) = G_{n-1}(t) = t^n + t^{-n}.$$

■

This was also proven in [1], who also prove the following identities.

Lemma 1.6 (Hunziker, Machiavelo, and Park). *Let $n = 2^k b$ where $b, k \in \mathbb{N}$. Then*

$$F_n(x) = x^{2^k-1} F_b(x)^{2^k}.$$

Proof. See [1] lemma 2.6.

■

Lemma 1.7. *The distinct roots of $F_n(x)$ are $\{\zeta + \zeta^{-1} : \zeta^n = 1, \zeta \neq 1\}$, plus 0 when n is even.*

Proof. Notice that $F_n(0) = F_{n-2}(0)$, so

$$F_n(0) = \begin{cases} 1 & n \text{ odd} \\ 0 & n \text{ even} \end{cases}.$$

Thus 0 is a root of $F_n(x)$ whenever n is even.

Let α be a non-zero root of F_n . Choose ζ satisfying

$$\zeta^2 + \alpha\zeta + 1 = 0.$$

Notice that ζ cannot be 0. Dividing by ζ ,

$$\begin{aligned} \zeta + \alpha + \zeta^{-1} &= 0 \\ \alpha &= \zeta + \zeta^{-1}. \end{aligned}$$

By, Lemma 1.5,

$$\begin{aligned} 0 &= \alpha F_n(\alpha) \\ &= \zeta^n + \zeta^{-n} \\ &= \zeta^{2n} + 1 \\ &= (\zeta^n + 1)^2 \\ &= \zeta^n + 1 \\ \zeta^n &= 1. \end{aligned}$$

Thus the order of ζ divides n . Notice too that $\zeta \neq 1$; otherwise, $\alpha = 1 + 1^{-1} = 0$, which we already showed as not possible.

Conversely, if $\zeta^n = 1$ and $\zeta \neq 1$, then let $\alpha = \zeta + \zeta^{-1}$. Then we see $\alpha \neq 0$ and

$$\alpha F_n(\alpha) = \zeta^n + \zeta^{-n} = 1 + 1^{-1} = 0.$$

Thus, α is a root of F_n . ■

2 Even Parity Covers

$\ker \Phi_G$ may also be understood via parity covers and dominating sets.

Definition 2.1. *An even dominating set of a graph $G = (V, E)$ is a subset $V' \subseteq V$ such that every vertex contains an even number of vertices from V' in its neighborhood.*

Odd dominating sets can be analogously defined. Sutner specifically studied odd dominating sets as the solution to *Lights Out* when all lights are initially on. He showed that such an odd dominating set always exists for any finite graph [4].

Lemma 2.1. *$V' \in \ker \Phi_G$ if and only if V' is an even dominating set of G .*

Proof. Let $V' \in \ker \Phi_G$. Let $N(v)$ be the neighborhood of vertex v : v and its immediate neighbors. Then for every $v \in V$,

$$|N(v) \cap V'| \text{ is even.}$$

Thus, V' is an even dominating set of G . We can apply the same steps in reverse to show the converse. ■

Lemma 2.2. *Φ_G is linear with respect to symmetric difference. That is, for all $V_1, V_2 \subseteq V$,*

$$\Phi_G(V_1 \triangle V_2) = \Phi_G(V_1) \triangle \Phi_G(V_2).$$

Proof.

$$\begin{aligned} \Phi_G(V_1 \triangle V_2) &= \{v \in V : |N(v) \cap (V_1 \triangle V_2)| \text{ is odd}\} \\ &= (\{v \in V : |N(v) \cap V_1| \text{ is odd}\} \cup \{v \in V : |N(v) \cap V_2| \text{ is odd}\}) \setminus \{v \in V : |N(v) \cap (V_1 \cap V_2)| \text{ is odd}\} \\ &= \{v \in V : |N(v) \cap V_1| \text{ is odd}\} \triangle \{v \in V : |N(v) \cap V_2| \text{ is odd}\} \\ &= \Phi_G(V_1) \triangle \Phi_G(V_2) \end{aligned}$$
■

Corollary 2.2.1. *If V_1 and V_2 have the same image under Φ_G if and only if $V_1 \triangle V_2 \in \ker \Phi_G$.*

Proof. Assume that V_1 and V_2 have the same image under Φ_G . Then

$$\begin{aligned} \Phi_G(V_1 \triangle V_2) &= \Phi_G(V_1) \triangle \Phi_G(V_2) \\ &= \Phi_G(V_1) \triangle \Phi_G(V_1) \\ &= \emptyset. \end{aligned}$$

Thus, $V_1 \triangle V_2 \in \ker \Phi_G$. Now assume $V_1 \triangle V_2 \in \ker \Phi_G$. Then

$$\begin{aligned} \Phi_G(V_1) &= \Phi_G(V_1) \triangle \emptyset \\ &= \Phi_G(V_1) \triangle \Phi_G(V_1 \triangle V_2) \\ &= \Phi_G(V_1) \triangle \Phi_G(V_1) \triangle \Phi_G(V_2) \\ &= \Phi_G(V_2). \end{aligned}$$

Thus V_1 and V_2 have the same image under Φ_G . ■

We may now use our geometric understanding of $\ker \Phi_G$ as even parity covers to construct a lower bound on $d(n)$.

Theorem 2.1. *For all $n, k \in \mathbb{N}$,*

$$d(nk - 1) \geq d(n - 1).$$

Proof. Let G_L be an $nk - 1 \times nk - 1$ grid and G_ℓ be an $n - 1 \times n - 1$ grid. Since $nk - 1 = k(n - 1) + (k - 1)$, an G may be broken into a $k \times k$ grid of $n - 1 \times n - 1$ grids, each separated by a horizontal or vertical strip of size 1.

Let Q be an even dominating set of G_ℓ . Let R be Q tiles k times horizontally and vertically, with size 1 separators between each tile, onto G_L where each tile is the reflection across its neighboring separators, as shown below.




Figure 2: Pattern Q tiled 5 times.

Consider an vertex v in G_L . There are 3 cases.

1. v is within a tile. Then $|N(v) \cap R|$ is even because Q being is an even parity cover.
2. v is on a separator between two tiles. Then only two elements in $N(v)$, the vertices within a tile, are also in R . Because neighboring tiles are reflections of each other across separators, these two vertices are either both in R or both not in R . In either case, $|N(v) \cap R|$ is even.
3. v is in the intersection of two separators. Then all of $N(v)$ is in a separator and thus not in R , so $|N(v) \cap R| = 0$, which is even.

Since for every vertex v of G_L , $|N(v) \cap R|$ is even, so R is an even dominating set of G_L . Thus, for every even dominating set of G_ℓ , there exists an even dominating set of G_L obtained through tiling. This tiling function is also injective because we can recover Q by restricting R to just the top left tile. Thus,

$$d(nk - 1) \geq d(n - 1),$$

as desired. ■

Sutner shows a similar result [3]. There is also a simple proof in terms of Fibonacci polynomials that also provides a divisibility relationship.

Proof. Since n divides nk , lemma 1.2 tells us $F_n(x)$ divides $F_{nk}(x)$. So,

$$\gcd(F_n(x), F_n(x+1)) \mid \gcd(F_{nk}(x), F_{nk}(x+1)).$$

Taking the degrees, we obtain $d(n-1)$ divides $d(nk-1)$, as desired. ■

3 Number Theory

We may also understand $d(n)$ and the Fibonacci polynomials with some results from number theory.

Lemma 3.1 (Ore). *Let a, b, c , and d be integers. Let (a, b) denote $\gcd(a, b)$. Then*

$$(ad, cd) = (a, c)(b, d) \left(\frac{a}{(a, c)}, \frac{d}{(b, d)} \right) \left(\frac{c}{(a, c)}, \frac{b}{(b, d)} \right).$$

Proof. See [6]. ■

Although Ore's result deals specifically with integers, both the integers and $\mathbb{F}_2[x]$ are Euclidean domains, so the result will still hold for our purposes.

Theorem 3.1. *For all $n \in \mathbb{N}$,*

$$d(2n+1) = 2d(n) + \delta_n,$$

where

$$\delta_n = \begin{cases} 2 & 3 \mid n+1 \\ 0 & \text{otherwise} \end{cases}.$$

Proof. Let (a, b) denote $\gcd(a, b)$. Applying lemmas 1.6 and 3.1,

$$\begin{aligned} d(2n+1) &= \deg(F_{2n+2}(x), F_{2n+2}(x+1)) \\ &= \deg(xF_{n+1}(x)^2, (x+1)F_{n+1}(x+1)^2) \\ &= \deg(x, x+1) (F_{n+1}(x)^2, F_{n+1}(x+1)^2) \left(\frac{x+1}{(x, x+1)}, \frac{F_{n+1}(x)^2}{(F_{n+1}(x)^2, F_{n+1}(x+1)^2)} \right) \left(\frac{x}{(x, x+1)}, \frac{F_{n+1}(x+1)^2}{(F_{n+1}(x)^2, F_{n+1}(x+1)^2)} \right) \\ &= \deg(F_{n+1}(x), F_{n+1}(x+1))^2 \left(x+1, \frac{F_{n+1}(x)^2}{(F_{n+1}(x), F_{n+1}(x+1))^2} \right) \left(x, \frac{F_{n+1}(x+1)^2}{(F_{n+1}(x), F_{n+1}(x+1))^2} \right) \\ &= 2d(n) + 2 \deg \left(x, \frac{F_{n+1}(x+1)^2}{(F_{n+1}(x), F_{n+1}(x+1))^2} \right) \end{aligned}$$

Thus, we have that

$$\delta_n = 2 \deg \left(x, \frac{F_{n+1}(x+1)^2}{(F_{n+1}(x), F_{n+1}(x+1))^2} \right).$$

So, δ_n is 2 exactly when $F_{n+1}(x)$ can be divided without remainder by $x+1$ more times than $F_{n+1}(x+1)$. When $n+1$ is not divisible by 3, corollary 1.2.1 tells us that $F_{n+1}(x)$ is not divisible by $x+1$, so $\delta_n = 0$. When $n+1$ is divisible by 3, let $n+1 = b2^k$ for $b, k \geq 0$ and b odd. Then b must be divisible by 3. Applying Lemma 1.6,

$$F_{n+1}(x) = x^{2^k-1} F_b(x)^{2^k} \quad F_{n+1}(x+1) = (x+1)^{2^k-1} F_b(x+1)^{2^k}.$$

Since b is an odd multiple of 3, corollary 1.2.1 tells us that $x + 1$ divides $F_b(x)$, but $x + 1$ does not divide $F_b(x + 1)$. So,

$$F_{n+1}(x) = x^{2^k-1}(x+1)^{2^k}g(x)^{2^k} \quad F_{n+1}(x+1) = (x+1)^{2^k-1}x^{2^k}g(x+1)^{2^k},$$

for some $g(x) \in \mathbb{F}_2[x]$, where $g(x)$ and $g(x+1)$ are both divisible by neither x nor $x+1$. So, we see that $F_{n+1}(x)$ can be divided without remainder by $x+1$ more times than $f_{n+1}(x+1)$, so $\delta_n = 2$. ■

This result resolves a conjecture of Sutner's [4].

We can repeatedly apply theorem 3.1 to obtain explicit formulas for $d(n)$.

Corollary 3.1.1. *Write odd n as $n + 1 = 2^s b$ for $s \geq 0$ and b odd. Then*

$$d(n) = 2^s d(b-1) + \begin{cases} 2(2^s - 1) & 3 \mid b \\ 0 & \text{otherwise} \end{cases}.$$

Proof. By induction on s . For $s = 0$,

$$\begin{aligned} n + 1 &= 2^0 b \\ n &= b - 1 \\ d(n) &= d(b - 1) \\ &= 2^0 d(b - 1) + \begin{cases} 2(2^0 - 1) & 3 \mid b \\ 0 & \text{otherwise} \end{cases}, \end{aligned}$$

as desired. Assume that our claim holds for $s \geq 0$. Then for $n + 1 = 2^{s+1}b$,

$$\begin{aligned} d(n) &= d(2^{s+1}b - 1) \\ &= d(2(2^s b - 1) + 1) \\ &= 2d(2^s b - 1) + \begin{cases} 2 & 3 \mid 2^s b \\ 0 & \text{otherwise} \end{cases} \\ &= 2 \left(2^s d(b - 1) + \begin{cases} 2(2^s - 1) & 3 \mid b \\ 0 & \text{otherwise} \end{cases} \right) + \begin{cases} 2 & 3 \mid 2^s b \\ 0 & \text{otherwise} \end{cases} \\ &= 2^{s+1} d(b - 1) + \begin{cases} 2(2^{s+1} - 1) - 2 & 3 \mid b \\ 0 & \text{otherwise} \end{cases} + \begin{cases} 2 & 3 \mid b \\ 0 & \text{otherwise} \end{cases} \\ &= 2^{s+1} d(b - 1) + \begin{cases} 2(2^{s+1} - 1) & 3 \mid b \\ 0 & \text{otherwise} \end{cases}, \end{aligned}$$

as desired. ■

We can now use this recurrence result to prove the impossibility of certain values of $d(n)$.

Theorem 3.2. *The smallest even k such that for all $n \in \mathbb{N}$, $d(n) \neq k$ is $k = 26$.*

Proof. Recall from lemma 1.4 that $d(n)$ is always even. Through calculation, we can find that

$$\begin{array}{llllll} d(5) = 2 & d(4) = 4 & d(11) = 6 & d(16) = 8 & d(29) = 10 & d(84) = 12 \\ d(23) = 14 & d(19) = 16 & d(101) = 18 & d(30) = 20 & d(59) = 22 & d(62) = 24. \end{array}$$

We will show that $d(n) = 26$ is impossible.

Assume for the sake of contradiction that there exists $n \in \mathbb{N}$ such that $d(n) = 26$. Since 26 is not divisible by 4, corollary 1.4.1 that n must be odd. So, we may write $n = 2m + 1$. By theorem 3.1,

$$26 = d(n) = d(2m + 1) = 2d(m) + \begin{cases} 2 & 3 \mid m + 1 \\ 0 & \text{otherwise} \end{cases}.$$

Since lemma 1.4 tells us that $d(m) = 13$ is not possible, we must have $d(m) = 12$ and $m + 1$ is divisible by 3. Let $m + 1 = 2^s b$ for odd b and $s \geq 0$. By corollary 3.1.1,

$$d(m) = 2^s d(b - 1) + 2(2^s - 1) = 12.$$

The only possible integer solutions here are $d(b - 1) = 12$ and $s = 0$ or $d(b - 1) = 5$ and $s = 1$. Since lemma 1.4 tells us that $d(b - 1) = 5$ is impossible, the only remaining possibility is that $d(m) = d(b - 1) = 12$ and $s = 0$. Thus, m must be even.

We know from lemma 1.1 that

$$d(m) = \deg(\gcd(F_{m+1}(x), F_{m+1}(x + 1))).$$

Since $m + 1$ is odd lemma 1.3 tells us that there exists some square-free $R_{m+1}(x)$ such that

$$F_{m+1}(x) = R_{m+1}(x)^2.$$

Define $H_{m+1}(x)$

$$H_{m+1}(x) = \gcd(R_{m+1}(x), R_{m+1}(x + 1)) \quad \gcd(F_{m+1}(x), F_{m+1}(x + 1)) = H(x)^2.$$

Since $R_{m+1}(x)$ is square-free, then $H_{m+1}(x)$ is too. Since $d(m) = 12$, $\deg H_{m+1}(x) = 6$. Notice too that $H_{m+1}(x) = H_{m+1}(x + 1)$, so our work in lemma 1.5 tells us that for $y = x^2 + x$, there exists square-free a cubic polynomial $h_{m+1}(y) \in \mathbb{F}_2[y]$ such that

$$H_{m+1}(x) = h_{m+1}(y).$$

There are only three possible square-free cubic polynomials in $\mathbb{F}_2[y]$:

1. $C_1(y) = y^3 + 1$
2. $C_2(y) = y^3 + y + 1$
3. $C_3(y) = y^3 + y^2 + 1$.

We will handle these case-by-case.

Define the “rank” of a polynomial P to be the smallest integer r such that P divides F_r . Calculating the rank of our three candidates for $h_{m+1}(x)$,

$$\begin{aligned} \text{rank}((x^2 + x)^3 + 1) &= 85 \\ \text{rank}((x^2 + x)^3 + (x^2 + x) + 1) &= 63 \\ \text{rank}((x^2 + x)^3 + (x^2 + x)^2 + 1) &= 63 \end{aligned}$$

If either rank-63 polynomials C_2 or C_3 divides $R_{m+1}(x)$, then 63 divides $m + 1$. However, by definition of rank, both polynomials would divide $R_{m+1}(x)$. C_2 and C_3 are coprime and each of degree 6 in x , so $\deg H_{m+1}(x) = \deg C_1(x) + \deg C_2(x) = 12$, contradicting our assumption that the degree was 6. If C_1 divides R_{m+1} , then 85 divides $m + 1$. Combining this with the fact that 3 divides $m + 1$ we showed earlier, we see that 255 must divide $m + 1$.

Define

$$P(x) = x^4 + x^3 + 1 \quad Q(x) = x^4 + x^3 + x^2 + x + 1.$$

P and Q are coprime to each other and C_1 . The ranks of $P(x)$ and $Q(x)$ are 17 and 15 respectively. Since both 17 and 15 divide 255, and 255 divides $m+1$, then both $P(x)$ and $Q(x)$ must divide $H_{m+1}(x)$. Therefore,

$$\deg H_{m+1}(x) \geq \deg C_1(x) + \deg P(x) + \deg Q(x) = 14,$$

which contradicts our assumption that the degree was 6.

Thus, there is no $3 \mid m+1$ such that $d(m) = 12$ and therefore no n such that $d(n) = 26$. ■

References

- [1] Markus Hunziker, António Machiavelo, and Jihun Park, *Chebyshev polynomials over finite fields and reversibility of σ -automata on square grids*, Theoretical Computer Science **320** (2004), no. 2, 465–483.
- [2] Donald Knuth, *The art of computer programming, volume 4A: Combinatorial algorithms, part 1*, Addison-Wesley, 2011, Section 7.1.3, Exercises 190-194.
- [3] Klaus Sutner, *On σ -automata*, Complex Systems **2** (1988), no. 1, 1–28.
- [4] ———, *Linear cellular automata and the Garden-of-Eden*, The Mathematical Intelligencer **11** (1989), no. 2, 49–53.
- [5] ———, *sigma-automata and chebyshev-polynomials*, Theoretical Computer Science **230** (1996), 49–73.
- [6] Øystein Ore, *Number theory and its history*, p. 109, McGraw-Hill, 1948, Section 5-4, Problem 2.