


Part of the Monistic Continuum Model (MCM)



Continuum-Based Spectral Physics (CBSP)

Spectral Signatures of Vortex Dynamics and Residual Stress in the
Continuum

Version V01 — August 16, 2026

DOI: [10.5281/zenodo.21963757](https://doi.org/10.5281/zenodo.21963757)

© 2026 Walter Moosbrugger — CC BY 4.0

Contents

1	Continuum-Based Spectral Physics	2
2	Scientific Question	2
3	Required Theoretical Structures	3
4	Mathematical Framework	4
4.1	Spectral State from Residual Stress	4
4.2	Diffuse Background from Vortex Dissolution	4
4.3	Nonlocal Spectral Modulation	4
4.4	Operator-Theoretic Definition of Spectral Structures	4
4.5	Concrete Model Example: Explicit Kernel and Fit-Based Spectral Form	6
5	Integration of Observational Data	8
5.1	Relevant Spectral Datasets	8
5.2	Observable CBSP Signatures	8
5.3	Comparison with Existing Data	9
5.4	Future Observational Tests	10
5.5	Conclusion	10
6	Physical Interpretation	10
7	Comparison of Spectral Paradigms	11
8	Consequences for the Theory	11
1	Mathematical Appendix: Operator Framework of Continuum-Based Spectral Physics	12
2	Functional Spaces and Basic Structures	12
3	Nonlocal Continuum Operator	12
4	Dark-Matter Spectral Operator	13
5	Background Radiation Operator	14
6	Spectral Projection Operator	14
7	Summary of Operator Framework	15
8	Citation Recommendation	16
8.1	APA Style	16
8.2	BibT _E X	16

1 Continuum-Based Spectral Physics

Introduction

The Monistic Continuum Model (MCM) is not intended as a replacement for established physical theories, nor as a competing cosmological framework. Instead, it is proposed as a conceptual and mathematical structure that organizes several known physical phenomena—including dark-matter signatures, background radiation components, and nonlocal spectral modulation—within a unified geometric-dynamical substrate.

The purpose of this document is therefore not to challenge the foundations of modern physics, but to offer an alternative perspective that may complement existing approaches. All standard cosmological and quantum-mechanical results remain valid within their respective domains; the MCM simply provides an additional layer of interpretation that connects these results through a continuum-based operator framework.

The mathematical structures introduced here are deliberately minimal and compatible with current observational practice. The model is fully falsifiable: if continuum-driven spectral modulation or dissolution-based background components are not observed, the MCM must be revised or rejected. In this sense, the framework is intended as an open scientific proposal rather than a definitive theory.

Readers are encouraged to approach the MCM incrementally. The operator definitions in Section 3 form the natural entry point, as they connect directly to measurable spectral quantities. The subsequent sections illustrate how these operators relate to observational datasets without requiring any departure from established physical principles.

The aim of this work is to invite collaboration, discussion, and critical evaluation. The MCM should be understood as a conceptual tool that may help structure existing knowledge and suggest new avenues for empirical investigation, not as a prescriptive or disruptive alternative to the successful theories of contemporary physics.

2 Scientific Question

The central scientific question addressed in this chapter is:

How does the Monistic Continuum Model (MCM) generate, modulate, and interpret spectral distributions, and in what precise way do these continuum-driven spectra differ from the baryonic and radiative spectra described in standard cosmology?

This question arises naturally from the MCM dark-matter theory: if transient vortex modes continuously form and dissolve, producing residual stress and diffuse radiation, then the continuum must imprint distinct spectral signatures that differ fundamentally from baryonic emission spectra.

3 Required Theoretical Structures

To answer the scientific question, the following theoretical concepts and operators must be defined:

1. The continuum field

$$C : M \times \Lambda \rightarrow V,$$

representing the geometric-dynamical substrate.

2. Vortex modes

$$W_n \in \mathcal{W},$$

classified into stable, metastable, and transient modes.

3. The residual stress operator

$$\sigma_{\text{res}}(x, \lambda),$$

generating dark-matter density via

$$\rho_{\text{DM}}(x) = \langle \sigma_{\text{res}}(x, \lambda) \rangle_{\lambda}.$$

4. The nonlocal continuum operator

$$(D_{\text{nl}}C)(x) = \int_M \mathcal{K}(x, y) C(y) d\mu(y),$$

responsible for continuum-driven spectral modulation.

5. The spectral projection operator

$$|\Psi_{\text{spec}}\rangle = \hat{\Pi}_{\text{spec}}(\sigma_{\text{res}}, W_n),$$

mapping geometric stress and vortex dynamics into observable spectral states.

4 Mathematical Framework

The continuum-based spectral physics requires the following mathematical relations:

4.1 Spectral State from Residual Stress

Residual stress contributes to spectral modulation through

$$|\Psi_{\text{DM}}(x)\rangle = \hat{\Pi}_{\text{DM}}(\langle\sigma_{\text{res}}(x, \lambda)\rangle_{\lambda}).$$

4.2 Diffuse Background from Vortex Dissolution

Transient vortex modes generate background radiation via

$$I_{\text{bg}}(\nu) = \mathcal{B}(W_{\text{transient}}),$$

where $\mathcal{B}$ is the background radiation operator.

4.3 Nonlocal Spectral Modulation

Continuum stress modulates spectra through

$$I_{\text{mod}}(\nu, x) = I_0(\nu) + \alpha [(D_{\text{nl}}C)(x)],$$

with α a coupling constant determined by continuum geometry.

4.4 Operator-Theoretic Definition of Spectral Structures

To formalize continuum-based spectral physics within the Monistic Continuum Model (MCM), we introduce the operator framework acting on well-defined functional spaces. This provides the mathematical backbone for spectral modulation, background radiation, and dark-matter related spectral signatures.

Functional Setting

Let (M, μ) be a smooth manifold equipped with a measure μ , representing the geometric continuum. Let Λ denote a parameter space of vortex modes. The continuum field is

$$C : M \times \Lambda \rightarrow V,$$

with V a finite-dimensional vector space. The residual stress field is

$$\sigma_{\text{res}} : M \times \Lambda \rightarrow \mathbb{R},$$

and the dark-matter density is defined by

$$\rho_{\text{DM}}(x) = \int_{\Lambda} \sigma_{\text{res}}(x, \lambda) w(\lambda) d\lambda,$$

where $w(\lambda)$ is a vortex-population weight.

Nonlocal Continuum Operator D_{nl}

For fixed $\lambda \in \Lambda$, assume $C(\cdot, \lambda) \in L^2(M, \mu; V)$. The nonlocal continuum operator is defined by

$$(D_{\text{nl}}C)(x, \lambda) = \int_M \mathcal{K}(x, y; \lambda) C(y, \lambda) d\mu(y),$$

with kernel

$$\mathcal{K} : M \times M \times \Lambda \rightarrow \text{End}(V).$$

This operator is linear and encodes continuum-driven spectral modulation:

$$I_{\text{mod}}(\nu, x) = I_0(\nu) + \alpha F[(D_{\text{nl}}C)(x, \lambda)],$$

where $F : V \rightarrow \mathbb{R}$ maps continuum configurations to observable intensity.

Dark-Matter Spectral Operator $\mathcal{D}_{\text{DM}}$

Let $\mathcal{H}_{\text{DM}}$ be a Hilbert space of spectral states. The dark-matter spectral operator is defined as

$$\mathcal{D}_{\text{DM}} : L^2(M \times \Lambda) \rightarrow \mathcal{H}_{\text{DM}}, \quad \sigma_{\text{res}} \mapsto |\Psi_{\text{DM}}\rangle,$$

with

$$|\Psi_{\text{DM}}\rangle = \int_M \int_{\Lambda} f(x, \lambda) \sigma_{\text{res}}(x, \lambda) |x, \lambda\rangle d\lambda d\mu(x),$$

where f encodes the coupling between residual stress and spectral representation.

Background Radiation Operator $\mathcal{B}$

Let $n_{\text{transient}}(\lambda)$ denote the density of transient vortex modes. The background radiation operator is

$$\mathcal{B} : \mathcal{W}_{\text{transient}} \rightarrow L^2(\mathbb{R}^+, d\nu), \quad W_{\text{transient}} \mapsto I_{\text{bg}}(\nu),$$

with

$$I_{\text{bg}}(\nu) = \int_{\Lambda} g(\nu, \lambda) n_{\text{transient}}(\lambda) d\lambda,$$

where g is an emission kernel describing dissolution-driven background radiation.

Spectral Projection Operator $\hat{\Pi}_{\text{spec}}$

For pairs $(\sigma_{\text{res}}, W_n)$ in

$$L^2(M \times \Lambda) \times \mathcal{W},$$

the spectral projection operator is defined by

$$\hat{\Pi}_{\text{spec}} : L^2(M \times \Lambda) \times \mathcal{W} \rightarrow \mathcal{H}_{\text{spec}},$$

$$|\Psi_{\text{spec}}\rangle = \hat{\Pi}_{\text{spec}}(\sigma_{\text{res}}, W_n) = \int_M \int_{\Lambda} h(x, \lambda; W_n) \sigma_{\text{res}}(x, \lambda) |x, \lambda\rangle d\lambda d\mu(x),$$

where h encodes the coupling between vortex modes and residual stress in the spectral representation.

Summary

The operator framework of the MCM consists of:

- D_{nl} — nonlocal continuum operator generating spectral modulation,
- $\mathcal{D}_{\text{DM}}$ — mapping residual stress to dark-matter spectral states,
- $\mathcal{B}$ — background radiation operator from transient vortex modes,
- $\hat{\Pi}_{\text{spec}}$ — projection operator producing observable spectral states.

These operators form the mathematical foundation of continuum-based spectral physics.

4.5 Concrete Model Example: Explicit Kernel and Fit-Based Spectral Form

To demonstrate how the operator framework of Continuum-Based Spectral Physics (CBSP) can be instantiated in a concrete, data-fit-ready model, we introduce an explicit form of the nonlocal

continuum kernel and derive a corresponding spectral intensity function suitable for observational comparison.

Explicit Kernel

We consider a Gaussian-type nonlocal kernel

$$\mathcal{K}(x, y; \lambda) = A(\lambda) \exp\left(-\frac{d(x, y)^2}{2 L(\lambda)^2}\right),$$

where

- $A(\lambda)$ is a mode-dependent amplitude,
- $L(\lambda)$ is a characteristic interaction length,
- $d(x, y)$ is the geodesic distance on the continuum manifold M .

This kernel represents a physically motivated, spatially decaying nonlocal interaction between continuum configurations.

Modelled Continuum Response

The nonlocal continuum operator becomes

$$(D_{\text{nl}}C)(x, \lambda) = A(\lambda) \int_M \exp\left(-\frac{d(x, y)^2}{2 L(\lambda)^2}\right) C(y, \lambda) d\mu(y).$$

For sufficiently smooth C , this operator acts as a mode-dependent smoothing and redistribution mechanism, encoding continuum geometry into spectral modulation.

Fit-Based Spectral Form

A fit-ready spectral intensity model is obtained by inserting the explicit kernel into the general modulation formula:

$$I_{\text{model}}(\nu, x) = I_0(\nu) + \alpha \int_{\Lambda} F \left[A(\lambda) \int_M \exp\left(-\frac{d(x, y)^2}{2 L(\lambda)^2}\right) C(y, \lambda) d\mu(y) \right] w(\lambda) d\lambda.$$

This expression provides a concrete spectral form that can be fitted to observational datasets such as CMB microstructure, diffuse X-ray backgrounds, or infrared background fluctuations.

Interpretation

The explicit kernel $\mathcal{K}(x, y; \lambda)$ introduces a tunable geometric scale $L(\lambda)$ and amplitude $A(\lambda)$, enabling direct comparison between continuum-driven spectral modulation and empirical data. This model bridges the abstract operator framework and measurable spectral signatures, bringing CBSP close to a fully testable theory.

5 Integration of Observational Data

Continuum-Based Spectral Physics (CBSP) provides a geometric-dynamical interpretation of spectral phenomena arising from vortex dynamics, residual stress, and dissolution processes in the Monistic Continuum Model (MCM). To connect the theoretical operator framework with empirical evidence, this section integrates relevant observational datasets and identifies specific signatures predicted by CBSP that can be tested with current and future missions.

5.1 Relevant Spectral Datasets

The following observational datasets provide direct or indirect access to spectral structures that CBSP aims to explain:

- **Cosmic Microwave Background (CMB)** High-precision measurements of temperature and polarization spectra from COBE, WMAP, and Planck.
- **Diffuse X-ray Background (XRB)** Measurements from Chandra, XMM-Newton, and eROSITA.
- **Extragalactic Gamma Background (EGB)** Data from Fermi-LAT and INTEGRAL.
- **Infrared and Submillimeter Backgrounds (IRB/SMB)** Observations from Herschel, Spitzer, and JWST.
- **Weak Lensing Surveys** Euclid, DES, KiDS, and LSST provide stress-correlated distortions of background spectra.

These datasets contain spectral features that can be compared directly with CBSP predictions.

5.2 Observable CBSP Signatures

The operator framework developed in CBSP predicts several observable spectral signatures:

1. **Non-thermal background components** Arising from dissolution of transient vortex modes via the operator

$$I_{\text{bg}}(\nu) = \mathcal{B}(W_{\text{transient}}).$$

2. **Spectral roughness and microstructure** Generated by continuum-driven modulation through

$$I_{\text{mod}}(\nu, x) = I_0(\nu) + \alpha F[(D_{\text{nl}}C)(x, \lambda)].$$

3. **Residual-stress-induced anisotropies** Encoded in the dark-matter spectral operator

$$|\Psi_{\text{DM}}\rangle = \mathcal{D}_{\text{DM}}(\sigma_{\text{res}}),$$

which predicts correlations between stress fields and spectral distortions.

4. **Frequency-dependent lensing-like distortions** Distinct from classical gravitational lensing because they arise from continuum geometry rather than spacetime curvature.
5. **Spectral excesses or deficits** In frequency bands where baryonic processes cannot account for observed intensities.

These signatures provide concrete targets for observational tests.

5.3 Comparison with Existing Data

Several known anomalies and features in observational datasets align with CBSP predictions:

- **CMB anomalies** Large-scale power deficits, hemispherical asymmetry, and non-Gaussian microstructure may reflect continuum-driven spectral modulation.
- **Diffuse X-ray and gamma backgrounds** Unresolved excesses in XRB and EGB spectra may correspond to dissolution-driven background radiation.
- **Infrared background fluctuations** IRB microstructure may indicate stress-correlated spectral modulation.
- **Weak lensing residuals** Frequency-dependent distortions in lensing maps may arise from residual stress fields.

These correlations do not constitute proof but provide strong motivation for targeted analysis.

5.4 Future Observational Tests

Upcoming missions offer opportunities to test CBSP predictions:

- **LiteBIRD** High-precision polarization spectra may reveal residual-stress anisotropies.
- **Euclid and LSST** Deep lensing surveys can detect continuum-driven spectral modulation.
- **JWST and Athena** High-resolution background spectra can identify dissolution-driven components.
- **SKA** Low-frequency background measurements may detect vortex-population signatures.

These missions can provide decisive evidence for or against continuum-based spectral physics.

5.5 Conclusion

The integration of observational data strengthens CBSP by linking its operator framework to empirical spectral structures. The predicted signatures—non-thermal components, spectral roughness, stress-induced anisotropies, and dissolution-driven backgrounds—are measurable with current and upcoming instruments. This establishes CBSP as a testable extension of the Monistic Continuum Model and provides a clear path toward empirical validation.

6 Physical Interpretation

The mathematical structures yield the following physical interpretation:

- **Baryonic spectra** arise from stable vortex modes and exhibit discrete lines, continuum radiation, and object-bound spectral features.
- **Residual-stress spectra** arise from the geometric stress field generated by transient vortex modes. They are delocalized, non-thermal, and imprint subtle modulations on background radiation.
- **Dissolution spectra** arise from the continuous decay of short-lived vortex modes, producing diffuse, weakly structured background radiation.
- **Continuum-driven modulation** produces frequency-dependent distortions, anisotropies, and spectral roughness not predicted by standard cosmology.

These interpretations unify dark matter and background radiation as two manifestations of the same continuum-dynamical process.

7 Comparison of Spectral Paradigms

The following TikZ graphic illustrates the conceptual difference between the spectrum in standard cosmology (top) and the continuum-based spectrum of the MCM (bottom). It is placed here because this section introduces the theoretical distinction that the graphic visualizes.




Figure 1: Comparison of spectral interpretation: standard cosmology (top) vs. MCM continuum-based spectral physics (bottom).

8 Consequences for the Theory

The continuum-based spectral physics has several consequences for the overall structure of the MCM:

1. It unifies dark matter and background radiation as two aspects of vortex dynamics.

2. It predicts spectral signatures that differ from Λ CDM, enabling observational tests.
3. It provides a geometric-dynamical interpretation of non-thermal background components.
4. It introduces spectral operators that integrate seamlessly with the MCM operator framework.
5. It establishes a continuum-driven mechanism for spectral modulation, extending classical lensing theory.

These consequences strengthen the MCM as a unified physical theory capable of explaining both visible and invisible components of the universe through a single geometric-dynamical substrate.

1 Mathematical Appendix: Operator Framework of Continuum-Based Spectral Physics

2 Functional Spaces and Basic Structures

Let (M, μ) be a smooth, σ -finite measure space representing the geometric continuum. Let Λ be a measurable parameter space of vortex modes. The continuum field is defined as

$$C : M \times \Lambda \rightarrow V,$$

where V is a finite-dimensional real or complex vector space.

The residual stress field is

$$\sigma_{\text{res}} : M \times \Lambda \rightarrow \mathbb{R},$$

with induced dark-matter density

$$\rho_{\text{DM}}(x) = \int_{\Lambda} \sigma_{\text{res}}(x, \lambda) w(\lambda) d\lambda,$$

where $w(\lambda)$ is a population weight.

We assume

$$C(\cdot, \lambda) \in L^2(M, \mu; V), \quad \sigma_{\text{res}}(\cdot, \cdot) \in L^2(M \times \Lambda, \mu \otimes d\lambda).$$

3 Nonlocal Continuum Operator

The nonlocal continuum operator

$$D_{\text{nl}} : L^2(M, \mu; V) \rightarrow L^2(M, \mu; V)$$

is defined by

$$(D_{\text{nl}}C)(x, \lambda) = \int_M \mathcal{K}(x, y; \lambda) C(y, \lambda) d\mu(y),$$

where

$$\mathcal{K} : M \times M \times \Lambda \rightarrow \text{End}(V)$$

is a measurable kernel.

Properties

- Linearity:

$$D_{\text{nl}}(aC_1 + bC_2) = aD_{\text{nl}}C_1 + bD_{\text{nl}}C_2.$$

- Boundedness (sufficient condition):

$$\sup_{x, \lambda} \int_M \|\mathcal{K}(x, y; \lambda)\| d\mu(y) < \infty.$$

- Symmetry (optional):

$$\mathcal{K}(x, y; \lambda) = \mathcal{K}(y, x; \lambda).$$

Spectral modulation is expressed as

$$I_{\text{mod}}(\nu, x) = I_0(\nu) + \alpha F[(D_{\text{nl}}C)(x, \lambda)],$$

with $F : V \rightarrow \mathbb{R}$ a measurable intensity map.

4 Dark-Matter Spectral Operator

Let $\mathcal{H}_{\text{DM}}$ be a Hilbert space of spectral states with generalized basis $\{|x, \lambda\rangle\}$.

The dark-matter spectral operator is defined as

$$\mathcal{D}_{\text{DM}} : L^2(M \times \Lambda) \rightarrow \mathcal{H}_{\text{DM}},$$

$$|\Psi_{\text{DM}}\rangle = \int_M \int_\Lambda f(x, \lambda) \sigma_{\text{res}}(x, \lambda) |x, \lambda\rangle d\lambda d\mu(x),$$

where $f : M \times \Lambda \rightarrow \mathbb{C}$ is a measurable coupling weight.

Norm Estimate

If

$$|f(x, \lambda)| \leq C_f,$$

then

$$\|\Psi_{\text{DM}}\|_{\mathcal{H}_{\text{DM}}} \leq C_f \|\sigma_{\text{res}}\|_{L^2(M \times \Lambda)}.$$

5 Background Radiation Operator

Let $n_{\text{transient}} : \Lambda \rightarrow \mathbb{R}_{\geq 0}$ denote the density of transient vortex modes.

The background radiation operator is

$$\mathcal{B} : L^1(\Lambda) \rightarrow L^2(\mathbb{R}^+, d\nu),$$

$$I_{\text{bg}}(\nu) = \int_\Lambda g(\nu, \lambda) n_{\text{transient}}(\lambda) d\lambda,$$

with emission kernel

$$g : \mathbb{R}^+ \times \Lambda \rightarrow \mathbb{R}.$$

Boundedness

If

$$\sup_\nu \int_\Lambda |g(\nu, \lambda)| d\lambda < \infty,$$

then $\mathcal{B}$ is bounded.

6 Spectral Projection Operator

Let $\mathcal{H}_{\text{spec}}$ be a Hilbert space of observable spectral states.

The spectral projection operator is defined as

$$\hat{\Pi}_{\text{spec}} : L^2(M \times \Lambda) \times \mathcal{W} \rightarrow \mathcal{H}_{\text{spec}},$$

$$|\Psi_{\text{spec}}\rangle = \int_M \int_{\Lambda} h(x, \lambda; W_n) \sigma_{\text{res}}(x, \lambda) |x, \lambda\rangle d\lambda d\mu(x),$$

where

$$h : M \times \Lambda \times \mathcal{W} \rightarrow \mathbb{C}$$

encodes the coupling between vortex modes and residual stress.

Interpretation

The operator $\hat{\Pi}_{\text{spec}}$ maps geometric-dynamical continuum structures into observable spectral states, combining:

- vortex-mode contributions,
- residual-stress fields,
- continuum geometry.

7 Summary of Operator Framework

The continuum-based spectral physics of the MCM is governed by the following operators:

- D_{nl} — nonlocal continuum operator generating spectral modulation,
- $\mathcal{D}_{\text{DM}}$ — mapping residual stress to dark-matter spectral states,
- $\mathcal{B}$ — background radiation operator from transient vortex modes,
- $\hat{\Pi}_{\text{spec}}$ — projection operator producing observable spectral states.

These operators form a mathematically rigorous foundation for interpreting baryonic, radiative, and dark-matter-related spectral signatures within the Monistic Continuum Model.

8 Citation Recommendation

This document represents the official reference version of *Continuum-Based Spectral Physics (CBSP): Spectral Signatures of Vortex Dynamics and Residual Stress in the Continuum*, published as Version V01 of the *Monistic Continuum Model (MCM)* series and archived with a permanent DOI on Zenodo.

The following citation formats may be used:

8.1 APA Style

Moosbrugger, W. (2026). *Continuum-Based Spectral Physics (CBSP): Spectral Signatures of Vortex Dynamics and Residual Stress in the Continuum*. Part of the *Monistic Continuum Model (MCM)*. Zenodo. <https://doi.org/10.5281/zenodo.21963757>

8.2 BibT_EX

```
@misc{moosbrugger2026_cbsp,  
  author    = {Moosbrugger, Walter},  
  title     = {Continuum-Based Spectral Physics (CBSP):  
    Spectral Signatures of Vortex Dynamics and  
    Residual Stress in the Continuum},  
  year      = {2026},  
  publisher = {Zenodo},  
  doi       = {10.5281/zenodo.21963757},  
  url       = {https://doi.org/10.5281/zenodo.21963757},  
  note      = {Version V01, Part of the Monistic Continuum Model (MCM)}  
}
```