


MCM — Master Document



Monistic Continuum Model (MCM)

Unified Geometric Ontology

DOI: [10.5281/zenodo.22128163](https://doi.org/10.5281/zenodo.22128163)

Version 07 — 27. August 2026

© 2026 Walter Moosbrugger

Licensed under CC BY 4.0

Contents

1	Introduction and Motivation	2
2	Ontological Entities in the MCM	3
3	Fundamental Axioms of the Medium	7
4	Emergence of Space from Bound Geometry	10
5	Mathematical Structure of the Medium	13
6	Dynamics of the Medium	15
7	Formalization of the Operators	18
8	Geometry from Stress	21
9	Matter as Bound Geometry	23
10	Vortex Geometry	25
11	Electromagnetism in the MCM	27
12	Quantization as a Geometric Effect	30
13	Cosmology of the Medium	32
14	Numerical Methods and Simulation	35
15	Experimental Accessibility	38
16	Applications in Applied Physics	41
17	Limitations of the MCM	43
18	Future Research Directions	46
19	Glossary of Terms	49
20	Conclusion	53
21	Citation Recommendation	55

Abstract

Das Monistic Continuum Model (MCM) stellt eine einheitliche geometrische Ontologie bereit, in der alle beobachtbaren Phänomene aus Stress, Rotation, Krümmung und interner Aktualisierung eines einzigen kontinuierlichen Mediums entstehen. Es werden keine physikalischen Primitiven wie Raum, Zeit, Felder, Teilchen oder Kräfte vorausgesetzt; sämtliches emergentes Verhalten wird durch interne geometrische Strukturen erzeugt, die in der prä-räumlichen Zustandsabbildung $C(\alpha, \Omega)$ kodiert sind und durch die Aktualisierungsrate τ reorganisiert werden.

Gebundene Geometrie besteht aus stabilen Konfigurationen von Stress, Rotation und Krümmung, die annähernde Aktualisierungsinvarianz erfüllen. Diese gebundenen Strukturen bilden diskrete Stabilitätsmodi und entsprechen materieähnlichem Verhalten. Ungebundene Geometrie bleibt kontinuierlich und unterstützt rotations-geometrische Ausbreitung, Krümmungsentwicklung und Domänenorganisation.

Das MCM liefert eine vollständig geometrische Beschreibung von Dynamik, bei der Entwicklung aus interner Reorganisation statt aus zeitlicher Evolution resultiert. Großskalige Muster – wie Clusterbildung, Krümmungssenken und rotationsgeometrische Kohärenz – entstehen aus Domäneninteraktionen ungebundener Geometrie. Das Modell bietet einen konsistenten Rahmen für numerische Simulation, analytische Untersuchung und die Übersetzung in physikalische Terminologie, ohne seine monistische geometrische Ontologie zu verlassen.

1 Introduction and Motivation

The Monistic Continuum Model (MCM) is developed in response to a fundamental conceptual limitation of established physical frameworks: the reliance on multiple primitive entities—space, time, fields, particles, and forces—that must be postulated independently and subsequently connected through additional rules. This multi-entity ontology produces structural discontinuities, requires external interaction laws, and obscures the geometric origin of observable phenomena. The MCM replaces this fragmented foundation with a single continuous medium whose internal geometry generates all emergent behavior.

The motivation for a monistic geometric ontology arises from three central observations. First, many physical phenomena traditionally treated as distinct (e.g., wave propagation, binding, curvature, rotational coherence) share a common geometric structure when expressed in continuum form. Second, stability patterns associated with matter, defects, and large-scale organization can be interpreted as bound geometric configurations rather than as properties of independent entities. Third, numerical and analytical methods reveal that stress, rotation, and curvature interact in ways that naturally produce localized binding, domain formation, and discrete stability modes without

requiring external forces or temporal evolution.

The MCM formalizes these insights by introducing a pre-spatial state map $C(\alpha, \Omega)$ and an update mechanism governed by the reactivity $\tau(\alpha, \Omega)$. The update index Ω is not a temporal parameter; it labels successive internal reorganizations of the medium. Dynamics are therefore geometric rather than temporal: development arises from changes in stress, rotation, and curvature induced by the constitutive operator $F(Vv, \sigma)$ and captured by the update derivative DC .

This geometric perspective resolves several conceptual tensions. It eliminates the need for independent field equations, avoids the dualism between particles and fields, and provides a unified explanation for bound and unbound behavior. Bound geometry corresponds to stable configurations that satisfy approximate update invariance, while unbound geometry describes continuous propagation and domain-level organization. Quantization emerges as a stability effect of bound geometry rather than as a fundamental principle. Large-scale patterns such as clustering, curvature sinks, and rotational coherence arise from domain-level interactions of unbound geometry. The purpose of this document is to establish the mathematical foundations of the MCM, formalize its operators, and demonstrate how stress, rotation, curvature, and updating generate the full spectrum of emergent behavior. The presentation is strictly geometric and fully compliant with the monistic ontology: no physical primitives are introduced, and all phenomena are derived from internal geometric structures of the medium.

The following sections develop the model systematically. After introducing the state map and geometric invariants, we formalize the update derivative and the constitutive operator, analyze bound and unbound geometry, examine rotational and curvature-based modes, and explore domain-level behavior. Numerical and analytical methods are outlined to support simulation and further development. The document concludes with conceptual implications and directions for future research within the monistic geometric framework.

2 Ontological Entities in the MCM

The Monistic Continuum Model (MCM) is based on a minimal set of ontological entities that describe the geometric, rotational, and stress-induced behavior of a single continuous medium. These entities replace the fundamental constructs of established physics—space, time, fields, particles, and forces—with purely geometric structures. The ontology is strictly monistic: all phenomena arise from internal relations of the medium.

The Medium

The medium is the sole fundamental entity of the MCM. It possesses no spatial or temporal structure. All observable structures—bound geometry (matter), rotational–geometric modes, curvature, and domains—arise from its internal stress, rotation, and curvature geometry.

Ontology: Only the medium.

Pre-Spatial State Map

The fundamental state map of the medium is

$$C(\alpha, \Omega) = \left(\rho(\alpha, \Omega), v_A(\alpha, \Omega), \sigma_{AB}(\alpha, \Omega), \tau(\alpha, \Omega) \right).$$

Here:

- α is a medium index (not a spatial coordinate),
- Ω is the global update index,
- all components are purely geometric quantities.

This map fully describes the medium prior to the emergence of space.

Density

$$\rho(\alpha, \Omega)$$

Geometric measure of local compression or expansion. Not a physical mass density. Determines the internal stress distribution.

Geometric Motion

$$v_A(\alpha, \Omega)$$

Motion induced by stress gradients. Not a physical velocity. Determines the reorganization of geometric patterns.

Stress

$$\sigma_{AB}(\alpha, \Omega)$$

Central ontological quantity of the MCM. Stress generates internal geometry, rotation, curvature, stability of bound geometry, and domain formation. All geometry in the MCM is stress geometry.

Update Rate

$$\tau(\alpha, \Omega)$$

Local reactivity of the medium. Not physical time. Modulates the strength of internal updating.

Update Derivative

$$DC(\alpha, \Omega)$$

Internal development of the medium. Measures the change of the state map under updating.

Three Stability Forms of Geometry

The geometry of the medium appears in three ontologically distinct stability forms:

(1) Open Geometry (Unbound Geometry)

$$DC(\alpha, \Omega) \neq 0$$

Pre-spatial form of the medium. Possesses no metric, no orientation, and no stability. Forms the geometric continuum from which space emerges, but is not space itself.

Open geometry is the *space above space*: the medium that carries and permeates emergent space. The cosmos consists almost entirely of open geometry; bound geometry is extremely rare.

(2) Unstable Bound Geometry

$$DC(\alpha, \Omega) \approx 0 \quad (\text{temporarily})$$

Short-lived bound configuration without permanent stability. Pre-material transitional form between open and stable bound geometry. Possesses temporary orientation and temporary metric.

Stable Bound Geometry (Matter)

Stable bound geometry forms the basis of matter and space. It is not a particle structure but localized stress, rotation, and curvature geometry. Stable bound geometry realizes itself as a vortex.

Only after the emergence of space do bound structures appear as material objects $M(x, \Omega)$.

Rotational–Geometric Modes

Propagating pre-spatial patterns in $L_{AB}(\alpha, \Omega)$. Correspond to electromagnetism-like behavior after spatial interpretation.

Curvature Domains

Large-scale regions of coherent internal curvature. Arise from global stress organization. Form the basis of cosmology-like behavior.

Ontological Minimality

The MCM uses the smallest possible set of ontological entities:

- the medium,
- the pre-spatial state map $C(\alpha, \Omega)$,
- stress,
- rotation,
- internal curvature,
- update rate,
- geometric motion.

All other phenomena—bound geometry (matter), waves, forces, fields, spacetime— are emergent geometric effects.

Summary

The ontological entities of the MCM define a strictly monistic geometric foundation. Stress, rotation, curvature, and update dynamics of a single medium generate all emergent phenomena. Space and matter arise only through stable bound stress geometry.

3 Fundamental Axioms of the Medium

The Monistic Continuum Model (MCM) is based on a minimal set of ontological axioms. These axioms define a single continuous medium without physical space, without physical time, without fields, without particles, and without forces. All observable phenomena arise from internal geometric relations of the medium.

In its pre-spatial state, the medium possesses only *unbound geometry*. Bound geometry emerges only when this geometry forms a stable configuration; in the MCM, bound geometry is identical to *matter*.

Axiom 1: The Medium

The medium is the sole ontological entity. It possesses no spatial or temporal structure. All structures arise from internal stress, rotation, and curvature.

Ontology: Only the medium.

Axiom 2: Pre-Spatial State Map

The state of the medium is described by the non-spatial map

$$C(\alpha, \Omega) = (\rho(\alpha, \Omega), v_A(\alpha, \Omega), \sigma_{AB}(\alpha, \Omega), \tau(\alpha, \Omega))$$

where:

- α is a medium index (not a spatial coordinate),
- Ω is a global update index,
- all components are purely geometric quantities.

This map fully describes the medium prior to the emergence of space.

Axiom 3: Stress-Induced Internal Geometry

Stress generates internal geometric relations:

$$\gamma_{AB}(\alpha, \Omega) = \delta_{AB} + \alpha_1 \sigma_{AB}(\alpha, \Omega) + \alpha_2 I_1(\alpha, \Omega) \delta_{AB},$$

with

$$I_1(\alpha, \Omega) = \text{tr}(\sigma(\alpha, \Omega)).$$

This geometry is pre-spatial and unbound.

Axiom 4: Internal Curvature

Curvature arises from the internal geometry:

$$\mathcal{R}_{ABCD}(\alpha, \Omega) = R(\gamma_{AB}(\alpha, \Omega)).$$

This is pure medium curvature, not spatial curvature.

Axiom 5: Rotational Behavior

Rotation is an internal geometric quantity:

$$L_{AB}(\alpha, \Omega) = \frac{1}{2}(\nabla_A v_B - \nabla_B v_A).$$

Rotation interacts with stress and curvature and generates pre-spatial wave-like patterns.

Axiom 6: Update Rate

The update rate is an ontological reactivity measure:

$$\tau(\alpha, \Omega) \in \mathbb{R}_{\geq 0}.$$

It is not time; it modulates the strength of internal updating.

Axiom 7: Internal Updating

The internal development of the medium is described by

$$DC(\alpha, \Omega)$$

Bound geometry arises precisely when

$$DC(\alpha, \Omega) \approx 0.$$

Stable bound geometry realizes itself as a vortex.

Axiom 8: Constitutive Dynamics

Stress reorganizes according to

$$D\sigma_{AB}(\alpha, \Omega) = F(Vv, \sigma)_{AB},$$

with internal transport

$$Vv(\sigma_{AB}) = v_C \nabla_C \sigma_{AB}.$$

All dynamics arise from internal geometric relations.

Axiom 9: Emergence of Space from Bound Geometry

Space arises exclusively from *bound geometry*. When a vortex becomes stable, a local map emerges:

$$x = X(\alpha, \Omega),$$

and the emergent metric is

$$g_{ij}(x, \Omega) = \gamma_{AB}(\alpha, \Omega) \frac{\partial X^A}{\partial x^i} \frac{\partial X^B}{\partial x^j}.$$

Space is the external appearance of bound geometry.

Axiom 10: Matter as Bound Space

Matter is not fundamental. Matter is *bound space*. It arises as stable binding of stress, rotation, and curvature:

$$M(x, \Omega) = \text{bound stress--rotation--curvature structures.}$$

The vortex is the only structure that fully realizes this binding.

Summary

The MCM possesses two levels:

$$\text{Medium: } C(\alpha, \Omega) \quad (\text{unbound geometry}),$$

Matter: $M(x, \Omega)$ (bound geometry = space).

Space arises from bound geometry. Matter is bound space. All phenomena are emergent geometric effects of a single medium.

4 Emergence of Space from Bound Geometry

In the Monistic Continuum Model (MCM), physical space is not a fundamental entity. The medium possesses an internal geometry induced by stress which, in its pre-spatial form, is *unbound* and does not constitute spatial structure. Only when patterns of stress, rotation, and curvature form a stable binding does *bound geometry* arise. In the MCM, bound geometry is identical to *matter*. Physical space is the external appearance of this bound geometry.

Unbound Geometry

The internal geometry of the medium is described by the stress-induced medium metric

$$\gamma_{AB}(\alpha, \Omega) = \delta_{AB} + \alpha_1 \sigma_{AB}(\alpha, \Omega) + \alpha_2 I_1(\alpha, \Omega) \delta_{AB},$$

with the first stress invariant

$$I_1(\alpha, \Omega) = \text{tr}(\sigma(\alpha, \Omega)).$$

This geometry is pre-spatial and unbound. As long as no stable binding structures exist, it remains dynamic, non-orientable, and non-measurable. Unbound geometry carries neither space nor matter.

Bound Geometry as the Origin of Space

Bound geometry emerges when stress, rotation, and curvature form a stable configuration, characterized by

$$DC(\alpha, \Omega) \approx 0.$$

Such a configuration is a *vortex defect*. In the MCM, matter is not a structure within space; it is the first form of bound geometry from which space emerges.

Bound geometry produces stable features such as:

- persistent stress centers,

- rotational cores,
- localized curvature domains.

These features form the geometric anchors that make physical space possible.

Formation of Spatial Axes

A stable vortex defect defines a local frame through:

- principal stress directions (eigenvectors of σ_{AB}),
- rotational axes (eigenvectors of L_{AB}),
- curvature gradients (principal directions of $\mathcal{R}_{ABCD}$).

These directions remain stable under updating and form the basis of emergent spatial orientation.

Emergence of Spatial Structure

Space emerges when bound geometry satisfies three conditions:

1. Persistence:

$$DC(\alpha, \Omega) \approx 0,$$

ensuring that the bound structure remains invariant under updating.

2. Orientation:

Stable eigenvectors of σ_{AB} , L_{AB} , and $\mathcal{R}_{ABCD}$ define persistent geometric axes.

3. Measurability:

The internal metric γ_{AB} becomes stable through binding, allowing a consistent spatial metric to be defined:

$$g_{ij}(x, \Omega) = \gamma_{AB}(\alpha, \Omega) \frac{\partial X^A}{\partial x^i} \frac{\partial X^B}{\partial x^j},$$

where $x = X(\alpha, \Omega)$ is the emergent spatial map.

Physical space is thus the stabilized, bound geometric domain of a vortex defect.

Space as a Derived Concept

Space is the stable, orientable geometric structure induced by bound geometry (matter):

Space = bound geometry anchored by matter.

Space is not fundamental; it arises exclusively through binding.

Absence of Space Before Matter

Before bound geometry forms, the medium exhibits:

- stress,
- geometric motion,
- rotation,
- internal curvature,
- update dynamics.

However, it lacks:

- stable axes,
- stable distances,
- stable orientation,
- measurable spatial structure.

Unbound geometry exists independently of space. Space emerges only through binding.

Summary

In the MCM, space is not fundamental. It emerges exclusively from bound geometry. Matter is bound space. Before bound structures arise, the medium possesses internal geometry but no space. This emergent view of spatial structure ensures a clear distinction between unbound geometry and bound, spatially interpretable behavior.

5 Mathematical Structure of the Medium

The Monistic Continuum Model (MCM) describes a single continuous medium through geometric and tensorial quantities that evolve under internal updating. The mathematical structure is entirely based on internal relations induced by stress, rotation, and geometric motion. Neither physical space nor physical time, nor fields, particles, or forces are assumed. All mathematical objects represent internal states of the medium and do not rely on external coordinates or physical measurement conventions.

In its pre-spatial form, the medium consists solely of *unbound geometry*. This geometry carries no matter and does not form physical space. Only through stable binding does bound geometry arise, which in the MCM is identical to matter.

State Map of the Medium

The pre-spatial state map is

$$C(\alpha, \Omega) = (\rho(\alpha, \Omega), v_A(\alpha, \Omega), \sigma_{AB}(\alpha, \Omega), \tau(\alpha, \Omega)),$$

where all components are purely geometric properties of the medium. The index α has no spatial meaning, and Ω is an update index without temporal interpretation.

Geometric Derivatives

All derivatives in the MCM are geometric. The update derivative

$$DC(\alpha, \Omega)$$

describes the internal development of the medium. Geometric derivatives replace spatial derivatives and operate on stress, motion, and rotation within the internal geometry. Bound geometry emerges when

$$DC(\alpha, \Omega) \approx 0.$$

Stress-Induced Internal Geometry

Stress generates an internal metric

$$\gamma_{AB}(\alpha, \Omega) = \delta_{AB} + \alpha_1 \sigma_{AB}(\alpha, \Omega) + \alpha_2 I_1(\alpha, \Omega) \delta_{AB},$$

with the first stress invariant

$$I_1(\alpha, \Omega) = \text{tr}(\sigma(\alpha, \Omega)).$$

This metric is pre-spatial and forms the basis of the later spatial metric $g_{ij}(x, \Omega)$ only through binding.

Internal Curvature

Curvature arises from derivatives of the internal metric:

$$\mathcal{R}_{ABCD}(\alpha, \Omega) = R(\gamma_{AB}(\alpha, \Omega)).$$

It describes geometric amplification or bending caused by stress gradients. This curvature is pure medium curvature; only bound curvature forms material cores.

Rotational–Geometric Structure

Rotation is encoded in the rotation tensor:

$$L_{AB}(\alpha, \Omega) = \frac{1}{2}(\nabla_A v_B - \nabla_B v_A),$$

which describes rotational–shear modes. These modes propagate through the internal geometry and form the basis of rotational–geometric patterns. Bound rotation produces a stable rotational core.

Constitutive Operator

The constitutive operator $F(Vv, \sigma)$ determines the reorganization of stress under updating:

$$D\sigma_{AB}(\alpha, \Omega) = F(Vv, \sigma)_{AB},$$

with geometric transport

$$Vv(\sigma_{AB}) = v_C \nabla_C \sigma_{AB}.$$

It contains rotation-related response terms, shear reorganization, curvature amplification, and geometric relaxation.

Absence of Physical Coordinates

All mathematical objects of the MCM are defined independently of physical space and physical time. The index α is not a spatial point but a medium index. Physical space arises only through bound geometry, which induces a map

$$x = X(\alpha, \Omega)$$

and stabilizes the spatial metric

$$g_{ij}(x, \Omega) = \gamma_{AB}(\alpha, \Omega) \frac{\partial X^A}{\partial x^i} \frac{\partial X^B}{\partial x^j}.$$

Summary

The mathematical structure of the MCM is based on internal geometric and tensorial quantities of a single medium. Stress generates internal geometry, curvature arises from stress gradients, rotational-geometric patterns propagate through the internal geometry, and the constitutive operator governs internal reorganization. This structure is pre-spatial and unbound. Only stable binding produces bound geometry (matter), whose external appearance is physical space.

6 Dynamics of the Medium

The dynamics of the Monistic Continuum Model (MCM) arise entirely from internal updating. Neither physical time nor external forces or field constructs are assumed. All evolution corresponds to changes in the geometric, rotational, and stress-induced structure of the medium. The update rate $\tau(\alpha, \Omega)$ describes the local reactivity of the medium and modulates the strength of this internal reorganization.

Internal Updating

The internal development of the medium is described by the update derivative

$$DC(\alpha, \Omega)$$

acting on the pre-spatial state map

$$C(\alpha, \Omega) = (\rho, v_A, \sigma_{AB}, \tau),$$

and reorganizing density, geometric motion, stress, and reactivity. As long as

$$DC(\alpha, \Omega) \not\approx 0,$$

the geometry remains unbound. Only approximate update invariance allows the formation of stable bound geometry.

Stress Gradients and Acceleration

Geometric motion v_A responds exclusively to stress gradients. Acceleration is given by a geometric operator G :

$$Dv_A(\alpha, \Omega) = G(\sigma(\alpha, \Omega), \nabla\sigma(\alpha, \Omega)).$$

No external forces exist; motion is induced purely geometrically.

Transport of Stress

Geometric motion transports stress through the medium. The transport operator is

$$Vv(\sigma_{AB}) = v_C \nabla_C \sigma_{AB}.$$

This transport is entirely pre-spatial and describes the movement of stress patterns within the internal geometry.

Constitutive Dynamics

The reorganization of stress under updating is governed by the constitutive operator:

$$D\sigma_{AB} = F(Vv, \sigma)_{AB}.$$

The operator contains rotation-related response terms, shear reorganization, curvature amplification, and geometric relaxation. It determines whether geometric configurations stabilize, propagate, or dissipate.

Rotational–Geometric Dynamics

Rotation is encoded in the rotation tensor

$$L_{AB} = \frac{1}{2}(\nabla_A v_B - \nabla_B v_A)$$

and rotational–geometric patterns propagate through the internal geometry, interacting with stress gradients. Bound rotation produces stable rotational cores within bound geometry.

Curvature Dynamics

Curvature arises from derivatives of the internal metric:

$$\mathcal{R}_{ABCD} = R(\gamma_{AB}).$$

Curvature dynamics describe how geometric amplification or bending evolves under updating. Curvature interacts with stress and rotation and influences the stability of bound structures.

Role of the Update Rate

The update rate $\tau(\alpha, \Omega)$ influences:

- the stabilization of bound geometry,
- the propagation of rotational–geometric patterns,
- the strength of geometric relaxation,
- the development of large-scale domains.

It is not a time coordinate but a material property of the medium.

Absence of Temporal Dynamics

The index Ω is not a temporal parameter. All dynamics are structural reorganization within the internal geometry of the medium.

Summary

The dynamics of the MCM arise from internal updating driven by stress gradients, rotation, and geometric coupling. Unbound geometry remains dynamic and non-material. Only when $DC(\alpha, \Omega) \approx 0$ holds does bound geometry emerge, whose external appearance is physical space. All dynamics are internal and geometric.

7 Formalization of the Operators

The Monistic Continuum Model (MCM) relies on two central operators that govern the internal evolution of the medium: the update derivative DC and the constitutive operator $F(Vv, \sigma)$. Both operators must be formulated without reference to physical fields, forces, or temporal dynamics. This section provides a mathematically consistent, invariant-based partial formalization of these operators, resolving all structural limitations identified in earlier chapters.

Update Derivative

The update derivative acts on the pre-spatial state map

$$C(\alpha, \Omega) = (\rho(\alpha, \Omega), v_A(\alpha, \Omega), \sigma_{AB}(\alpha, \Omega), \tau(\alpha, \Omega)).$$

It is defined component-wise:

$$DC(\alpha, \Omega) = (D\rho, Dv_A, D\sigma_{AB}, D\tau).$$

The operator D is not a temporal derivative; it describes structural reorganization of the medium. Its algebraic properties are:

- D is linear on scalar and tensor quantities.
- D commutes with contractions:

$$D(T_{AB}S_{AB}) = (DT_{AB})S_{AB} + T_{AB}(DS_{AB}).$$

- D acts on covariant derivatives via

$$D(\nabla_A T_{BC}) = \nabla_A(DT_{BC}) + K_{A,BC},$$

where $K_{A,BC}$ is a geometric correction induced by the update of the stress-generated metric.

- D preserves geometric invariants of the medium.

A discrete approximation is

$$C(\alpha, \Omega + \Delta\Omega) \approx C(\alpha, \Omega) + \Delta\Omega DC(\alpha, \Omega),$$

where $\Delta\Omega$ is an update increment without temporal meaning.

Constitutive Operator

The constitutive operator governs the update of the stress tensor:

$$D\sigma_{AB} = F(Vv, \sigma)_{AB}.$$

Since the MCM is purely geometric, F must be formulated using only internal quantities of the medium. A full specification is not required; however, mathematical consistency demands an invariant-based partial formalization. The operator is decomposed into four geometric contributions:

$$F(Vv, \sigma)_{AB} = F_{AB}^{(\text{rot})} + F_{AB}^{(\text{shear})} + F_{AB}^{(\text{curv})} + F_{AB}^{(\text{relax})}.$$

Rotational Contribution

Rotation is encoded in

$$L_{AB} = \frac{1}{2}(\nabla_A v_B - \nabla_B v_A).$$

The rotational contribution is

$$F_{AB}^{(\text{rot})} = \beta_1 L_{AC} \sigma_{CB} + \beta_2 \sigma_{AC} L_{CB}.$$

Shear Contribution

Shear transport is described by

$$Vv(\sigma_{AB}) = v_C \nabla_C \sigma_{AB}.$$

The shear contribution is

$$F_{AB}^{(\text{shear})} = \gamma_1 Vv(\sigma_{AB}) + \gamma_2 \nabla_A v_C \sigma_{CB} + \gamma_3 \sigma_{AC} \nabla_C v_B.$$

Curvature Contribution

The stress-induced metric

$$\gamma_{AB} = \delta_{AB} + \alpha_1 \sigma_{AB} + \alpha_2 I_1 \delta_{AB}$$

generates internal curvature

$$\mathcal{R}_{ABCD} = R(\gamma_{AB}).$$

Curvature contributes via

$$F_{AB}^{(\text{curv})} = \delta_1 \mathcal{R}_{ACBD} \sigma_{CD} + \delta_2 \mathcal{R}_{AB}.$$

Relaxation Contribution

Relaxation smooths stress concentrations:

$$F_{AB}^{(\text{relax})} = -\lambda_1 \sigma_{AB} - \lambda_2 I_1 \delta_{AB}.$$

Combined Operator

The invariant-based partial formalization of the constitutive operator is

$$\begin{aligned} D\sigma_{AB} = & \beta_1 L_{AC} \sigma_{CB} + \beta_2 \sigma_{AC} L_{CB} + \gamma_1 Vv(\sigma_{AB}) \\ & + \gamma_2 (\nabla^A v_C) \sigma_{CB} + \gamma_3 \sigma_{AC} (\nabla_C v_B) \\ & + \delta_1 \mathcal{R}_{ACBD} \sigma_{CD} + \delta_2 \mathcal{R}_{AB} - \lambda_1 \sigma_{AB} - \lambda_2 I_1 \delta_{AB} \end{aligned}$$

This representation is:

- geometrically consistent,
- invariant-based,
- compatible with the stress-induced metric,
- fully pre-spatial,
- mathematically sufficient for numerical implementation,
- compliant with the V08 ontology.

Summary

The update derivative DC and the constitutive operator $F(Vv, \sigma)$ are now partially formalized in a mathematically consistent and invariant-based manner. This resolves all structural limitations identified in earlier sections and provides a solid foundation for numerical simulation, analytical development, and further refinement of the MCM.

8 Geometry from Stress

In the Monistic Continuum Model (MCM), geometry is fundamental and arises directly from the stress state of the medium. In its pre-spatial form, the medium consists solely of *unbound geometry*, which carries no matter and forms no spatial structure. Only when patterns of stress, rotation, and curvature form a stable binding does *bound geometry* emerge. In the MCM, bound geometry is identical to matter and forms the foundation of emergent physical space.

Stress-Induced Internal Metric

The internal geometric structure of the medium is determined by the stress-induced metric

$$\gamma_{AB} = \delta_{AB} + \alpha_1 \sigma_{AB} + \alpha_2 I_1 \delta_{AB},$$

with the first stress invariant

$$I_1 = \text{tr}(\sigma),$$

This metric is pre-spatial and describes internal relations of unbound geometry. Only through stable binding does it become the basis of the later spatial metric $g_{ij}(x, \Omega)$.

Curvature from Stress Gradients

Curvature arises from derivatives of the internal metric:

$$\mathcal{R}_{ABCD} = R(\gamma_{AB}),$$

and represents geometric amplification or bending caused by stress gradients. This curvature is pure medium curvature; only bound curvature forms material cores.

Interaction of Stress, Rotation, and Curvature

Stress interacts with rotation and curvature to produce complex geometric patterns. Rotation is encoded in the tensor

$$L_{AB} = \frac{1}{2}(\nabla_A v_B - \nabla_B v_A)$$

and propagates through the internal geometry. The interplay of these quantities determines:

- the formation of bound geometry,

- the propagation of rotational–geometric patterns,
- the stabilization of geometric domains,
- the emergence of spatial structure.

Geometric Stability

Geometric stability arises when internal updating becomes approximately invariant:

$$DC(\alpha, \Omega) \approx 0.$$

Stable configurations form bound geometry. Bound geometry is matter, and physical space is the external appearance of this bound structure.

Geometry Without Space

Before bound geometry forms, the medium exhibits:

- stress-induced internal metric,
- internal curvature patterns,
- rotational–geometric behavior,
- geometric motion.

However, it lacks:

- stable axes,
- stable distances,
- stable orientation,
- measurable spatial structure.

Unbound geometry is pre-spatial. Only through binding does matter arise, and only matter generates space.

Role of the Update Rate

The update rate $\tau(\alpha, \Omega)$ influences geometric stability by determining how quickly stress reorganizes. High reactivity leads to rapid change, while low reactivity promotes stability. It therefore plays a central role in the formation of bound geometry and thus physical space.

Summary

Geometry in the MCM arises from stress and is shaped by rotation and curvature. Unbound geometry is pre-spatial and carries no matter. Only stable binding produces bound geometry, whose external appearance is physical space. The update rate controls stability and influences the emergence of spatial domains.

9 Matter as Bound Geometry

In the Monistic Continuum Model (MCM), matter is not a fundamental entity. It arises only when stress, rotation, and curvature form a stable binding. This binding produces *bound geometry*, which in the MCM is identical to matter. Matter is therefore not a structure within space, but bound internal geometry from which physical space emerges.

Definition of Bound Geometry

Bound geometry exists when internal updating becomes approximately invariant:

$$DC(\alpha, \Omega) \approx 0.$$

In this state, stress, rotation, and curvature remain stable under updating and form persistent, localized structures. This stability is the ontological basis of matter.

Stress–Rotation Binding

Bound geometry arises through strong coupling between stress and rotation. The rotation tensor

$$L_{AB} = \frac{1}{2}(\nabla_A v_B - \nabla_B v_A)$$

interacts with the stress tensor σ_{AB} and produces localized rotational–geometric cores. These cores stabilize the surrounding region and generate persistent curvature.

The stable binding configuration is the *vortex*. It forms the elementary structure of bound geometry and thus the foundation of all matter.

Localization of Curvature

Curvature becomes strongly localized around binding cores:

$$\mathcal{R}_{ABCD} = R(\gamma_{AB}).$$

This localization enhances geometric stability and prevents dissipation. Bound curvature is a characteristic feature of material cores.

Update Stability

The stability of bound geometry depends on the update rate $\tau(\alpha, \Omega)$. High reactivity destabilizes binding cores, while moderate reactivity promotes persistence. The update rate therefore determines whether bound structures can exist long-term.

Emergent Properties of Bound Geometry

Bound geometry exhibits emergent properties:

- localized stress concentration,
- persistent rotational–geometric cores,
- stable curvature domains,
- geometric anchoring of spatial axes,
- resistance to geometric dissipation.

These properties correspond to mass-like, charge-like, and interaction-like behavior, but arise solely from bound geometry.

Bound Geometry as the Basis of Spatial Structure

Bound geometry stabilizes the internal structure of the medium and enables the spatial mapping

$$x = X(\alpha, \Omega).$$

The representation of bound geometry in emergent space is

$$M(x, \Omega).$$

Matter therefore forms the basis of spatial structure. Space is the external appearance of bound geometry.

Summary

Matter is not fundamental in the MCM. It arises as bound geometry stabilized through stress, rotation, and curvature. Bound geometry satisfies approximate update invariance and forms the foundation of physical space as well as all observable phenomena.

10 Vortex Geometry

In the Monistic Continuum Model (MCM), the vortex is the first stable form of *bound geometry*. A vortex is not a structure within space, but a pre-spatial binding of stress, rotation, and curvature that remains stable under internal updating. Since bound geometry in the MCM is identical to matter, the vortex constitutes the elementary structure of material defects.

A vortex defect is a localized configuration that satisfies the condition of update invariance:

$$DC(\alpha, \Omega) \approx 0.$$

Only a single geometric realization satisfies all axiometric requirements of the MCM; other known topological or field-theoretic models (Skyrmion, string model) do not meet these conditions and therefore cannot form bound geometry.

Axiometric Requirements for an MCM Vortex

A valid vortex in the MCM must satisfy the following conditions:

1. **Rotational Core:** The rotation tensor is nontrivial:

$$L_{AB} \neq 0.$$

2. **Stress Localization:** The stress tensor is concentrated in the core:

$$\sigma_{AB} \text{ localized.}$$

3. **Bound Curvature Domain:** Internal curvature forms a closed, stable structure:

$$\mathcal{R}_{ABCD} \text{ stable.}$$

4. **Topological Stability:** The structure possesses a well-defined linking number.

5. **Update Invariance:** The configuration remains stable under updating:

$$DC(\alpha, \Omega) \approx 0.$$

Only the MCM vortex satisfies all five conditions and forms bound geometry. Thus, it is the only structure capable of generating matter.

Why Skyrmions and String/Knot Models Are Not MCM Vortices

Skyrmion: A Skyrmion possesses no true rotational core, no stress-induced geometry, and no localized curvature domain. It satisfies neither

$$L_{AB} \neq 0, \quad \sigma_{AB} \text{ localized}, \quad DC \approx 0,$$

and therefore cannot form bound geometry.

String/Knot Models: Knot structures possess topology but no rotation, no stress-induced geometry, and no bound curvature domain. They satisfy none of the geometric axioms of the MCM and therefore cannot generate matter.

Conclusion: Both structures are purely topological objects without geometric binding and cannot form material defects in the MCM.


The Stable MCM Vortex

The stable vortex possesses a closed, intertwined rotation–curvature structure that satisfies all axiometric requirements:

- closed rotation lines (nontrivial L_{AB}),
- localized stress concentration in the core,
- stable bound curvature domain with defined linking number,
- discrete stability modes under D ,
- unique topological charge.

Thus, the vortex is the elementary form of bound geometry and the foundation of all material structures in the MCM.

Representation of a Vortex



Geometric parameters: curvature, spiral pitch, compression
 Topological charge q_W , winding number, linking index
 Discrete stability modes W_n , spectral gap ΔW
 Local internal dynamics $D\sigma_{AB}$, coupling of $L_{AB}, \mathcal{R}_{ABCD}$
 Stability norm $S(W_n)$, $DC(\alpha, \Omega) \approx 0$
 Cluster operator C : aggregation of bound defects
 MCM embedding: $\sigma_{AB}(\alpha, \Omega)$, $C(\alpha, \Omega)$
 Emergent domains: bound geometry $\rightarrow M(x, \Omega)$

Vortex as the Basis of Emergent Matter

Since the stable vortex satisfies all axioms of the MCM, it is the sole candidate for stable material defects. Its intertwined rotation–curvature structure produces:

- discrete stability modes (quantization),
- defined topological charges,
- stable stress anchors,
- persistent geometric axes,
- robust cluster formation.

The vortex is the only structure capable of generating bound geometry. It therefore forms the geometric foundation of all emergent material structures in the MCM.

11 Electromagnetism in the MCM

In the Monistic Continuum Model (MCM), electromagnetic behavior does not arise from physical fields or charges, but from rotational–geometric modes of the medium. These modes are pre-spatial structures of unbound geometry and become stable and spatially interpretable only through bound geometry (matter). The ontology remains fully geometric: electromagnetism is a rotational mode of the medium, not an external field.

Rotational–Geometric Modes

Rotational–geometric modes are encoded in the rotation tensor

$$L_{AB} = \frac{1}{2}(\nabla_A v_B - \nabla_B v_A).$$

These modes propagate through the stress-induced internal geometry and produce signatures that, in physical theories, are interpreted as electromagnetic waves. In the MCM they correspond to:

- transverse rotational–geometric patterns,
- stress-coupled rotational propagation,
- curvature-modulated rotational behavior,
- defect-induced rotational scattering.

They already exist in unbound geometry but become stable and measurable only through bound geometry.

Stress–Rotation Coupling

Electromagnetic behavior arises from coupling between stress and rotation. The stress tensor σ_{AB} interacts with the rotation tensor L_{AB} and produces propagating rotational–geometric structures. Stability of these modes requires approximate update invariance:

$$DC(\alpha, \Omega) \approx 0.$$

Only bound geometry (matter) enables persistently propagating rotational modes.

Curvature Interaction

The stress-induced internal metric

$$\gamma_{AB} = \delta_{AB} + \alpha_1 \sigma_{AB} + \alpha_2 I_1 \delta_{AB},$$

generates internal curvature

$$\mathcal{R}_{ABCD} = R(\gamma_{AB}).$$

Rotational–geometric modes interact with this curvature and produce effects such as:

- curvature gradients,
- rotation–curvature coupling,
- defect-induced curvature sinks,
- curvature-modulated propagation.

These effects are pre-spatial and become spatially interpretable only through bound geometry.

Defect Level of Electromagnetic Behavior

Bound geometry produces local rotational–geometric behavior. The internal structure of a stable vortex generates:

- rotational–geometric cores,
- stress-induced rotational anchoring,
- bound curvature localization,
- discrete rotational stability modes.

These features correspond to charge-like and dipole-like behavior in physical interpretations, but in the MCM they arise solely from bound geometry.

Quantized Rotational Modes

Rotational–geometric modes exhibit discrete stability patterns because only certain rotational configurations remain invariant under updating. This discreteness corresponds to quantized electromagnetic behavior. It arises from:

- rotational stability conditions,
- stress-induced mode selection,
- curvature-induced mode filtering,
- constraints imposed by the update rate.

Quantization is therefore an effect of bound geometry.

Summary

Electromagnetism in the MCM arises from rotational–geometric behavior of the medium. Stress, rotation, and curvature interact to produce pre-spatial rotational modes. Only through bound geometry (matter) do these modes become stable, measurable, and spatially interpretable. Electromagnetic phenomena are geometric effects and require no physical fields or charges.

12 Quantization as a Geometric Effect

In the Monistic Continuum Model (MCM), quantization is not a fundamental principle. It arises solely from the stability conditions of *bound geometry*. Unbound geometry is continuous and not quantized. Only stable binding of stress, rotation, and curvature produces discrete stability modes. Bound geometry is identical to matter in the MCM; quantization is therefore an effect of this bound structure.

Update-Invariant Geometric Modes

Quantization arises from the requirement that bound configurations satisfy approximate update invariance:

$$DC(\alpha, \Omega) \approx 0.$$

Only specific bound patterns satisfy this condition. They form discrete stability modes of stress, rotation, and curvature. The medium therefore supports a finite set of stable bound configurations, each representing a quantized geometric state.

Rotational Stability Modes

Rotational–geometric behavior is encoded in the rotation tensor:

$$L_{AB} = \frac{1}{2}(\nabla_A v_B - \nabla_B v_A).$$

Stable rotational modes arise only in bound geometry. The rotational–geometric structure of a Hopf vortex remains invariant under updating and forms discrete families because only certain rotational patterns maintain geometric stability. This discreteness corresponds to spin-like behavior in physical interpretations.

Stress-Induced Quantization

Stress contributes to quantization through the stress tensor σ_{AB} . Stable stress configurations require that both magnitude and orientation of the stress field remain invariant under updating. This produces discrete stress patterns that represent quantized bound configurations.

The stress-induced internal metric

$$\gamma_{AB} = \delta_{AB} + \alpha_1 \sigma_{AB} + \alpha_2 I_1 \delta_{AB},$$

supports only certain stable bound curvature structures. These structures correspond to quantized curvature modes.

Curvature-Based Quantization

Curvature arises from the internal metric:

$$\mathcal{R}_{ABCD} = R(\gamma_{AB}).$$

Only specific curvature patterns remain invariant under updating. Stable bound curvature configurations form discrete families and correspond to quantized geometric behavior.

Defect Level of Quantization

Material defects exhibit quantized behavior because their bound geometric structure supports only certain stable configurations. Defects therefore possess:

- discrete rotational–geometric cores,
- discrete curvature sinks,
- discrete stress patterns,
- discrete stability modes.

These discrete geometric features correspond to mass-like, charge-like, and spin-like quantization in physical interpretations. The discreteness is an effect of bound geometry.

Update Rate and Quantization

The update rate $\tau(\alpha, \Omega)$ influences which bound modes remain stable. High reactivity destabilizes many configurations and reduces the number of stable modes. Moderate reactivity allows a richer

variety of quantized patterns. Quantization therefore depends directly on the reactivity of the medium.

Summary

Quantization in the MCM arises from bound geometric stability. Only specific stress, rotation, and curvature patterns remain invariant under internal updating and form discrete families of stable bound configurations. Material defects exhibit quantized behavior because their bound geometric structure supports only specific stability modes. Quantization is therefore a geometric effect of bound geometry and not a fundamental principle.

13 Cosmology of the Medium

The cosmology of the Monistic Continuum Model (MCM) does not describe the evolution of a physical universe in space and time. It characterizes the large-scale behavior of *unbound and bound geometry* under internal updating. Global stress reorganization, domain formation, and clustering of bound geometry generate emergent structures that correspond, in physical interpretations, to cosmological phenomena. The ontology remains entirely geometric and independent of physical space or physical time.

Global Stress Structure

The medium possesses a global stress distribution described by the pre-spatial stress tensor σ_{AB} . Large-scale behavior arises from slow reorganization of this stress distribution. Global stress gradients generate:

- large-scale internal curvature fields,
- rotational–geometric patterns,
- tendencies toward clustering of bound geometry,
- expansion or contraction of unbound domains.

These effects correspond to cosmological expansion, contraction, and structure formation, but are purely geometric phenomena.

Domain Formation

The stress-induced internal metric γ_{AB} supports the formation of large-scale geometric domains. Domains arise when regions of the medium develop coherent stress, rotation, and curvature patterns that remain approximately stable under updating:

$$DC(\alpha, \Omega) \approx 0.$$

Domains exhibit:

- coherent internal curvature profiles,
- stable rotational–geometric behavior,
- persistent stress gradients,
- clustering of bound geometry (vortices).

These domains correspond to cosmic structures such as filaments, voids, and clusters, but are pre-spatial geometric regions.

Defect Clustering

Bound geometry acts as a geometric anchor. Its stability condition $DC \approx 0$ enables clustering in regions of compatible stress and compatible curvature. Clustering arises through:

- curvature sinks,
- rotational–geometric attraction,
- alignment of stress gradients,
- modulation by the update rate.

This clustering corresponds to large-scale structure formation, but is a pre-spatial reorganization of bound geometry.

Rotational–Geometric Cosmology

Large-scale rotational–geometric patterns propagate through the medium. They are encoded in the rotation tensor L_{AB} and generate domain-level coherence. Rotational patterns can:

- align domains,

- amplify rotational flows through curvature modulation,
- stabilize geometric domains.

These effects correspond to cosmic rotation, polarization, and anisotropy.

Curvature Evolution

Internal curvature arises from the stress-induced metric:

$$\mathcal{R}_{ABCD} = R(\gamma_{AB}).$$

Large-scale curvature evolves under updating and determines:

- expansion or contraction of geometric domains,
- patterns of clustering of bound geometry,
- propagation of rotational–geometric patterns,
- global geometric stability.

These effects correspond to gravitational structure formation and curvature evolution in physical interpretations.

Role of the Update Rate

The update rate τ influences cosmological behavior by determining how quickly large-scale geometric structures reorganize. High reactivity leads to rapid domain development; low reactivity promotes long-term stability.

Cosmology therefore depends on the reactivity of the medium, not on physical time.

Summary

Cosmology in the MCM arises from the large-scale behavior of unbound and bound geometry. Global stress reorganization, domain formation, clustering of bound geometry, rotational–geometric propagation, and curvature evolution generate emergent structures that correspond to cosmological phenomena. Cosmology is a geometric effect of the medium and does not rely on physical space, physical time, or field constructs.

14 Numerical Methods and Simulation

Numerical simulation in the Monistic Continuum Model (MCM) serves to approximate the internal updating of the medium. Since the model uses neither physical space, physical time, nor field constructs, numerical methods operate exclusively on pre-spatial quantities: stress, rotation, curvature, density, and update rate. Simulation therefore corresponds to iterating the pre-spatial state map

$$C(\alpha, \Omega) = (\rho(\alpha, \Omega), v_A(\alpha, \Omega), \sigma_{AB}(\alpha, \Omega), \tau(\alpha, \Omega))$$

under discrete approximations of the update derivative.

Discrete Update Approximation

The internal update operator D is approximated by a discrete increment:

$$C(\alpha, \Omega + \Delta\Omega) \approx C(\alpha, \Omega) + \Delta\Omega DC(\alpha, \Omega),$$

where $\Delta\Omega$ is a numerical computation parameter. It has no temporal meaning; it controls the resolution of the updating process.

Approximation of Geometric Derivatives

Geometric derivatives such as $\nabla_A v_B$ and $\nabla_C \sigma_{AB}$ are approximated using geometric finite-difference schemes defined on the stress-induced internal geometry of the medium. These schemes operate on the internal metric

$$\gamma_{AB} = \delta_{AB} + \alpha_1 \sigma_{AB} + \alpha_2 I_1 \delta_{AB},$$

and ensure that numerical derivatives respect the pre-spatial geometry.

Evaluation of the Constitutive Operator

The constitutive operator

$$D\sigma_{AB} = F(Vv, \sigma)_{AB}$$

is evaluated numerically by computing:

- rotation-related response terms from L_{AB} ,

- shear reorganization from $\nabla_C \sigma_{AB}$,
- curvature amplification from $\mathcal{R}_{ABCD}$,
- geometric relaxation from stress–rotation coupling.

All terms are approximated using finite-difference or finite-volume methods.

Simulation of Rotational–Geometric Modes

Rotational–geometric modes, encoded in

$$L_{AB} = \frac{1}{2}(\nabla_A v_B - \nabla_B v_A),$$

are simulated by iterating rotation-related update equations. Numerical stability requires approximate update invariance:

$$DC(\alpha, \Omega) \approx 0.$$

Only under this condition do rotational–geometric waves propagate without numerical dissipation.

Curvature Computation

Internal curvature is computed from the stress-induced metric:

$$\mathcal{R}_{ABCD} = R(\gamma_{AB}).$$

Curvature computation is essential for:

- formation of bound geometry (defects),
- development of geometric domains,
- curvature-modulated rotational propagation,
- stress-gradient-driven reorganization.

Defect Simulation

Bound geometry satisfies approximate update invariance:

$$DC(\alpha, \Omega) \approx 0.$$

Numerical simulation identifies bound geometry by regions in which stress, rotation, and curvature patterns remain stable over many steps. Stability is monitored through:

- stress localization,
- persistence of rotational–geometric cores,
- curvature amplification,
- influence of the update rate.

Domain-Level Simulation

Large-scale geometric domains arise from global stress reorganization. Simulation tracks:

- domain curvature profiles,
- clustering of bound geometry,
- rotational–geometric coherence,
- development of large-scale stress gradients.

These processes correspond to cosmology-like behavior but remain pre-spatial geometry.

Numerical Stability

Numerical stability requires appropriate choice of $\Delta\Omega$. Instability occurs when:

- stress gradients exceed numerical resolution,
- rotational–geometric modes are under-resolved,
- curvature localization becomes too pronounced,
- the update rate varies strongly.

Stable simulation requires adaptive increments and geometric refinement.

Summary

Numerical simulation in the MCM approximates the internal updating of geometric, rotational, and stress-induced structures. Geometric derivatives, constitutive operators, rotational–geometric modes, curvature, defects, and domains are simulated using geometric finite-difference methods. All simulation behavior arises from the internal geometry of the medium and is independent of physical space or physical time.

15 Experimental Accessibility

The Monistic Continuum Model (MCM) describes a continuous medium whose internal dynamics are geometric rather than physical. Although the model assumes neither physical space, physical time, fields, nor particles, its bound geometric structures generate signatures that can be interpreted through physical measurement frameworks. Experimental accessibility therefore arises only when *bound geometry* is present and a spatial interpretation becomes possible.

Bound geometry satisfies approximate update invariance:

$$DC(\alpha, \Omega) \approx 0,$$

and enables the spatial mapping

$$x = X(\alpha, \Omega),$$

through which geometric structures become measurable.

Defect-Based Observables

Bound geometry produces stable, localized structures that appear as physical observables. These include:

- localized stress concentrations,
- persistent rotational–geometric cores,
- bound curvature sinks,
- discrete stability modes.

These signatures correspond to mass-like, charge-like, and spin-like behavior.

Rotational–Geometric Observables

Rotational–geometric modes, encoded in

$$L_{ij} = \frac{1}{2}(\nabla_i v_j - \nabla_j v_i),$$

produce wave-like signatures that are interpreted as electromagnetism-like behavior. Observable effects include:

- transverse rotational–geometric patterns,
- curvature-modulated propagation,
- rotational–geometric coherence across domains,
- defect-induced rotational scattering.

Curvature Observables

The spatial metric arises from bound internal geometry:

$$g_{ij} = \gamma_{AB} \frac{\partial X^A}{\partial x^i} \frac{\partial X^B}{\partial x^j},$$

with spatial curvature tensor

$$R_{ijkl} = R(g_{ij}).$$

Observable curvature signatures include:

- curvature gradients around bound cores,
- domain-level curvature profiles,
- curvature-modulated rotational behavior,
- curvature amplification during binding.

These correspond to gravity-like behavior.

Observables of the Update Rate

The update rate $\tau(x, \Omega)$ determines the reactivity of bound geometry. Variations produce observable effects such as:

- changes in defect stability,
- shifts in propagation speed of rotational–geometric modes,
- geometric relaxation behavior,
- domain expansion or contraction.

These effects appear dynamic, even though τ is not time.

Domain-Level Observables

Large-scale geometric domains arise from global stress reorganization. Observable signatures include:

- clustering of bound geometry,
- rotational–geometric coherence,
- domain curvature profiles,
- stress-gradient-driven domain evolution.

These correspond to cosmological structures.

Measurement Translation

Geometric structures are translated into observables through physical interpretation frameworks. Typical correspondences include:

- stress localization → mass-like behavior,
- rotational–geometric cores → spin-like behavior,
- bound curvature sinks → gravity-like behavior,
- rotational–geometric waves → electromagnetism-like behavior.

These translations make geometric effects experimentally accessible without altering the ontology of the MCM.

Summary

Experimental accessibility arises only through bound geometry. Defects, rotational modes, curvature, variations of the update rate, and domain behavior generate observable signatures when interpreted through physical measurement frameworks. All observables are geometric effects of bound geometry and do not rely on physical fields, particles, or forces.

16 Applications in Applied Physics

Although the Monistic Continuum Model (MCM) assumes neither physical space, physical time, fields, particles, nor forces, its geometric structures can be translated into physical interpretative frameworks. In this way, the model provides conceptual and structural insights relevant to applied physics. Applications arise from stable or propagating configurations of bound and unbound geometry once a spatial interpretation becomes possible.

Material Stability and Defect Dynamics

Bound geometry satisfies approximate update invariance:

$$DC(\alpha, \Omega) \approx 0.$$

Its stability provides insights into material-relevant phenomena such as:

- formation and annihilation of bound structures,
- stress localization and amplification,
- persistence of rotational–geometric cores,
- curvature-driven structural stability.

These mechanisms correspond to fracture dynamics, defect propagation, and stability analysis in materials science.

Wave Propagation and Signal Transmission

Rotational–geometric modes, encoded in

$$L_{AB} = \frac{1}{2}(\nabla_A v_B - \nabla_B v_A),$$

produce wave-like behavior when spatially interpreted. Applications include:

- rotational–geometric waveguides,
- curvature-modulated signal transmission,
- defect-induced scattering,
- domain-level coherence.

These correspond to electromagnetic-like propagation, optical behavior, and signal transmission.

Curvature-Driven Structural Effects

Internal curvature arises from the stress-induced metric γ_{AB} with curvature tensor

$$\mathcal{R}_{ABCD} = R(\gamma_{AB}).$$

Curvature-driven behavior provides insights into:

- structural reinforcement,
- geometric buckling,
- curvature-modulated stability,
- domain-level deformation.

These correspond to gravity-like structural effects in physical interpretations.

Dynamic Response and Reactivity

The update rate τ determines the reactivity of bound and unbound geometry. Applications include:

- dynamic stability analysis,
- reactivity-driven structural adaptation,
- modulation of rotational–geometric propagation,
- domain-level relaxation behavior.

These correspond to dynamic effects, even though τ is not time.

Domain-Level Behavior

Large-scale geometric domains arise from global stress reorganization. Their behavior provides insights into:

- large-scale structural coherence,
- clustering of bound geometry,
- curvature-driven domain evolution,
- rotational–geometric alignment.

These correspond to cosmology-like behavior and can be used for structural modeling.

Translation into Applied Frameworks

Applications rely on translation schemes that connect geometric structures of the MCM with physical constructs:

- stress localization → mass-like behavior,
- rotational–geometric cores → charge-like behavior,
- bound curvature sinks → gravity-like behavior,
- rotational–geometric waves → electromagnetism-like behavior,
- discrete stability modes → quantum-like behavior.

These translations enable physical interpretation without altering the monistic ontology of the MCM.

Summary

Applications of the MCM in applied physics arise from geometric, rotational, and stress-induced structures of the medium. Bound geometry, rotational modes, curvature, reactivity, and domain behavior correspond to physical phenomena when interpreted through appropriate translation schemes. The model provides conceptual insights into stability, propagation, structural reinforcement, and large-scale behavior.

17 Limitations of the MCM

The Monistic Continuum Model (MCM) provides a unified geometric ontology in which all observable phenomena arise from stress, rotation, curvature, and internal updating of a single continuous medium. Despite its conceptual coherence, the model possesses limitations resulting from its monistic structure, its pre-spatial formulation, and the absence of physical fundamental entities. These limitations mark areas that require further development.

Absence of Physical Fundamental Entities

The MCM assumes neither physical space, physical time, fields, particles, nor forces. This ensures ontological consistency but complicates direct comparison with physical theories that rely on these constructs. Translation schemes are required to map pre-spatial geometric structures to physical observables. These translations are not unique and may vary depending on the physical framework.

Dependence on Geometric Interpretation

All behavior in the MCM arises from internal geometric quantities:

$$\sigma_{AB}, \quad L_{AB}, \quad \mathcal{R}_{ABCD}, \quad \tau.$$

This complete geometric dependence limits the model's ability to describe phenomena that are formulated non-geometrically in established physics. The model must rely on geometric analogies rather than direct physical constructs.

Lack of Closed Constitutive Relations

The constitutive operator

$$D\sigma_{AB} = F(Vv, \sigma)_{AB}$$

is conceptually defined but not fully specified in closed form. This limits predictive capability and requires numerical or phenomenological approximations. Without explicit constitutive relations, certain behaviors cannot currently be derived analytically, including:

- stability conditions of bound geometry,
- transitions between unbound and bound geometry,
- large-scale domain development.

Complexity of Numerical Simulation

Simulation of the MCM requires geometric finite-difference or finite-volume methods operating on the stress-induced internal geometry. Numerical challenges include:

- resolving strongly localized bound curvature,
- maintaining rotational–geometric stability,
- handling stress-gradient singularities,
- adaptive selection of update increments $\Delta\Omega$,
- ensuring geometric consistency across domains.

These challenges limit resolution and scale, particularly during formation of bound geometry (vortices).

Limited Analytical Tools

Because the MCM uses neither physical coordinates nor temporal evolution, many analytical tools from classical physics, quantum theory, and relativity cannot be applied directly. New geometric tools must be developed to analyze:

- stability of bound geometry,
- rotational–geometric propagation,
- curvature evolution,
- behavior of large-scale domains.

These tools are still under development.

Interpretation Ambiguity

Translation schemes linking geometric structures to physical observables are not unique. For example:

- bound stress localization $\rightarrow$ mass-like behavior,
- rotational–geometric cores $\rightarrow$ charge-like behavior,
- bound curvature sinks $\rightarrow$ gravity-like behavior.

Different physical frameworks may interpret the same geometric structure differently. This leads to ambiguity in physical interpretation and complicates direct experimental correspondence.

Lack of Empirical Calibration

The MCM has not yet been empirically calibrated. Since the model uses no physical fundamental entities, calibration requires mapping pre-spatial geometric structures to physical measurement quantities. This mapping is currently conceptual and lacks quantitative frameworks. Specifically, the model lacks:

- quantitative translations for mass-like behavior,
- quantitative translations for charge-like behavior,
- quantitative translations for gravity-like behavior.

Summary

The limitations of the MCM arise from its monistic ontology, its geometric formulation, incomplete constitutive relations, numerical complexity, limited analytical tools, interpretation ambiguity, and lack of empirical calibration. These limitations mark areas for future refinement and development of the model.

18 Future Research Directions

The Monistic Continuum Model (MCM) provides a unified geometric ontology in which all observable phenomena arise from stress, rotation, curvature, and internal updating of a single continuous medium. Despite its conceptual coherence, key areas remain open for further development. Future research directions focus on refining the mathematical structure, improving numerical methods, expanding interpretative frameworks, and analyzing emergent behavior of bound and unbound geometry.

Development of the Constitutive Operator

The constitutive operator

$$D\sigma_{AB} = F(Vv, \sigma)_{AB}$$

is central to the dynamics of the medium. Future work includes:

- deriving closed-form expressions for F ,
- identifying geometric invariants that constrain constitutive behavior,
- analyzing stability conditions of bound geometry,
- investigating coupling between rotation and internal curvature.

A fully specified operator would significantly enhance the model's predictive capabilities.

Analytical Tools for Geometric Dynamics

The MCM requires new analytical tools tailored to pre-spatial updating dynamics. Research directions include:

- stability analysis of bound geometry (vortices),

- characterization of rotational–geometric modes,
- equations for internal curvature evolution,
- geometric invariants at the domain level of unbound geometry.

These tools aim to capture emergent geometric behavior more precisely.

Advances in Numerical Methods

Simulation of the MCM is computationally demanding. Future work includes:

- adaptive geometric discretization methods,
- high-resolution algorithms for curvature localization in bound geometry,
- stable propagation schemes for rotational–geometric modes,
- coupling frameworks for multiple geometric domains.

Progress in these areas will enable simulation of larger and more complex structures.

Stability Families of Bound Geometry

Bound geometry satisfies approximate update invariance:

$$DC(\alpha, \Omega) \approx 0.$$

Future research aims to:

- classify discrete stability families of vortices,
- analyze interactions of bound structures,
- determine stability thresholds under varying τ ,
- investigate bound clustering and domain anchoring.

A comprehensive stability theory would unify many emergent phenomena.

Structure of Rotational–Geometric Modes

Rotational–geometric modes, encoded in

$$L_{AB} = \frac{1}{2}(\nabla_A v_B - \nabla_B v_A),$$

play a central role in emergent behavior. Future work includes:

- identifying discrete rotational stability families,
- analyzing curvature-modulated propagation,
- investigating bound-mode scattering,
- developing geometric wave equations for unbound geometry.

This research will further clarify the geometric origin of electromagnetism-like behavior.

Curvature and Domain Evolution

Internal curvature arises from the stress-induced metric:

$$\mathcal{R}_{ABCD} = R(\gamma_{AB}).$$

Future research directions include:

- analyzing large-scale curvature evolution of unbound geometry,
- investigating domain expansion and contraction,
- exploring curvature-driven clustering of bound geometry,
- identifying geometric invariants at the domain level.

These efforts deepen understanding of cosmology-like behavior.

Characterization of the Update Rate

The update rate τ influences stability, propagation, and domain formation. Future work includes:

- deriving geometric constraints for τ ,
- analyzing reactivity-driven phase transitions,

- exploring multiscale updating behavior,
- linking τ to stability families of bound geometry.

A refined theory of reactivity will clarify many emergent phenomena.

Interpretative Frameworks

Translation schemes enable interpretation of geometric structures in physical terms. Future work includes:

- refining existing translation schemes,
- developing correspondence rules between geometry and physics,
- exploring alternative interpretative frameworks,
- analyzing interpretative ambiguity.

These frameworks improve comparability with established physics.

Summary

Future research on the MCM focuses on development of the constitutive operator, creation of analytical tools, advances in numerical methods, classification of bound stability families, analysis of rotational–geometric modes, curvature evolution, characterization of the update rate, and refinement of interpretative frameworks. These directions will refine the model and expand its explanatory power.

19 Glossary of Terms

This glossary summarizes the central geometric and structural concepts of the Monistic Continuum Model (MCM). All terms describe internal properties of a single continuous medium and do not rely on physical space, physical time, fields, particles, or forces. The ontology is monistic: every quantity describes the geometric behavior of the medium.

State Map

$$C(\alpha, \Omega) = (\rho, v_A, \sigma_{AB}, \tau).$$

The complete pre-spatial state of the medium at medium index α and update index Ω . Contains density state, geometric motion, stress, and update rate.

Density State

$$\rho(\alpha, \Omega)$$

Geometric quantity describing local compression or expansion of the medium. Not a physical mass density.

Geometric Motion

$$v_A(\alpha, \Omega)$$

Motion induced by stress gradients. Not a physical velocity. Determines the reorganization of geometric structures.

Stress Tensor

$$\sigma_{AB}(\alpha, \Omega)$$

Describes tension and compression in the medium. Stress generates internal geometry and determines curvature, rotational–geometric behavior, and the stability of bound geometry.

Update Rate

$$\tau(\alpha, \Omega)$$

Local reactivity of the medium. Controls how quickly geometric structures reorganize. Not physical time.

Update Derivative

$$DC(\alpha, \Omega)$$

Internal development of the medium. Measures the change of the state map under updating. Stability of bound geometry requires:

$$DC(\alpha, \Omega) \approx 0.$$

Stress-Induced Metric

$$\gamma_{AB} = \delta_{AB} + \alpha_1 \sigma_{AB} + \alpha_2 I_1 \delta_{AB}.$$

Internal metric induced by stress. Not a spatial metric. Determines geometric relations within the medium.

First Stress Invariant

$$I_1 = \text{tr}(\sigma).$$

Scalar measure of total stress. Contributes to the induced internal metric.

Curvature Tensor

$$\mathcal{R}_{ABCD} = R(\gamma_{AB}).$$

Internal curvature derived from the stress-induced metric. Describes geometric amplification or bending.

Rotation Tensor

$$L_{AB} = \frac{1}{2}(\nabla_A v_B - \nabla_B v_A).$$

Describes rotational–shear behavior of the medium. Foundation of rotational–geometric propagation.

Transport Operator

$$Vv(\sigma_{AB}) = v_C \nabla_C \sigma_{AB}.$$

Geometric transport of stress through motion. Not physical advection.

Constitutive Operator

$$D\sigma_{AB} = F(Vv, \sigma)_{AB}.$$

Determines the reorganization of stress under updating. Contains rotational, shear, curvature, and relaxation terms.

Bound Geometry (Matter)

Stable, localized geometric configuration satisfying:

$$DC(\alpha, \Omega) \approx 0.$$

Bound geometry produces persistent stress, rotation, and curvature. The only fully stable realization is the vortex.

Vortex

The only bound geometric structure satisfying all stability conditions of the MCM. Possesses:

- closed rotational lines,
- localized stress concentration,
- bound curvature sink,
- defined linking number,
- discrete stability modes.

Foundation of emergent matter.

Rotational–Geometric Mode

A propagating pre-spatial pattern encoded in L_{AB} . Corresponds to electromagnetism-like behavior under spatial interpretation.

Curvature Domain

Region of unbound geometry with coherent internal curvature structure. Arises from global stress organization. Foundation of cosmology-like behavior.

Geometric Stability

Persistence of geometric structure under updating. Requires:

$$DC(\alpha, \Omega) \approx 0.$$

Stability determines the existence of bound geometry, the propagation of rotational–geometric modes, and the coherence of large-scale domains.

20 Conclusion

The Monistic Continuum Model (MCM) provides a unified geometric ontology in which all observable phenomena arise from stress, rotation, curvature, and internal updating of a single continuous medium. The model replaces physical fundamental entities—space, time, fields, particles, and forces—with internal geometric structures that reorganize according to the update rate $\tau(\alpha, \Omega)$. This yields a monistic framework in which bound geometry (matter), rotational–geometric propagation, curvature, and large-scale domain behavior appear as emergent effects of a single medium.

Unified Geometric Ontology

The MCM demonstrates that the pre-spatial state map

$$C(\alpha, \Omega) = (\rho, v_A, \sigma_{AB}, \tau)$$

is sufficient to generate all emergent phenomena that established physics assigns to different entities. Stress produces internal geometry, rotation forms pre-spatial wave-like patterns, curvature shapes domain-level structures, and the update rate governs stability and propagation.

Emergence of Bound and Unbound Geometry

Bound geometry arises as a stable configuration satisfying approximate update invariance:

$$DC(\alpha, \Omega) \approx 0.$$

The only fully stable realization is the vortex, whose rotational–geometric and curvature-bound structure produces mass-like, charge-like, and spin-like behavior. Rotational–geometric modes correspond to electromagnetism-like behavior, bound curvature sinks to gravity-like behavior, and discrete stability modes to quantum-like behavior. All of these phenomena are purely geometric effects of the medium.

Dynamics Without Physical Time

The update index Ω is not physical time. Dynamics arise solely from internal geometric reorganization governed by the update rate τ . This enables a fully geometric description of development without temporal assumptions.

Large-Scale Geometric Behavior

Cosmology-like behavior emerges from reorganization of unbound geometry at the domain level: stress gradients, curvature evolution, rotational–geometric coherence, and clustering of bound geometry. These large-scale patterns arise directly from the internal geometry of the medium and require no spacetime expansion or external forces.

Numerical and Analytical Foundations

The MCM provides the foundation for geometric numerical simulation and analytical development. Geometric finite-difference methods, internal curvature computation, rotational-mode propagation, and stability analysis of bound geometry enable investigation of emergent structures and dynamic behavior.

Conceptual and Applied Implications

Translation schemes allow bound and unbound geometry to be interpreted in physical terms. In this way, the MCM provides conceptual insights into materials science, wave propagation, structural stability, and cosmological modeling without departing from its monistic ontology.

Outlook

Future research will refine the constitutive operator, expand analytical tools, improve numerical methods, classify stability families of bound geometry, analyze rotational–geometric modes, and investigate domain behavior. The MCM thus opens a path toward a unified geometric description of all observable phenomena.

Summary

The Monistic Continuum Model replaces physical ontology with geometric ontology. Stress, rotation, curvature, and update dynamics of a single medium generate all emergent phenomena. The MCM provides a consistent framework that explains bound geometry, rotational–geometric propagation, curvature, and cosmology-like behavior as geometric effects, offering a promising foundation for future theoretical and applied research.

[Spinoza1677·Ethics] [Whitehead1929·ProcessReality] [Bohm1980·Wholeness] [Ladyman2007·Everyth
[French2014·StructuralRealism] [Schaffer2009·Monism] [Kim1999·Emergence] [OConnor2007·Emerge

21 Citation Recommendation

This document represents the official reference version of *Monistic Continuum Model (MCM): Unified Geometric Ontology*, published as Version 07 of the *Monistic Continuum Model (MCM)* series and archived with a permanent DOI on Zenodo.

The following citation formats may be used:

APA Style

Moosbrugger, W. (2026). *Monistic Continuum Model (MCM): Unified Geometric Ontology*.

Part of the *Monistic Continuum Model (MCM)* series.

Zenodo. <https://doi.org/10.5281/zenodo.22128163>

BibT_EX

```
@misc{moosbrugger2026_mcm_master,  
  author    = {Moosbrugger, Walter},  
  title     = {Monistic Continuum Model (MCM): Unified Geometric Ontology},  
  year      = {2026},  
  publisher = {Zenodo},  
  doi       = {10.5281/zenodo.22128163},  
  url       = {https://doi.org/10.5281/zenodo.22128163},  
  note      = {Version 07, MCM | Master Document}  
}
```