

Command	Explanation
<code>pnorm(<math>x</math>)</code>	$P(Z \leq x)$ for $Z \sim \mathcal{N}(0, 1)$
<code>pt(<math>x, n - 1</math>)</code>	$P(T_{n-1} \leq x)$ for $T_{n-1} \sim T(n - 1)$
<code>qnorm(<math>p</math>)</code>	gives $x$ such that $P(Z \leq x) = p$ for $Z \sim \mathcal{N}(0, 1)$
<code>qt(<math>p, n - 1</math>)</code>	gives $x$ such that $P(T_{n-1} \leq x) = p$ for $T_{n-1} \sim T(n - 1)$
<code>pnorm(<math>x</math>, lower.tail = FALSE)</code>	$P(Z \geq x)$ for $Z \sim \mathcal{N}(0, 1)$
<code>pt(<math>x, n - 1</math>, lower.tail = FALSE)</code>	$P(T_{n-1} \geq x)$ for $T_{n-1} \sim T(n - 1)$
<code>qnorm(<math>p</math>, lower.tail = FALSE)</code>	gives $x$ such that $P(Z \geq x) = p$ for $Z \sim \mathcal{N}(0, 1)$
<code>qt(<math>p, n - 1</math>, lower.tail = FALSE)</code>	gives $x$ such that $P(T_{n-1} \geq x) = p$ for $T_{n-1} \sim T(n - 1)$

Notice that the `lower.tail = FALSE` option switches around the inequality in the probabilities. Hence it considers the upper tail instead of the lower tail, explaining its name.




FIGURE 1: The green area is given by `pt(-1, n-1)`, and the pink area is given by `pt(-1, n-1, lower.tail = FALSE)`.




FIGURE 2: The number  $p$  is the number such that the probability of being below it is 0.84, and likewise the probability of being above it is 0.16. Apropos R commands, `qt(0.84, n-1)` and `qt(0.16, n-1, lower.tail = FALSE)`.