Table 14.3.11.1 Prior Medications (Safety Analysis Set)

	Arm I (Nab-paclitaxel and gemcitabine) (N=xx)	Arm II (VCN-01 with Nab-paclitaxel and gemcitabine) (N=xx)
Any prior medications	xx (xx.x%)	xx (xx.x%)
X, xxxxxxxxxxxxx	xx (xx.x%)	xx (xx.x%)
X0N, xxxxxxxxxxxxx	xx (xx.x%)	xx (xx.x%)
X0NXX, xxxxxxxxxxx	xx (xx.x%)	xx (xx.x%)
xxxxxxxxxxxxx	xx (xx.x%)	xx (xx.x%)
X0NXX, xxxxxxxxxxx	xx (xx.x%)	xx (xx.x%)
xxxxxxxxxxxx	xx (xx.x%)	xx (xx.x%)
…		
…		

Prior medications/medical procedures are defined as any medication taken/any medical procedures performed within 28 days of the Screening visit.
Data are summarized by: Anatomical main group (ATC level 1); Therapeutic subgroup (ATC level 2); Therapeutic/pharmacological subgroup (ATC level 3); Chemical/therapeutic/pharmacological subgroup (ATC level 4); Preferred term.
Percentage is the number of patients in the category (n) divided by the number of patients within the column (N).
WHO-DDE version <XXXX>.

Source: Listing 16.2.12
Path\Filename

Programming Note: This layout also applies to Table 14.3.11.2 Concomitant Medications (Safety Analysis Set) [Footnote: Concomitant medications are defined as any medication taken at or after the time of the Screening visit.]