

Sum of Matrix After Queries

You are given an integer n and a **0-indexed 2D array** `queries` where `queries[i] = [typei, indexi, vali]`.

Initially, there is a **0-indexed** $n \times n$ matrix filled with `0`'s. For each query, you must apply one of the following changes:

- if `typei == 0`, set the values in the row with `indexi` to `vali`, overwriting any previous values.
- if `typei == 1`, set the values in the column with `indexi` to `vali`, overwriting any previous values.

Return the sum of integers in the matrix after all queries are applied.

Example 1:

Initial Matrix	Query 0	Query 1	Query 2	Query 3																																													
<table><tr><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td></tr></table>	0	0	0	0	0	0	0	0	0	<table><tr><td>1</td><td>1</td><td>1</td></tr><tr><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td></tr></table>	1	1	1	0	0	0	0	0	0	<table><tr><td>1</td><td>1</td><td>2</td></tr><tr><td>0</td><td>0</td><td>2</td></tr><tr><td>0</td><td>0</td><td>2</td></tr></table>	1	1	2	0	0	2	0	0	2	<table><tr><td>1</td><td>1</td><td>2</td></tr><tr><td>0</td><td>0</td><td>2</td></tr><tr><td>3</td><td>3</td><td>3</td></tr></table>	1	1	2	0	0	2	3	3	3	<table><tr><td>4</td><td>1</td><td>2</td></tr><tr><td>4</td><td>0</td><td>2</td></tr><tr><td>4</td><td>3</td><td>3</td></tr></table>	4	1	2	4	0	2	4	3	3
0	0	0																																															
0	0	0																																															
0	0	0																																															
1	1	1																																															
0	0	0																																															
0	0	0																																															
1	1	2																																															
0	0	2																																															
0	0	2																																															
1	1	2																																															
0	0	2																																															
3	3	3																																															
4	1	2																																															
4	0	2																																															
4	3	3																																															

Input: $n = 3$, `queries = [[0,0,1],[1,2,2],[0,2,3],[1,0,4]]`

Output: 23

Explanation: The image above describes the matrix after each query. The sum of the matrix after all queries are applied is 23.

Example 2:

Initial Matrix	Query 0	Query 1	Query 2	Query 3																																													
<table><tr><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td></tr></table>	0	0	0	0	0	0	0	0	0	<table><tr><td>4</td><td>4</td><td>4</td></tr><tr><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td></tr></table>	4	4	4	0	0	0	0	0	0	<table><tr><td>4</td><td>4</td><td>4</td></tr><tr><td>2</td><td>2</td><td>2</td></tr><tr><td>0</td><td>0</td><td>0</td></tr></table>	4	4	4	2	2	2	0	0	0	<table><tr><td>1</td><td>4</td><td>4</td></tr><tr><td>1</td><td>2</td><td>2</td></tr><tr><td>1</td><td>0</td><td>0</td></tr></table>	1	4	4	1	2	2	1	0	0	<table><tr><td>1</td><td>4</td><td>4</td></tr><tr><td>1</td><td>2</td><td>2</td></tr><tr><td>3</td><td>3</td><td>3</td></tr></table>	1	4	4	1	2	2	3	3	3
0	0	0																																															
0	0	0																																															
0	0	0																																															
4	4	4																																															
0	0	0																																															
0	0	0																																															
4	4	4																																															
2	2	2																																															
0	0	0																																															
1	4	4																																															
1	2	2																																															
1	0	0																																															
1	4	4																																															
1	2	2																																															
3	3	3																																															
Query 4																																																	
<table><tr><td>1</td><td>4</td><td>1</td></tr><tr><td>1</td><td>2</td><td>1</td></tr><tr><td>3</td><td>3</td><td>1</td></tr></table>					1	4	1	1	2	1	3	3	1																																				
1	4	1																																															
1	2	1																																															
3	3	1																																															

Input: $n = 3$, queries = $[[0,0,4],[0,1,2],[1,0,1],[0,2,3],[1,2,1]]$

Output: 17

Explanation: The image above describes the matrix after each query. The sum of the matrix after all queries are applied is 17.

Constraints:

- $1 \leq n \leq 10^4$
- $1 \leq \text{queries.length} \leq 5 * 10^4$
- $\text{queries}[i].\text{length} == 3$
- $0 \leq \text{type}_i \leq 1$
- $0 \leq \text{index}_i < n$
- $0 \leq \text{val}_i \leq 10^5$