

ASSIGNMENT 2: MAXIMUM LIKELIHOOD



Institute for Machine Learning

Contact

Heads:

Wei Lin

Florian Sestak

Institute for Machine Learning

Johannes Kepler University

Altenberger Str. 69

A-4040 Linz

E-Mail: {lin,sestak}@ml.jku.at

[Institute Homepage](#)

External Lecturers

Botond Fazekas

Guilherme Aresta

Taha Emre

Teresa Araújo

Antoine Rivail

Jürgen Plasser

E-Mail: `botond.fazekas@jku.at`

`guilherme.aresta@jku.at`

`taha.emre@jku.at`

`teresa.araujo@jku.at`

`antoine.rivail@meduniwien.ac.at`

`juergen_richard.plasser@jku.at`

Copyright statement:

This material, no matter whether in printed or electronic form, may be used for personal and non-commercial educational use only. Any reproduction of this material, no matter whether as a whole or in parts, no matter whether in printed or in electronic form, requires explicit prior acceptance of the authors.

Agenda

- Basics of Maximum Likelihood
- Properties of the Maximum Likelihood Estimator
- Further Literature:
 - Lecture notes (Hochreiter, 2014)
 - Mathematics for Machine Learning (Deisenroth et al., 2018)

Recap: Maximum Likelihood Estimator

- **Given:** n samples x_i from a distribution $p(x; w)$, where $w = \{w_1, w_2, \dots, w_N\}$ represents the parameter(s) of the distribution.
- **Task:** find the parameter(s) w that were most likely used to produce this data.
- **Idea:** how likely was a given w used to produce the dataset?

$$\mathcal{L}(\{x\}; w) = \prod_{i=1}^n p(x_i; w)$$

- **Solution:** find the w that maximizes $\mathcal{L}(\{x\}; w)$ or $\ln \mathcal{L}(\{x\}; w)$ (which is equivalent):

$$\hat{w} = \operatorname{argmax}_w \ln \mathcal{L}(\{x\}; w)$$

Recap: Maximum Likelihood Estimator

Properties of the Maximum Likelihood Estimator (MLE) \hat{w} :

- the MLE is invariant under parameter change
 - Model and data stay the same: Inference for two sets of parameters w and μ should be the same, if μ is a one-to-one transformation $\mu = f(w)$, i.e. $\hat{\mu} = f(\hat{w})$

Recap: Maximum Likelihood Estimator

Properties of the Maximum Likelihood Estimator (MLE) \hat{w} :

- the MLE is invariant under parameter change
- the MLE is asymptotically unbiased
 - Bias goes to 0 as sample size goes to infinity, i.e.
$$\lim_{n \rightarrow \infty} E(\hat{w}) = w$$

Recap: Maximum Likelihood Estimator

Properties of the Maximum Likelihood Estimator (MLE) \hat{w} :

- the MLE is invariant under parameter change
- the MLE is asymptotically unbiased
- the MLE is asymptotically efficient, i.e. asymptotically optimal

□ Reaches CRLB as sample size goes to infinity, i.e.

$$\lim_{n \rightarrow \infty} \text{Covar}(\hat{w}) = \mathbf{I}_F^{-1}(w)$$

Recap: Maximum Likelihood Estimator

Properties of the Maximum Likelihood Estimator (MLE) \hat{w} :

- the MLE is invariant under parameter change
- the MLE is asymptotically unbiased
- the MLE is asymptotically efficient, i.e. asymptotically optimal
- the MLE is consistent for zero Cramer-Rao lower bound
 - If variance can go to 0 (i.e. zero CRLB) and MLE is asymptotically unbiased and asymptotically efficient, this means the estimated parameter will be the true parameter as the sample size goes to infinity, i.e. $\lim_{n \rightarrow \infty} \hat{w} = w$