A Machine Learning Approach to Affine Conditional Moment Restriction Models

Wonjun Choi*

November 2, 2026

[Most recent version]

Abstract

This paper develops a general framework for estimation and inference in a class of affine conditional moment restriction models. Empirical applications of these models often involve a small number of endogenous regressors but a large set of instruments. Exploiting the affine structure, we leverage machine learning methods to develop an estimation procedure that is computationally fast and scales well to high dimensional instrumental variables. Using an orthogonalized minimum distance criterion, we characterize the requirement on the machine learners through their prediction errors, without restricting the analysis to a particular class of learners. Under this requirement, we develop a Wald statistic for hypotheses concerning smooth functionals. We also provide lower-level verification of our requirement for selected machine learning methods, as well as a Gaussian coupling of the estimated sieve coefficients useful for various inference problems. We examine the finite sample performance of our method through simulation studies and illustrate its use in an empirical application to migration in Tanzania.

*Department of Economics, Emory University. The author acknowledges the use of OpenAI's ChatGPT for language editing and editorial formatting.

1 Introduction

Conditional moment restriction (CMR) models provide a flexible framework for structural and reduced-form analysis (Compiani, 2022; Denteh and Liebert, 2026) and have been extensively studied with sieve and kernel methods (Newey and Powell, 2003; Ai and Chen, 2003; Hall and Horowitz, 2005; Carrasco, Florens, and Renault, 2007; Darolles et al., 2011, to name a few). An important subclass of CMR models arises when the residual function is affine in the parameter of interest. Despite its simple structure, affine CMR models encompass a broad range of empirically relevant models, including nonparametric instrumental variable (NPIV) models, partially linear models, and many others (Blundell, Chen, and Kristensen, 2007; Florens and Sokullu, 2017; Escanciano et al., 2021; Bennett et al., 2023). Meanwhile, modern empirical research utilizes increasingly rich datasets, and in particular, many instruments are often available for identifying the structural function in many fields of economics. For example, Bastos, Silva, and Verhoogen (2018) estimate the effect of export destination income on firms' input prices using 100 instruments based on exchange rates and firms' initial export destinations, while Deryugina et al. (2019) estimate the effects of fine particulate pollution on health outcomes using 300 instruments based on wind directions and geographic groups.

While nonparametric methods provide a flexible tool for capturing those rich information contained in modern data, a major obstacle to the empirical use of nonparametric methods is their computational challenge. It is well known that the computational burden of classical methods such as kernel and sieve estimators increases exponentially with the dimension of the variables used. Perhaps for this reason, empirical researchers often rely on parametric specifications (Bastos, Silva, and Verhoogen, 2018; Deryugina et al., 2019; Alfaro, Bloom, and Lin, 2024) or low-dimensional summaries of a larger set of instruments (Autor, Dorn, and Hanson, 2013; Burchardi, Chaney, and Hassan, 2019) even when hundreds of instrumental variables are available. This leaves a gap between the flexibility offered by nonparametric methods and the rich information available in empirical applications.

This paper addresses this gap by utilizing machine learning methods that can

handle high-dimensional variables effectively. Empirical applications of affine CMR models often involve a small number of endogenous variables with a large set of instrumental variables. Hence, we build on the sieve minimum-distance framework of Chen and Pouzo (2015), where the estimation step involves two stages: the first stage conditional moment estimation and the second-stage sieve coefficient estimation. We begin by approximating the structural parameter $\theta(x)$ with a linear sieve. Under the affine structure, the conditional moment reduces to

$$m(Z, \theta_n(X)) = \mathbb{E}[c(O)|Z] + \mathbb{E}[q_n(O)|Z]^\top \beta_n, \tag{1}$$

where $c(O)$ and $q_n(O)$ are known transformations of the observed variables $O = (Y, X, Z)$, X is a vector of inputs of the structural function $\theta(\cdot)$ which can contain endogenous variables, Z is a vector of instruments, Y is a vector of other variables, and β_n is the vector of sieve coefficients. Thus, the first stage conditional moment estimation becomes a nonparametric conditional mean prediction problem with many instruments, where modern machine learning methods are well suited.

Our core contribution is to provide a simple but general framework for estimation and inference in affine CMR models. We characterize the first stage requirements through prediction errors, so that the subsequent theory depends on the first stage learner only through its predictive performance. Under these conditions, we derive an asymptotic linear representation of the estimated sieve coefficients. Building on this representation, we provide Wald inference for smooth functionals and a Gaussian coupling of the sieve coefficients that can work as a building block for various inference problems. A technical challenge is that the dimension of the first stage prediction target grows with the sieve dimension. We therefore provide lower-level verification of the prediction-error requirements for several machine learning methods using error bounds commonly available in the machine learning literature.

Our framework combines practical advantages of minimum distance and GMM estimation. Minimum distance estimation works directly with conditional moment restrictions, instead of a set of unconditional moments implied by the conditional moment. Switching the conditional moment to unconditional ones involves the choice

of instrument functions and the number of unconditional moments, which can affect GMM performance (Domínguez and Lobato, 2004; Donald, Imbens, and Newey, 2009; Bennett and Kallus, 2023). On the other hand, GMM provides a natural standard error estimator once appropriate moments are specified, whereas the conventional minimum distance method may require first stage learner-specific variance analysis (Chen and Pouzo, 2015). By orthogonalizing the plug-in minimum distance criterion function, we obtain a first-order representation and standard error estimator whose forms are agnostic to the choice of first stage learner. This allows empirical researchers to flexibly choose machine learning methods best-suited for their applications at hand.

Our framework also provides a computationally attractive estimator by avoiding iterative training of the nested first stage problem. In CMR models, the conditional moment is a function of the structural parameter, so evaluating the second-stage criterion function at a new parameter point during optimization may require iterative training of the first stage conditional moment. This can be avoided by utilizing a fixed object that can be reused throughout the second-stage optimization, such as sieve space (Chen, Chen, and Tamer, 2023; Chen, Liao, and Wang, 2025), conditional mean operator (Darolles et al., 2011), conditional density/distribution (Hartford et al., 2017; Shao et al., 2025), RKHS embeddings (Singh, Sahani, and Gretton, 2019). In our case, we exploit the conditional mean of the known transformation of the observations. Several papers iteratively solve the nested problem, despite the computational burden, for adversarial or minimax formulations (Lewis and Syrgkanis, 2018; Bennett, Kallus, and Schnabel, 2019; Dikkala et al., 2020). This computational advantage can be particularly valuable when employing machine learning methods that are often costly to train.

In the simulation study, we examine the finite sample performance of our Wald tests for point evaluations and average derivatives. We find that our method works well with different first stage learners and across our simulation designs. The tests maintain size close to the nominal level under the null. The power decreases as the number of instruments increases, particularly in smaller samples, but improves with larger sample sizes as expected. We also find that the performance of our method is

particularly sensitive to noise the endogenous regressor that is not explained by the instruments, especially when the number of instruments is large.

In the empirical application, we revisit the study of migration and economic mobility in Tanzania by Beegle, De Weerdt, and Dercon (2011). We apply nonparametric methods in both stages of one of their 2SLS specifications to estimate the relationship between migration distance and consumption growth. The application involves six excluded instruments, making conventional tensor-product sieve estimation computationally expensive. We find that the positive relationship the authors found is actually close to zero over much of the migration-distance distribution but increases sharply at longer distances.

This paper contributes to a growing literature incorporating machine learning methods into CMR models. Existing approaches built around particular learners include neural networks (Hartford et al., 2017; Bennett, Kallus, and Schnabel, 2019; Shao et al., 2025), forests (Athey, Tibshirani, and Wager, 2019; Oprescu, Syrgkanis, and Wu, 2019), and reproducing kernel Hilbert space (Singh, Sahani, and Gretton, 2019; Muandet et al., 2020), among many others. Much of this literature emphasizes flexible second-stage estimation of the structural function. In contrast, our focus is on accommodating flexible machine learning methods in the first stage. Related literature studies machine learning for instrument selection (Belloni et al., 2012) and the construction of instruments through flexible first stage prediction (Chen, Chen, and Lewis, 2020). Our work complements this line of research by exploiting the affine structure to formulate first stage estimation as conditional mean prediction.

Our work also contributes to the literature on Neyman-orthogonalization and debiased learning (Chernozhukov et al., 2018, 2022; Foster and Syrgkanis, 2023). Drawing on Foster and Syrgkanis (2023) who studied the orthogonalization of general loss functions, we orthogonalize the sieve minimum distance criterion function. This orthogonalization can be equivalently performed by setting the functional of interest as the criterion function in the framework of Chernozhukov et al. (2022). On top of the usual rate gain from the Neyman orthogonality, this construction allows us to derive an asymptotic linear representation whose form is agnostic to the choice of first stage learner in the minimum distance framework.

A closely related paper is Bruns-Smith (2025), which also emphasizes flexible first stage estimation in NPIV models. Bruns-Smith constructs a data-adaptive RKHS from a reduced-form prediction of the outcome and uses this learned representation for second-stage estimation. Our approach differs in several aspects. First, we consider the broader class of affine CMR models, where a natural reduced-form problem could be unclear. Second, we directly estimate the conditional means with any learner, while it is important in the framework of Bruns-Smith (2025) that the chosen machine learning method can produce a sufficiently rich RKHS. Finally, Bruns-Smith (2025) allows machine learning methods in both stages, whereas we retain a linear sieve in the second stage. This compromises the computational scalability of our second stage, but allows us to access the extensive literature on challenging inference problems built on sieve methods. Bruns-Smith (2025) instead provides inference for regular functionals through a separate debiasing step.

The remainder of the paper is organized as follows. Section 2 introduces affine CMR models with examples. Section 3 develops the proposed estimator and Wald inference for smooth functionals. Section 4 examines first stage machine learning methods and sufficient rate requirements for inference. Section 5 presents Monte Carlo simulations, and Section 6 provides an empirical application. Appendix A establishes a Gaussian coupling of the sieve coefficients. Appendices B and C contain the proofs and lower-level verification of the first stage conditions, respectively.

Notation. For a measurable function f, let $\|f\|_{P,2} = (\mathbb{E}_P[f^2])^{1/2}$ and $\|f\|_{P,\infty} = \inf_t\{t \geq 0 : P(|f(X)| > t) = 0\}$. For vectors, $\|\cdot\|_2$ denotes the Euclidean norm, and for matrices, $\|\cdot\|_{\text{op}}$ denotes the operator norm. We write $a_n \lesssim b_n$ if $a_n \leq Cb_n$ for some constant $C > 0$, and $a_n \asymp b_n$ if both $a_n \lesssim b_n$ and $b_n \lesssim a_n$. When uniformity over $P \in \mathcal{P}$ is asserted, the constants are also independent of P.

2 Affine CMR models

In this section, we formally introduce the affine CMR model and discuss its reduction in our problem with examples. The residual function in (**??**) is affine in the structural

parameter θ if

$$\rho(O_i, \theta) \quad = \quad c(O_i) + \mathcal{L}[\theta; O_i]$$

for some known function $c(\cdot)$ and a known linear functional $\mathcal{L}[\cdot; o] : \mathcal{H} \to \mathbf{R}$. We assume the residual function $\rho(\cdot, \theta)$ is P-measurable for each $\theta \in \Theta \subset \mathcal{H}$ and $P \in \mathbf{P}$ where $\mathcal{H}$ is an Hilbert space equipped with a norm $\|\cdot\|_{\mathcal{H}}$ and Θ is a set of admissible parameter space.

We parametrize Θ with a linear sieve. That is, for some choice of sieve basis functions $\{p_{nl}(x)\}_{l=1,...,k_n} \subset \mathcal{H}$, we define a linear span $\mathcal{V}_n = \text{span}\{p_{n1}(x), ..., p_{nk_n}(x)\}$ and define the sieve space $\Theta_n \subset \Theta \cap \mathcal{V}_n$ as

$$\Theta_n \quad := \quad \{p_n(x)^\top \beta : \beta \in \mathcal{B}_n \subset \mathbf{R}^{k_n}\},$$

which grows dense in Θ. Here, $\mathcal{B}_n$ is a set of feasible coefficients, which can reflect, for example, shape constraints or regularizations. For convenience, after replacing all nonparametric parts with its sieve counterparts, we rewrite the residual function as a linear function of c_n and $q_n := \mathcal{L}[p_n(\cdot); O_i]$ thanks to the linearity of $\mathcal{L}$ as follow:

$$\rho(O_i, \theta_n) = c(O_i) + q_n(O_i)^\top \beta_n. \tag{2}$$

Taking the conditional expectation with respect to instruments Z yields the moment equation in Equation (1).

Remark 1. It is straightforward to extend the above formulation to a fixed-dimensional Euclidean parameter $b \in \mathbf{R}^{d_b}$ and a fixed number of nonparametric components $h_j \in \mathcal{H}_j$, $j = 1, ..., J$. Let $\theta = (b, h_1, \ldots, h_J) \in \mathbb{R}^{d_b} \times \mathcal{H}_1 \times \cdots \times \mathcal{H}_J$. The residual function takes the form

$$\rho(O, \theta) = c(O) + a(O)^\top b + \sum_{j=1}^{J} L_j[h_j; O]$$

where $L_j[\cdot; O]$, $j = 1, ..., J$ are known linear functionals. Approximating each h_j by

$p_{j,n}^\top \gamma_{j,n}$ and stacking the sieve coefficients and the corresponding basis functions again yields the expression in Equation (2) with $k_n = d_b + \sum_{j=1}^{J} k_{j,n}$ unknown Euclidean parameters. See, for example, Chen and Pouzo (2015) for the precise product-space construction of semi/nonparametric conditional moment models.

The following are examples of affine CMR models and their reduction to the expression in Equation (1).

Example 2.1 (NPIV)**.** The recent work of Compiani (2022) and Denteh and Liebert (2026) respectively estimate the demand of strawberry markets and the conditional local average treatment effect of a randomized policy by using NPIV and partially linear models. NPIV model specifies an unknown structural function h_0 through the moment equation $\rho(Y, X, h_0) = Y - h_0(X)$ where Y is an outcome, X a vector of regressors, and Z a vector of instruments. After replacing the structural function h_0 with a sieve approximate $h_n(x) = p_n(x)^\top \beta_n$, the residual function fits the formulation in (2) with $c_n(O) = Y$ and $q_n(O) = -p_n(X)$.

Example 2.2 (Linear CMR)**.** Bennett et al. (2023) consider the conditional moment restriction that is linear in the structural function of interest h

$$\mathbb{E}[g_1(O)h_0(X) \mid Z] = \mathbb{E}[g_2(O) \mid Z],$$

where g_1 and g_2 are known functions of the observed data and h_0 is unknown. Hence, we may set $c(O) = -g_2(O)$ and $q_n(O) = g_1(O)p_n(X)$ after substituting $h_n(x) = p_n(x)^\top \beta_n$ in the residual equation. Bennett et al. (2023) discuss proximal causal inference with unmeasured confounding and missing-not-at-random data with shadow variables as applications of this conditional moment restriction.

Example 2.3 (Transformation IV)**.** Florens and Sokullu (2017) consider nonparametric models on both sides of the equations. In their motivational model, a magazine serves readers and advertisers, and readership demand satisfies

$$H_0(Y) = \varphi_0(D) + X^\top b_0 + U, \qquad \mathbb{E}[U \mid X, W] = 0,$$

where Y is the magazine's readership market share, D is the number of adver-

tisements, which is endogenous, H_0 is the inverse readership demand, and φ_0 is the unknown function of advertising. The vector X contains exogenous magazine characteristics, W is a vector of instruments, and U captures unobserved magazine characteristics. Replacing sieve approximates $H_n(y) = p_H(y)^\top \beta_H$ and $\varphi_n(d) = p_\varphi(d)^\top \beta_\varphi$, and setting $Z = (X, W)$, Equation (1) is met with $c(O) = 0$, $q_n(O) = (-X^\top, p_H(Y)^\top, -p_\varphi(D)^\top)^\top$, and $\beta_n = (b^\top, \beta_H^\top, \beta_\varphi^\top)^\top$.

Example 2.4 (Euler equation)**.** Escanciano et al. (2021) consider the consumption based asset pricing Euler equation

$$\mathbb{E}[b_0 R_{t+1} g_0(Z_{t+1}) \mid Z_t] - g_0(Z_t) = 0, \qquad Z_t = (C_t, V_t),$$

where C_t is consumption, V_t contains other utility-relevant variables, R_{t+1} is a gross asset return, b_0 is the discount factor, and g_0 is marginal utility. For a fixed discount factor, the residual is linear in g. When both b and g are unknown, we may introduce $f = bg$ and write $\theta = (f, g)$. Then the residual becomes $\rho(O_t, \theta) = R_{t+1} f(Z_{t+1}) - g(Z_t)$, which is again linear in (f, g). The original model is retained by imposing $f = bg$ for some $b \in (0, 1)$ as a restriction. With the sieve approximates $f_n(z) = p_n(z)^\top \gamma$ and $g_n(z) = p_n(z)^\top \beta$, we can set $q_n(O_t) = (R_{t+1} p_n(Z_{t+1})^\top, -p_n(Z_t)^\top)^\top$ and $\beta_n = (\gamma^\top, \beta^\top)^\top$. We leave this example for future research, as it requires time-series generalization. See, for example, Chen, Liao, and Wang (2025) for time-series generalization of (nonlinear) sieve minimum distance estimators.

3 Estimation and Inference

This section develops estimation and inference for affine CMR models. We focus on the affine model with a single nonparametric function for ease of exposition. We first introduce an orthogonalized sieve minimum distance estimator and establish an asymptotic linear representation of its coefficients under high-level first stage conditions. Building on this representation, we develop Wald inference for smooth functionals of the structural parameter.

3.1 Orthogonalized Sieve Minimum Distance Estimator

The conventional (population) plug-in minimum distance criterion function is

$$Q_P^{plug-in}(\theta) \quad = \quad \mathbb{E}_P\Big[\frac{m(Z,\theta)^2}{\sigma^2(Z)}\Big],$$

where $\sigma^2(z)$ is a weighting function. Since the true structural function θ_P satisfies the moment condition $m(Z,\theta_P)=0$, it is a natural minimizer of $Q_P(\theta)$. By the law of iterated expectations, the criterion function can be equivalently expressed as

$$Q_P(\theta) \quad = \quad \mathbb{E}_P[\frac{2\rho(Z,\theta)m(Z,\theta)-m(Z,\theta)^2}{\sigma^2(Z)}].$$

The above criterion function is orthogonal in the sense of Foster and Syrgkanis (2023). That is, the gradient of the criterion function is Neyman-orthogonal to perturbations in the estimated nuisance, which is the conditional moment in our case. As a result, the first stage prediction error only appears in second-order and the estimators enjoy the same rate gain as in Chernozhukov et al. (2018). Through out the analysis, we treat the weighting function $\sigma^2(\cdot)$ as known[1] and leave the optimal weighting scheme to future work.

We denote the sieve minimizer of the orthogonalized criterion function over the sieve space as $\theta_{P,n} := \text{argmin}_{\Theta_n} Q_P$ and the associated sieve coefficients as $\beta_{P,n}$, so that $\theta_{P,n}(x) = p_n(x)^\top \beta_{P,n}$. With the estimated conditional moment function $\hat{m}_n(z,\theta)$ from the first stage, denote the sample counterpart of the criterion function as $\hat{Q}_n$. We define the ortogonalized sieve minimum distance estimator as

$$\hat{\theta}_n \quad := \quad \underset{\Theta_n}{\text{argmin}}\, \hat{Q}_n(\theta) = \underset{\Theta_n}{\text{argmin}} \left\{ \frac{1}{n}\sum_{i=1}^{n} \frac{2\hat{m}_n(Z_i,\theta)\rho(Z_i,\theta) - \hat{m}_n(Z_i,\theta)^2}{\sigma^2(Z_i)} \right\}.$$

Likewise, we denote the sieve coefficients as $\hat{\beta}_n$, so that $\hat{\theta}_n(x) = p_n(x)^\top \hat{\beta}_n$.

Remark 2 (Orthogonal projection first stage). Suppose the conditional moment is

[1] A typical choice of $\sigma^2(z)$ is $\text{Var}_P(\rho(O,\theta_P)|z)$. It can be estimated by $\text{Var}_n(\rho(O_i,\tilde{\theta}_n)|Z_i)$ with a preliminary estimator $\tilde{\theta}_n$, for instance, without weighting.

estimated by orthogonal projection methods such as sieve projections (Chen and Pouzo, 2015). For example, consider a sieve IV space spanned by a basis functions $\{b_1, \ldots, b_{J_n}\}$. Write $b^{J_n}(z) := (b_1(z), \ldots, b_{J_n}(z))^\top$, $P := \left(b^{J_n}(Z_1), \ldots, b^{J_n}(Z_n)\right)^\top$, and $\Pi := P(P^\top P)^- P^\top$ where $(P^\top P)^-$ denotes the Moore–Penrose inverse. For the residual vector $R = (\rho(O_1, \theta), \ldots, \rho(O_n, \theta))^\top$, ignoring weighting and sample-splitting for simplicity,

$$\hat{Q}_n(\theta) = \frac{1}{n}\{2R^T(\Pi R) - R^\top \Pi^\top \Pi R\} = \frac{1}{n} R^\top \Pi^\top \Pi R = \frac{1}{n}\sum_i \hat{m}_n(Z_i, \theta)^2,$$

due to the idempotency of Π. Thus, the orthogonalized criterion reduces to the conventional plug-in criterion function. However, such reduction may not extend to general first stage learners.

We now introduce the assumptions underlying our asymptotic analysis and discuss each component of the estimator in detail. We begin with regularity conditions on the data-generating process and the sieve approximation, followed by conditions on first stage estimation and the population criterion.

Assumption 3.1 (DGP)**.** (i) $\{O_i\}_{i=1}^n$ are i.i.d., $O_i \sim P \in \mathbf{P}$. (ii) There exists a constant $C_\rho, C_c < \infty$ such that $\sup_P \sup_\theta \|\mathbb{E}_P[\rho(O,\theta)^2|Z]\|_{P,\infty} \leq C_\rho$ and $\sup_{P\in\mathcal{P}} \|E_P[c(O)^2|Z]\|_{P,\infty} \leq C_c$. (iii) For every $P \in \mathbf{P}$, there exists $\theta_P \in \Theta$ such that $\mathbb{E}[\rho(O;\theta_P)|Z] = 0$ for P-almost every Z. (iv) There exists a constant $C_\mathcal{L} < \infty$, such that for every $P \in \mathbf{P}$, $\|\mathcal{L}[h;\cdot]\|_{P,2} := \{E_P[\mathcal{L}[h;\cdot]^2]\}^{1/2} \leq C_\mathcal{L}\|h\|_\mathcal{H}$ for all $h \in \mathcal{H}$ and $\|\mathcal{L}[h;\cdot]\|_{P,\infty} := ess\sup_o |\mathcal{L}[h;\cdot]| \leq C_\mathcal{L}\|h\|_\infty$ for all $h \in \mathcal{H}$. (v) For some constant $M < \infty$, $\sup_{\theta\in\Theta} \|\theta\|_\mathcal{H} \leq M$. (vi) There exist constants $0 < c \leq C \leq \infty$ such that for every $P \in \mathbf{P}$, $c\|h\|^2_{L_2(P_X)} \leq \|h\|^2_\mathcal{H} \leq C\|h\|^2_{L_2(P_X)}$ for all $h \in \mathcal{H}$.

Assumption 3.2 (Sieve)**.** (i) Let the Gram matrix be $G_n := \mathbb{E}_P[p_n(X)p_n(X)^\top]$. There exist constants $\underline{\lambda}$ and $\bar{\lambda}$ such that for every n and $P \in \mathbf{P}$, $0 < \underline{\lambda} \leq \lambda_{\min}(G_n) \leq \lambda_{\max}(G_n) < \overline{\lambda} < \infty$. (ii) There exists a sequence ξ_n such that for every n and $P \in \mathbf{P}$, $\operatorname{ess\,sup}_x \|p_n(x)\|_{l_2} \leq \xi_n$. (iii) For every $\theta \in \Theta$, $\inf_{\theta_n\in\Theta_n} \|\theta - \theta_n\|_\mathcal{H} = O(r_{s,n}) = o(1)$.

Assumption 3.1(iv) translates the primitive Assumption 3.2(ii) on the sieve basis functions to $q = \mathcal{L}[p;O]$, which is actually used in analysis. In the NPIV example,

Assumption 3.1(iv) trivially holds with constant 1 for the sup-norm and $c^{-1/2}$ for the L_2 norm due to Assumption 3.1(vi). Assumption 3.1(v) is mild and standard in the literature. In particular, we assume the constant M is large enough, so that the minimizers of the criterions are always interior solutions. A convenient sufficient condition for Assumption 3.1(vi) is that the density of X is bounded both from below and above.

For theoretical convenience, we orthonormalize the sieve basis with respect to $\|\cdot\|_{\mathcal{H}}$ without loss of generality. Together with Assumption 3.1(vi), we may obtain $C^{-1}I_{k_n} \preceq \tilde{G}_n \preceq c^{-1}I_{k_n}$ where $\tilde{G}_n$ is the $\|\cdot\|_{\mathcal{H}}$-orthonormalized Gram matrix. In Assumption 3.2(ii), ξ_n is $O(\sqrt{k_n})$ for many of the popular choices of sieves, including B-splines. Assumption 3.2(iii) is a standard sieve approximation assumption and the rate is readily available for popular choices of sieve (e.g., Chen, 2007).

By the affine representation in Equation (1), we can construct the conditional moment at any sieve coefficient β_n from the conditional means $\mu_c(z) = \mathbb{E}_P[c(O)|Z = z]$ and $\mu_q(z) = \mathbb{E}_P[q_n(O)|Z = z]$. We write $\mu_{n,P}(z) := (\mu_c(z), \mu_q(z)^\top)^\top \in \mathbb{R}^{k_n+1}$ and let $\hat{\mu}_n^{(-\ell)}$ collect the corresponding conditional mean estimates. We estimate these conditional means using flexible machine learning methods with L-fold cross-fitting. That is, we partition the sample into L disjoint folds and estimate the conditional means for each fold using observations from the remaining folds (Chernozhukov et al., 2018). For each observation i in fold ℓ, we evaluate the conditional moment as

$$\hat{m}_n(Z_i, \beta) = \hat{\mu}_c^{(-\ell)}(Z_i) + \hat{\mu}_q^{(-\ell)}(Z_i)^\top \beta,$$

with the conditional means $\hat{\mu}^{(-\ell)}(\cdot)$ are learned on the training folds $\mathcal{T}_\ell$ that excludes ℓ. The conditional mean estimators are always obtained by cross-fitting, so we often omit the superscript $(-\ell)$ when the context is clear.

Assumption 3.3 (first stage)**.** For some deterministic sequences $r_{\mu,n} \to 0$ and $r_{m,n} \to 0$, $\mathbb{E}_P[\|\hat{\mu}_n^{(-\ell)}(Z) - \mu_{n,P}(Z)\|_2^2 | \mathcal{T}_\ell] = O_P(r_{\mu,n}^2)$ and $\mathbb{E}_P[\{\hat{m}_n^{(-\ell)}(Z, \beta_{P,n}) - m_{n,P}(Z, \beta_{P,n})\}^2 | \mathcal{T}_\ell] = O_P(r_{m,n}^2)$.

Although we obtain the conditional moment $\hat{m}_n(z, \cdot)$ through the estimated con-

ditional means $\hat{\mu}_n(z)$, the regularization on $m(z,\cdot)$ can significantly reduce the rate $r_{m,n}$ relative to $r_{\mu,n}$. Hence, we state the rates separately for future generalization. We discuss the verification of the above assumption using finite sample error bounds in the literature in Section 4.

Remark 3. We mainly concern machine learning methods that can predict vector outcomes. However, k_n-many separate scalar learners can still be used. Suppose individual learners satisfy $\mathbb{E}_P[\int |\hat{\mu}_{n,j}^{(-\ell)}(z) - \mu_{n,P,j}(z)|^2 dP_Z|\mathcal{T}_\ell] \leq b_{n,j}^2$ for $j = 1, ...k_n$. Summing up the componentwise errors, the vector mean squared error is bounded by $\sum_{j=0}^{k_n} b_{n,j}^2$. If there is an a_n such that $b_{n,j}^2 \leq Ca_n^2$ for each $j = 0, ..., k_n$ accounting for $k_n \to \infty$, then $r_{\mu,n}$ can be taken as $\sqrt{k_n+1}a_n$.

In population, the orthogonalized criterion function coincides with the conventional minimum distance criterion by the law of iterated expectations. Hence, the orthogonalization does not alter identification. The identification in affine model is particularly transparent. For $h \in \mathcal{H}$, define the weak norm as

$$\|h\|_w^2 := \mathbb{E}_P\left[\frac{\{\mathbb{E}_P[\mathcal{L}[h;O] \mid Z]\}^2}{\sigma^2(Z)}\right].$$

Thanks to the affinity of the moment and the moment condition $m(\cdot,\theta_P) = 0$, it holds that $m(\cdot,\theta) = m(\cdot,\theta-\theta_P)$. Hence, the population criterion satisfies

$$Q_P(\theta) = \mathbb{E}_P\left[\frac{m(Z,\theta)^2}{\sigma^2(Z)}\right] = \|\theta-\theta_P\|_w^2.$$

Assumption 3.4(i) below imposes point identification of θ_P. In the affine model, this condition is equivalent to injectivity of the conditional mean operator $h \mapsto \mathbb{E}_P[\mathcal{L}[h;O] \mid Z]$ on the set of admissible deviations $\{\theta - \theta_P : \theta \in \Theta\}$. In the NPIV model, this reduces to the usual completeness condition (Newey and Powell, 2003).

Assumption 3.4 (Criterion function)**.** (i) For every $P \in \mathbf{P}$ and $\theta \in \Theta$, $Q_P(\theta) = 0$ if and only if $\|\theta - \theta_P\|_{\mathcal{H}} = 0$. (ii) There is a sequence $r_{s,n}$ and $\Pi_n\theta_P \in \Theta_n$ such that $\|\theta_P - \Pi_n\theta_P\|_{\mathcal{H}} = O(r_{s,n}) = o(1)$. (iii) For each $P \in \mathbf{P}$, there is a sequence $r_{w,P,n}$ such that $Q_P(\Pi_n\theta_P) = \|\Pi_n\theta_P - \theta_P\|_w^2 \leq O(r_{w,P,n}^2) = o(1)$. (iv) There exist constants

$0 < \underline{\sigma}^2 \leq \bar{\sigma}^2 < \infty$, such that for every $P \in \mathbf{P}$, $\underline{\sigma}^2 \leq \sigma^2(Z) \leq \bar{\sigma}^2$ P-a.s..

Assumption 3.4(ii) follows from the sieve approximation condition in Assumption 3.2(iii). Assumption 3.4(ii) also implies Assumption 3.4(iii) with $r_{w,P,n} = r_{s,n}$ by Jensen's inequality, Assumptions 3.1(iv), and 3.4(iv). We state the weak approximation rate separately to allow for sharper bounds. Notice Assumption 3.4(ii)(iii) are on the sieve approximation of the true structural function $\Pi_n\theta_0$, which in general differ from the sieve minimizer $\theta_{P,n}$.

Following Chen and Pouzo (2015), we define the sieve measure of ill-posedness. For a sieve direction $v = p_n^\top b \in \mathcal{V}_n$, let

$$H_n := \mathbb{E}_P\left[\frac{\mu_q(Z)\mu_q(Z)^\top}{\sigma^2(Z)}\right], \qquad \tau_n^2 := \sup_{b\neq 0} \frac{b^\top G_n b}{b^\top H_n b},$$

where G_n is defined in Assumption 3.2(i). On the sieve space, the weak norm satisfies $\|v\|_w^2 = b^\top H_n b$ and $\|v\|_{L_2(P_X)} \leq \tau_n \|v\|_w$. If the conditional moment operator is injective on $\mathcal{V}_n$, then H_n is positive definite for each n, although its smallest eigenvalue may converge to zero as $n \to \infty$. Because the conditional moment is affine in θ, its directional derivative and the matrix H_n do not depend on the evaluation point. See Newey and Powell (2003), Carrasco, Florens, and Renault (2007), Chen and Christensen (2018), and Chen, Christensen, and Kankanala (2025) for treatments of ill-posedness in NPIV and CMR models.

Now we derive the asymptotic linear representation of the sieve coefficients, which is useful for subsequent inference. Define the score vector

$$\psi_n(O;\theta) := \frac{\rho(O,\theta)\mu_q(Z) + m(Z,\theta)\{q_n(O) - \mu_q(Z)\}}{\sigma^2(Z)}. \tag{3}$$

For $\theta_n = p_n^\top \beta$, differentiating the (half of) the orthogonalized criterion function with

respect to β gives the gradient and Hessian matrix

$$\begin{aligned}\Psi_{n,P}(\theta_n) &:= \frac{1}{2}\nabla_\beta Q_P(\theta_n) = \mathbb{E}_P[\psi_n(O;\theta_n)],\\ \nabla_\beta \Psi_{n,P}(\theta_n) &= \mathbb{E}_P\left[\frac{\rho(O,\theta_n)\mu_q(Z) + m(Z,\theta_n)\{q_n(O) - \mu_q(Z)\}}{\sigma^2(Z)}\right].\end{aligned}$$

Notice the population Hessian matrix coincide with H_n defined above in the weak norm. Similarly, differentiating the cross-fitted sample criterion function yields

$$\hat{\Psi}_n(\beta) := \frac{1}{n}\sum_{i=1}^n \hat{\psi}_{ni}(\beta), \quad \text{where} \quad \hat{\psi}_{ni}(\beta) := \frac{\rho(O_i,\beta)\hat{\mu}_q(Z_i) + \hat{m}_n(Z_i,\beta)(q_n(O_i) - \hat{\mu}_q(Z_i))}{\sigma(Z_i)^2}$$

and the sample Hessian matrix

$$\hat{H}_n := \nabla_\beta \hat{\Psi}_n(\beta) = \frac{1}{n}\sum_{i=1}^n \frac{q_n(O_i)\hat{\mu}_q(Z_i)^\top + \hat{\mu}_q(Z_i)(q_n(O_i) - \hat{\mu}_q(Z_i))^\top}{\sigma(Z_i)^2}.$$

Remark 4 (Comparison to double/debiased machine learning). As expected from Foster and Syrgkanis (2023), the first order condition of the orthogonalized criterion function can work as the orthogonal moment condition for GMM (Chernozhukov et al., 2018, 2022). Hence, the orthogonalized minimum distance estimator here and the GMM estimator of double machine learning can coincide with a particular choice of weighting functions. However, they in general differ (Newey and McFadden, 1994).

Our first order expansion relies on the inverse of the estimated Hessian $\hat{H}_n$. Lemma B.2 shows that $\hat{H}_n$ is indeed positive definite (and hence invertible) with probability approaching one. A sufficient rate requirement for this is

$$r_{H,n} := \xi_n\sqrt{\frac{\log k_n}{n}} + \frac{r_{\mu,n}\xi_n}{\sqrt{n}} + r_{\mu,n}^2 = o(\tau_n^{-2}). \tag{4}$$

Provided the inverse of $\hat{H}_n$ and the interior solution $\hat{\beta}_n$, the following holds without

remainder, due to the affinity of the moment.

$$\hat{\beta}_n - \beta_{P,n} = \hat{H}_n^{-1}\{\hat{\Psi}_n(\hat{\beta}_n) - \hat{\Psi}_n(\beta_{P,n})\} = -\hat{H}_n^{-1}\hat{\Psi}_n(\beta_{P,n}). \tag{5}$$

The last equality uses $\hat{\Psi}_n(\hat{\beta}_n) = 0$ and $\hat{H}_n$ is positive definite. Hence, we derive a natural linear representation of the sieve coefficients with the population version of Equation (5) as a leading term.

Lemma 3.1. Suppose Assumptions 3.1, 3.2, 3.3, and 3.4(iii)(iv) hold. Moreover, suppose $r_{H,n} = o(\tau_n^{-2})$. Then,

$$\sqrt{n}(\hat{\beta}_n - \beta_{P,n}) = -H_{n,P}^{-1}\frac{1}{\sqrt{n}}\sum_i \psi_n(O_i, \theta_{P,n}) + R_n$$

where $\|R_n\|_2 = O_p(a_n + b_n)$ with

$$a_n = \tau_n^2\{\xi_n\sqrt{\frac{\log k_n}{n}} + \frac{r_{\mu,n}\xi_n}{\sqrt{n}} + r_{\mu,n}^2\}\{\tau_n\sqrt{k_n} + \tau_n^2\xi_n r_{w,n}\}, \tag{6}$$

$$b_n = \tau_n^2\{r_{\mu,n} + \xi_n r_{m,n} + \sqrt{n}r_{m,n}r_{\mu,n}\}. \tag{7}$$

The rate requirement $\tau_n^2 r_{H,n} = o(1)$ is a sufficient condition for the Hessian inversion. It is mild and dominated by the rate requirements implied by a_n and b_n. The rates in a_n and b_n are the convergence rates of the remainders from replacing the estimated term in Equation (5) to population terms. Hence, they can be captured by the prediction error rate $r_{\mu,n}$ in Assumption 3.3.

Remark 5 (Comparison to CP score). Evaluating ψ_n at the true structural parameter gives $\psi_n(O;\theta_P) = \rho(O,\theta_P)\mu_q(Z)/\sigma^2(Z)$, which corresponds to the leading stochastic term in Chen and Pouzo (2015, Equation (3.11)). The distinction lies in how this leading term is obtained from the sample criterion. Specifically, the gradient of the plug-in minimum distance criterion is $n^{-1}\sum_{i=1}^n \hat{m}_i(\theta)\hat{\mu}_{q,i}/\sigma^2(Z_i)$, so replacing $\hat{m}_i$ with ρ_i requires a separate justification. The orthogonalized criterion places ρ_i directly in the gradient, and its difference from the population score can be controlled through the prediction errors for any first stage learners.

Building on the linear representation in Lemma 3.1, we provide a Wald statistic for testing smooth functionals in the following subsection. In Appendix A, we provide a Gaussian coupling of the linear representation, which is useful for various inference problems.

3.2 Inference for smooth functionals

We provide a Wald statistic for smooth functionals ϕ that covers important inference problems in empirical research. Particularly, we modify the Wald statistic of Chen and Pouzo (2015) to our setting. Consider the null hypothesis

$$H_0 : \phi(\theta_0) = 0.$$

For smooth functionals $\phi(\cdot)$, their first order behavior on the sieve space admits a convenient expression in terms of coefficient loadings l_n (Newey, 1997), as shown in the following examples.

Example 2.1 (NPIV, continued)**.** Suppose an empirical researcher estimates a demand function using the NPIV model $Y = \theta_0(X) + U$, where Y is (log) quantity and X contains (log) prices. A researcher may be interested in demand at price x_0 and its confidence interval. Hence, define the testing functional $\phi : \mathcal{H} \to \mathbf{R}$

$$\phi(\theta) = \theta(x_0).$$

On the sieve space, the testing functional is written as $\phi(\theta_{P,n}) = \theta_{P,n}(x_0) = p_n(x_0)^\top \beta_{P,n}$. Hence, the loading l_n is taken as $p_n(x_0)$.

Example 2.1 (NPIV, continued)**.** Continuing from the previous example, a researcher may be interested in the (weighted) average price elasticity of its own price. Let x_1 denote the log of the own price. Then, for some known weights $w(x)$, define

$$\phi(\theta) = \int \frac{\partial \theta(x)}{\partial x_1} w(x)\, dx.$$

On the sieve space, $\phi(\theta_{P,n}) = \left\{\int (\partial p_n(x)/\partial x_1) w(x)\, dx\right\}^\top \beta_{P,n}$ Hence, the loading l_n can be taken as the expression in the curly bracket.

Example 2.1 (NPIV, continued)**.** Continuing from the previous example, a researcher may instead be interested in demand in levels at x_0 and define the testing functional as $\phi(\theta) = \exp\{\theta(x_0)\}$. Since $\phi(\cdot)$ is smooth, it can be approximated by the Tayler expansion near $\beta_{P,n}$ on the sieve space (e.g., Magnus and Neudecker, 2019, Chapter 6, Theorem 12):

$$\phi(\theta_n) = \exp\{p_n(x_0)^\top \beta_{P,n}\} + \nabla_\beta \exp\{p_n(x_0)^\top \beta_{P,n}\}(\beta_n - \beta_{P,n}) + O(\|\beta - \beta_{P,n}\|_2^2).$$

Since $\beta_{P,n}$ is unknown, one may estimate the loading as $l_n = \nabla_\beta \exp\{p_n(x_0)^\top \hat{\beta}_n\}$.

Provided the coefficient loading l_n and the linear expansion in Lemma 3.1, we can write the plug-in test statistic as

$$\sqrt{n}(\phi(\hat{\theta}_n) - \phi(\theta_P)) \;\approx\; -l_n^\top H_n^{-1} \frac{1}{\sqrt{n}} \sum_i \psi_n(O_i) + l_n^\top R_{n,P}. \tag{8}$$

This suggests the sieve Wald score and its estimator

$$\begin{aligned} S_{ni} &= l_n^\top H_n^{-1} \frac{\rho_i(\beta_{P,n})\mu_q(Z_i) + m(Z_i, \beta_{P,n})(q(X_i) - \mu_q(Z_i))}{\sigma^2(Z_i)}, \\ \hat{S}_{ni} &= l_n^\top \hat{H}_n^{-1} \frac{\rho_i(\hat{\beta}_n)\hat{\mu}_q(Z_i) + \hat{m}(Z_i, \hat{\beta}_n)(q(X_i) - \hat{\mu}_q(Z_i))}{\sigma^2(Z_i)}. \end{aligned}$$

We define the standard error of the Wald statistic and its estimator

$$V_{n,P} := \mathrm{Var}_P[S_{ni}], \qquad \hat{V}_{n,P} := \mathrm{Var}_n[\hat{S}_{ni}],$$

where $\mathrm{Var}_n(X) = \mathbb{E}_n[X^2] - (\mathbb{E}_n X)^2$. Equivalently, they can be written as $V_{n,P} = l_n^\top H_n^{-1} \Omega_n H_n^{-1} l_n$ where $\Omega_n := E_P[\psi_{n,P}(O)\psi_{n,P}(O)^\top]$, and $\hat{V}_n = l_n^\top \hat{H}_n^{-1} \hat{\Omega}_n \hat{H}_n^{-1} l_n$ where $\hat{\Omega}_n = \mathbb{E}_n[\hat{\psi}_i \hat{\psi}_i^\top]$. The consistency of $\hat{V}_n$ is provided in Lemma B.3. As the dimension of the sieve grows, $\|l_n\|_2$ may diverge depending on the testing functional. For details

on such *irregular* functionals, see Chen and Liao (2014) and Chen and Pouzo (2015).

Following Chen and Pouzo (2015), we impose the following assumptions.

Assumption 3.5 (Testing functional ϕ)**.** (i) There exists $\mathcal{N}_{n,P} \subset \Theta_n$ and a linear functional $\dot{\phi} : \mathcal{V}_n \to \mathbf{R}$ such that $P\{\hat{\theta}_n \in \mathcal{N}_{n,P}\} \to 1$ and $\sup_{\theta \in \mathcal{N}_{n,P}} \frac{\sqrt{n}}{\sqrt{V_n}} |\phi(\theta) - \phi(\theta_{P,n}) - \dot{\phi}[\theta - \theta_{P,n}]| = o(1)$. (ii) $\frac{\sqrt{n}}{\sqrt{V_n}} |\phi(\theta_{P,n}) - \phi(\theta_P)| = o(1)$.

Once $\dot{\phi}$ is obtained, the loading can be formally defined as $l_n := (\dot{\phi}[p_{n1}], \cdots, \dot{\phi}[p_{nk_n}])^\top$. If ϕ is linear, then the absolute value in Assumption 3.5(i) is exactly 0 and $\mathcal{N}_{n,P}$ can be taken as the entire function space. Otherwise, one may construct a local neighborhood near $\beta_{P,n}$ after establishing consistency and control the estimation and linearization errors across the local neighborhood. Assumption 3.5(ii) is a combined requirement on the sieve approximation bias and the testing functional. It sets the lower bound on the sieve complexity k_n.

Under current assumptions, $\beta_{P,n}$ is always an interior minimizer. Hence, it always holds that $\mathbb{E}_P[S_{ni}] = 0$ for any (population) loading l_n. If the parameter space is restricted, we would require the centering condition $\mathbb{E}_P[S_{ni}] = o(\sqrt{V_n/n})$ to maintain the current Wald inference Chen and Pouzo (2015).

Assumption 3.6 (Sieve score)**.** For every $\epsilon > 0$, $E_P[\frac{S_{ni}^2}{V_n} 1\{|S_{ni} > \epsilon \sqrt{n V_n}\}] \to 0$.

A convenient sufficient condition for Assumption 3.2 is the Lyapunov condition $E_P[|S_{ni}|^{2+\delta}/n^{\delta/2} V_n^{1+\delta/2}] \to 0$, which in turn can be satisfied by the $2+\delta$ moments of the residual, sieve basis, and conditional means.

The following theorem establishes the asymptotic distribution of the Wald statistic.

Theorem 3.1 (Wald)**.** Suppose Assumptions 3.1, 3.2, 3.3, 3.4(iii)(iv), 3.5, and 3.6 hold. Moreover, suppose the rate conditions $r_{H,n} = o(\tau_n^{-2})$, $\|l_n\|_2 (a_n + b_n)/\sqrt{V_n} = o(1)$ hold. Then,

$$\frac{\sqrt{n}}{\sqrt{\hat{V}_n}} \{\phi(\hat{\theta}_n) - \phi(\theta_P)\} \rightsquigarrow N(0,1).$$

The first rate requirement $\tau_n^2 r_{H,n} = o(1)$ for Hessian convergence is mild. The second rate requirement is to control the remainder term in Equation (8), and is stronger than necessary. A convenient sufficient condition for the second rate requirement is $a_n + b_n = o(1)$, although the rate requirement can be relaxed up to τ_n^2. We defer the discussion of the rate requirement to the following section.

4 first stage learners and the rate requirements

This section surveys several machine learning methods and discuss the verification of rate requirement in Assumption 3.3 and its implication to the Wald inferences.

4.1 First stage learners and their error bounds

Recall Assumption 3.3 requires bounds on the prediction errors of the estimated conditional means and the resulting conditional moment. However, such bounds are often not directly available in the theory of machine learning methods. Therefore, we verify Assumption 3.3 for selected machine learning methods by adapting their derivation of the error bounds. The full proofs are provided in Appendix C.

The first class of learners consists of local-smoothing learners such as kernel estimators. These methods exploit the smoothness of conditional means on shrinking neighborhoods of the conditioning point, while the neighborhoods are constructed differently across learners. Their convergence theory often provides pointwise prediction-error bounds. We impose additional regularity conditions when necessary and integrate these bounds over P_Z to verify Assumption 3.3.

Example 4.1 (Generalized random forest)**.** Athey, Tibshirani, and Wager (2019) propose generalized random forest, which uses the grown trees to obtain kernel-like weights α_i. For an evaluation point z and in each tree, observations whose Z_i falls in the same leaf as z recieves a unit weight. Aggregating those weights over trees yields the final weight $\alpha_i(z)$. Our first stage conditional means then can be estimated as

the weighted average

$$\hat{\mu}_n(z) = \sum_i \alpha_i(z)(c_n(O_i), q_n(O_i)^\top)^\top.$$

Their Lemma 7 provides pointwise bias and variance bounds of $\hat{\mu}(z)$, which does not depend on z. Under some regularity conditions, such as the bounded second moment and the Lipschitz continuity of the conditional means, we may bound the first stage error by aggregating the pointwise bound and obtain

$$\mathbb{E}_P[\|\hat{\mu}_n^{(-\ell)}(Z) - \mu_{n,P}(Z)\|_2^2 \mid \mathcal{T}_\ell] = O_P\Big(L_{\mu,n}^2 s_n^{-2\gamma} + (1+\xi_n^2)\frac{s_n}{n}\Big).$$

Hence, we may take $r_{\mu,n}$ as the square root of the sequence in $O_p(\cdot)$ term. Here, s_n is the subsampling size, which is a tuning parameter, $L_{\mu,n}$ is the Lipschitz constant of $\mu_{n,P}$, which may grow with n, ξ_n is the sieve envelope introduced in Assumption 3.2, and $\gamma > 0$ is a constant that depends on the dimension of instruments d_z and other tuning parameters. These constants can be taken independent of P under appropriate conditions.

Example 4.2 (Subsampled nearest neighbors)**.** Khosravi, Lewis, and Syrgkanis (2022) suggest subsampled nearest neighbors that is adaptive to the covariates' intrinsic dimension. Similarly to the previous example, the conditional mean estimator is

$$\hat{\mu}_n(z) = \sum_i \alpha_i(z)(c_n(O_i), q_n(O_i)^\top)^\top$$

where the wegiths $\alpha_i(z)$ is obtained from independently drawn B_n subsamples of size s_n. Within each subsample, weight $1/\kappa$ is assigned to the κ-nearest neighbors of z. Averaging these weights over B_n subsamples gives $\alpha_i(z)$.

Their Equation (8) in Theorem 1 provides a finite sample root mean squared error bound at each fixed z. Under the extra conditions in Lemma C.1, we make the

pointwise bound uniform[2] in z and bound the first stage error by

$$\mathbb{E}_P[\|\hat{\mu}_n^{(-\ell)}(Z) - \mu_{n,P}(Z)\|_2^2 \mid \mathcal{T}_\ell] = O_P\Big(L_{\mu,n}^2 s_n^{-2/d_z} + (1+\xi_n^2)\frac{s_n}{n}\Big),$$

where $L_{\mu,n}$ is the Lipschitz constant of $\mu_{n,P}$, which may grow with n, and ξ_n is the sieve envelope. Hence, we may take $r_{\mu,n}^2$ as the sequence in the O_p term.

The bounds in Examples 4.1 and 4.2 exploit only Lipschitzness of the conditional mean of the first stage outcomes. Faster prediction rates under stronger regularity conditions are available for other subsampling-based learners such as two-scale distributional nearest neighbors (Demirkaya et al., 2024). These results suggest first stage learners with potentially sharper error bounds.

The second class of learners exploits stronger smoothness assumptions on the conditional means. For example, Schmidt-Hieber (2020) derives prediction error bounds for sparse deep ReLU networks for a scalar outcome, with rates depending on the Hölder smoothness of the conditional mean and the number of instruments. Cattaneo, Klusowski, and Underwood (2026) obtain similar rates for debiased Mondrian forests under smoothness conditions on the conditional mean and the instrument density. Under these bounds, root mean squared prediction error for a scalar outcome can converge faster than $n^{-1/4}$ for any fixed d_Z, provided sufficient smoothness and the remaining regularity conditions. We leave the verification of Assumption 3.3 for these learners to future work.

The last class of learners is the modern conditional distribution estimators, such as conditional diffusion and Gaussian mixture models. The theory of these estimators typically provides convergence rates under probability metrics such as total variation distance, Kullback-Leibler divergence, Wasserstein distance, and more. See Gibbs and Su (2002) for the review of such metrics. Hence, we may translate these bounds to verify the L_2 prediction error bound in Assumption 3.3.

Example 4.3 (Conditional diffusion models)**.** Separate O into (V, Z), where $V \in$

[2]This should not be confused with the convergence in sup-norm or convergence as a function.

$\mathbb{R}^{d_V}$ and d_V, d_Z are fixed. Conditional diffusion models learn the conditional distribution of V given Z by adding Gaussian noise to V and estimating the conditional score function from the noisy version of V and the conditioning variable Z (Song et al., 2021; Ho and Salimans, 2022). Provided the estimated conditional distribution function, we can define

$$\widehat{\mu}_n^{(-\ell)}(z) = \int (c(v,z), q_n(v,z)^\top)^\top d\widehat{P}_n^{(-\ell)}(v \mid z),$$

and estimate the conditional means by Monte Carlo integration.

The theoretical result in Fu et al. (2024) provides a rate of convergence of the (integrated) total variation distance between the estimated and true conditional law. Hence, we may apply the Cauchy-Schwarz inequality and bound the prediction error by using the Hellinger distance. Then, Assumption 3.3 can be verified by

$$\mathbb{E}_P\big[\|\widehat{\mu}_n^{(-\ell)}(Z) - \mu_{n,P}(Z)\|_2^2 \mid \mathcal{T}_\ell\big] = \tilde{O}_P\big((1+\xi_n^2) n^{-2s/(2s+d_V+d_Z)}\big) \tag{9}$$

uniformly over P. Here $\tilde{O}_P(\cdot)$ refers to up to a logarithmic factor and s is a smoothness parameter of the conditional density.

Gaussian mixture models provide another approach to conditional distribution estimation. Hartford et al. (2017) and Shao et al. (2025) employ mixture density networks to estimate the parameters in conditional Gaussian mixtures. Convergence rates for mixture density networks and related Gaussian mixture models are also available under several measures (Zhou et al., 2023; Ho, Yang, and Jordan, 2022; Nguyen, Nguyen, and Ho, 2023).

4.2 Discussion on the first stage requirement

We briefly discuss the implications of the rate conditions on the first stage prediction error rates. For a benchmark, we consider simple sufficient conditions that the rates in Lemma 3.1 satisfy $\tau_n^2 r_{H,n} = o(1)$ and $a_n + b_n = o(1)$. To simplify the rate conditions, suppose $\xi_n \asymp \sqrt{k_n}$, $\tau_n r_{w,n} = O(1)$, and $r_{m,n} \asymp r_{\mu,n}$. The first condition holds for many commonly used sieves. The second condition is a conservative benchmark,

but imposing a faster rate for $r_{w,n}$ does not improve the sufficient rate conditions derived below. The last condition is a natural choice when no additional structure is imposed to reduce $r_{m,n}$ relative to $r_{\mu,n}$. Under these conditions, the sufficient rate requirements can be obtained as

$$\tau_n^3 k_n \sqrt{\frac{\log k_n}{n}} = o(1), \qquad r_{\mu,n} = o(\tau_n^{-1} n^{-1/4}). \tag{10}$$

The first one sets the upper bound on the growth of k_n. Together with the lower bound on k_n implied by Assumption 3.5(ii), they set the admissible range of sieve complexity for the statistical inference. The second one is the familiar faster-than-$n^{-1/4}$ requirement, augmented by the ill-posedness of the inverse problem.

For illustration, suppose further $L_{\mu,n} = O(1)$, which can be achieved under strong assumptions on the conditional density. Balancing the subsampling sizes in Examples 4.1 and 4.2 gives, respectively,

$$r_{\mu,n} \lesssim \left(\frac{k_n}{n}\right)^{\gamma_{\mathrm{RF}}/(1+2\gamma_{\mathrm{RF}})}, \qquad r_{\mu,n} \lesssim \left(\frac{k_n}{n}\right)^{1/(\tilde{d}_Z+2)}.$$

Since $\gamma_{\mathrm{RF}} < 1/2$ for any d_Z, the available GRF bound does not meet the benchmark first stage requirement. For subsampled nearest neighbors, the bound can meet the requirement when the intrinsic dimension is one, provided k_n and τ_n grows sufficiently slowly. However, we stress that these results are based on the upper bounds, whose theory often intentionally does not exploit strong regularities for more general applications. Therefore, the failure to meet the $o(n^{-1/4})$ rate of the upper bounds does not establish that these methods cannot be used in the first stage.

Other learners that exploit additional smoothness or structure have sharper prediction-error bounds that can satisfy the benchmark requirement with larger d_Z. For example, LASSO can exploit approximate sparsity and yield a rate that depends on the sparsity level and scales logarithmically in d_Z (Belloni et al., 2012). Two-scale distributional nearest neighbors can achieve rates faster than $n^{-1/4}$ for scalar outcomes when $d_Z \leq 7$ under additional regularity conditions (Demirkaya et al., 2024). Neural networks and Mondrian forests can achieve rates faster than $n^{-1/4}$ for any fixed

d_Z under sufficiently strong smoothness and other regularity conditions (Schmidt-Hieber, 2020; Cattaneo, Klusowski, and Underwood, 2026). We leave verification of Assumption 3.3 for these methods to future work.

Finally, the rate requirements can be relaxed by putting additional structure on the CMR problems. In the first stage, some sieve basis functions may have natural shape restrictions. These may potentially reduce the required covering or bracketing numbers of the first stage function class, and could result in sharper bounds. In the second stage, additional structure on the structural function, such as approximate sparsity, bounded variation, or economically motivated shape restrictions, may reduce the rate $r_{m,n}$ relative to $r_{\mu,n}$. We leave these relaxations for future work.

5 Simulation Study

In this section, we study how the performance of our method changes with the number of instruments, sample size, and noise in the endogenous regressor.

DGP We begin with a first stage design that is implied by a certain neural network architecture as an optimistic benchmark (Chen, Chen, and Tamer, 2023). We vary the number of instruments while keeping the distribution of the underlying predictive signal fixed by aggregating the instruments. This allows us to control the instruments' dimensionality and their joint relevance separately.

We first independently generate

$$Z \sim N(0, I_{d_Z}), \quad V \sim N(0,1), \quad \epsilon \sim N(0,1).$$

Then we partition the instruments into five equally sized groups I_r, and define

$$S_r = \frac{\sum_{j \in I_r} Z_j}{\sqrt{|I_r|}}, \quad g(Z) = \frac{\sum_{r=1}^{5} \tanh(S_r)}{\sqrt{5\nu}},$$

where $\nu = \mathbb{E}[\tanh(N(0,1))^2] \approx 0.39$. Note $S_r \sim N(0,1)$ regardlessly of d_Z and r.

Finally, we generate the endogenous regressor and outcome as

$$\begin{aligned} X &= \Phi\left(\frac{g(Z)+\sigma_X V}{\sqrt{1+\sigma_X^2}}\right)-\frac{1}{2}, \quad U=\sigma_U\{\eta V+\sqrt{1-\eta^2}\epsilon\}, \\ Y &= \theta_0(X)+U=2\sin(\pi X)+U. \end{aligned}$$

There are three DGP parameters $(\sigma_X, \sigma_U, \eta)$. The parameter σ_X controls the amount of variation in the endogenous regressor that is not explained by the instruments. The parameter σ_U determines the standard deviation of the structural error, while η controls its dependence on the shared shock V and hence the degree of endogeneity. By construction, $\mathrm{Corr}(U, V) = \eta$.

First stage learners We approximate θ_0 with quadratic and cubic B-splines having three or five interior knots. The first stage jointly predicts the conditional mean of the outcome and spline basis functions. We use the generalized random forests (GRF; Athey, Tibshirani, and Wager, 2019), feedforward neural networks (NN), and conditional diffusion estimator as first stage learners. We do 5-fold cross-fitting.

For GRF, we fix the subsampling proportion as 0.4 in all simulations[3]. We tried different subsampling sizes in pilot simulations and found that GRF in general works better with a larger subsampling size in our first stage. We fix other tuning parameters, such as minimum split portion and variable selections, to the default in the authors' R package.

For the neural networks, we consider an architecture with three hidden layers with sixteen neurons each. We use the ReLU activation function and Adam as a training algorithm with a batch size of 128. We try different learning rates $\{0.001, 0.003, 0.01\}$ and L_2 regularization parameters $\{0.001, 0.01\}$ in the Python package scikit-learn. For each DGP configuration, we generate three extra datasets and decide the tuning parameters by 5-fold out-of-sample prediction. In actual Monte Carlo simulations, we fixed the chosen tuning parameters.

[3]Theoretically, the subsampling size needs to satisfy $s_n/n \to 0$, so the proportion has to decrease, but we fix it at 0.4 due to computational burden.

For the conditional diffusion model, we separately estimate the conditional distribution of Y and X given Z. For X, we apply a logit transformation on $X + 1/2$ to make its support a real line. We use three hidden layers of 32 neurons and the SiLu activation function. We use the Adam algorithm with learning rate 0.001 and a batch size of 256. In each training fold, we reserved 20% of the training data for validation of early stopping. After fitting the model conditional distribution, we sampled 1024 Monte Carlo draws for numerical integration.

Simulation settings We do 200 Monte Carlo repetitions for neural networks and 50 repetitions for GRF due to computational burden. The Monte Carlo repetitions of the diffusion model are currently 50. We set as baseline $\sigma_X = \sqrt{0.1} \approx 0.31$, $\sigma_U = 0.5$, and $\eta = 0.9$. In the first experiment, we set the DGP parameters to the baseline and vary the number of instruments $d_Z \in \{5, 10, 20\}$ and the sample size $n \in \{2500, 5000, 10000\}$. In the second experiment, we fix $n = 5000$, $\sigma_U = 0.5$, $\eta = 0.9$, and vary $\sigma_X \in \{0.2, 0.31, 0.5\}$ and $d_Z \in \{5, 10, 20\}$.

In the baseline design, $\text{Corr}(X, U) \approx 0.26$. To quantify instrument relevance, we define $R_X^2 := \text{Var}(\mathbb{E}[X|Z])/\text{Var}(X)$, which measures the proportion of variation in X explained by the whole instruments jointly. We note that the conditional mean $\mathbb{E}[X|Z]$ is for reference (useful in linear IV), and it is not the object directly used in our first stage[4]. In the baseline design, R_X^2 is approximately 90.4%, which is very high in empirical settings. In the second experiment with noise, $R_X^2 \approx 0.96$ when $\sigma_X = 0.2$ and $R_X^2 \approx 0.79$ when $\sigma_X = 0.5$. We set it deliberately high because we find that the finite sample performance of our inference deteriorates substantially with the noise in instruments.

We perform two Wald inferences with nominal level 5%. One is the point evaluation at $x = 0$, and the other is the average derivative. For point evaluation, the null value ϕ_0 is 0. For average derivative, we trim the top and bottom 5 percentiles of the observations, and test for $\mathbb{E}[\theta_0'(X)|X_{(0.05)} \leq X \leq X_{0.95}]$ where $X_{(\alpha)}$ refers to

[4]For the conditional mean of Y and the quadratic B-spline (with three knots) basis functions, the analogue measures are $R_Y^2 = 0.69$ and $R_p^2 = (0.82, 0.73, 0.62, 0.62, 0.73, 0.82)$, respectively, in the baseline design.

α-quantile of X. The null value is fixed by Monte Carlo at $\phi_0 = 4.34$. As a performance report, we provide local power curves. To obtain the curve, we fix the DGP and instead shift the null value by $\phi_0 + \kappa\sqrt{V_n/n}$ for $\kappa \in \{0, 0.1, \cdots, 4\}$. V_n is fixed from separate Monte Carlo simulations. At $\kappa = 0$, the rejection frequency indicates the (empirical) size of the test, so we do not separately report size.

Results Figure 1 and 2 respectively illustrate the local power of the point evaluation and average derivative Wald test by sample size and instrument dimension. The reported second-stage sieve is a quadratic B-spline with three interior knots. In both tests and all configurations, the size was kept close to the nominal level of 5%. In Figure 1, when the sample size was small ($n = 2500$), the performance decreased substantially as the number of instruments grew. This is expected because modern machine learning methods can accommodate many variables computationally, but they need not eliminate the exponential growth in required sample sizes. Nevertheless, the power recovered fast in our simulations as the sample size gets larger. For the neural net with $d_Z = 20$, the rejection frequency at $\kappa = 4$ increased from 30% to 71% to 93% as n increases. The corresponding average derivative rejection rates were 50%, 98%, and 99%. The average derivative testing achieved nearly ideal rejection rates in all scenarios when $n = 10000$.

Although the upper bound in Example 4.1 does not deliver a first stage convergence rate faster than $n^{-1/4}$, we find that GRF performs well when the number of instruments is small. Its performance is less sensitive to tuning parameters, and it outperforms neural networks in some low-dimensional configurations. However, GRF performance deteriorates rapidly as the number of instruments increases, whereas neural networks perform better with moderately many instruments.

Figure 3 examines the sensitivity of local power to the noise level σ_X, holding the sample size ($n = 5000$) and other DGP parameters fixed. The results show that increased noise in the endogenous regressor can substantially reduce rejection frequencies, particularly when the number of instruments is large. The rejection frequencies of the neureal net at $\kappa = 4$ were 99%, 99%, 45% for $\sigma_X = 0.20, 0.31, 0.50$, resepectively.




Figure 1: Local power of point evaluation Wald test by sample size n and the instrument dimension d_Z. Nominal significance level is 5%. The dotted curve represents the Gaussian reference rejection probability, and the horizontal line indicates the nominal significance level.




Figure 2: Local power of average derivative Wald test by sample size n and the instrument dimension d_Z. Nominal significance level is 5%. The dotted curve represents the Gaussian reference rejection probability, and the horizontal line indicates the nominal significance level.

We may expect that greater noise in the endogenous regressor may affect the performance in multiple channels: aggravating population (sieve) ill-posedness and making the first stage regressions more challenging by reducing their signal-to-noise ratio. These effects can reinforce each other, as weaker population curvature amplifies first stage estimation errors and increases the risk of instability in the estimated Hessian. In DGP1, increasing the number of instruments does not change population sieve ill-posedness, but makes the conditional means harder to estimate. The bottom-right panel of Figure 3 suggests this potential vulnerability to noise could be pronounced in high-dimensional settings. The rejection rate of the neural net was 26% when $\kappa = 4$ in the plot. Also, while we find neural networks are highly sensitive to tuning, we only tried a few primitive tunings. Hence, the performance could improve with finer tuning.

6 Empirical Application

In this section, we revisit the empirical work of Beegle, De Weerdt, and Dercon (2011), who studied migration and economic mobility in Tanzania. The authors investigate whether geographic mobility improves individuals' living standards with the Kagera Health and Development Survey data, which tracks individuals from their original households in northwestern Tanzania between 1991 and 2004. While the original paper examines several aspects of migration and economic mobility, we focus on one of their instrumental variable regression, which estimates the effect of migration distance on consumption growth. We extend the linear IV specification by nonparametrically estimating the effect of migration distance on consumption growth.

Particularly, we consider the 2SLS in Table 9, column 4 of their study. Let Y_i denote the change in log real consumption per capita between 1991 and 2004, and let $X_i = \log(1 + D_i)$, where D_i is migration distance in kilometers. The original 2SLS model considers

$$Y_i = \beta X_i + C_i^\top \gamma + \alpha_{H_i} + U_i,$$




Figure 3: Local power of average derivative Wald test by noise σ_X and the instrument dimension d_Z. Nominal significance level is 5%. The dotted curve represents the Gaussian reference rejection probability, and the horizontal line indicates the nominal significance level.

where C_i contains twenty eight individual and family characteristics controls and α_{H_i} is an origin-household fixed effect. The six (excluded) instruments are household-position indicators and interactions involving age rank, distance to the regional capital, and the original rainfall-shock measure. The data contain 3227 individuals and 681 origin households. Of these individuals, 2147 have zero recorded migration distance, and 1080 have positive distance. In the 2SLS regression, the authors find the slope of X is 0.104 with strong significance.

To allow for a nonlinear relationship between migration distance and consumption growth, we consider the following partially linear NPIV specification while retaining the original controls, origin-household fixed effects, and excluded instruments:

$$Y_i = h(X_i) + C_i^\top \gamma + \alpha_{H_i} + U_i, \qquad \mathbb{E}[U_i \mid Z_i, C_i, H_i] = 0.$$

We approximate h with quadratic B-splines with five equally spaced knots. Without loss of generality, we normalize $h(0) = 0$ by subtracting its estimated value at zero and absorbing the differences into the intercept in C. Hence, the redefined function $h(x) = h^{raw}(x) - h(0)$ measures the effect of migration distance relative to staying in the original location. In the first stage, we employ GRF and neural networks. For GRF, we set the subsampling proportion to 0.4 and retain all other tuning parameters as the R package default. For neural networks, we use three layers of sixteen nodes with ReLU activation function and the Adam optimizer with learning rate and L_2 regularization to 0.001. We used a batch size of 128 and used 500 training epochs. We set aside 15% of the training samples as validation samples and use early stopping with patience of 25 epochs and tolerance of 10^{-4}. The first stage inputs include the excluded instruments, original controls, and origin-household indicators. We do 5-fold cross-fitting[5].

Figure 4 presents the estimated structural functions alongside the linear IV estimate. Both GRF and neural network estimates suggest that the relationship between

[5]The individuals that share the same household can be related. We tried several cross-fittings with and without preserving household relationships, but the results were similar. We report the one that considers household relationships in both origin (year 1991) and destination (year 2004) households.




Figure 4: Consumption-growth curves against $X = \log(1 + \text{distance in km})$. The B-spline curves use quadratic B-splines with five interior knots. The dotted line is the original linear IV fit. All curves are normalized to $h(0) = 0$. The scatter points show residualized outcomes, with colors matching the corresponding fitted curves. Vertical lines mark deciles (gray), P95 (red), and P98 (purple).

	Estimate	SE	Estimate	SE
Comparison in whole range	GRF		NN	
Linear IV slope	0.104**	0.043	-	-
Average derivative, P68-98	0.128***	(0.050)	0.054	(0.034)
Average derivative, P71-95	0.114**	(0.050)	0.008	(0.045)
Heterogeneity in distance	GRF		NN	
Average derivative, P70-90	-0.026	(0.064)	-0.053	(0.050)
Average derivative, P90-95	0.582***	(0.112)	0.396***	(0.104)
Average derivative, P90-98	0.564***	(0.099)	0.434***	(0.075)
Difference (P90-95)-(P70-90)	0.607***	(0.142)	0.450***	(0.124)

Table 1: Migration distance and consumption growth in Tanzania. The table reports the linear IV estimate and average derivatives of the estimated structural function over selected percentile ranges of migration distance. GRF and NN denote the first stage learners. Standard errors are reported in the adjacent columns. P70–90 denotes the 70th to 90th percentiles of the transformed migration distance. *** and ** respectively indicate 1% and 5% significance levels.

migration distance and consumption growth is relatively flat over most of the distribution, but increases sharply at higher migration distances. We therefore compare average derivatives across different ranges of migration distance to investigate potential heterogeneity in their marginal effect.

Before presenting the inference results, we note three caveats. First, our current theory assumes iid samples, so our standard errors do not account for the clustered error within households. We leave this generalization to future work. Second, the number of households increases with sample size. Although our theory does not preclude an increasing number of first stage inputs, the theory for employed machine learning methods may not allow it. In particular, the theory underlying GRF assumes a fixed number of inputs, although it can be relatively large. Finally, around two-thirds of observations are concentrated at the zero boundary of migration distance, while nonparametric methods sometimes have unstable behaviors near the boundary. This concentration may affect the stability of the sieve estimation. Also, the interesting findings are concentrated near the right boundary.

Table 1 provides suggestive evidence of heterogeneity. GRF estimates the average

derivative in the whole range, after trimming top and bottom 2% and 5% of observations with positive moving distance, similarly to the linear IV slope. Both GRF and NN estimates show that the average derivative is close to zero between the 70th and 90th percentiles but significantly larger between the 90th and 95th percentiles. The differences between the average derivatives are also statistically significant. De Weerdt, Christiaensen, and Kanbur (2021) also find larger consumption gains among migrants to distant cities than among those moving to nearby towns in Tanzania by distinguishing migrant destinations in the analysis.

7 Conclusion

This paper develops an orthogonalized sieve minimum distance framework for affine CMR models, allowing flexible machine learning methods in the first stage. Our analysis points to several directions for future research. First, the orthogonalized criterion function can be extended to smooth nonlinear CMR models, provided estimators of the conditional moment and its derivative. Second, incorporating penalization and economically motivated shape constraints may alleviate the demanding first-stage rate requirements. Third, the special structure of the first-stage prediction outcomes motivates the development of machine learning methods and theory that can exploit the relationships among them. Finally, our simulation results suggest the need for methods that are more robust to noise in the endogenous regressor and the amplification of first-stage prediction errors through ill-posedness.

References

Ai, C. and X. Chen (2003): "Efficient Estimation of Models with Conditional Moment Restrictions Containing Unknown Functions," *Econometrica*, 71, 1795–1843.

Alfaro, I., N. Bloom, and X. Lin (2024): "The Finance Uncertainty Multiplier," *Journal of Political Economy*, 132, 577–615.

Athey, S., J. Tibshirani, and S. Wager (2019): "Generalized Random Forests," *The Annals of Statistics*, 47, 1148–1178.

Autor, D. H., D. Dorn, and G. H. Hanson (2013): "The China Syndrome: Local Labor Market Effects of Import Competition in the United States," *American Economic Review*, 103, 2121–2168.

Bastos, P., J. Silva, and E. Verhoogen (2018): "Export Destinations and Input Prices," *American Economic Review*, 108, 353–392.

Beegle, K., J. De Weerdt, and S. Dercon (2011): "Migration and Economic Mobility in Tanzania: Evidence from a Tracking Survey," *Review of Economics and Statistics*, 93, 1010–1033.

Belloni, A., D. Chen, V. Chernozhukov, and C. Hansen (2012): "Sparse Models and Methods for Optimal Instruments with an Application to Eminent Domain," *Econometrica*, 80, 2369–2429.

Belloni, A., V. Chernozhukov, D. Chetverikov, and K. Kato (2015): "Some New Asymptotic Theory for Least Squares Series: Pointwise and Uniform Results," *Journal of Econometrics*, 186, 345–366.

Bennett, A. and N. Kallus (2023): "The variational method of moments," *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 85, 810–841.

Bennett, A., N. Kallus, X. Mao, W. Newey, V. Syrgkanis, and M. Uehara (2023): "Inference on Strongly Identified Functionals of Weakly Identified Functions," in *Proceedings of the Thirty Sixth Conference on Learning Theory*, vol. 195 of *Proceedings of Machine Learning Research*, 2265.

Bennett, A., N. Kallus, and T. Schnabel (2019): "Deep Generalized Method of Moments for Instrumental Variable Analysis," in *Advances in Neural Information Processing Systems*, vol. 32.

Blundell, R., X. Chen, and D. Kristensen (2007): "Semi-Nonparametric IV Estimation of Shape-Invariant Engel Curves," *Econometrica*, 75, 1613–1669.

Bruns-Smith, D. (2025): "Two-Stage Machine Learning for Nonparametric Instrumental Variable Regression," Working paper, Stanford University.

Burchardi, K. B., T. Chaney, and T. A. Hassan (2019): "Migrants, Ancestors, and Foreign Investments," *Review of Economic Studies*, 86, 1448–1486.

Carrasco, M., J.-P. Florens, and E. Renault (2007): "Linear Inverse Problems in Structural Econometrics: Estimation Based on Spectral Decomposition and Regularization," in *Handbook of Econometrics*, ed. by J. J. Heckman and E. E. Leamer, Amsterdam: Elsevier, vol. 6B, 5633–5751.

Cattaneo, M. D., J. M. Klusowski, and W. G. Underwood (2026): "Inference with Mondrian Random Forests," *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 88, 1060–1085.

Chen, J., D. L. Chen, and G. Lewis (2020): "Mostly Harmless Machine Learning: Learning Optimal Instruments in Linear IV Models," arXiv:2011.06158.

Chen, J., X. Chen, and E. Tamer (2023): "Efficient Estimation of Average Derivatives in NPIV Models: Simulation Comparisons of Neural Network Estimators," *Journal of Econometrics*, 235.

Chen, X. (2007): "Large Sample Sieve Estimation of Semi-Nonparametric Models," in *Handbook of Econometrics*, ed. by J. J. Heckman and E. E. Leamer, Amsterdam: Elsevier, vol. 6B, 5549–5632.

Chen, X., T. Christensen, and S. Kankanala (2025): "Adaptive Estimation and Uniform Confidence Bands for Nonparametric Structural Functions and Elasticities," *Review of Economic Studies*, 92.

Chen, X. and T. M. Christensen (2018): "Optimal Sup-Norm Rates and Uniform Inference on Nonlinear Functionals of Nonparametric IV Regression," *Quantitative Economics*, 9, 39–84.

Chen, X., Y. Liao, and W. Wang (2025): "Inference on Time Series Nonparametric Conditional Moment Restrictions Using Nonlinear Sieves," *Journal of Econometrics*, 249, 105920.

Chen, X. and Z. Liao (2014): "Sieve M Inference on Irregular Parameters," *Journal of Econometrics*, 182, 70–86.

Chen, X. and D. Pouzo (2015): "Sieve Wald and QLR Inferences on Semi/Nonparametric Conditional Moment Models," *Econometrica*, 83, 1013–1079.

Chernozhukov, V., D. Chetverikov, M. Demirer, E. Duflo, C. Hansen, W. Newey, and J. Robins (2018): "Double/Debiased Machine Learning for Treatment and Structural Parameters," *The Econometrics Journal*, 21, C1–C68.

Chernozhukov, V., J. C. Escanciano, H. Ichimura, W. K. Newey, and J. M. Robins (2022): "Locally Robust Semiparametric Estimation," *Econometrica*, 90, 1501–1535.

Chernozhukov, V., S. Lee, and A. M. Rosen (2013): "Intersection Bounds: Estimation and Inference," *Econometrica*, 81, 667–737.

Compiani, G. (2022): "Market counterfactuals and the specification of multiproduct demand: A nonparametric approach," *Quantitative Economics*, 13, 545–591.

Darolles, S., Y. Fan, J.-P. Florens, and E. Renault (2011): "Nonparametric Instrumental Regression," *Econometrica*, 79, 1541–1565.

De Weerdt, J., L. Christiaensen, and R. Kanbur (2021): "When Distance Drives Destination, Towns Can Stimulate Development," Policy Research Working Paper 9622, World Bank.

Demirkaya, E., Y. Fan, L. Gao, J. Lv, P. Vossler, and J. Wang (2024): "Optimal Nonparametric Inference with Two-Scale Distributional Nearest Neighbors," *Journal of the American Statistical Association*, 119, 297–307.

Denteh, A. and H. Liebert (2026): "Who increases emergency department use? New insights from the Oregon health insurance experiment," *Journal of the Royal Statistical Society Series A: Statistics in Society*, 189, 1378–1427.

Deryugina, T., G. Heutel, N. H. Miller, D. Molitor, and J. Reif (2019): "The Mortality and Medical Costs of Air Pollution: Evidence from Changes in Wind Direction," *American Economic Review*, 109, 4178–4219.

Dikkala, N., G. Lewis, L. Mackey, and V. Syrgkanis (2020): "Minimax Estimation of Conditional Moment Models," in *Advances in Neural Information Processing Systems*, vol. 33.

Domínguez, M. A. and I. N. Lobato (2004): "Consistent estimation of models defined by conditional moment restrictions," *Econometrica*, 72, 1601–1615.

Donald, S. G., G. W. Imbens, and W. K. Newey (2009): "Choosing Instrumental Variables in Conditional Moment Restriction Models," *Journal of Econometrics*, 152, 28–36.

Escanciano, J. C., S. Hoderlein, A. Lewbel, O. Linton, and S. Srisuma (2021): "Nonparametric Euler Equation Identification and Estimation," *Econometric Theory*, 37, 851–891.

Fang, Z. and J. Seo (2021): "A Projection Framework for Testing Shape Restrictions That Form Convex Cones," *Econometrica*, 89, 2439–2458.

Florens, J.-P. and S. Sokullu (2017): "Nonparametric Estimation of Semiparametric Transformation Models," *Econometric Theory*, 33, 839–873.

Foster, D. J. and V. Syrgkanis (2023): "Orthogonal Statistical Learning," *The Annals of Statistics*, 51, 879–908.

Fu, H., Z. Yang, M. Wang, and M. Chen (2024): "Unveil Conditional Diffusion Models with Classifier-Free Guidance: A Sharp Statistical Theory," arXiv:2403.11968.

Gibbs, A. L. and F. E. Su (2002): "On Choosing and Bounding Probability Metrics," *International Statistical Review*, 70, 419–435.

Hall, P. and J. L. Horowitz (2005): "Nonparametric Methods for Inference in the Presence of Instrumental Variables," *The Annals of Statistics*, 33, 2904–2929.

Hartford, J., G. Lewis, K. Leyton-Brown, and M. Taddy (2017): "Deep IV: A Flexible Approach for Counterfactual Prediction," in *Proceedings of the 34th International Conference on Machine Learning*, vol. 70 of *Proceedings of Machine Learning Research*, 1414–1423.

Ho, J. and T. Salimans (2022): "Classifier-Free Diffusion Guidance," arXiv:2207.12598.

Ho, N., C.-Y. Yang, and M. I. Jordan (2022): "Convergence Rates for Gaussian Mixtures of Experts," *Journal of Machine Learning Research*, 23, 1–81.

Khosravi, K., G. Lewis, and V. Syrgkanis (2022): "Non-Parametric Inference Adaptive to Intrinsic Dimension," in *Proceedings of the First Conference on Causal Learning and Reasoning*, vol. 177 of *Proceedings of Machine Learning Research*.

Lewis, G. and V. Syrgkanis (2018): "Adversarial Generalized Method of Moments," arXiv:1803.07164.

Magnus, J. R. and H. Neudecker (2019): *Matrix Differential Calculus with Applications in Statistics and Econometrics*, Hoboken, NJ: Wiley, third ed.

Muandet, K., A. Mehrjou, S. K. Lee, and A. Raj (2020): "Dual Instrumental Variable Regression," in *Advances in Neural Information Processing Systems*, vol. 33.

Newey, W. K. (1997): "Convergence Rates and Asymptotic Normality for Series Estimators," *Journal of Econometrics*, 79, 147–168.

Newey, W. K. and D. McFadden (1994): "Large Sample Estimation and Hypothesis Testing," in *Handbook of Econometrics*, ed. by R. F. Engle and D. L. McFadden, Amsterdam: Elsevier, vol. 4, 2111–2245.

Newey, W. K. and J. L. Powell (2003): "Instrumental Variable Estimation of Nonparametric Models," *Econometrica*, 71, 1565–1578.

Nguyen, H., T. Nguyen, and N. Ho (2023): "Demystifying Softmax Gating Function in Gaussian Mixture of Experts," in *Advances in Neural Information Processing Systems*, vol. 36.

Oprescu, M., V. Syrgkanis, and Z. S. Wu (2019): "Orthogonal Random Forest for Causal Inference," in *Proceedings of the 36th International Conference on Machine Learning*, vol. 97 of *Proceedings of Machine Learning Research*, 4932–4941.

Pollard, D. (2002): *A User's Guide to Measure Theoretic Probability*, Cambridge: Cambridge University Press.

Portier, F. and J. Segers (2019): "Monte Carlo Integration with a Growing Number of Control Variates," *Journal of Applied Probability*, 56, 1168–1186.

Schmidt-Hieber, J. (2020): "Nonparametric Regression Using Deep Neural Networks with ReLU Activation Function," *The Annals of Statistics*, 48, 1875–1897.

Shao, D., A. Soleymani, F. Quinzan, and M. Kwiatkowska (2025): "Double Machine Learning for Conditional Moment Restrictions: IV Regression, Proximal Causal Learning and Beyond," arXiv:2506.14950.

Singh, R., M. Sahani, and A. Gretton (2019): "Kernel Instrumental Variable Regression," in *Advances in Neural Information Processing Systems*, vol. 32.

Song, Y., J. Sohl-Dickstein, D. P. Kingma, A. Kumar, S. Ermon, and B. Poole (2021): "Score-Based Generative Modeling through Stochastic Differential Equations," in *International Conference on Learning Representations*.

Yurinskii, V. V. (1977): "On the Error of the Gaussian Approximation for Convolutions," *Theory of Probability and Its Applications*, 22, 236–247.

Zhou, Y., C. Shi, L. Li, and Q. Yao (2023): "Testing for the Markov Property in Time Series via Deep Conditional Generative Learning," *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 85, 1204–1222.

A Gaussian coupling of the sieve coefficients

In this section, we provide a Gaussian coupling of the linear representation in Lemma 3.1. The coupling can be used as a building block for various nonstandard inference problems.

Assumption A.1. (i) Assumptions 3.3 and Assumption 3.4(iii) hold uniformly over $P \in \mathcal{P}$, with constants and deterministic rate sequences independent of P. (ii) Suppose there is a deterministic sequence τ_n^2 such that $\sup_P \|G_{n,P}^{1/2} H_{n,P}^{-1/2}\|_{op} \leq \tau_n$ and $\tau_n^2 r_{H,n} = o(1)$. (iii) For a given deterministic sequence $D_{n,P}$ for each $P \in \mathbf{P}$, $\sup_{P\in\mathcal{P}} \mathbb{E}_P\left[\left\|D_{n,P} H_{n,P}^{-1} \psi_{n,P}(O)\right\|_2^3\right] < \infty$. (iv) Define $\Delta_{D,P} := n^{-1/2}\mathbb{E}_P[\|D_{n,P} H_{n,P}^{-1}\psi_{n,P}(O)\|_2^3]$. For a deterministic sequence $c_n \to 0$, suppose $c_n^{-3} k_n \sup_P \Delta_{D,P} \to 0$ and $(a_n + b_n)\sup_P \|D_{n,P}\|_{op} = o(c_n)$.

$D_{n,P}$ is a problem specific choice. For example, $D_{n,P}$ can be taken as $n^{-1/2} r_n I_{k_n}$ to verify Assumption 3.2(i) in Fang and Seo (2021), or can be taken as $(H_{n,P}^{-1}\Omega_{n,P} H_{n,P}^{-1})^{-1/2}$ to verfify Condition C in Chernozhukov, Lee, and Rosen (2013). (i) For Assumption 3.3, the examples provided yield bounds with the constants that do not depend on P. Requiring Assumption 3.4(iii) uniformly over P restricts the class of DGP so that the sieve approximation error measured in weak norm is bounded by a common deterministic rate $r_{w,n}$. Thus, the rate is chosen to accommodate the largest weak norm approximation error over $P \in \mathcal{P}$. (ii) imposes an upper envelope on the sieve measure of ill-posedness uniformly over P (Chen and Christensen, 2018). (iii) is the third momoment condition for Yurinskii's coupling. (iv) c_n is a tolerance of approximation a researcher chose. For instance, it can be taken as $\log(n)$ or n^{-a} for some $a > 0$.

Lemma A.1. Suppose Assumption A.1 holds. For a deterministic sequence of matrix $D_{n,P}$, there exists Gaussian random vectors $W_{n,P}^D \sim N(0, D_{n,P}\Sigma_{n,P} D_{n,P}^\top)$ such that

$$\|\sqrt{n} D_{n,P}(\hat{\beta}_n - \beta_{P,n}) - W_{n,P}^D\|_2 = o_P(c_n)$$

uniformly over $P \in \mathbf{P}$.

The proof is a direct application of the Yurinskii coupling (Yurinskii, 1977).

B Proofs

B.1 Proofs for Section 3

To relieve notational burden, we write $f_i = f(A_i)$ for any function f and random variable A_i.

Lemma B.1 (Hessian estimation error)**.** Suppose Assumptions 3.1(i)(iv)(vi), 3.2(i)(ii), 3.3, and 3.4(iv) hold.

$$\|\hat{H}_n - H_n\|_{op} = O_p(\xi_n\sqrt{\frac{\log k_n}{n}} + \frac{r_{\mu,n}\xi_n}{\sqrt{n}} + r_{\mu,n}^2).$$

Proof of Lemma B.1. Denote $\tilde{H}_n := \mathbb{E}_n[\sigma_i^{-2}\{\mu_{qi}q_i^\top + (q_i - \mu_{qi})\mu_{qi}^\top\}]$ and $e(Z_i) := \hat{\mu}_q^{(-\ell)}(Z_i) - \mu_q(Z_i)$ for $i \in I_\ell$.

$$\begin{aligned}\hat{H}_n &\equiv \mathbb{E}_n\Big[\frac{1}{\sigma^2}\Big\{\hat{\mu}_{qi}q_i^\top + (q_i - \hat{\mu}_{qi})\hat{\mu}_{qi}^\top\Big\}\Big] \\ &= \tilde{H}_n + \mathbb{E}_n\Big[\frac{1}{\sigma^2}\Big\{e(q-\mu)^\top + (q-\mu)e^\top - ee^\top\Big\}\Big].\end{aligned}$$

By Assumption 3.4(iv), $\mathbb{E}_P[\|e(Z_i)\|_2^2|\mathcal{T}] = O_p(r_{\mu,n}^2)$ where $\mathcal{T}$ denotes the training fold(s). Meanwhile, by Assumption 3.1(iv) and Assumption 3.2(ii), $\|q_n(O)\|_2 \lesssim \xi_n$ P-a.s.. By Jensen's inequality, $\|\mu_q(Z)\|_2 \lesssim \xi_n$ P-a.s., hence, by triangle inequality $\|q - \mu_q\|_2 \lesssim \xi_n$ P-a.s.. Together with Assumption 3.4(iv), in each fold ℓ,

$$\begin{aligned}\mathbb{E}\Big[\Big\|\frac{1}{n_\ell}\sum_{i\in I_\ell}[e_i(q_i-\mu_i)^\top/\sigma_i^2]\Big\|_{op}^2\Big|\mathcal{T}_\ell\Big] &\leq \mathbb{E}\Big[\Big\|\frac{1}{n_\ell}\sum_{i\in I_\ell}[e_i(q_i-\mu_i)^\top/\sigma_i^2]\Big\|_F^2\Big|\mathcal{T}_\ell\Big] \\ &= \frac{1}{n_\ell}\mathbb{E}\Big[\frac{\|e_i(q_i-\mu_i)^\top\|_F^2}{\sigma_i^4}\Big|\mathcal{T}_\ell\Big] \\ &= \frac{1}{n_\ell}\mathbb{E}\Big[\frac{\|e_i\|_2^2\|q_i-\mu_i\|_2^2}{\sigma_i^4}\Big|\mathcal{T}_\ell\Big] \\ &\lesssim O_p\big(\frac{r_{\mu,n}^2\xi_n^2}{n_\ell}\big) = O_p\big(\frac{r_{\mu,n}^2\xi_n^2}{n}\big).\end{aligned}$$

Aggregating across ℓ, by the law of iterative expectation and the Markov inequality,

$\left\|\mathbb{E}_n\left[\frac{e_i(q_i-\mu_i)^\top}{\sigma_i^2}\right]\right\|_{op} = O_p(\frac{r_{\mu,n}\xi_n}{\sqrt{n}})$. The second remainder has the same bound. Similarly, the last remainder term is $O_p(r_{\mu,n}^2)$ by Assumption 3.4(iv), the Cauchy-Schwarz inequality, and the Markov inequality. Hence,

$$\|\hat{H}_n - \tilde{H}_n\|_{op} = O_p(\frac{r_{\mu,n}\xi_n}{\sqrt{n}} + r_{\mu,n}^2).$$

Notice $\tilde{H}_n = \mathbb{E}_n[\frac{1}{\sigma^2}(q_i q_i^\top - (q_i - \mu_{qi})(q_i - \mu_{qi})^\top)]$. We may apply Lemma 6.2 of Belloni et al. (2015) separately to each term. From the bounds on $\|q_i\|_2$ and $\|q_i - \mu_{qi}\|_2$ above, the envelope condition holds with $M = \xi_n^2/\underline{\sigma}^2$ and $M = 4\xi_n^2/\underline{\sigma}^2$, respectively. Then, provided $\xi_n\sqrt{\frac{\log k_n}{n}} = o(1)$,

$$\|\tilde{H}_n - H_n\|_{op} = O_p\Big(\frac{\xi_n^2 \log k_n}{n} + \xi_n\sqrt{\frac{\log k_n}{n}}\Big) = O_p\Big(\xi_n\sqrt{\frac{\log k_n}{n}}\Big).$$

Finally, by the triangle inequality,

$$\|\hat{H}_n - H_n\|_{op} = O_p\Big(\xi_n\sqrt{\frac{\log k_n}{n}} + \frac{r_{\mu,n}\xi_n}{\sqrt{n}} + r_{\mu,n}^2\Big).$$

□

Lemma B.2 is a standalone lemma that collects several established results in the literature. See, for instance, Appendix F.1 of Chen and Christensen (2018) and Appendix A.1 of Belloni et al. (2015). A proof is provided for completeness.

Lemma B.2. Let H_n be a deterministic symmetric positive-definite $k_n \times k_n$ matrix. Suppose $\lambda_{\min}(H_n) \geq c_n > 0$ for some deterministic sequence $c_n > 0$ that can go to 0. Let $\hat{H}_n$ be an estimator of H_n, which is symmetric and $k_n \times k_n$. Define $\Delta_{H,n} := H_n^{-1/2}(\hat{H}_n - H_n)H_n^{-1/2}$ and suppose that $\|\Delta_{H,n}\|_{op} = o_p(1)$ and $\xi_n\sqrt{\frac{\log k_n}{n}} = o(1)$. Then, $\hat{H}_n$ is positive definite (and hence invertible[6]) with probability approaching to 1. Moreover, $\|\hat{H}_n^{-1}\|_{op} = O_p(c_n^{-1})$ and $\|\hat{H}_n^{-1} - H_n^{-1}\|_{op} = O_p(\|\hat{H}_n - H_n\|_{op}/c_n^2)$.

[6]Define $\hat{H}_n^{-1}$ as any matrix, for instance 0 matrix, if $\hat{H}_n$ is singular.

The second half of the result might not be useful, if we take care of the product terms separately as in Fang and Seo (2021).

Proof of Lemma B.2. By definition of $\Delta_{H,n}$, $\hat{H}_n = H_n^{1/2}(I_{k_n} + \Delta_{H,n})H_n^{1/2}$. Fix $\epsilon > 0$. On the event $\mathcal{A}_n(\epsilon) = \{\|\Delta_{H,n}\|_{op} \leq \epsilon\}$, for any nonzero $u \in \mathbf{R}^{k_n}$, let $v := H_n^{1/2}u$. Then,

$$\begin{aligned} u^\top \hat{H}_n u &= u^\top H_n^{1/2}(I_{k_n} + \Delta_{H,n})H_n^{1/2}u \\ &= \|v\|_2^2 + v^\top \Delta_{H,n} v \\ &\geq \|v\|_2^2 - \|\Delta_{H,n}\|_{op}\|v\|_2^2 \\ &\geq (1-\epsilon)\|v\|_2^2 \\ &= (1-\epsilon)u^\top H_n u \\ &> 0. \end{aligned}$$

By assumption, $Pr(\mathcal{A}_n(\epsilon)) \to 1$. Hence, $Pr(\hat{H}_n > 0) \geq Pr(\mathcal{A}_n(\epsilon)) \to 1$.

Define $\hat{H}_n^{-1} = H_n^{-1/2}(I_{k_n} + \Delta_{H,n})^{-1}H_n^{-1/2}$. For the same reason, $(I_{k_n} + \Delta_{H,n})$ is positive definite on $\mathcal{A}_n(\epsilon)$ and hence invertible. By construction, $\hat{H}_n^{-1}$ is the inverse of $\hat{H}_n$. Again, on the event $A_n(\epsilon)$, from the definition of eigenvalue,

$$\|\hat{H}_n^{-1}\|_{op} = \frac{1}{\lambda_{\min}(\hat{H}_n)} \leq \frac{1}{(1-\epsilon)\lambda_{\min}(H_n)} \leq \frac{1}{(1-\epsilon)c_n}. \tag{11}$$

The last inequality follows from $\|H_n^{-1}\|_{op} = 1/\lambda_{\min}(H_n) \leq 1/c_n$.

Finally, on the same event, both inverses exists, and

$$\begin{aligned} \|\hat{H}_n^{-1} - H_n^{-1}\|_{op} &= \|\hat{H}_n^{-1}(H_n - \hat{H}_n)H_n^{-1}\|_{op} \\ &\leq \|\hat{H}_n^{-1}\|_{op}\|\hat{H}_n - H_n\|_{op}\|H_n^{-1}\|_{op} \\ &\leq \frac{(1-\epsilon)^{-1}\|\hat{H}_n - H_n\|_{op}}{c_n^2} \\ &= O_p(\frac{\|\hat{H}_n - H_n\|_{op}}{c_n^2}). \end{aligned}$$

□

Proof of Lemma 3.1 (ALR). Define $\tilde{\Psi}_n = \mathbb{E}_n[\psi_i(\beta_{P,n})]$ where ψ_i is defined in Equation (3). Recall we assume $\beta_{P,n}$ is an interior solution. We may decompose as

$$\begin{aligned}\sqrt{n}(\hat{\beta}_n - \beta_{P,n}) &= -H_n^{-1}\frac{1}{\sqrt{n}}\sum_i \psi_i + (H_n^{-1} - \hat{H}_n^{-1})\frac{1}{\sqrt{n}}\sum_i \psi_i - \hat{H}_n^{-1}\sqrt{n}(\hat{\Psi}_n - \tilde{\Psi}_n) \\ &:= -H_n^{-1}\mathbb{G}_n\psi_i + R_{H,n} + R_{\psi,n} \qquad (12)\end{aligned}$$

The first term is the leading term in the lemma. It is centered due to the interior solution and the first order condition.

For the first remainder $R_{H,n}$, we first bound $\|H_n^{-1}\mathbb{G}_n\psi_i\|_2$. Recall $\psi_i = (\rho_i - m_i)\mu_{qi}/\sigma_i^2 + m_i q_i/\sigma_i^2$. The first term is by independence, LIE, Assumption 3.1(ii), Assumption 3.4(iii),

$$\begin{aligned}\mathbb{E}_P\Big[\Big\|H_n^{-1}\frac{1}{\sqrt{n}}\sum_i \frac{(\rho_i - m_i)\mu_{qi}}{\sigma_i^2}\Big\|_2^2\Big] &\lesssim \mathbb{E}_P[\frac{\mu_q^T H_n^{-2}\mu_q}{\sigma_i^2}] \\ &= \mathrm{tr}(H_n^{-1}E_P[\frac{\mu_{qi}\mu_{qi}^\top}{\sigma_i^2}]H_n^{-1}) \\ &= \mathrm{tr}(H_n^{-1}) \lesssim \tau_n^2 k_n. \qquad (13)\end{aligned}$$

The second term is by independence, the Cauchy-Schwarz inequality, Assumption 3.4(iii),

$$\begin{aligned}\mathbb{E}_P\Big[\|H_n^{-1}\frac{1}{\sqrt{n}}\sum_i \frac{m_i q_i}{\sigma_i^2}\|_2^2\Big] &= \mathbb{E}_P[\frac{m_i^2}{\sigma_i^4}q_i^\top H_n^{-2}q_i] \\ &\le \|H_n^{-1}\|_{op}^2 E_P[\frac{m_i^2\|q_i\|_2^2}{\sigma_i^4}] \lesssim \tau_n^4\xi_n^2 r_{w,n}^2. \qquad (14)\end{aligned}$$

The last inequality holds from $\mathbb{E}_P[m_i^2/\sigma_i^2] = Q_P(\theta_{P,n}) \le Q_P(\Pi_n\theta_P) = O(r_{w,n}^2)$.

Hence, by Equations (13), (14), and Lemma B.1,

$$\begin{aligned}
\|R_{H,n}\|_2 &= \Big\|\hat{H}_n^{-1}(\hat{H}_n - H_n)H_n^{-1}\frac{1}{\sqrt{n}}\sum_i \psi_i\Big\|_2 \\
&\leq \|\hat{H}_n^{-1}\|_{op}\|(\hat{H}_n - H_n)\|_{op}\Big\|H_n^{-1}\frac{1}{\sqrt{n}}\sum_i \psi_i\Big\|_2 \\
&= O_p(\tau_n^2\{\xi_n\sqrt{\frac{\log k_n}{n}} + \frac{r_{\mu,n}\xi_n}{\sqrt{n}} + r_{\mu,n}^2\}\{\tau_n\sqrt{k_n} + \tau_n^2\xi_n r_{w,n}\}). \qquad (15)
\end{aligned}$$

For the second remainder $R_{\psi,n}$, we first derive the estimation error of the $\hat{\Psi}_n$ term. Denoting $e_\mu = \hat{\mu}_q - \mu_q$ and $e_m = \hat{m}(\cdot, \beta_{P,n}) - m(\cdot, \beta_{P,n})$,

$$\begin{aligned}
\hat{\Psi}_n(\beta_{P,n}) &:= \frac{1}{n}\sum_i \frac{1}{\sigma_i^2}\Big\{\rho_i\hat{\mu}_i + \hat{m}_i(q_i - \hat{\mu}_i)\Big\} \\
&= \frac{1}{n}\sum_i \frac{1}{\sigma^2}\Big\{\rho(\mu + e_\mu) + (m + e_m)(q - \mu - e_\mu)\Big\} \\
&= \tilde{\Psi}_n(\beta_{P,n}) + \frac{1}{n}\sum_i \frac{1}{\sigma^2}\Big\{(\rho - m)e_\mu + e_m(q - \mu) - e_m e_\mu\Big\}
\end{aligned}$$

Similarly to the proof of Lemma B.1, in each fold ℓ, by indendency, Assumption 3.1(ii), the Cauchy-Schwarz inequality, and Assumption 3.4(iv), 3.3,

$$\begin{aligned}
\mathbb{E}\Big[\Big\|\frac{1}{n_\ell}\sum_{i\in I_\ell}[(\rho - m)e_\mu/\sigma^2]\Big\|_2^2\Big|\mathcal{T}_\ell\Big] &= \frac{1}{n_\ell}\mathbb{E}[\sigma_i^{-4}\|(\rho_i - m_i)e_{\mu,i}\|_2^2|\mathcal{T}_\ell] \\
&\lesssim \frac{1}{n_\ell}\mathbb{E}\big[\|e_{\mu,i}\|_2^2|\mathcal{T}_\ell\big] = O_p(\frac{r_{\mu,n}^2}{n})
\end{aligned}$$

Similarly,

$$\mathbb{E}\Big[\Big\|\frac{1}{n}\sum_i[e_m(q - \mu)/\sigma^2]\Big\|_2^2\Big|\mathcal{T}_\ell\Big] \leq \frac{1}{n_\ell}\mathbb{E}\Big[\frac{e_m^2\|q - \mu_q\|_2^2}{\sigma^4}\Big|\mathcal{T}_\ell\Big] = O_p(\frac{\xi_n^2 r_{m,n}^2}{n})$$

The last $e_m e_\mu$ term is by the Cauchy-Schwarz and Markov inequalities, $O_p(r_{\mu,n}r_{m,n})$.

Hence,

$$\|\hat{\Psi}_n(\beta_{P,n}) - \tilde{\Psi}_n(\beta_{P,n})\|_2 = O_p(\frac{r_{\mu,n}}{\sqrt{n}} + \frac{\xi_n r_{m,n}}{\sqrt{n}} + r_{m,n} r_{\mu,n}). \tag{16}$$

Combining Equation (12), Lemma B.2, (16)

$$\begin{aligned} \|R_{\psi,n}\|_2 &\leq \sqrt{n}\|\hat{H}_n^{-1}\|_{op}\|\hat{\Psi}_n - \tilde{\Psi}_n\|_2 \\ &= O_p(\tau_n^2 r_{\mu,n} + \xi_n r_{m,n} + \sqrt{n} r_{m,n} r_{\mu,n}) = O_p(b_n). \end{aligned} \tag{17}$$

Finally, define $R_n = R_{H,n} + R_{\psi,n}$, then

$$\|R_n\|_2 \leq \|R_{H,n}\|_2 + \|R_{\psi,n}\|_2 = O_p(a_n + b_n).$$

□

Proof of Theorem 3.1. Suppose V_n is known. By Assumption 3.5 and 3.6,

$$\begin{aligned} \frac{\sqrt{n}}{\sqrt{V_n}}(\phi(\hat{\theta}_n) - \phi(\theta_P)) &= \frac{\sqrt{n}}{\sqrt{V_n}}(\phi(\hat{\theta}_n) - \phi(\theta_{P,n})) + \frac{\sqrt{n}}{\sqrt{V_n}}(\phi(\theta_{P,n}) - \phi(\theta_P)) \\ &= \frac{\sqrt{n}}{\sqrt{V}_n} l_n^\top(\hat{\beta}_n - \beta_{P,n}) + o_p(1). \end{aligned}$$

By Lemma 3.1,

$$\begin{aligned} \frac{\sqrt{n}}{\sqrt{V}_n} l_n^\top(\hat{\beta}_n - \beta_{P,n}) &= -\frac{1}{\sqrt{V}_n} l_n^\top \frac{1}{\sqrt{n}} \sum_i \psi_{ni} + \frac{1}{\sqrt{V}_n} l_n^\top (R_{H,n} + R_{\psi,n}) \\ &= -\frac{1}{\sqrt{n}} \sum_i \frac{S_{ni}}{\sqrt{V_n}} + \frac{1}{\sqrt{V}_n} l_n^\top (R_{H,n} + R_{\psi,n}). \end{aligned}$$

By the triangle inequality, the Cauchy-Schwarz inequality, and the assumption that $\|l_n\|_2(a_n + b_n)/\sqrt{V_n} = o_p(1)$,

$$\Big|\frac{1}{\sqrt{V_n}}(l_n^\top R_{H,n} + l_n^\top R_{\psi,n})\Big| \leq \frac{\|l_n\|_2}{\sqrt{V_n}}(\|R_{H,n}\|_2 + \|R_{\psi,n}\|_2) = O_p(\frac{\|l_n\|_2(a_n + b_n)}{\sqrt{V_n}}) = o_p(1).$$

Define $X_{ni} = \frac{S_{ni}}{\sqrt{V_n}}$ and apply the Lindeberg-Feller CLT with Assumption 3.6. Then, conclude by Lemma B.3 and Slutsky's lemma. □

Lemma B.3 (Wald variance estimator)**.** Suppose $\tau_n^2 r_{H,n} = o(1)$ and $\|l_n\|_2(a_n + b_n)/\sqrt{V_n} = o(1)$ (the same assumptions as Theorem 3.1). Then, $\hat{V}_n/V_{n,P} \xrightarrow{P} 1$.

Proof. Denote $\|f\|_{n,2} := (\mathbb{E}_n\|f_i\|_2^2)^{1/2}$. Recall $\hat{V}_n = \mathrm{Var}_n(\hat{S}_{ni})$. We first establish $\mathbb{E}_n\|\hat{S}_{ni} - S_{ni}\|_2/\sqrt{V}_n = o_p(1)$, then show $|\mathbb{E}_n S_{ni}^2/V_n - 1| = o_p(1)$.

$$\hat{S}_{ni} - S_{ni} = l_n^\top(\hat{H}_n^{-1} - H_n^{-1})\psi_{ni} + l_n^\top \hat{H}_n^{-1}(\hat{\psi}_{ni} - \psi_{ni}) + l_n^\top \hat{H}_n^{-1}\{\hat{\psi}_{ni}(\hat{\beta}_n) - \hat{\psi}(\beta_{P,n})\}.$$

The first term is by the Cauchy-Schwarz inequality and Equation (15),

$$\|l_n^\top(\hat{H}_n^{-1} - H_n^{-1})\psi_{ni}\|_{n,2} \leq \|l_n\|_2\|\hat{H}_n^{-1}\|_{op}\|\hat{H}_n - H_n\|_{op}\|H_n^{-1}\psi_{ni}\|_{n,2} = O_p(\|l_n\|_2 a_n) \quad (18)$$

Similarly, $\|l_n^\top \hat{H}_n^{-1}(\hat{\psi}_{ni} - \psi_{ni})\|_{n,2} = O_p(\|l_n\|_2 b_n)$ from the second part of Equation (17) and the Cauchy-Schwarz inequality. To bound the last term,

$$\begin{aligned}
\|\hat{\psi}_{ni}(\hat{\beta}_n) - \hat{\psi}(\beta_{P,n})\|_{n,2}^2 &= \left\|\frac{\hat{\mu}_{qi}q_i^\top + (q_i - \hat{\mu}_{qi}\hat{\mu}_{qi})^\top}{\sigma_i^2}(\hat{\beta}_n - \beta_{P,n})\right\|_{n,2}^2 \\
&:= \frac{1}{n}\sum_i \left\|\hat{H}_{ni}(\hat{\beta}_n - \beta_{P,n})\right\|_2^2
\end{aligned}$$

For any $v \in \mathbf{R}^{k_n}$, by Assumption 3.4(iii), the Cauchy-Schwarz inequality,

$$\begin{aligned}
\|\hat{H}_{ni}v\|_2^2 &= \|\frac{1}{\sigma_i^2}\big(\mu_{qi}q^\top v + (q_i - \mu_{qi})\mu_q^\top v + e_\mu(q_i - \mu_{qi})^\top v + (q_i - \mu_{qi})e_\mu^\top v - e_\mu e_\mu^\top v\big)\|_2^2 \\
&\lesssim \|\mu_{qi}q^\top v\|_2^2 + \|(q_i - \mu_{qi})\mu_q^\top v\|_2^2 + \|e_\mu(q_i - \mu_{qi})^\top v\|_2^2 + \|(q_i - \mu_{qi})e_\mu^\top v\|_2^2 + \|e_\mu e_\mu^\top\|_2^2 \\
&\leq \xi_n^2|q_i^\top v|^2 + 4\xi_n^2|\mu_q^\top v|^2 + \|e_\mu\|_2^2|(q_i - \mu_{qi})^\top v|^2 + 4\xi_n^2|e_\mu^\top v|^2 + \|e_\mu\|_2^2|e_\mu^\top v|^2 \\
\mathbb{E}_n\|\hat{H}_{ni}v\|_2^2 &\lesssim \xi_n^2 v^\top \mathbb{E}_n[q_i q_i^\top]v + 4\xi_n^2 v^\top \mathbb{E}_n[\mu_{qi}\mu_{qi}^\top]v + O_p(\xi_n^2 r_{\mu,n}^2\|v\|_2^2) + O_p(\xi_n^2 r_{\mu,n}^2\|v\|_2^2) \\
&\quad + \mathbb{E}_n\|e_\mu\|_2^4\|v\|_2^2.
\end{aligned}$$

The first two terms are $O_p(\xi_n^2\|v\|_2^2)$ because $\mathbb{E}_n[q_i q_i^\top]$ and $\mathbb{E}_n[\mu_{qi}\mu_{qi}^\top]$ have a bounded operator norm due to Lemma 6.2 of Belloni et al. (2015) as in Lemma B.1. To

bound the last term, since $(\mathbb{E}_n\|e_\mu\|_2^2)^2 = n^{-1}\mathbb{E}_n\|e_\mu\|_2^4 + 2n^{-2}\sum_{i>j}\|e_{\mu,i}\|_2^2\|e_{\mu,j}\|_2^2 \geq n^{-1}\mathbb{E}_n\|e_\mu\|_2^4$, multiplying n both sides yields

$$\frac{1}{n}\sum_i \|e_\mu\|_2^4 \leq n(\mathbb{E}_n\|e_\mu\|_2^2)^2 = O_p(nr_{\mu,n}^4).$$

The above bound on $\|\hat{H}_{ni}v\|_2^2$ holds for any $v \in \mathbf{R}^{k_n}$. Therefore, $\|\hat{\psi}_{ni}(\hat{\beta}_n) - \hat{\psi}_{ni}(\beta_{P,n})\|_{n,2} = O_p(\{\xi_n + \xi_n r_{\mu,n} + \sqrt{n}r_{\mu,n}^2\})\|\hat{\beta}_n - \beta_{P,n}\|_2$. From Equations (13), (14), and Lemma 3.1, $\|\hat{\beta}_n - \beta_{P,n}\|_2 = O_p(n^{-1/2}\{\tau_n^2\sqrt{k_n + \xi_n^2 r_{w,n}^2} + a_n + b_n\})$. Combining all three terms,

$$\begin{aligned}\|l_n^\top \hat{H}_n^{-1}\{\hat{\psi}_{ni}(\hat{\beta}_n) - \hat{\psi}_{ni}(\beta_{P,n})\}\|_{n,2} &= O_p\Big(\|l_n\|_2\tau_n^2\big(\frac{\xi_n}{\sqrt{n}} + \frac{\xi_n r_{\mu,n}}{\sqrt{n}} + r_{\mu,n}^2\big)\big(\tau_n\sqrt{k_n} + \tau_n^2\xi_n r_{w,n} + a_n + b_n\big)\Big) \\ &= O_p(\|l_n\|_2(a_n + b_n)),\end{aligned}$$

where the last step uses the assumption that $\tau_n^2 r_{H,n} = o(1)$.

Finally, under the assumption that $\|l_n\|_2(a_n+b_n)/\sqrt{V}_n = o_p(1)$, $\|\hat{S}_{ni} - S_{ni}\|_{n,2}/\sqrt{V_n} = o_p(1)$.

To conclude the proof, under Assumption 3.6, apply, for example, Proposition 1 of Portier and Segers (2019) with $Y_{ni} = \frac{S_{ni}^2}{V_n}$. Then, $\mathbb{E}_n Y_{ni} \xrightarrow{p} 1$, hence, $|S_{ni}^2/V_n - 1| \xrightarrow{p} 0$ as desiered. □

B.2 Proofs for Appendix A

Proof of Lemma A.1 (Gaussian coupling). Let $\mathbb{G}_n\psi_{n,P}(O_i) := \frac{1}{\sqrt{n}}\sum_i\{\psi_{n,P}(O_i) - \mathbb{E}_P[\psi_{n,P}(O)]\}$. From the interior solution assumption, $\mathbb{E}_P[\psi_{n,P}] = 0$. Multiplying $D_{n,P}$ to the linear expansion in Lemma 3.1,

$$\|\sqrt{n}D_{n,P}(\hat{\beta}_n - \beta_{P,n}) - W_{n,P}^D\|_2 \leq \| - D_{n,P}H_{n,P}^{-1}\mathbb{G}_n\psi_{n,P} - W_{n,P}^D\|_2 + \|D_{n,P}R_{n,P}\|_2.$$

We may apply Theorem 10.10 of Pollard (2002) to the empirical process term and show the remainder is $o(c_n)$. Choose $\epsilon_n \to 0$ sufficiently slowly so that $\frac{k_n}{\epsilon_n^3 c_n^3}\sup_P \Delta_{D,P} \to$

0. Applying the theorem with $\delta = \epsilon_n c_n$ and $B = \frac{k_n \Delta_{D,P}}{\epsilon_n^3 c_n^3}$,

$$P(\|D_{n,P} H_{n,P}^{-1} \mathbb{G}_n \psi_{n,P} - W_{n,P}^D\|_2 > 3\epsilon_n c_n) \leq C_0 B\{1 + \frac{|\log(1/B)|}{k_n}\} \to 0$$

for some universal constant C_0 uniformly over $P \in \mathcal{P}$. Hence, for any fixed $\eta > 0$, whenever n is sufficiently large,

$$P(\|D_{n,P} H_{n,P}^{-1} \mathbb{G}_n \psi_{n,P} - W_{n,P}^D\|_2 > \eta c_n) \leq P(\|D_{n,P} H_{n,P}^{-1} \mathbb{G}_n \psi_{n,P} - W_{n,P}^D\|_2 > 3\epsilon_n c_n) \to 0.$$

Hence, $\|D_{n,P} H_{n,P}^{-1} \mathbb{G}_n \psi_{n,P} - W_{n,P}^D\|_2 = o_P(c_n)$ uniformly over P. The remainder term is $o_P(c_n)$ uniformly over P by Assumption A.1(iv). □

C Verification of Assumption 3.3

Lemma C.1 (Verification of Assumption 3.3 for KLS)**.** Suppose the following conditions hold. (i) $Z \in [0,1]^{dz}$ and the marginal density is bounded away from 0 uniformly over P. (ii) $\|\mu_{n,P}(z) - \mu_{n,P}(z')\|_2 \leq L_{\mu,n}\|z - z'\|_2$, for any $z, z' \in \mathcal{Z}_P$ for every $P \in \mathcal{P}$ with the Lipschitz constant independent of P. (iii) There exists constatns $C_c, C_q < \infty$ independent of n, P, and a deterministic sequence ξ_n independent of P, such that $\|\mathrm{Var}_P(c_n(O)|Z)\|_{P,\infty} \leq C_c^2$ and $\|q_n(O)\|_2 \leq C_q \xi_n$ P-a.s.. (iv) Subsample size $s_n \to \infty$, $s_n/n \to 0$, κ is fixed, and the number of subsampling $B_n \gtrsim n/s_n$.

Then, for an evaluation draw $Z \sim P_Z$ independent of $\mathcal{T}_\ell$,

$$\mathbb{E}_P[\|\hat{\mu}_n^{(-\ell)}(Z) - \mu_{n,P}(Z)\|_2^2 | \mathcal{T}_\ell] = O_P(L_{\mu,n}^2 s_n^{-2/d_z} + (1 + \xi_n^2)\frac{s_n}{n}).$$

Condition (ii) depends on the choice of the sieve dictionary. The Lipschitz constant $L_{\mu,n}$ may grow with n, as the number of the sieve dictionary grows. Condition (iii) is direct from the envelop condition in the main text.

Proof of Lemma C.1. Fix the fold ℓ, the evaluation index set I_ℓ, the training index set J_ℓ, and an evaluation point z. Let $n_J := |J| = n - |I|$. Let A denote all the

auxiliary learner randomness (such as subsampling, tie-breaking randomness), which is independent of the observations. Define the sigma field $\mathcal{D} = \sigma(\{Z_j\}_{j\in J_\ell}, A)$. By construction, the weights depend only on the point z, the conditioning variables $\{Z_j, j \in J\}$, and the auxiliary randomness A. Hence, the weights are $\mathcal{D}$-measurable and considered as fixed real numbers conditional on $\mathcal{D}$. Decompose the prediction error as

$$\hat{\mu}_n^{(-\ell)}(z) - \mu_{n,P}(z) = \sum_{j\in J_\ell} \alpha_j^{(-\ell)}(z)\{\begin{pmatrix} c_n(O_j) \\ q_n(O_j) \end{pmatrix} - \mu_{n,P}(Z_j)\} + \sum_{j\in J_\ell} \alpha_j^{(-\ell)}(z)\{\mu_{n,P}(Z_j) - \mu_{n,P}(z)\}.$$

Conditional on $\mathcal{D}$, the first term has mean 0, and the second term is fixed. Hence, the cross-product between the two terms has mean 0. Then,

$$\begin{aligned} \mathbb{E}_P[\|\hat{\mu}_n^{(-\ell)}(z) - \mu_{n,P}(z)\|_2^2|\mathcal{D}] &= \|\sum_{j\in J_\ell} \alpha_j^{(-\ell)}(z)\{\mu_{n,P}(Z_j) - \mu_{n,P}(z)\}\|_2^2 \\ &\quad + \sum_{j\in J_\ell} \alpha_j^{(-\ell)}(z)^2 \mathbb{E}_P[\|(c_n(O_j), q_n(O_j))^\top - \mu_{n,P}(Z_j)\|_2^2|Z_j]. \end{aligned}$$

By the Jensen inequality and the Lipschitzness assumption,

$$\begin{aligned} \|\sum_{j\in J_\ell} \alpha_j^{(-\ell)}(z)\{\mu_{n,P}(Z_j) - \mu_{n,P}(z)\}\|_2^2 &\leq \sum_{j\in J_\ell} \alpha_j^{(-\ell)}(z)\|\mu_{n,P}(Z_j) - \mu_{n,P}(z)\|_2^2 \\ &\leq L_{\mu,n}^2 \sum_{j\in J_\ell} \alpha_j^{(-\ell)}(z)\|Z_j - z\|_2^2 \\ &= \frac{L_{\mu,n}^2}{B_n} \sum_b \sum_{j\in J_\ell} \alpha_{b,j}^{(-\ell)}(z)\|Z_j - z\|_2^2 \\ &\leq \frac{L_{\mu,n}^2}{B_n} \sum_b R_\kappa(z, S_b)^2. \end{aligned}$$

where $R_\kappa(z, S_b)$ denotes the distance from z to the κ-th nearest neighbor in subsample

S_b. Meanwhile, by the bounded conditional second moment assumption,

$$\begin{aligned}\mathbb{E}_P[\|(c_n(O_j), q_n(O_j))^\top - \mu_{n,P}(Z_j)\|_2^2 | Z_j] &= \mathrm{Var}_P(c_n(O_j)|Z_j) + \mathbb{E}_P[\|q_n(O_j)\|_2^2|Z_j] - \|\mu_q(Z_j)\|_2^2 \\ &\leq C_c^2 + C_q^2 \xi_n^2.\end{aligned}$$

Combining the bounds and taking expectation over $\mathcal{D}$,

$$\mathbb{E}_P[\|\hat{\mu}_n^{(-\ell)}(z) - \mu_{n,P}(z)\|_2^2] \leq L_{\mu,n}^2 \mathbb{E}_P[R_\kappa(z,S)^2] + (C_c^2 + C_q^2\xi_n^2)\mathbb{E}_P[\sum_{j \in J_\ell} \alpha_j^{(-\ell)}(z)^2].$$

Now we bound the two expectation terms on the right-hand side. Under the condition (i), for small $t \in (0, \frac{1}{2})$,

$$P_Z\{B(z,t)\} = \int_{B(z,t)\cap[0,1]^{d_z}} f_{P,Z}(u)du \geq \underline{f} 2^{-d_z} v_{d_z} t^{d_z}$$

where v_{dz} is the volume of d_z-dimensional unit ball. Then, their Lemma 2 and the tail bound in Appendix H.2 imply that there exist positive constants c and C_0 independent of P, z, ℓ, such that, for sufficiently large n, for $t \in [C_0 s_n^{-1/d_z}, r_0]$,

$$P\{R_\kappa(z,S) > t\} \leq 2\exp(-cs_n t^{d_z}).$$

Following their integration argument in their Appendix H.2 on Page 27[7] with the identity $\mathbb{E}[X^2] = \int_0^\infty 2tPr(X > t)dt$ and $t_n = C_0 s_n^{-1/d_z}$, we obtain $\mathbb{E}_P[R_\kappa(z,S)^2] \leq C_R s_n^{-2/d_z}$ for some $C_R > 0$ uniformly over P and P-almost every z. We show the mean of the sum of squared weights is $O(\frac{1}{\kappa}(\frac{1}{B_n} + \frac{(B_n-1)s_n}{B_n n_J}))$. Hence, for sufficiently

[7]where they obtain the first moment of R_κ in their analysis, while we need the second moment.

large B_n, it is $O(\frac{s_n}{\kappa n_J})$.

$$\begin{aligned}\mathbb{E}_P[\sum_{j\in J_\ell}(\alpha_j^{(-\ell)}(z))^2|Z_j, j\in J_\ell] &= \frac{1}{B_n^2}\sum_{b=1}^{B_n}\sum_{j\in J_\ell}\mathbb{E}_P[(\alpha_{b,j}^{(-\ell)}(z))^2|Z_j, j\in J_\ell] \\ &\quad +\frac{1}{B_n^2}\sum_{b\neq b'}^{B_n}\sum_{j\in J_\ell}\mathbb{E}_P[\alpha_{b,j}^{(-\ell)}(z)\alpha_{b',j}^{(-\ell)}(z)|Z_j, j\in J_\ell] \\ &= \frac{1}{\kappa B_n}+\frac{B_n(B_n-1)}{B_n^2}\sum_{j\in J_\ell}\mathbb{E}_P[\alpha_{b,j}^{(-\ell)}(z)|Z_j, j\in J_\ell]^2\end{aligned}$$

because b and b' are independent, and $\alpha_{b,j}^{(-\ell)}(z) = \frac{1}{\kappa}$ if j is in κ-nearest neighbor in subsample S_b, and 0 otherwise. Meanwhile, $\mathbb{E}_P[\alpha_{b,j}^{(-\ell)}(z)|Z_j, j\in J_\ell]$ has natural worst-case and best-case bounds 0 and $s_n/\kappa n_J$, and sums up to 1 by taking the conditional expectation to the identity $\sum_{j\in J_\ell}\alpha_{b,j}^{(-\ell)}(z) = 1$. Hence,

$$\begin{aligned}\sum_{j\in J_\ell}\mathbb{E}_P[\alpha_{b,j}^{(-\ell)}(z)|Z_j, j\in J_\ell]^2 &\leq (\max_{j\in J_\ell}\mathbb{E}_P[\alpha_{b,j}^{(-\ell)}(z)|Z_j, j\in J_\ell])\sum_{j\in J_\ell}\mathbb{E}_P[\alpha_{b,j}^{(-\ell)}(z)|Z_j, j\in J_\ell] \\ &\leq \frac{s_n}{\kappa n_J}.\end{aligned}$$

Therefore, for a fixed point z and sufficiently large $B_n \geq n_J/s_n$, we obtain the explicit finite sample bound

$$\mathbb{E}_P[\|\hat{\mu}_n^{(-\ell)}(z) - \mu_{n,P}(z)\|_2^2] \leq (C_c^2 + C_q^2\xi_n^2)\frac{s_n}{\kappa n_J} + C_R L_{\mu,n}^2 s_n^{-2/d_z} = O(r_{\mu,n}^2),$$

where $r_{\mu,n}^2 = (C_c^2 + C_q^2\xi_n^2)\frac{s_n}{\kappa n} + L_{\mu,n}^2 s_n^{-2/d_z}$. We use $n \asymp n_J$ in the last equality. Note that the above bound holds for z-a.s., with a constant independent of P, z, and the fold ℓ.

Finally, to obtain the desired result in Assumption 3.3, let $Z \sim P_Z$ be an inde-

pendent evaluation draw. Then,

$$\begin{aligned}\mathbb{E}_P\Big[\mathbb{E}_P[\|\hat{\mu}_n^{(-\ell)}(Z)-\mu_{n,P}(Z)\|_2^2|\mathcal{T}_\ell]\Big] &= \mathbb{E}_P[\int_Z \|\hat{\mu}_n^{(-\ell)}(Z)-\mu_{n,P}(Z)\|_2^2 dP_Z] \\ &= \int_Z \mathbb{E}_P[\|\hat{\mu}_n^{(-\ell)}(Z)-\mu_{n,P}(Z)\|_2^2] dP_Z \\ &\leq Cr_{\mu,n}^2,\end{aligned}$$

for some fixed $C > 0$. By the Markov inequality $\mathbb{E}_P[\|\hat{\mu}_n^{(-\ell)}(Z)-\mu_{n,P}(Z)\|_2^2|\mathcal{T}_\ell] = O_p(r_{\mu,n}^2)$ uniformly over $P \in \mathcal{P}$. □

Lemma C.2 (Verification of Assumption 3.3 for ATW (GRF))**.** Suppose the observations are i.i.d., the training folds satisfy $|\mathcal{T}_\ell| \asymp n$, and the following conditions hold uniformly over $P \in \mathcal{P}$. (i) $Z \in [0,1]^{d_z}$ for a fixed d_z and the density of Z is bounded above and away from zero by constants independent of P. (ii) $\|\mu_{n,P}(z)-\mu_{n,P}(z')\|_2 \leq L_{\mu,n}\|z-z'\|_2$ for some deterministic sequence $L_{\mu,n}$. (iii) There exist constants $C_c, C_q < \infty$, independent of n, P, such that $\operatorname{ess\,sup}_z \mathbb{E}_P[c_n(O)^2 \mid Z = z] \leq C_c^2, \|q_n(O)\|_2 \leq C_q\xi_n \quad P$-a.s. (iii) The forest is honest and regular in the sense of Specification 1 of Athey, Tibshirani, and Wager (2019), with a fixed leaf size. (iv) Subsample size $s_n \to \infty$, the number of trees $B_n \gtrsim n/s_n$.

Then, for $Z \sim P_Z$ independent of $\mathcal{T}_\ell$,

$$\mathbb{E}_P\left[\|\hat{\mu}_n^{(-\ell)}(Z)-\mu_{n,P}(Z)\|_2^2|\mathcal{T}_\ell\right] = O_P\Big(L_{\mu,n}^2 s_n^{-2\gamma} + (1+\xi_n^2)\frac{s_n}{n}\Big)$$

uniformly over $P \in \mathcal{P}$. Here, s_n is a tuning parameter and γ is a constant[8] decided from some other tuning parameters and d_z.

Proof of Lemma C.2. Fix a fold ℓ and an evaluation point z. Let the training index set be J_ℓ and write $n_J = |J_\ell|$. Let $J_{\ell b} \subset J_\ell$ be the subsample for $b = 1, ..., B_n$. For

[8] $\gamma := \frac{\pi}{2d_z}\frac{\log((1-w)^{-1})}{\log(w^{-1})}$ where π is a constant from 'each coordinate is selected for splitting with probability at least π/d_z', w is a minimum split fraction. Stronger balance w might exclude useful split – learn data less, but prevent worst outcome. It affects the bias term (larger w means larger γ. The default choice of 'mtry' in GRF package does not guarantee a specific π. One may force it by using a different feature selection rule, e.g., uniformly choose one among d_z instruments, so that $\pi = 1$. We stick to the default 'mtry' in our implementation.

honest split, let $J_{\ell b} = J_{\ell b1} \cup J_{\ell b2}$, where $J_{\ell b1}$ is the evaluation split and $J_{\ell b2}$ is the partitioning split. Let A contain all auxiliary randomness used for subsampling, honest splitting, feature selection, and tie-breaking. A is independent of the observations. Define a sigma field $\mathcal{D}_\ell := \sigma(\{Z_j, O_j : j \in J_\ell\}, A)$. Let index b denote an individual tree. Write the tree prediction

$$T_b(z) = \sum_{i \in J_\ell} \alpha_{b,i}(z) \begin{pmatrix} c_n(O_i) \\ q_n(O_i) \end{pmatrix}, \qquad \hat{\mu}_n^{(-\ell)}(z) = \frac{1}{B_n} \sum_{b=1}^{B_n} T_b(z),$$

where $\alpha_{b,i}(z) \geq 0$ and $\sum_i \alpha_{b,i}(z) = 1$. See Athey, Tibshirani, and Wager (2019) for the exact definition. Define the finite and infinite forest

$$\hat{\mu}_n^{(-\ell)}(z) = \frac{1}{B_n} \sum_{b=1}^{B_n} T_b(z), \qquad \hat{\mu}_{\infty,n}^{(-\ell)}(z) := \mathbb{E}_A[T_b(z) | O_j, j \in J_\ell].$$

where the expectation is taken over tree randomness A_b. Note $\mathbb{E}_{P,A}[\hat{\mu}_n^{(-\ell)}(z)] = \mathbb{E}_P[\hat{\mu}_\infty^{(-\ell)}(z)]$.

Decomposing the mean squared error,

$$\begin{aligned}
&\mathbb{E}_{P,A} \|\hat{\mu}_n^{(-\ell)}(z) - \mu_{n,P}(z)\|_2^2 \\
&= \mathbb{E}_{P,A} \left\| \{\mathbb{E}_{P,A}[\hat{\mu}_n^{(-\ell)}(z)] - \mu_{n,P}(z)\} + \{\hat{\mu}_{\infty,n}^{(-\ell)}(z) - \mathbb{E}_P[\hat{\mu}_{\infty,n}^{(-\ell)}(z)]\} + \{\hat{\mu}_n^{(-\ell)}(z) - \hat{\mu}_{\infty,n}^{(-\ell)}(z)\} \right\|_2^2 \\
&= \|\mathbb{E}_{P,A}[\hat{\mu}_n^{(-\ell)}(z)] - \mu_{n,P}(z)\|_2^2 + \mathbb{E}_P \|\hat{\mu}_{\infty,n}^{(-\ell)}(z) - \mathbb{E}_P[\hat{\mu}_{\infty,n}^{(-\ell)}(z)]\|_2^2 + \mathbb{E}_{P,A} \|\hat{\mu}_n^{(-\ell)}(z) - \hat{\mu}_{\infty,n}^{(-\ell)}(z)\|_2^2.
\end{aligned}$$

The first term in the decomposition is deterministic, the second term has mean 0, and the last term has mean 0 conditionally on $\{(Z_i, O_i) : i \in J_\ell\}$. Hence, the cross product terms are 0 by iterative expectation, and we obtain the last equality.

The first term is $\|\mathbb{E}_P[\hat{\mu}_{\infty,n}^{(-\ell)}(z)] - \mu_{n,P}(z)\|_2^2 \leq C L_{\mu,n}^2 s_n^{-2\gamma}$ by the Lipschitz condition and the localization bound in Equation (26) of ATW18. The second term satisfies

$$\mathbb{E}_P \|\hat{\mu}_{\infty,n}^{(-\ell)}(z) - \mathbb{E}_P[\hat{\mu}_{\infty,n}^{(-\ell)}(z)]\|_2^2 = \operatorname{tr}(\operatorname{Var}_P(\hat{\mu}_{\infty,n}^{(-\ell)}(z))) \leq \frac{s_n}{n_J}(C_c^2 + C_q^2 \xi_n^2),$$

where the inequality follows by applying the first ineqaulity in Equation (27) of

ATW18 to our conditional mean score $\psi_{\mu_{n,P}(z)}(O_i) = (c_n, q_n^\top)^\top - \mu_{n,P}(z)$, Jensen's inequality, honesty, and the condition (ii) in Lemma C.2. For the last term, by LIE conditioning on $\{Z_i, O_i : i \in J_\ell\}$, conditional independency and mean 0 of the trees,

$$\begin{aligned}\mathbb{E}_{P,A}\big[\|\hat{\mu}_n^{(-\ell)}(z) - \hat{\mu}_{\infty,n}^{(-\ell)}(z)\|_2^2\big] &= \frac{1}{B_n}\mathbb{E}_{P,A}\big[\|T_1(z) - \hat{\mu}_{\infty,n}^{(-\ell)}(z)\|_2^2\big] \\ &\leq \frac{1}{B_n}\mathbb{E}_{P,A}\|T_1(z)\|_2^2 \leq \frac{C_c^2 + C_q^2\xi_n^2}{B_n}.\end{aligned}$$

The last inequality follows from Jensen's inequality, LIE conditioning on $\{A_1, (Z_i, O_i) : i \in J_{\ell 11}\}$, honesty, condition (ii), and $\sum_i \alpha_{1,i}(z) = 1$.

Substituting the three bounds into the mean squared error decomposition and taking $B_n \geq n_J/s_n$, we obtain

$$\mathbb{E}_{P,A}\|\hat{\mu}_n^{(-\ell)}(z) - \mu_{n,P}(z)\|_2^2 \leq 2(C_c^2 + C_q^2\xi_n^2)\frac{s_n}{n_J} + CL_{\mu,n}^2 s_n^{-2\gamma}.$$

The constants are independent of P, z, ℓ.

Finally,

$$\begin{aligned}\mathbb{E}_{P,A}\left[\mathbb{E}_P\left[\|\hat{\mu}_n^{(-\ell)}(Z) - \mu_{n,P}(Z)\|_2^2 \mid \mathcal{D}_\ell\right]\right] &= \int \mathbb{E}_{P,A}\|\hat{\mu}_n^{(-\ell)}(z) - \mu_{n,P}(z)\|_2^2 \, dP_Z(z) \\ &= O_P(L_{\mu,n}^2 s_n^{-2\gamma} + (1+\xi_n^2)\frac{s_n}{n}),\end{aligned}$$

where the last equality uses $n_J \asymp n$. The Markov's inequality yields

$$\mathbb{E}_P\big[\|\hat{\mu}_n^{(-\ell)}(Z) - \mu_{n,P}(Z)\|_2^2 \mid \mathcal{D}_\ell\big] = O_P(L_{\mu,n}^2 s_n^{-2\gamma} + (1+\xi_n^2)\frac{s_n}{n})$$

uniformly over $P \in \mathcal{P}$. □

Lemma C.3 (Verification of the first stage assumption for conditional diffusion)**.** Write $O = (V, Z)$, where $V \in \mathbb{R}^{d_V}$ and d_V, d_z are fixed, and write $c(O) = c(V, Z)$ and $q_n(O) = q_n(V, Z)$. Suppose the following additional conditions hold, with constants independent of P, n, and the fold.

Suppose Assumptions 3.1(i)(ii),(iv)–(vi) and 3.2(i)(ii) hold. Moreover, suppose

the following conditions hold. (i) Let $Z \in [0,1]^{d_z}$. Let φ_λ be the density of $N(0, \lambda^{-1} I_{d_V})$ for a fixed $\lambda > 0$. The conditional density of V given $Z = z$ satisfies $f_{0,P}(v \mid z) = \varphi_\lambda(v) g_P(v, z)$, where g_P is bounded above and away from zero and belongs to a s-Hölder ball on $\mathbb{R}^{d_V} \times [0,1]^{d_z}$. (ii) The architecture of the neural network and tuning parameters satisfy the specification in Theorems 3.4 and 4.2 of Fu et al. (2024). (iii) The estimated conditional distribution satisfies $\operatorname{ess\,sup}_z \int c(v,z)^2 \, d\hat{P}_n^{(-\ell)}(v \mid z) = O_P(1)$ uniformly over $P \in \mathbf{P}$.

Define

$$\widehat{\mu}_n^{(-\ell)}(z) = \int \begin{pmatrix} c(v,z) \\ q_n(v,z) \end{pmatrix} d\widehat{P}_n^{(-\ell)}(v \mid z).$$

Then, for $Z \sim P_Z$ independent of $\mathcal{T}_\ell$,

$$\mathbb{E}_P\big[\|\widehat{\mu}_n^{(-\ell)}(Z) - \mu_{n,P}(Z)\|_2^2 \mid \mathcal{T}_\ell\big] = O_P\big((1 + \xi_n^2)\varepsilon_n^2\big) \tag{19}$$

uniformly over P where $\varepsilon_n^2 = n^{-2s/(2s+d_V+d_z)} (\log n)^{\max\{19, s+2\}}$.

In Condition (i), the support of Z is for convenience and can be relaxed to unbounded distributions with small tails (Fu et al., 2024, Appendix C). Condition (i) requires the conditional density to be a smooth, uniformly bounded positive multiple of a fixed Gaussian density. It allows non-Gaussian conditional distributions but imposes restrictive support and tail conditions.