

	Training (n=300)	Testing (n=128)	P-value
OS			
Mean (SD)	894 (811)	882 (729)	0.881
Median [Min, Max]	672 [14.0, 4500]	706 [6.00, 4230]	
age			
Mean (SD)	66.9 (13.0)	66.1 (13.3)	0.525
Median [Min, Max]	69.0 [31.0, 90.0]	67.0 [34.0, 90.0]	
status			
0	234 (78.0%)	100 (78.1%)	1
1	66 (22.0%)	28 (21.9%)	
gender			
male	159 (53.0%)	69 (53.9%)	0.914
female	141 (47.0%)	59 (46.1%)	
preoperative.CEA			
Mean (SD)	35.4 (218)	44.8 (168)	0.699
Median [Min, Max]	2.91 [0.200, 2610]	3.80 [0.00, 1390]	
Missing	115 (38.3%)	41 (32.0%)	
stage			
i	55 (18.3%)	18 (14.1%)	0.682
ii	114 (38.0%)	54 (42.2%)	
iii	87 (29.0%)	39 (30.5%)	
iv	44 (14.7%)	17 (13.3%)	
lymphatic.invasion			
no	165 (55.0%)	71 (55.5%)	0.731
yes	103 (34.3%)	48 (37.5%)	
Missing	32 (10.7%)	9 (7.0%)	
perineural.invasion			
no	81 (27.0%)	48 (37.5%)	0.0665
yes	34 (11.3%)	9 (7.0%)	
Missing	185 (61.7%)	71 (55.5%)	
venous.invasion			
no	193 (64.3%)	90 (70.3%)	0.285
yes	66 (22.0%)	23 (18.0%)	
Missing	41 (13.7%)	15 (11.7%)	

Note:Chi-sq test for categorical variables,continuity correction is only used in the 2-by-2 case, and t-test for continous variables.