

	alive (n=343)	dead (n=98)	P-value
Overall survival (days)			
Mean (SD)	943 (803)	711 (690)	0.00523
Median [Min, Max]	742 [6.00, 4500]	425 [29.0, 3040]	
Disease free survival (days)			
Mean (SD)	893 (667)	923 (763)	0.86
Median [Min, Max]	739 [6.00, 4500]	820 [29.0, 2530]	
Missing	83 (24.2%)	76 (77.6%)	
Age (years)			
Mean (SD)	65.9 (12.8)	69.5 (13.4)	0.0174
Median [Min, Max]	67.0 [31.0, 90.0]	72.5 [34.0, 90.0]	
Gender			
male	182 (53.1%)	54 (55.1%)	0.733
female	161 (46.9%)	44 (44.9%)	
preoperative.CEA (ng/ml)			
Mean (SD)	21.5 (99.0)	114 (416)	0.122
Median [Min, Max]	2.80 [0.00, 1290]	6.20 [1.09, 2610]	
Missing	114 (33.2%)	47 (48.0%)	
Stages			
i	70 (20.4%)	4 (4.1%)	<0.001
ii	141 (41.1%)	28 (28.6%)	
iii	96 (28.0%)	31 (31.6%)	
iv	30 (8.7%)	31 (31.6%)	
Missing	6 (1.7%)	4 (4.1%)	
person.neoplasm.status			
tumor free	260 (75.8%)	22 (22.4%)	<0.001
with tumor	72 (21.0%)	61 (62.2%)	
Missing	11 (3.2%)	15 (15.3%)	
lymphatic.invasion			
no	203 (59.2%)	42 (42.9%)	0.0085
yes	108 (31.5%)	44 (44.9%)	
Missing	32 (9.3%)	12 (12.2%)	
perineural.invasion			
no	107 (31.2%)	24 (24.5%)	0.144
yes	32 (9.3%)	13 (13.3%)	
Missing	204 (59.5%)	61 (62.2%)	
venous.invasion			
no	241 (70.3%)	51 (52.0%)	<0.001
yes	56 (16.3%)	34 (34.7%)	
Missing	46 (13.4%)	13 (13.3%)	

Note: Categorical variable using Chi-Square tests and T-test for continuous variable.