

Table S. Clinical features in the training and validation group

	Training group (n=300)	Validation group (n=128)	p-value
OS			0.689
Mean (sd)	880 (752)	915 (864)	
Median [Min, Max]	676 [14.0, 4500]	686 [6.00, 4230]	
Age			0.12
Mean (sd)	66.1 (13.3)	68.1 (12.3)	
Median [Min, Max]	68.0 [31.0, 90.0]	71.0 [36.0, 90.0]	
Status			1
Alive	234 (78.0%)	100 (78.1%)	
Dead	66 (22.0%)	28 (21.9%)	
Gender			0.449
Male	156 (52.0%)	72 (56.2%)	
Female	144 (48.0%)	56 (43.8%)	
Preoperative CEA (ng/ml)			0.555
Mean (sd)	33.6 (197)	50.4 (218)	
Median [Min, Max]	3.20 [0.200, 2610]	2.85 [0.00, 1390]	
Missing	106 (35.3%)	50 (39.1%)	
Stage			0.988
i	51 (17.0%)	22 (17.2%)	
ii	117 (39.0%)	51 (39.8%)	
iii	90 (30.0%)	36 (28.1%)	
iv	42 (14.0%)	19 (14.8%)	
Lymphatic invasion			0.909
No	167 (55.7%)	69 (53.9%)	
Yes	106 (35.3%)	45 (35.2%)	
Missing	27 (9.0%)	14 (10.9%)	
Perineural invasion			0.513
No	98 (32.7%)	31 (24.2%)	
Yes	35 (11.7%)	8 (6.2%)	
Missing	167 (55.7%)	89 (69.5%)	
venous invasion			0.685
No	201 (67.0%)	82 (64.1%)	
Yes	61 (20.3%)	28 (21.9%)	
Missing	38 (12.7%)	18 (14.1%)	