

## JANUARY 2024 PRELIMINARY EXAM

**Problem 1.** Suppose  $\{a_n\}$  and  $\{b_n\}$  are two complex sequences such that

$$\lim_{n \rightarrow \infty} a_n b_n = 0.$$

Show that at least one of  $\{a_n\}$  and  $\{b_n\}$  has a subsequence that converges to zero.

**Problem 2.** Suppose  $f : [a, b] \rightarrow \mathbb{R}$  is bounded, and moreover  $f$  is Riemann integrable on  $[a, c]$  for all  $a < c < b$ . Show that  $f$  is Riemann integrable on  $[a, b]$ .

**Problem 3.** Suppose  $f : \mathbb{R} \rightarrow \mathbb{R}$  is differentiable and  $\lim_{x \rightarrow \infty} f'(x) = 0$ . Show that if the sequence  $\{f(n)\}_{n \in \mathbb{N}}$  converges, then the limit  $\lim_{x \rightarrow \infty} f(x)$  exists.

*Proof.* Let  $L := \lim_{n \in \mathbb{N}, n \rightarrow \infty} f(n)$ . We will show  $\lim_{x \rightarrow \infty} f(x) = L$ .

Let  $\epsilon > 0$ . Then there exists an  $N \in \mathbb{N}$  such that  $|L - f(m)| < \epsilon/2$  for all  $m \geq N$ . Similarly, there exists an  $M \in \mathbb{N}$  such that  $|f'(p)| < \epsilon/2$  for all  $p \geq M$ . Set  $P = \max\{N, M\}$ . Then for all  $x \geq P$  set  $\tilde{x} = \lceil x \rceil$  so that  $\tilde{x} \in \mathbb{N}$  and  $\tilde{x} \geq P$  as well. Then

$$\begin{aligned} |L - f(x)| &= |L - f(x) - f(\tilde{x}) + f(\tilde{x})| \\ &\leq |L - f(\tilde{x})| + |f(\tilde{x}) - f(x)| \text{ by the triangle inequality,} \\ &< \epsilon/2 + |f(\tilde{x}) - f(x)| \text{ since } \tilde{x} \geq P. \end{aligned}$$

It remains to show that  $|f(\tilde{x}) - f(x)| < \epsilon/2$ . Since  $f$  is differentiable, by the Mean Value Theorem, there exists  $c \in (x, \tilde{x})$  such that

$$\frac{|f(\tilde{x}) - f(x)|}{|\tilde{x} - x|} = |f'(c)| < \epsilon/2$$

where the inequality holds since  $c \geq P$ . Then,

$$|f(\tilde{x}) - f(x)| < |\tilde{x} - x|\epsilon/2.$$

Putting it altogether, we get the required bound

$$|L - f(x)| < \epsilon/2 + |f(\tilde{x}) - f(x)| < \epsilon/2 + \epsilon/2 = \epsilon.$$

This shows that the limit  $\lim_{x \rightarrow \infty} f(x)$  equals  $L$ , and hence, exists. □

**Problem 4.** A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is called Lipschitz if there exists  $M > 0$  such that

$$|f(x) - f(y)| \leq M|x - y|$$

for all  $x, y \in \mathbb{R}$ . Show that every Lipschitz function on  $\mathbb{R}$  is uniformly continuous on  $\mathbb{R}$ , but not every uniformly continuous function on  $\mathbb{R}$  is necessarily Lipschitz on  $\mathbb{R}$ .