

Unit 103 - Field Data Review

Mohawk Valley West Slope demonstration. Method **Tractor**, 53.4 ac, BAF 20 variable-radius cruise on a 300 ft grid, 27 plots, 276 trees tallied. Simulated data with planted recording errors; see docs/methods.md.

Table 1. Stand summary from the BAF 20 cruise, per acre

STAND DENSITY		STAND INFO	
Basal area	204 sq ft/ac (1 SE of mean, BA 3.5 %)	Type	Sierran mixed conifer
Trees per acre	108	CWHR size / density	4 / D (cover 80 %)
QMD	18.6 in	SDI	273 (47 % of max 585)
SAWTIMBER (>= 10 in)		BIOMASS (< 10 in)	
BA QMD	196 sq ft/ac 20.6 in	BA QMD	8 sq ft/ac 8.0 in
TPA	85	TPA	23
Volume	30.3 MBF/ac (6028 cu ft)	Volume	2.7 green tons/ac (118 cu ft)
TREATMENT TARGET		CRUISE DESIGN	
Leave	35 % of max SDI = ~153 sq ft/ac BA	Sampling error (volume)	14.9 % at 95 % (CV 38 %, t = 2.06, n = 27)
Species (BA)	WF 65, PP 58, DF 37, IC 27, SP 18	Stratum standard	40 % of volume - MEETS; plots for standard: 4
		Sampling error (basal area)	22 % at 95 % (CV 18 %); secondary, not the standard
		Plot count	27 vs 20-plot local practice, not a handbook standard: ok

Cubic volume = total-stem CVTS by species (PNW-FIA California equations, MacLean and Berger 1976, CARB 2011 compendium) net of defect; Scribner board feet at 5.02 bf/cu ft (Keegan et al. 2010, table 2, California 2000-2006); green tons at PP 45, WF 47, DF 38, SP 49, IC 45 lb/cu ft of wood (Miles and Smith 2009, NRS-38 table 1A). SDI in the summation form; maximum is the basal-area-weighted FVS Western Sierra species maximum (table 3.5.1, rev. 2025-09-23). The Keegan ratio is Scribner log scale per cubic foot of delivered sawlog fiber, so applied to total-stem CVTS it overstates sawlog volume; merchantable cubic volume would be the consistent pairing. Sampling error at 95 % confidence with Student's t on plot net cubic volume per acre, the quantity the handbook standards apply to (FSH 2409.12 ch. 40, 41.1(5)(a) and (b)); stratum standard 40 % of volume for tree-measurement sales; the basal-area error is secondary. Stand-summary BA counts every tallied tree (plot count x BAF); the stand and stock tables exclude records flagged by QA, such as an out-of-range DBH, so the two can differ by a few tenths. Sources and URLs: cruise_data.xlsx, Standards and assumptions.

Table 2. Stand table by 4-inch DBH class, per acre

DBH class, in	Trees	TPA	BA	cu ft	MBF
4-7.9	4	10.6	3.0	44	0.2
8-11.9	19	26.0	14.1	216	1.1
12-15.9	36	24.8	26.7	581	2.9
16-19.9	41	17.7	30.4	727	3.6
20-23.9	46	13.0	34.1	1001	5.0
24-27.9	32	6.5	23.7	739	3.7
28-31.9	26	4.0	19.3	591	3.0
32-35.9	17	2.0	12.6	497	2.5
36-39.9	15	1.4	11.1	460	2.3
40-43.9	12	1.0	8.9	335	1.7
44-47.9	8	0.5	5.9	270	1.4
48-51.9	10	0.5	7.4	319	1.6
52-55.9	2	0.1	1.5	80	0.4
56-59.9	3	0.1	2.2	103	0.5
60-63.9	5	0.2	3.7	182	0.9
All	276	108.4	204.4	6146	30.9

Table 3. Stock table by species, per acre

Species	Trees	TPA	BA	QMD	cu ft	MBF
WF	88	33.8	65.2	18.8	2087	10.5
PP	78	33.6	57.8	17.8	1818	9.1
DF	50	18.3	37.0	19.2	1076	5.4
IC	36	13.6	26.7	19.0	663	3.3
SP	24	9.1	17.8	18.9	502	2.5
All	276	108.4	204.4	18.6	6146	30.9

Trees = trees tallied in the cruise; TPA, BA (sq ft/ac), cubic feet and Scribner MBF per acre by BAF expansion. Species: PP ponderosa pine, WF white fir, DF Douglas-fir, SP sugar pine, IC incense-cedar.

Unit 103: plots on the 300 ft grid (orange = QA flag)




Figure 1. Plot locations with QA status.

QA findings: no issues found; unit accepted

Table 4. Plot list

Plot	Trees in	BA sq ft/ac	Slope %	Status
103-01	9	180	25	ok
103-02	11	220	19	ok
103-03	14	280	32	ok
103-04	10	200	33	ok
103-05	12	240	36	ok
103-06	8	160	28	ok
103-07	12	240	33	ok
103-08	10	200	22	ok
103-09	7	140	32	ok
103-10	9	180	29	ok
103-11	11	220	37	ok
103-12	8	160	26	ok
103-13	10	200	25	ok
103-14	13	260	30	ok
103-15	11	220	27	ok
103-16	9	180	22	ok
103-17	10	200	23	ok
103-18	12	240	25	ok
103-19	9	180	22	ok
103-20	7	140	16	ok
103-21	7	140	19	ok
103-22	10	200	35	ok
103-23	11	220	18	ok
103-24	12	240	16	ok
103-25	12	240	21	ok
103-26	12	240	14	ok
103-27	10	200	5	ok